RIEMANNIAN GEOMETRY DURING THE SECOND HALF OF THE TWENTIETH CENTURY

Marcel BERGER

CERN LIBRARIES, GENEVA

CM-P00066872

Institut des Hautes Études Scientifiques
35, route de Chartres
91440 – Bures-sur-Yvette (France)

Mars 1997

IHES/M/97/26

RIEMANNIAN GEOMETRY DURING THE SECOND HALF OF THE TWENTIETH CENTURY

Marcel Berger

Table of contents

-1. INTRODUCTION

0. RIEMANNIAN GEOMETRY UP TO 1950

- A. Gauss, Riemann, Christoffel and Levi-Civita
- B. Hadamard, Elie Cartan and Heinz Hopf
- C. Synge, Myers, Preissmann : the use of geometric tools
- D. Hodge harmonic forms and Bochner technique : the use of Analysis
- E. Allendoerfer, Weil and Chern
- F. Existing tools and a brief look at the new ones
- G. Existing examples and a brief look at the new ones

1. COMMENTS ON THE MAIN TOPICS I, II, III, IV, V UNDER CONSIDERATION

An important digression : curvature is a very complicated
invariant, how does one see curvatures today

I. CURVATURE AND TOPOLOGY

A. Pinching problems

- 1. Introduction
- 2. The positive pinching
A digression : comparison theorems
- 3. Pinching around zero
- 4. The negative pinching

B. Curvature of a given sign

- 1. The positive side : sectional curvature
- 2. The positive side : Ricci curvature
- 3. The positive side : scalar curvature
- 4. The negative side : sectional curvature
- 5. The negative side : Ricci curvature

C. Finiteness, compactness, collapsing and the space of all Riemannian structures

- 1. Finiteness results
A digression : does the curvature determines the metric ?
- 2. Compactness, convergence results
- 3. The set of all Riemannian metrics : collapsing

II. THE GEOMETRIC HIERARCHY OF RIEMANNIAN MANIFOLDS :
SPACE FORMS

- A. The constant curvature case
- B. Space forms of rank one
- C. Space forms of symmetric spaces of rank larger than one

III. THE SET OF RIEMANNIAN STRUCTURES ON A GIVEN COMPACT
MANIFOLD : IS THERE A BEST METRIC ?

- A. The problem
- B. The minimal volume and $\text{Min}\|R\|^{d/2}$
- C. The case of Einstein manifolds
Digression : the Yamabe problem
- D. Some topological closures

IV. THE SPECTRUM, THE EIGENFUNCTIONS

V. PERIODIC GEODESICS, THE GEODESIC FLOW

- A. Periodic geodesics
A digression : geodesics joining two points
- B. The geodesic flow (geometry and dynamics)
Digression : the entropies in Riemannian geometry

TOP. SOME OTHER RIEMANNIAN GEOMETRY TOPICS OF INTEREST

- 1. Volumes
 - A. Bishop's theorem
 - B. The isoperimetric profile
 - C. The embolic volume
 - D. The minimal volume
 - E. The systolic story
- 2. Isometric embedding
- 3. Holonomy groups and special metrics : another (very restricted)
Riemannian hierarchy
 - A. Holonomy groups
 - B. Kähler manifolds
- 4. Cut-loci
- 5. Non compact manifolds
- 6. Bundles over Riemannian manifolds
 - A. Exterior differential forms (and some others)
 - B. Spinors
 - C. Various other bundles
- 7. Harmonic maps between Riemannian manifolds
- 8. Low dimensional Riemannian geometry
- 9. Some generalizations of Riemannian geometry
- 10. Submanifolds
 - A. The case of surfaces in $\mathbb{R}^3$
 - B. Higher dimensions
 - C. Geometric measure theory and pseudo-holomorphic
curves

-1. INTRODUCTION

From the 50's to the 70's the task of describing the evolution, the techniques and the results of Riemannian Geometry would have been quite easy enough. But Riemannian Geometry underwent since the 70's such a blowing up that this task becomes now almost impossible. This not only for volume, dimension reasons but also because of the difficulties of an ordering. In fact today's results, techniques, examples are interweaving each other enormously. This implies that we have been forced to take several biased steps.

First : not to be exhaustive. Second we have adopted quite an artificial classification. As a consequence this text is not really a more or less complete survey. A contrario we will refer to existing partial surveys as soon as we know their existence. Finally we will mention briefly at the end topics which are important but were not treated at large or even explicitly lack of space. We still hope that we will be clear and fair enough.

There is also the problem to define "**Riemannian geometry**". For obvious reasons of dimensions and competence we restrict it quite strongly to **what's really happening downstairs on the manifold itself and as Riemannian, in particular as a metric space**. In particular we are going to ignore completely most of the topics of **differential geometry**, even **Riemannian bundles**. This is a field which today underwent tremendous developments, including among others : differential topology, fiber bundles and connections, singularities and transversality, contact manifolds, symplectic geometry, minimal surfaces and generalizations of them, Yang-Mills (gauge) theory, twistors, foliations, submanifolds of $\mathbf{R}^d$, conformal geometry. For example, despite their great interest among various Riemannian bundles, Yang-Mills fields are not considered at least here as Riemannian geometry but of course are an important part of differential geometry. We will of course mention them but only briefly in **TOP. 6**. The tangent bundle, in particular the unit tangent bundle, will be used but typically in connection with the geodesic flow. Some people can find our "definition" of Riemannian geometry too narrow. They might be right, because it reflects the (biased) "elementary geometry" temper of the present author. As well as his liking for results simple to state. Because of their basic use in Riemannian geometry and the way they are constructed, we will detail a little more exterior differential forms and spinors in **TOP. 6. A and B**. One conclusion is the desire that quite soon will appear

some "Handbook of differential geometry", for the moment one can consult the three volumes of (Greene & Yau, 1993).

We also made one more gamble. That to be at the same time historical and to tell what is the state of affairs today. This will not take too much room since we are using an "author-date" reference system, which will hence give credits automatically. Concerning references we will be far from being exhaustive, since we are not presenting a complete survey and also because our bibliography would be unreasonably numerous. In most case what we will do is to give a reference which is good with respect to date and will enable at the same time the reader to go backwards using one or more bibliographies by induction. This will be specially in vigour for the briefly mentionned topics.

**manifold (unless otherwise stated) =
Riemannian manifold = a (the) "Riemannian metric"
= metric**

**most manifolds will be COMPACT
and never with boundary (closed)
and if non compact then always complete**

Acknowledgments are very important for the present text. First to the editors of a book to come on mathematics during the second half of the century who asked me to write such a text for the differential geometry part. I soon started writing it and found myself very fast embarked in a length exceeding from very far (as the reader can see now) the twenty pages allowance I was offered. So that I decided to write a larger object, the one you have now.

I was able to write such a text with enthusiasm thanks to the Rome University La Sapienza, the Indian Institute of Technology at Powai-Bombay, the University of Pennsylvania and the Zürich Polytechnicum who all invited me to give lectures, Roma in 1992, Bombay in 1993, Penn in the fall 94 and Zürich in the Wintersemester 95-96. Most important is the fact that both these two departments permitted me to give lectures entitled "Topics in Riemannian geometry" where I was covering a lot but without almost any proof, only sketching ideas and ingredients. I want to thank those very much these four mathematics departments for having accepted me to give lectures not entering in a classical frame.

Last, not least. A preliminary version was sent around to fifty colleagues. I got many answers, all important. I will now give a succession of names and for obvious reasons do this without trying to give the importance of their help. Those who helped me a lot and whom I even bothered will recognize themselves. From all of them help was important : Stephanie Alexander, Michael Anderson, Ivan Babenko, Pierre Bérard, Lionel Bérard Bergery, Gérard Besson, Jean-Benoît Bost, Jean-Pierre Bourguignon, Robert Bryant, Dan Burghelea, Peter Buser, Jeff Cheeger, Shiing-shen Chern, Tobias Colding, Alain Connes, Yves Colin de Verdière, Thibault Damour, Kenji Fukaya, Jacques Gasqui de Saint-Joachim, Paul Gauduchon, Robert Greene, Mikhael Gromov, Nancy Hingston, Dominique Hulin, Mikhail Katz, Ruth Kellerhals, Bruce Kleiner, Horst Knörrer, Jean-Louis Koszul, Jacques Lafontaine, Rémi Langevin, Blaine Lawson, André Lichnerowicz, Wolfgang Meyer, René Michel, Robert Osserman, Pierre Pansu, Peter Petersen, Hans-Bert Rademacher, Takashi Sakai, Katsuhiko Shiohama, Shanta Shrinivasan, Alain-Sol Sznitman, Iskander Taimanov, Domingo Toledo, Lieven Vanhecke, Takao Yamaguchi, Wolfgang Ziller.

Finally I do hope that, whatever its imperfections are, the present text will be of help for Riemannian geometers of any age and might be even pleasant to read in part or in totality for non experts.

0. RIEMANNIAN GEOMETRY UP TO 1950

A. Gauss, Riemann, Christoffel and Levi-Civita

In 1827 Gauss was the first, for ordinary surfaces in $\mathbf{R}^3$, to decouple the intrinsic metric from their embedding. In fact he knew that result much before, see (Dombrowski, 1979). More precisely he proved that the total curvature, namely the product of the two principal curvatures, could be computed only with the intrinsic metric (theorema egregium).

In 1854 Riemann was defining surfaces with a "Riemannian metric" in a completely abstract way, making them free of any embedding or immersion. Moreover he was doing so in any dimension for all manifolds, even if his notion of manifold was somewhat a little bit imprecise. See in the volume II of (Spivak, 1970) a perfect analysis of Riemann founding paper. With the now

solidly established notion of a differentiable manifold one can now define a general Riemannian manifold (M, g) as a differentiable manifold M endowed with a Riemannian metric g , i.e. one is given at every point m of M some positive definite quadratic form $g(m)$ on the tangent space $T_m M$ to M at m . One asks moreover the application $m \rightarrow g(m)$ to be differentiable. Riemann defined the metric $d(p, q)$ as the infimum of the length of all the curves joining p to q , the length making sense precisely because of g . He also defined the curvature tensor as well as the sectional curvature. We will come back below to this very hard object to tackle in the digression at the end of **COMMENTS on TOPIC I**.

Thereafter Christoffel succeeded partially to understand what was going on, namely the, still at that time obscure, notion of connection. This was circa 1865, see (Klingenberg & Pinl, 1981). Then at the turn and the beginning of the century one knew that Riemannian manifolds own a **golden triangle** which is the primary tool in Riemannian geometry. This was due to Levi-Civita and Ricci : (Levi-Civita & Ricci, 1901) and (Levi-Civita, 1917). The triangle is made of the three vertices

Let us be more precise. The urge is to be able to develop on a Riemannian manifold a differential calculus of any order. In affine spaces the tangent spaces at various points can be all identified with a single vector space by translations and that is enough. But this is not possible a priori in a smooth manifold since there is no way to compare tangent spaces at different points. The discovery was that such a comparison is possible infinitesimally in a unique way if one demands that the euclidean structures of the tangent spaces to be preserved and moreover the second differential of a function to be symmetric. The precise notion of such a **connection** was seen in various ways, mostly in coordinate writing. Elie Cartan was using mobile frames. One can also use, in the tangent bundle to the manifold, the notion of horizontal subspace. Since Koszul the best writing for the canonical connection (as well as for connections in various bundles with ad hoc modifications) is to define it as an operation $D_X Y$ on vector fields which is linear in X

and a derivation on Y . Namely if f is any function then $D_f X Y = f D_X Y$ but $D_X(f.Y) = X(f).Y + f.D_X Y$. Koszul introduced this concept in it 1951, it was used soon in (Nomizu, 1954) and finally presented in a systematic expository way in (Koszul, 1960). Preserving the metric is insured by $X\langle Y, Z \rangle = \langle D_X Y, Z \rangle + \langle Y, D_X Z \rangle$ for every X, Y, Z and the symmetry asked for the second differential will be insured if and only if the connection is without torsion, i.e. $D_X Y - D_Y X = [X, Y]$ for every pair of vector fields.

The connection can be integrated along any curve (but not on a surface in general) because it is then a linear first order differential equation, the result is the Levi-Civita parallelism. And then the tangent spaces along the curve can be identified with a single one of them : one can now take derivatives of any order along a curve. Important to remark is the fact that the parallel transport along a loop is in general different from the identity. This lack of "holonomy" is the starting point for **TOP. 3**.

Let us now comment on the arrows between the three vertices of the golden triangle. Note first that the metric comes from it because shortest paths (called **segments**) between points are supported by geodesics, which can be defined as the self-parallel transported curves (for their speed vectors). The curvature tensor is given by the derivative of the parallel transport along infinitesimal parallelograms and in writing it is nothing but the defect of commutation in $D_X \circ D_Y$ namely $R(X, Y) = D_X \circ D_Y - D_Y \circ D_X$. So that the curvature tensor is an antisymmetric differential form with values in the endomorphisms of the tangent spaces.

Now the absolute (covariant) derivative $D_v S$ of a form (or more generally of any tensor S) is obtained by taking the ordinary derivative at 0 of the values taken by the form on a set of vector which are parallel transported along a curve whose initial speed vector is v .

The third arrow concerns the relation between curvature and covariant derivatives of tensors. In a differentiable manifold the differential of a function is well defined, but no longer the second differential because the chain rule formula is not valid for the second derivatives when one looks at coordinates changes. The Riemannian canonical Levi-Civita connection precisely yields such an invariant second derivative, called the **Hessian** of the function : it is a bilinear symmetric differential form (this is Schwarz lemma for Riemannian manifolds). And one can keep going on with derivatives (covariant) of any order. But then typically the third

differential of a function is no longer symmetric but precisely the defect of this symmetry is given by the curvature tensor. The same for the second derivative of 1-forms, etc. This is basic in many places below and can be seen as the revenge of the Riemannian geometer. The explicit formulas for those defects are called **Ricci commutation formulas**, see their first use in **D** below. In Koszul's writing Ricci formulas are straightforward.

The **curvature tensor**, known to Riemann, is introduced above, as a tensor of type (3,1), namely the exterior 2-form $R(X,Y)$ with values in the endomorphisms of the tangent spaces. But alternatively (using the duality given by the metric) as a differential form of degree **four** (quadrilinear), enjoying moreover some symmetries : $R(x,y,z,t) = \langle R(x,y)z,t \rangle$. We will see below in various places that such an object, even pointwise from the point of view of the sole linear algebra, remains quite mysterious today, see the digression in **1. COMMENTS on I**).

B. Hadamard, Elie Cartan and Heinz Hopf

In (Hadamard, 1898) it was already known that negative curvature and simply connectedness for surfaces embedded in $\mathbf{R}^3$ force uniqueness of geodesics joining two points, implying that moreover any piece of geodesic is a shortest path (**segment**). But there is a long way toward the modern statement : any complete abstract Riemannian manifold of negative (in fact nonpositive and of any dimension) curvature is the quotient of its universal covering by a discrete group of isometries. This is especially needed for studying space forms of negative curvature, see **II** and **I. B. 4**.

Missing as well defined notions were : abstract manifolds, completeness of Riemannian manifolds, this in any dimension. Hopf was interested to solidly establish the meaning of **completeness** (with a good notion of manifolds) in relation with the prolongability of geodesics. This is done in (Hopf & Rinow, 1931), surprisingly enough only for dimension 2. It was remarked only in (Myers, 1935a) that Hopf-Rinow proof applies to any dimension without any non trivial change.

In the second edition (Cartan, 1946) of his fundamental book (Cartan, 1928), Elie Cartan was interested in the technical tools (and ignored systematically what is precisely a manifold). He developed a dual version of the today second order differential

equation called **Jacobi fields** (known to Gauss for surfaces). A Jacobi field is, by definition, the transverse derivative to a one-parameter family of geodesics. In vector writing (and using parallel transport to work with only "ordinary derivatives") the equation for Jacobi fields reads $Y'' + R(Y) = 0$ where R is the $R(\gamma', \cdot)\gamma'$ part of the curvature tensor given by the speed γ' of the geodesic γ one is working along. For the curvature tensor, a frightening object, see various places below. Cartan's tool was "repère mobile" but this is equivalent. He obtained the fact that the curvature tensor and the parallel transport together determine the metric (see at the end of **TOP. 4** for the global problem : Ambrose's problem). Remark that one needs here to know the geodesics through a point and not only the curvature, see the digression in **I. C. 1**. From his "philosophy" (chapter X of (Cartan, 1946)) Cartan deduced that a real analytic Riemannian manifold was determined locally by the values, at any given point, of the curvature tensor and of all its covariant derivatives. In fact Cartan knew that philosophy much before, when he studied the symmetric spaces in the middle 20's (**II. C**), since in that case the curvature tensor has a vanishing covariant derivative and then the metric is known as soon as the curvature tensor is known at a single point. As a corollary he showed that nonpositive curvature implies (simply connectedness is asked) an inequality which says that triangles in this case are "larger than equal to the corresponding euclidean ones" :

$$a^2 \geq b^2 + c^2 - 2 b c \cos \alpha$$

In (Cartan, 1929b), section 16, this inequality is used to prove the existence and uniqueness of the center of mass in those manifolds (see **I. C. 1**). From it he deduced that compact maximal subgroups of semi-simple Lie groups are always conjugate, this because symmetric spaces of nonpositive curvature are nothing but the quotient of a non compact Lie group by one of its maximal compact subgroup.

Parallel to that and with obvious interactions he developed also the complete theory of symmetric spaces (see **G** below) and complete mastering of their geodesic behavior.

One can now see that the simply connected manifolds of constant sectional curvature are unique : they are the "round" spheres in the positive case, the euclidian spaces for the zero case and the hyperbolic spaces **Hyp^d** for the negative case. Remember here that

the first correct and complete definition of Hyp^d appeared for the first time in Riemann 1854 text. Moreover the other space forms are "just" quotients of those by some suitable group of isometries. These quotients will be studied quite in detail in II. In the case $K = 0$, which is equivalent to locally euclidean, one uses the wording flat.

C. Synge, Myers, Preissmann : the use of geometric tools

Back in the late 20's Heinz Hopf was working very hard on the links between curvature and topology, see the landmark paper (Hopf, 1932). He put many students on the subject. The first one was Myers. In particular in (Myers, 1941) the second variation formula is used to prove that the diameter can be bounded (optimally) when the manifold has a positive lower bound for the Ricci curvature (see 1. COMMENTS on I). The striking corollary is that the fundamental group is finite as soon as one has a positive lower bound for Ricci curvature (just look at the universal covering with the lifted up metric).

In the case of surfaces the second variation formula has already been exploited in (Bonnet, 1855) to show the diameter property for surfaces of positive (total, Gauss) curvature which coincides with Ricci curvature in dimension 2.

The second variation formula was again used in (Synge, 1936), together with the parallel transport along a periodic geodesic, to prove that positive sectional curvature, compactness and even dimensionality force the fundamental group to be zero (or $\mathbb{Z}_2$ in the non orientable case). A beautifully simple idea : move by parallel transport vectors along a smallest curve in any free homotopy class (which is necessarily a periodic geodesic). When coming back to the origin point the vector is moved by an orientation preserving orthogonal transformation, which in odd dimension has always at least one non zero fixed vector. This yields a strip which will furnish smallest near by curves

(Preissmann, 1942-43) is a somewhat turning point paper. Except the theory of symmetric spaces, it contains in 40 pages all global results available in Riemannian geometry in those years. Preissmann contribution was the first to give a solid proof of the second variation formula. Then he used Jacobi fields to rediscover the above triangle inequality of Cartan for nonpositive curved

manifolds. Moreover he used this triangle comparison to show that if the curvature is negative then abelian subgroups of the fundamental groups have to be necessarily cyclic. This was generalized only in (Lawson and Yau 1972), see I. B. 4 .

D. Hodge harmonic forms and Bochner technique : the use of Analysis

All the above results were obtained by geometric tools. A historical event, still of basic influence, was (Bochner, 1946). At that time was already available de Rham theorem which links cohomology and exterior differential forms : (de Rham, 1931). Note that this theorem is differential topology, for the moment Riemannian geometry does not enter at all in it. The precise statement of de Rham theorem appeared first in (Cartan, 1928b). Elie Cartan suspected this result when working to compute the topology of symmetric spaces, see (Cartan, 1928b) and more generally that of homogeneous spaces, see (Cartan, 1929a) (or his Complete works : the same applies, books excepted, to all other Elie Cartan citations below).

But moreover Hodge theory of harmonic forms was available : (Hodge, 1941), although it was solidly founded later, see (de Rham, 1954). It says that (via de Rham theorem) the elements of the real cohomology of a compact Riemannian manifold are represented by exterior harmonic forms, and this in a unique way : for an exterior form ω harmonic means $\Delta\omega = 0$, where we have as usual denoted by Δ the Laplacian operating on numerical functions and on exterior forms, see IV and TOP. 6 . This is on such an exterior form a strong Riemannian condition. The relation with cohomology is insured by de Rham theorem.

Let now use Ricci commutation formulas for permuting the order of (covariant) derivatives : as seen in B the defect of the Schwarz lemma in classical differential calculus can here completely computed with the help of the curvature tensor. One gets finally a formula evaluating the Laplacian of the square norm of such a harmonic p-form ω :

$$-\frac{1}{2} \Delta(\|\omega\|^2) = \|D\omega\|^2 + \text{Curv}_p(\omega, \omega)$$

where $D\omega$ is the covariant derivative of ω and Curv_p some universal algebraic term involving only the form ω (quadratically)

and the curvature tensor (linearly). . We now integrate over the compact manifold and see that such a ω cannot exist if Curv_p is positive definite, because the integral of a Laplacian is always zero by Stokes formula. A short way to write the formula above is

$$\Delta = d \circ d^* + d^* \circ d = D^*D + \text{Curv}_p .$$

This was first used in a global setting for $p=1$, i.e. for 1-forms, in (Bochner, 1946), and very soon underwent a tremendous development, first for p -forms thereafter for various vector bundles and elliptic operators. The story is still not finished, see the surveys (Wu, 1987), (Bérard, 1988) , (Bourguignon, 1988) and their bibliographies. We will meet applications of analogous formulas in many places below for this **vanishing technique** . It was remarked in (de Rham, 1954), page 131, that the above formula for Δ with Curv_p was already in fact in (Weitzenböck, 1923), so that in fact Bochner in (Bochner, 1946) (see also (Bochner, 1948)) rediscovered it. We detail now the initial Bochner case.

This Curv_p term is specially nice when $p = 1$, since its value is $\text{Curv}_1(\omega, \omega) = \text{Ricci}(\omega, \omega)$. As a conclusion the positiveness of the Ricci curvature implies the vanishing of the real 1-cohomology since the two other terms are nonnegative This is certainly less strong than Myers. Yes, but it is such a completely new technique ! Moreover one can also immediately use it to take care of the nonnegative case : the positivity forces $D\omega = 0$, and a parallel transported form (i.e. with a zero covariant derivative) forces a local Riemannian product decomposition. See the digression in **I. COMMENTS on I.**

Thereafter people tried to see what results one can get for other degrees p , see among others (Bochner & Yano, 1953). But it is only in (Meyer, 1971) that it was made clear that it is the **curvature operator** which completely governs the term Curv_p : see the end of section **I. B. 1** . For Curv_p a very lucid exposition is in (Lawson & Michelsohn, 1989), theorem 8.6 .

It is important to see the philosophy here. The Laplacian Δ on exterior forms is what is called a **natural operator**, because it finally involves only Riemannian invariants : D the covariant differentiation , D^* its adjoint (defined by the global scalar product) and the curvature tensor . More generally for any operator S the product S^*S is always nonnegative by definition. Then one has vanishing of $D\omega=0$ as soon as $\Delta\omega=0$. This scheme can and will be

used for more general differential operators on various bundles, e.g. in **I. B. 3** and **TOP. 6**.

E. Allendoerfer, Weil and Chern

In the late 20's Hopf was wondering about extending the Blaschke-Gauss-Bonnet formula which says that for a compact surface M and its curvature K the Euler-Poincaré characteristic is given by

$$\chi(M) = \frac{1}{2\pi} \int_M K(m) \, dm$$

Consequently if K has a given sign then χ has the same sign. All books seem to call this formula "Gauss-Bonnet theorem". This is quite surprising since the notion $\chi(\cdot)$ did not exist then. This formula as such first appeared partially in (Blaschke, 1921), page 109 and completely in (Blaschke, 1930), page 167. We owe this historical information to Robert Osserman.

Hopf was asking for higher dimensions : when the sectional curvature (see **1. COMMENTS on I**) has a given sign, does necessarily χ have the expected sign ? Hopf managed to prove it in the very special case of space forms (**II. A**) in (Hopf, 1925a). And was probably guessing that in the general case it would follow from a generalization of Blaschke formula to higher dimension. Such a formula was indeed obtained in (Allendoerfer & Weil, 1943). In dimension 4 it reads

$$8\pi^2 \chi(M) = \int_M (\|R\|^2 - \|\text{Ricci}\|^2 - \frac{\text{scal}}{4} \cdot g^2) \, dm$$

in invariant form. But in a nice basis the integrand becomes $K_{12}K_{34} + K_{13}K_{42} + K_{14}K_{23} + R^2_{1234} + R^2_{1342} + R^2_{1423}$ (with the obvious notations for the sectional curvature). This implies Hopf's conjecture as said in (Chern, 1955). The invariant writing will be used in **I. B. 1** and in **III. B**. But starting dimension 6 we will see that the formula is so complicated that it cannot say too many good things in general (cf. (Bourguignon & Polombo, 1981) , mostly only some non-zero (vanishing) results : see **TOP.1. D** . Starting dimension 6 Hopf's question is still open, see the end of **I. B**.

In (Chern, 1944) Chern was providing Riemannian geometers with a very conceptual proof of the generalized Blaschke formula. Then of course he tried to do the same for the Stiefel-Whitney classes.

But because they come from (real) Stiefel manifolds, they are torsion classes so inaccessible directly via exterior forms. Reading Ehresmann he learned that complex Stiefel manifolds have no torsion and this lead him to discover the "Chern classes" which are cohomology invariants defined for any complex vector bundle over a manifold, generalizing Euler-Poincaré characteristic and which for "real" manifold are the Pontryagin classes $P_{4k}(M)$: (Chern, 1946). His proof was yielding integral formulas for those classes . The integrands are involving universal polynomials in the curvature tensor. As above for the Euler-Poincaré characteristic the formula for P_1 in dimension 4 is useful, see III. C for Einstein manifolds. Higher dimensions yield only general results of "non-zero" type but, using algebraic examples of the curvature tensor, it is proved in (Bourguignon & Polombo, 1981) that those formulas are "useless" or at least disappointing for the Riemannian geometer when using only pointwise estimates of the curvature tensor.

It is important to know that there are **no other** such universal Riemannian integral formulas yielding topological invariants. This was a query of Gelfand, proved finally in (Gilkey, 1974) (see also (Abrahamov, 1951)). See (Gilkey, 1995) for an up to date text and also **TOP. 6** .

This does not prevent Chern classes to be basic. Recall that they are defined for any bundle, not only the tangent one : (Chern, 1946). They are today a building tool in algebraic geometry and in the heat kernel. But what also counts is the fact that there is an integral formula whose writing is **polynomial and universal** in the curvature. See (Berline, Getzler, & Vergne, 1992), (Gilkey, 1995) and **TOP. 6** .

A basic formula fact discovered in the first edition (in 1956) of (Hirzebruch, 1966)) . A suitable combination of Pontryagin classes yields the **signature** for $4k$ -dimensional manifolds. Read via de Rham theorem this signature is the linear algebra signature of the quadratic form which is defined by the cup product of cohomology $2k$ -classes. Hence this important fact : there is an integral formula involving the curvature for the signature. See essentially (Gilkey, 1995) and (Berline, Getzler & Vergne, 1992). See also **TOP. 6** for secondary characteristic classes and the η -invariant.

F. Existing tools and a brief look at the new ones

From the above one sees that the existing tools in 1950 were only the following ones. The golden triangle and its rules and the equation for Jacobi fields, which is the equation giving the transverse field to a one-parameter family of geodesics. The basic fact is that equation is linear (of second order) and involving only the curvature tensor, the time being the equation for geodesics themselves is definitively not linear in general.

Then one had also the first and the second variation formulas. The first is given by angles of vectors at the extremities, the second needs an integral along a geodesic where enters a sectional curvature term. The first variation was used to prove that there exists at least one periodic geodesic in every non zero free homotopy class of a compact manifold. It seems that was folklore, for example it is used in (Hadamard, 1898), where moreover uniqueness is proved in the negative curvature case. The second variation formula gives for a geodesic the second derivative of the lengths of a family of curves as in integral along the geodesic which mixes the sectional curvature and the defect of the family to be a parallel strip. From Analysis were available Hodge theory and Bochner technique.

The very geometric notion of convexity came up quite recently for both subsets and functions in a Riemannian manifold, see its use in the non compact situation in B. 1 and B. 4.

For the geodesic behavior, a basic tool is Morse theory (Morse, 1934) of the global calculus of variations, see the digression in V. A. Before 1950 this extremely general and powerful theory was still not yielding substantial results in Riemannian geometry. This is not because of the theory in itself but since algebraic topology was not advanced enough.

More about tools . We will not make a systematic exposition, merely will indicate them at the time they are used. Tools used by Riemannian geometers are of two kinds. Those they created themselves for their need and the preexisting ones that they borrowed and had in some instances to be adapted.

Riemannian geometry being "differentiable", one can risk to say caricaturally that "half of it is Analysis". The reader might have noticed that many names mentioned are those of analysts : Jacobi,

Hadamard, Poincaré, Elie Cartan, Morse, Chern, Bochner, Rauch, among others . This fact can also be remarked in the sequel. We mention now important tools coming from Analysis which appeared since the 50's. We are certainly biased and again insist on the fact that we are really sticking to Riemannian geometry inside the huge world of differential geometry. The reader could consult the following surveys : (Yau, 1987), (Schoen, 1991).

One basic tool is Geometric Measure Theory (GMT) . The starting idea is that of minimal surfaces, but GMT was the first to offer general enough results. It will be used for example **TOP. 1. B** , **TOP. 10. C** and **I. B. 3** . The major impetus came from (Federer & Fleming, 1960) and things cristallized in (Federer, 1969). See more in **TOP. 10. C**.

There is also the control of functions of special type, like distance functions, harmonic functions, eigenfunctions of the Laplacian (in particular the first one). The two basic facts are : in polar coordinates the formula giving the Laplacian of a function depending only of the polar radius involves only the logarithmic derivative of the angular volume element θ ("solid angle") : in particular if d is a distance function (it can be a Busemann one from infinity) its Laplacian is $\Delta d = -\frac{\theta'}{\theta}$. And this is upper bounded when Ricci curvature is lower bounded by Bishop's theorem (**TOP. 1. A**). The second is Bochner's formula above as applied to any function : wrote in full details it reads $-\frac{1}{2}\Delta(\|df\|^2) = \|\text{Hessian}(f)\|^2 + \langle D\Delta f, Df \rangle + \text{Ricci}(df, df)$. One can apply it to a distance function, this will give more control with the Ricci curvature on the differential of its Laplacian $d(\Delta d)$, and/or to some suitably chosen harmonic functions. We will see that this two facts are heavily used, not only for the spectrum, but also to obtain metric comparison theorems with only a lower Ricci bound : see e.g. the digression in **I. A. 2**.

Other tools are existence theorems for harmonic maps. The topic started with (Eells & Sampson, 1964). Today it is of basic use, see **TOP. 7** .

Harmonic coordinates were systematically introduced in (Jost & Karcher, 1982) and are now incontournable : see the digression in **I. C** and **1. COMMENTS on I**. We note that harmonic coordinates have been used a long time ago by theoretical physicists, see

(Einstein, 1916) and (Lanczos, 1922). See the formula for Δg_{ij} in the digression of **1. COMMENTS on I**, then **III. C** for the use of harmonic coordinates to prove that Einstein manifolds are real analytic and the basic need of them in **I. C. 1 and 2**.

Very new are **deformations** techniques. This means on a manifold a partial differential equation, like the heat equation, and describing an **evolution**. They are used to get existence of desired objects. The technique is to prove first that the deformation equation (typically a partial differential equation of parabolic type) has a solution for all time t and second that it is converging to "the" (or "a" desired object) when t goes to infinity. See **TOP. 7** for harmonic maps and **5.A** for obtaining periodic geodesics. If one deforms suitably chosen functions in the complex case this is the technique suggested in (Calabi, 1954) to obtain Kähler Einstein manifolds. These deformation equations are quite natural : they consist in deforming an initial object in the best way for reducing the **defect** with respect to the imposed condition.

Calabi's equation was a particular case of the more recent and more abstract technique consisting in deformations in the space of all Riemannian metric on a given manifold, see **III. C**. In the general case the deformation is really for Riemannian metrics. Very often one uses the **Ricci flow**. That is to say one considers a one-parameter t and a family of metrics $g(t)$ and the PDE :

$$\frac{\partial(g(t))}{\partial t} = - 2 \text{ Ricci}(g(t))$$

This equation appears under slightly different forms where the right hand term is normalized (in various ways according to authors) by adding the scalar curvature and playing on both coefficients of Ricci and scalar, the point being to keep the volume fixed. The difficulty is that this equation is not really parabolic because the right side is only "elliptic degenerate". We will meet various results later on, a partial is the section 4 of (Besse, 1994). The Ricci flow was first used by Hamilton in dimensions 3 and 4, see the results in **I. B. 1 and 2**.

See also **TOP. 3. A** for results on manifolds with holonomy G_2 and $\text{Spin}(7)$ which are obtained by such deformation technique but with completely different equations. See also **I. C**.

G. Existing examples and a brief look at the new ones

Another very important aspect of Riemannian geometry is that of **examples**. This might not be obvious but some people even say that to feel, to understand what is a Riemannian manifold, cannot be achieved without being familiar with quite a few examples and we will mention them in time even if briefly. The following books are specially insisting on examples in an expository way : (Gallot, Hulin, & Lafontaine, 1990), (Besse, 1987), (Petersen, 1997b). Here are the examples which were known in 1950 and whose important geometric properties were more or less understood.

First following Riemann, the hyperbolic spaces Hyp^d in any dimension. On the compact side first of course the spheres. Then the complex projective spaces $\mathbb{C}P^n$ with their Fubini-Study metric : (Fubini, 1903) and (Study, 1905)).

Those $\mathbb{C}P^n$ were put by Elie Cartan in the general frame of all symmetric spaces (see section II.C). This includes first the symmetric spaces of rank one, which are on the compact side the spheres, the various $\mathbb{K}P^n$ for $\mathbb{K} = \mathbb{C}$ or $\mathbb{H}$ (the quaternion field) and for the Cayley numbers $\mathbb{Ca}$ only $\mathbb{Ca}P^1 = S^8$ and $\mathbb{Ca}P^2$ are permitted. The Cayley projective plane $\mathbb{Ca}P^2$ is beautiful, we like to call it the **panda** of Riemannian geometry. The projective lines are 8-dimensional spheres and verify the projective plane axioms. Cartan suspected it in (Cartan, 1939), page 354. One had to wait Freundenthal in 1951 for a solid projective construction of it which, by the way, was never published. For this intricate history of the panda, see the chapter 3 of (Besse, 1978).

On the non compact side Cartan exhibited the dual generalized hyperbolic spaces $\text{Hyp}\mathbb{K}P^n$. Higher rank symmetric spaces were also completely classified by him, they include in particular all the Grassmann manifolds $\text{Grass}_{\mathbb{K}}(p,q)$ as well as their nonpositive curvature analogues, plus a very restricted list. For the non simply connected quotients of the above, see the entire chapter II. Except for the spherical case and for surfaces compact quotients were not known to exist in 1950 besides the dimension 3 hyperbolic examples of (Löbell, 1931).

This complete theory of symmetric spaces (geometry, classification, see more in II. C) was built up by Elie Cartan in the very short

interval 1926-27 , in a series of paper from (Cartan, 1926b) to (Cartan, 1927). See more in **II. C.**

The geometry of the geodesic flow was understood : for surfaces of revolution (Clairaut circa 1730), for ellipsoids (Jacobi in 1832, see an exposition in (Klingenberg, 1982), section 3), (these geodesic flows are in the modern jargon called "integrable"), for symmetric space in (Cartan, 1927). One should add Morse's theorem quoted in **V. A** . But one still does not have a complete mastering of the metric of ellipsoids, see the end of **TOP. 4** .

A trivial but of basic need is the **product** of two Riemannian manifolds. The relation with holonomy groups (see **TOP. 3**) and the global product result in the complete simply connected case are in (de Rham, 1952) and (Borel & Lichnerowicz, 1952) (we make this an exception to the 1950 rule). In the case of symmetric spaces this notion of product and holonomy group was known to Cartan and was a basic tool in his work we mentioned above.

This hides the difficulty, sometimes extremely hard, one has to exhibit a Riemannian manifold having imposed properties. By this we mean most often the control of the curvature or/and of the basic geometric invariants : volume, diameter, injectivity radius (**I. A. 1**).

There is quite a variety of techniques to build up examples. It starts with Algebra. First to come are **homogeneous** spaces G/H of Lie groups with H compact : they always admit invariant (called homogeneous) metrics and there are explicit formulas for computing the curvature see **III. C** . But their discrete quotients are much much harder to built up. For example building up general space forms was in the air only in the 50 and moreover one had to wait (Borel, 1963) to get a general result. Note that it was more Number Theory than geometry.

Then comes the theory of **Riemannian submersions**. This means a submersion $M \rightarrow N$ between two Riemannian manifolds which should fulfill the following condition. The metric in M determines at every point an orthogonal complement (called horizontal) of the space tangent to the fiber : at every point the restriction of the projection to this horizontal space is asked to be an euclidean isometry between it and the tangent space at its image in the basis. Here the formulas of (O'Neill, 1966b) enables us to compute the curvature of the total space if one knows : the curvatures of the

basis and of the fibres, how the fibration is twisted and finally how the fibres are embedded (the best case is when the fibers are totally geodesic) . For example topological surgery can be riemannianly controlled by these formulas. A now classical example is for the positive scalar curvature : **I. B. 3** . Particularly important Riemannian submersions are the **warped products**. Definitions here vary with authors. When the basis is one-dimensional, in coordinates the metric will appear as $g = dt^2 + f(t).g_{d-1}(t)$. If g_{d-1} is independent of t this is a rotationnally symmetric metric, a generalization of surfaces of revolution. O'Neill formulas then become much more simple. More recent are even the **double warped products**. For these formulas in book form see (Cheeger & Ebin, 1975), (Besse, 1987), (Sakai, 1996) and (Petersen, 1997b). Warped products are of basic use in (Cheeger & Colding, 1996d) (and subsequent works of the authors) where they serve as smooth rigid models for studying the structure of manifolds with a lower bound on Ricci curvature.

Other examples are those obtained by **conformal** changes : g becomes $f.g$ where f is a numerical positive function. Although extremely special when $d \geq 3$ those changes are very useful sometimes and the formulas for the curvatures are extremely simple.

Mixing these two techniques one can compute the curvature of the Riemannian manifolds of **high cohomogeneity**. This means manifolds which are almost homogeneous, i.e. the isometry group is large in the sense that its generic orbits have low codimension. The main difficulty is that of the singularities which occur at the places where the dimension of the orbit is not the maximal one (the case of the poles of surfaces of revolution is typical). This was used for example in (Page, 1979) and (Bérard Bergery, 1982) to built up Einstein manifolds. As one will see below this kind of examples are now more and more numerous and often used as counterexamples.

Deep analysis, in particular strong results for various types of partial differential equations, is basic. See **III** for example and what was said about the new tools from Analysis.

Another approach is quite recent. It turned out in the 60's that some results (of algebraic topology, then for complex manifolds and even more recently of Riemannian geometry) on a manifold cannot be obtained by just working on the manifold itself. One had to study various **bundles** over that manifold, principally vector

bundles. Some bundles are canonically attached to a Riemannian manifolds, like those of exterior forms but it is also important to build various "twisted" bundles. This consideration has become an basic tool, we will devote **TOP 6** to it. A special mention should be given to **spinors**. A quite complete reference is (Lawson & Michelsohn, 1989), see also (Gilkey, 1995) and (Berline, Getzler & Vergne, 1992).

Last, not least : "pure" geometry. One can see two parts in it. The first is geometry on the manifold itself. Most often one needs to adapt in the Riemannian context techniques coming from differential topology, surgery in particular. In this case one needs to control the curvature, an example is (Gromov and Lawson 1980b) used in **I. B. 3**. One finds also various geometric controls of various volumes during surgery and other topological constructions in relation with **TOP. 1. E** . We will mention other examples when going on, but Gromov's construction of a sphere with almost negative curvature (see details in (Buser & Gromoll, 1988) and a generalization in (Bavard, 1987b) show how high can be the degree of sophistication when trying to build up examples with only purely geometric tools. See the end of **I. B. 4**.

Complexity theory is entering slowly in Riemannian geometry, see 8. 11 of (Semmes, 1996) for the case of bounded geometry (**TOP. 5**).

1. COMMENTS ON THE MAIN TOPICS UNDER CONSIDERATION

As already explained above we made a choice of five main topics, namely

- I** : curvature and topology ((Hopf's urge)
 - II** : looking for constructing and classifying space forms
 - III** : looking for distinguished metrics, in particular Einstein
ones
 - IV** : study of the eigenvalues and the eigenfunctions of the
Laplacian (in brief : study of the spectrum)
 - V** : study periodic geodesics, more generally the geodesic flow
- TOP** : at the end we will briefly mention interesting
important topics, but not as much in detail

THE TOPIC I started with Hadamard, was pursued by H. Hopf, Synge, Myers, Preissmann and Bochner. The subject blew up with Rauch in 1952 and did not stop since. We will devote a large part of our text to it. It was a very strong incentive. Considering the huge harvest of results we had a very hard time organizing the material. We will make a first division in three parts : pinching, curvature of a given sign, finiteness theorems (and more). The reader will judge if that division is too artificial. Inside each part the idea is also threesome, with sectional, Ricci and scalar curvatures. But moreover we will at the same time follow the strong stream of today research in Riemannian geometry : obtain results with less and less hypothesis, the simplification being of having less invariants in the statement and/or to replace pointwise norms by integral norms.

In particular recently one obtained more and more deep results with only Ricci curvature bounds. The motivation is this : in dimension d Ricci curvature has as many parameters as a metric : $d(d+1)/2$. The time being the sectional curvature depends on $d^2(d^2-1)/12$ parameters. For $d=2$ this is not enough information , but for $d \geq 4$ this is "too much" information . We saw above in 1. F one basic formula involving only Ricci curvature, the second is in the digression below.

If one were asked to sum up in one question what remains to be done today in the field we owe to Gromov the following one : "how many positive curvature can a manifold carry"?

An important digression : curvature is a very hard invariant, how
does one see curvatures today

The curvature tensor appears naturally as endomorphisms $R(x,y)$ of the tangent space attached to couples (x,y) of tangent vectors. Naïvely simple is the **sectional curvature** attached (from the curvature tensor) to a 2-dimensional tangent space P (a **tangent plane**) which is the scalar $K(P) = K(x,y) = R(x,y,x,y)$ when x,y is any orthonormal basis of P . Note that the form $R(x,y,z,t)$ is multilinear of degree 4 ("quadrilinear" or "biquadratic") and enjoys the following symmetries : it is anticommuting in x,y and in z,t , one has $R(x,y,z,t) + R(x,z,t,y) + R(x,t,y,z) = 0$ and $R(x,y,z,t) = R(z,t,x,y)$. Geometrically the geometric interpretation of the sectional curvature is wonderful : compute the length of the small metric

circles generated by the geodesics starting with a speed vector in P . Then $K(P)$ measures exactly the defect of that length to be the euclidean one, precisely it gives the first non trivial term in the limited expansion in function of the radius.

But despite the symmetries of the curvature tensor the function K as a numerical function defined on the Grassmann manifold of the planes at a point is still almost a complete mystery : where are the critical points (planes), what are its critical values and in particular its extremal ones ? Things are trivial in dimensions 2 and 3 but starting dimension 4 one has only one partial result in (Singer & Thorpe, 1969) for 4-dimensional Einstein manifolds. A notation like $K < a$ for a Riemannian manifold will mean that the sectional curvature is smaller than the scalar a at every point and this for every tangent plane at that point.

The fact that the knowledge of the curvature tensor and the knowledge of the sectional curvature are equivalent is important theoretically but it is hard to use it explicitly. Important to know is the dimension of the space of curvature tensors (pointwise) : $d^2(d^2-1)/12$. It is also very important to realize that the curvature tensor, in general, does not determine the metric, even with that huge dimension (when $d > 3$) as compared to the number, equal to $d(d+1)/2$ of parameters for a metric : see the digression in I. C. 1. Opposite the sectional curvature almost always does it.

Computing explicitly the curvature tensor in a coordinate chart $\{x_i\}$ involves the two first derivatives of the metric (say the $g_{ij} = g(\partial/\partial x_i ; \partial/\partial x_j)$) but the expression is very complicated. Very roughly speaking the curvature (tensor or sectional) can be seen as "the Hessian of the metric". This is more apparent at the origin of the normal coordinates of Riemann, those which are given by the geodesics issuing from a given point as an origin. Then the curvature involves only the second derivatives of the g_{ij} . This does not explain but makes very plausible the comparison theorems of I.2 : one can control a function when one has a control on its second derivative. But there is a long way to really state the results and prove them.

It turned out (this will be seen in detail in I. C. 1) that those very geometric and natural normal coordinates are in fact badly adapted for coordinates changes when one controls only curvature. This is because the equation for Jacobi fields in 0. B looks of second order but this is only after having integrated the equation for geodesics.

The discovery of (Jost & Karcher, 1982) is that harmonic coordinates (O. F) are the well adapted ones. With them one does not lose derivatives. They are quite simple to define and build up, being just nicely chosen linearly independent harmonic functions (see I. C. 2 and IV).

If the reader does not like biquadratic forms as algebraic objects and the still not unveiled sectional curvature, he should be immediately more enthusiastic about **Ricci curvature**. It is nothing but the trace (with respect to the metric) of the sectional curvature : for a unit vector x its Ricci curvature $\text{Ricci}(x)$ is the sum of the sectional curvatures $K(x, x_i)$ where $\{x, x_2, \dots, x_d\}$ is any orthonormal basis completing x . One can also consider it as the mean value of the sectional curvature through a given vector. This is a quadratic form and then has the same number of parameters as the metric itself. It has been a vital lead in Riemannian geometry to deduce as much as possible on the metric from its Ricci curvature. This will be seen amply in the sequel. One thing which really helped to unwrap the situation was that in **harmonic coordinates** (O. F) the Laplacian Δ of the metric (see V for the laplacian) is exactly the Ricci curvature up to terms Q of lower order (which are quadratic) :

$$\Delta g_{ij} + Q(g, \partial g) = -\text{Ricci}(\partial/\partial x_i ; \partial/\partial x_j)$$

Then one has at his disposal all the classical Analysis to get information on functions and their Hessian from the knowledge of their Laplacian. For distance functions and Ricci curvature see O. F.

Another main point about the Ricci curvature was in fact discovered in (Bishop, 1963) : a lower bound on Ricci curvature gives an upper (optimal) bound on the volume of metric balls. This will turn out to be basic for topological implications but, even more, for convergence and precompactness results. The proof is quite simple : Ricci curvature is a trace and traces are derivatives of determinants (volumes). For all this see TOP. 1. A and B. In fact Ricci curvature yields the first non trivial term in the asymptotic expansion of the volume form along a geodesic (solid angle). Bishop's result is that can be integrated with a lower bound. Oppositely it definitively cannot be integrated with an upper bound as soon as $d \geq 4$.

Scalar curvature is the trace of Ricci curvature, it is a scalar (at every point). Infinitesimally it gives the first non trivial term in the asymptotic expansion in function of the radius of the volume of balls centered at a given point . But this time it cannot be integrated under a lower nor an upper bound.

THE TOPIC II is in fact the most natural one : study the Riemannian hierarchy. Riemann's idea was to generalize euclidean geometry. So that we look next for geometries which are successively less and less close to euclidean ones. Otherwise stated : relax more and more the euclidean axioms. This is the **space form problem**. It goes back to Killing who convinced Klein of its interest in a lecture. This explained partly the classical phrasing of "Clifford-Klein spaces".

It is the place here to comment on two volets (wings) of Riemannian geometry, namely those of positive and negative curvature. We will see more than once below how much those two volets differ, often paradoxically. In the case of space forms here is the point. Briefly speaking space forms of positive curvature are "childish" to build up and "easily" classified. Negative space forms are very hard to build up, they always need Number Theory even if in disguise. In exchange negative curvature space forms, once they are built up, enjoy many optimal results, see **I. B. 4** .

Oppositely positive curvature space forms are still very mysterious. For example there are almost no reasonable invariant either explicitly known for them or characterizing them : see **III. B** for the minimal volume, **TOP. 1. C** for the embolic volume. And from **III. C** : there are non-uniqueness results for Einstein metrics on them in some cases, but not a single case of uniqueness is known. It starts with S^4 .

THE TOPIC III is satisfactory only for surfaces this thanks to conformal representation. For higher dimension it started with a question of Calabi in 1950. The first general answer to that question came only in (Yau, 1978). Among the few results obtained since, let us mention (Hamilton, 1982). In particular the number one question "does or not any compact manifold of dimension more than 4 admit an Einstein metric" is completely untouched and with no guess from experts. One has only local non deformations results.

THE TOPIC IV started seriously only with the conjunction of (Milnor, 1964) and (McKean & Singer, 1967). But it was then a real blowing up. The links between IV and V are well developed since (Colin de Verdière, 1973), a fact which could be surprising considering what comes next. The point in IV can be described as : build up a Fourier analysis on Riemannian manifolds and study it. This is basic in some pure geometric questions. Let us mention in classical geometry the use of spherical harmonics to study the kissing number of spheres, see the chapter 13 of (Conway & Sloane, 1993) and in our text the control of distance functions in the digression in I. A. 2..

THE TOPIC V remains in a miserable state of affairs. Results started to appear with Poincaré and Hadamard, but despite the initial efforts of Birkhoff, and even more Morse, the today final harvest is still very meager. The case of surfaces is a little better understood. One had to wait (Lusternik & Fet, 1951) for a first general result and the very recent (Rademacher, 1994). It is only for the negative curvature that one has good, in fact excellent, results : as soon (Hopf, 1939) for surfaces and (Anosov, 1967) for the general case.

I : CURVATURE AND TOPOLOGY

Since Heinz Hopf in the late 20's this topic was and remains the strongest incentive for research in Riemannian geometry. Most books on Riemannian geometry give some results in this topic but none is covering it completely, so that we will rely on surveys and references. Books which give special attention to this topic are , after (Cheeger & Ebin, 1975), (Sakai, 1996) and (Petersen, 1997b).

A. Pinching problems

1. INTRODUCTION

Recall that the manifolds of constant sectional curvature are locally isometric to spheres, to euclidean spaces or to hyperbolic ones. So that, if compact, they are compact quotients of the sphere, the euclidean space or the hyperbolic space. For more on these **space forms** see **II. A** . To establish this solidly was one of the tasks of Heinz Hopf in the 30's and one of the motivations for the Hopf-Rinow theorem. Natural is the following : assume a compact manifold has a sectional curvature varying not too much (one will say that the manifold is "**pinched**"). Can one deduce from this that the underlying manifold is, topologically (or more - differentiably), identical with one of the above space forms ? After normalization we are left with three cases : the pinching question around $\sigma = +1, 0, -1$. We assume that the **sectional curvature satisfies** (everywhere and for every tangent plane) $|K - \sigma| \leq \varepsilon$ and look for an ε which will lead to the conclusion that we are on a topological (or more a differentiable) space form. Note that, for the zero case, some normalization is needed, which is done usually by asking the diameter to be not too large. This because stretching the metric by a large factor will make the whole curvature going to zero. We will detail quite a lot the positive pinching problem because of its historical importance.

Back to the general pinching problem we will see that the answer has surprisingly different answers in the three cases. This will start to illustrate the basic differences between space forms of different signs. There is an answer covering the three cases together, but one needs, besides the pinching of the sectional curvature, both an upper bound on the diameter and a lower bound

on the volume. This was obtained in an unified way in (Fukaya, 1990), see theorem 15.1 and is covering prior results : the positive case was solved in 1951 by Rauch, see below, the zero case was solved in (Gromov, 1978a) and the negative case in (Gromov, 1978b).

There are more general types of pinching, the most general being for symmetric space forms. Such a programm was started by Rauch, for references and intermediate results see (Min-Oo & Ruh, 1979; Min-Oo & Ruh, 1981). Informative texts are (Gromov, 1990b) and (Petersen, 1996e). For Ricci-pinching see the very end of 4 below.

2. THE POSITIVE PINCHING

Natural is the following question, repeatedly asked around by H. Hopf, in particular when Rauch (an analyst, expert in Riemann surfaces) visited Zürich in the late 40's. Since the standard sphere is the only simply connected manifold of constant positive sectional curvature and having in mind some heuristic principle of semi-continuity, one can hope to be able to prove that if the sectional curvature is close to a positive constant, the underlying manifold (simply connected) is still the sphere. This was indeed proved in (Rauch, 1951) with a pinching constant (i.e. the ratio between the lower and the upper bounds of the sectional curvature) about $3/4$. Rauch' paper was seminal in two respects. It was first giving a control of the metric on both sides : if the curvature is lower than Δ , then the metric (in a sense of course to be precised) is larger than the metric in the sphere of constant curvature equal to Δ (at least in regions small enough). And mutatis mutandis for the curvature larger than δ . The case where $\Delta = 0$ was known to Cartan and Preissmann as we saw in O. B and O.C. Second, to get a global result, Rauch made a subtle geometric study where he proved that, under the pinching assumption, one can build (by a sort of analytic prolongation using exponential maps) a covering of the manifold by the sphere.

We comment now on Rauch's result and those of his followers. First about the tools, thereafter what assumptions are made on the curvatures (sectional or "only" Ricci), and finally about optimality. We will mainly follow history, a survey is (Abresch & Meyer, 1997).

The basic role of the **injectivity radius** (as linked to the cut-locus: **TOP. 4**) was precised in (Klingenberg, 1959). The injectivity radius $\text{Inj}(g)$ of a Riemannian manifold (M, g) is the largest real number such that every pair of points whose distance is less than it are always joined by a unique shortest geodesic (segment). In particular every open metric ball of radius $\text{Inj}(g)$ (or smaller) is diffeomorphic to $\mathbb{R}^d$. By the elementary foundations of Riemannian geometry for every point there is such a positive number (called the injectivity radius at that point). By the foundations this number is continuous as a function of the point and in particular is positive if the manifold is compact. We will see below how basic is the injectivity radius. In the positive curvature case two results of Klingenberg tell us this : in even dimension and with simply connectedness the double hypothesis $0 < K \leq 1$ implies always that $\text{Inj}(g) \geq \pi$ (think of the sphere). The proof uses Synge's result. This injectivity radius bound is definitively false in odd dimension, see below. But for odd dimensions it remains valid if moreover $1/4 < K \leq 1$. This remains true even for $1/4 \leq K \leq 1$ but the proof is more delicate. The key idea is due to Klingenberg, but some gaps had to be filled. This was done in (Cheeger & Gromoll, 1980), who circulated a preprint as soon as 1972. See also (Klingenberg & Sakai, 1980). For both results the use of Morse theory is basic, see the "long homotopy" below. One mixes it with comparison theorems for Jacobi fields. We will see below in section C. 1 that a lower control of the injectivity radius needs in all generality definitively **four** conditions : an upper and a lower bound for the sectional curvature, an upper bound on the diameter and a lower bound on the volume. Remark that Rauch's proof implies, under his pinching constant, Klingenberg's injectivity radius lower bound.

A digression : comparison theorems. The best possible local control for the metric from $\Delta = \sup K$ and $\delta = \inf K$ follows directly (and for Δ or δ of any sign) from Rauch's estimate for Jacobi fields. Even if other techniques are now available, see below, the geometric metric two-way bounds gotten for triangles (large or small) from $\sup K$ and $\inf K$ are basic. Then Rauch by integration along curves obtained bounds for (small) triangles, i.e. within the injectivity radius. One call these results **RCT** (Rauch comparison theorems).

They are to be distinguished from **Toponogov's** theorem : (Toponogov, 1959). The remarkable fact in Toponogov's theorem is that, for a lower bound on the sectional curvature, one can "ignore

the injectivity radius or the cut-locus" (TOP. 4). This definitively cannot be the case for an upper Toponogov's result : just look at a very small periodic geodesic in a flat torus. However this is possible in the negative curvature case if moreover one has simply connectedness : see Cartan's result in O. B and I. B. 4 below. Toponogov theorem is simple to state : for $K \geq 0$ it just says that every triangle is infra-euclidean : $a^2 \leq b^2 + c^2 - 2bc \cos \alpha$. For $K \geq k$ it is the analogous inequality but as applied to the classical formulas for triangles in the space of constant curvature equal to k , namely the spheres (resp. hyperbolic spaces) when $k > 0$ (resp. $k < 0$). The relevant formulas are the classical ones of spherical or hyperbolic trigonometry.

One should be very careful not to expect too much from these comparison theorems. They are great in the sense that a double control on the sectional curvature gives us a double control on the metric. For example one recovers the classical fact that, locally, spaces of constant sectional curvature are isometric to the standard ones. But for variable curvature things are different. Here are for example two cases where to exert carefulness. The first is that the curvature tensor does not determine the metric in general, the case of constant curvature is completely exceptional. This is a whole topic in itself and which started only quite recently. We will come back to it in the digression of C. 1. The second is the pinching around zero, see 3 below.

Rauch's proof of RCT was quite painful, as well as the proof of Toponogov's theorem. It was discovered in (Gromov, Lafontaine, & Pansu, 1981a) (the "little green book") that the good concept was the control of the distance functions (distance to a point, to some hypersurface, to some periodic geodesic or from the infinite : Busemann functions in B. 2). The key idea of Gromov was to look at the ODE for the second fundamental form of the level hypersurfaces : this is a Riccati equation (but not surprising since any second order linear equation (in particular the Jacobi field equation) can be reduced to one of Riccati type. Things are well written in (Karcher, 1989), thereafter in the lectures notes (Eschenburg, 1994), also in the books (Sakai, 1996) and (Petersen, 1997b). But heuristically, if one thinks of the sectional curvature as "the Hessian of the metric" (see the digression in the 1. COMMENTS on I), such results are not finally too surprising.

Toponogov triangle theorem needs essentially a lower bound for the sectional curvature. But these days one can enjoy more and more metric type results, which were obtained previously using

sectional curvature, now valid under bounds only on Ricci curvature. We saw in O. F why such a hope was not too unrealistic. Among landmarks are : (Calabi, 1958), (Cheeger & Gromoll, 1971), (Abresch & Gromoll, 1990a). Quite astonishing for the reader can be (Colding 1996c). One finds there a statement yielding both an upper and a lower bound for triangles and this with only a lower bound for the Ricci curvature (of any sign). The result is of L^2 -type and implies that most of the (thin) triangles are OK for what one wants to do with them. We state it explicitly :

"in a Riemannian manifold of dimension d with nonnegative Ricci curvature, for any $\varepsilon > 0$ there exists some $\eta(d, \varepsilon) > 0$ such that for any $R > 0$ then for any $r \leq \eta R$ and any points p, q with $d(p, q) > 2R$ one has

$$\frac{1}{\text{Vol}(U(B(q, R)))} \int_{v \in B(q, R)} |d(\gamma_v(r), p) - d(\gamma_v(0), p) - r \langle v, \text{grad}(d(p, \cdot)) \rangle| dv \leq \varepsilon R$$

where $U(B(q, R))$ denotes the unit tangent bundle over the ball $B(q, R)$ ".

There are analogous formulas when the lower bound for Ricci curvature is any real number. Two remarks are now in order, the first is that the triangles should have at least one side which is large with respect to another. But in exchange, they are controlled on both sides. The reader could find surprising to control both sides with only a lower bound for Ricci curvature. The heuristic explanation is this : from the second variation formula and Myer's trick (see O. C and below) one can feel that a long segment forbids along it too large sectional curvatures : a segment being an absolute minimum the second variation along it cannot be negative. This two side control with only a lower ones appeared in fact in the rigidity theorems in B. 2 below for non compact manifolds with nonnegative sectional and Ricci curvature, where infinite segments (lines) were used. Colding likes to think of his result as a predictability one. Given a point p , when one is given the initial velocity of a geodesic starting close enough to q and not too long, then one knows with a good probability, i.e. in a local integral fashion, where is the end of this geodesic as seen from p .

Here the tools are those explained in O. F : playing Analysis, in particular with distance and suitably chosen harmonic functions. Colding's formula introduced a new view point which is put in a

very general frame in various texts quoted below, see in particular for a systematic exposition (Cheeger & Colding, 1997a), (Cheeger & Colding, 1997c), (Cheeger & Colding, 1998). The philosophy could be this : the interplay between the Ricci curvature lower bound and the Laplacian on functions plays a role analogous to that of **RCT** and Toponogov theorems in the case of a lower bound for sectional curvature.

We come now to the optimal constant in the (positive) pinching problem. As soon as (Klingenberg, 1961) one knew that a strict $(1/4)$ -pinching was enough to ensure that the manifold is the sphere. This is called in the literature the **sphere theorem**. And cannot do better since the KP^n (with $K \neq R$) have a curvature between $1/4$ and 1 . The classical proof consists simply in covering the manifold with two topological balls (which are metric balls in the construction, but beware that metric balls are not always topological ones when their radius is too large, typically after the injectivity radius). But only topological spheres can be covered by two balls (Reeb's theorem, see (Milnor, 1963), page 25). To find such two balls one takes two points realizing the diameter and applies Toponogov theorem to prove that the two balls centered at those points and of radius just a little more than half the diameter do have a radius smaller than the injectivity radius. In fact the proof can be precised in order to prove directly the desired homeomorphism, see (do Carmo, 1992), page 287. For "sphere theorems and below", beside what follows, see the survey (Abresch & Meyer, 1997).

An interesting proof is due to Gromov and explained in (Eschenburg, 1986). It is in fact very close in spirit to Rauch's original proof spirit. Look at balls centered at a fixed point : then just after $\pi/2$ the boundary becomes concave (this was proved and used by Rauch). With some work and using positive curvature one shows that the outside of that concave ball can be contracted to a point just by "following the normal field". This proof does not use Toponogov's theorem nor Klingenberg's injectivity radius bound, this bound being in fact a corollary of the proof.

The sphere theorem was triggering geometers to go further. We will now mention the essential developments. Surveys on the topic are (Shiohama, 1990) and (Wu, 1990).

In the sphere theorem as stated above there is a dramatic proviso. A covering with only two balls characterizes spheres but only topological spheres. Now there exist topological spheres with **differentiable** structures different from the standard one (so-called "exotic"). Moreover some exotic spheres are precisely obtained by gluing two half spheres along their equator, but with a weird identification. So that we are led to the "differentiable pinching problem" (since the simply connected manifolds of constant positive curvature are the standard spheres, see 0. B).

There is also the problem to see one can be done for the non simply connected case. For sure, looking at the universal covering, one is sure that the manifold downstairs is a quotient of a sphere, but this does not say that the topology is that of a space form. The problem is the **equivariant** pinching question. It was solved in (Im Hof & Ruh, 1975). For the non-simply connected case and the diameter see (Flach, 1994).

But the pinching constant is still under work, the latest reference is (Suyama, 1995) with a pinching independent of the dimension : 0.654 . See also the references there for intermediate results. Nobody knows if $1/4$ is possible or not. Worst : there is none exotic sphere known which has positive curvature, only nonnegative ones are shown to exist in (Gromoll and Meyer 1974). However in (Weiss, 1993) it is shown that many exotic spheres cannot be strictly $1/4$ -pinched.

Historically the differentiable pinching problem was solved by Calabi (unpublished) and in (Shikata, 1967) by methods close to those which will appear in the section C below, but yielding no explicit pinching constants. Then (Gromoll, 1966) did this : looking back at the proof "with two balls" centered at diametrically opposite points, one uses Toponogov and an extra control of parallel transport to show that the gluing of the two balls along their intermediate $(d-1)$ -sphere is closed to the identity. Then differential topology implies a total diffeomorphism. Gromoll pinching constants were explicit, but going to 1 with the dimension going to infinity.

The method of (Ruh, 1971) was completely different : the sphere was seen "from outside". The fact that the curvature is very pinched enables us to construct a line bundle on the manifold M^d which resembles enough to the normal bundle of the standard sphere in $\mathbb{R}^{d+1}$ to yield finally an embedding from M into $\mathbb{R}^{d+1}$. This embedding yields a sphere closed to the standard one and

hence diffeomorphic to it. There are other techniques, in (Otsu, 1993) for example one uses thin triangles. For lower Ricci and volume bounds, see below.

There is also the recent "pinching" result (Grove & Wilhelm, 1995) yielding manifolds diffeomorphic to the standard sphere. It uses in the hypothesis a pure metric invariant called $\text{pack}_{d+1}(g)$. For a Riemannian manifold (M^d, g) it is defined as one half of the maximum of the minimum non zero mutual distances between $d+1$ points : the maximum is to be taken over all such sets of points in M . When one consider only two points instead of $d+1$ one has the definition of (half) the diameter. The result is then : if the sectional curvature is such that $K \geq 1$ and $\text{pack}_{d+1} > \pi/4$ then we have diffeomorphism with the standard sphere. The proof uses Alexandrov geometry (see TOP. 9) and is also a method "from outside" as Ruh's one, except it just uses the metric to get a suitable embedding. A very informative survey of this kind of results (and of some other below involving the radius) is (Grove, 1992). Another type of result was (Otsu, Shiohama, & Yamaguchi, 1989): if $K \geq 1$ and the volume of the manifold close enough to that of the standard sphere, then one has diffeomorphy. But now one can obtain this with only Ricci curvature, see below.

Natural now is to ask what is happening just at $1/4$ or a little below $1/4$? There is a survey : (Abresch & Meyer, 1997). We only give the essential points. First for $1/4$ exactly, if we are not on a sphere, then we are necessarily on a KP^n and moreover with the standard metric : (Berger, 1960). This kind of theorem is called a **rigidity** result. One can think of it as a susceptibility result : the smallest touch to the standard metric of KP^n (with K not $\mathbb{R}$) will forces the curvature range to be strictly outside of the interval $[1/4, 1]$.

But today one can go a little bit below $1/4$. There is an $\varepsilon(d) > 0$ such that a $(1/4 - \varepsilon)$ -pinching implies that one is on a sphere or on a KP^n : (Berger, 1983). The proof was not only incomplete, see the survey, but moreover the ε was in the air since gotten by the compactness result in C. 2 and contradiction. In (Abresch & Meyer, 1994b) and (Abresch & Meyer, 1996b) one has explicit $\varepsilon(d)$ pinching constants. But it yields the optimal topological answer only in odd dimensions. For even dimension one gets only a strong topology information which is yielding "almost only" the KP^n . See the references for more details. The proof is very geometric, using Toponogov control of triangles on one hand and Morse theory and

parallel transport on the other to control the injectivity radius. The proof is as follows. If the injectivity radius were too small it will exist some very small periodic geodesic. The simply connectedness enables us to deform it through a point by a "long critical homotopy". The homotopy has to be long because the mountain pass point is a conjugate point, and of index exactly 1. But the curvature conditions yields finally a contradiction. The best $\varepsilon(d)$ is unknown, the present one is of the order of 10^{-5} but in view of B. 1 any guess is senseless.

We have to stop here for the pinching constant. In fact, to go further below $1/4$ will mean merely look for a classification of all manifolds (since any manifold has some Riemannian structure) or at least for manifolds of positive curvature : see B. 1 and C.

But the game is far from being finished, because one would like to get results with weaker hypothesis. An important event was (Grove & Shiohama, 1977). The result is that one still gets a sphere when asking only the curvature to be lower bounded by 1 and the diameter to be larger than $\pi/2$. The upper bound for the curvature disappears. This is perhaps not too surprising, as we saw in Colding's comparison theorem in the digression above, because of Myers's type results. The technique was new, it still used Toponogov for working, but the main point is to be able to go **beyond** the injectivity radius. The trick is to use the distance function $d(p, \cdot)$ from a point p . One says that a point q is **critical** for $d(p, \cdot)$ if for any direction v at q there exists a segment (a shortest geodesic) from p to q whose speed vector w at q makes with v an angle not larger than $\pi/2$. Heuristically this means that in any direction one can get closer to p . Two extreme examples of critical pairs are : first when the distance between p and q is equal to the diameter. This was exactly the idea of the proof of the sphere theorem above. The second case is when the two points are antipodal on a periodic geodesic.

If q is not critical then a contrario one can find some direction in which $d(p, \cdot)$ is strictly increasing. Then, as above, one succeeds to cover the manifold with two topological balls : like in Morse theory one can "push things without changing the topology" as soon as one does not meet a critical point.

The study of what is happening for the limiting case where the diameter is equal to $\pi/2$ was strongly advanced in (Gromoll & Grove, 1987) but is still not finish.

The **critical point** technique is now a basic tool in many instances. It was put in quite general a frame in (Greene & Shiohama, 1981a) and (Greene & Shiohama, 1981b). Surveys of this technique and its applications are : (Cheeger, 1991), (Grove, 1985), (Meyer, 1989), (Grove, 1990), (Greene, 1990), (Karcher, 1989). See also various places below.

In fact Grove-Shiohama result has a flavor close to Myers theorem ((Myers, 1941), see O. C) which say this. Assume the Ricci curvature is lower bounded by $d-1$. Then the diameter is bounded from above by π . Then one can ask for a Myers-type diameter pinching result. The first thing even to do is to look if the equality characterizes the standard sphere. This can be done in many ways, the first appearance is (Cheng, 1975). In (Anderson, 1990a) is was shown that such a pinching result **cannot** exist, see also (Otsu, 1991). This was achieved by constructing suitable examples with $\text{Ricci} \geq d-1$ and diameter closer and closer to π , with tools from Riemannian submersions. Oppositely, with an extra positive lower sectional curvature bound, one can get the conclusion : (Perelman, 1995b).

But if the diameter is replaced by the **volume** then there came first a result (Shiohama, 1983) : there exists $\varepsilon(d)$ such that $\text{Ricci} \geq d-1$ and $\text{volume} > \beta(d) - \varepsilon(d)$ imply homeomorphism with the sphere, but with an extra lower (negative) bound for the sectional curvature (of course here $\beta(d)$ denotes the volume of the standard sphere S^d). In (Perelman, 1994a) homeomorphism with the sphere is obtained without any extra condition. The proof is very geometrical. In brief it mixes techniques of critical point theory, contractibility of balls in concentric ones of larger radius and playing smartly with algebraic topology (see B. 1). Gromov's view of it is "non trivial homology generates large **shadows**", hence restricts too much the volume . Needless to say that the injectivity radius is ignored. If one still uses the full force of $K \geq 1$ and $\text{volume} > \beta(d) - \varepsilon(d)$ then the same conclusion was proved in (Otsu, Shiohama & Yamaguchi, 1989), this with an explicit ε . The technique is to build up an embedding into $\mathbf{R}^{d+1}$, it is an "outer" method. But all these results are now superseded by (Cheeger & Colding, 1997a). In its appendix 1 diffeomorphism with the

standard sphere is proved under a lower Ricci bound and a volume closed to the canonical one (moreover the volume bound can be explicitly computed). This is only the emerged part of an iceberg, still not finished to be fathomed. We will see below other results of that vein. We suggest the reader to read the introduction and appendix 2 of (Cheeger & Colding, 1997a) for a survey of ideas, results and a programme concerning Ricci curvature. Among other tools Colding L^2 -Toponogov type theorem mentioned in the digression above is basic. The fact that "thin" triangles are useful, as they were first for the splitting theorem in B. 2 below, can be heuristically explained if one sees sphere theorems as "splitting ones" (into two balls). A systematic tool introduced by the authors for the general study of manifolds with a lower bound on Ricci curvature is that of suitable warped products (O. G).

A recent pinching with "distances" result is (Colding, 1996a). There the diameter is replaced by a very close but still stronger metric invariant, the **radius**. The radius looks very similar to the diameter but is surprisingly more powerful. It is the minimum of the radius of a metric ball which is containing (and hence equal to) the whole manifold. Think of a very thin ellipsoid : then the radius is close to half the diameter. It is proved in (Grove & Petersen, 1993) that $K \geq 1$ and $\text{radius} > \pi/2$ implies homeomorphy to a sphere. Colding result is that $\text{Ricci} \geq d-1$ and radius close enough to π imply homeomorphism with the sphere (how close enough is demanded can be made explicit). The idea of the proof is to show that the Gromov-Hausdorff distance (C. 2) between the manifold and the standard sphere can be made as small as wanted. This is done by squeezing suitable balls in the manifold and defining with them a map to the sphere. Then one uses the predictability result (see the digression in A. 2) of Colding as applied to the unit tangent bundles of these balls. Using again and again Bishop-Gromov estimate for the volume of balls (TOP. 1. A), volume arguments finish the proof.

One should mention also the **pointwise pinching** problem : the curvature is asked to be squeezed in a given ratio at every point, but the global sup and inf need not to be squeezed. Lack of space we just refer to the almost final result of (Margerin, 1991, 1993, 1994), and to it for detailed statements and the very subtle control of the deformation technique. A nice survey on the **deformation technique** along the Ricci flow (see 0. F) is the section 4 of (Besse, 1994). See the solution of the (1/4)-case in B.1 . For an exposition in book form of some of these results see (Hebey, 1997).

And one would like again weaker hypotheses. The drive started with sectional curvature, then one tried to replace it by Ricci curvature and now finally by **integral norms** for various curvatures. We give only some references : (Gallot 1987), (Anderson and Cheeger 1991a), (Gao 1990a), (Gao 1990b), (Petersen & Wei, 1996c), (Anderson, 1990d) and in the book (Petersen, 1997b). Note that the work to do here is more Analysis than geometry. See also C. 1.

3. PINCHING AROUND ZERO

The space forms with zero curvature are flat tori or finite quotients of them (see II). Can one prove the following : there exists some $\varepsilon > 0$ such that if a manifold has a diameter bounded by 1 and a curvature bounded both sides by ε and $-\varepsilon$, then this manifold admits a flat structure? Oppositely to the positive case the answer in the present case is **no**. There are other manifolds known to be "almost flat", namely the so-called "nilmanifolds". They can be defined as compact quotients of nilpotent Lie groups. Topologically they are nothing but successive fibrations whose fibers are always a circle and ending in a point. They admit, for any $\varepsilon > 0$, a metric with curvature between $-\varepsilon$ and ε and diameter equal to 1. Heuristically, this almost flatness is possible because a circle (a curve) has no curvature. Then one just applies the formulas for Riemannian submersions (O. G). See also C. 3.

The basic paper (Gromov, 1978a) proved the converse : the solution of the pinching problem is true, but the answer are not flat manifolds but nilpotent ones (exactly in fact the so-called *infranilmanifolds* namely the finite quotients of nilmanifolds). The complete statement was in (Ruh, 1982). It says that there exists some universal $\varepsilon(d) > 0$ such that this is the conclusion for manifolds with $-1 < K < 1$ and diameter $< \varepsilon(d)$. A detailed proof of Gromov's result is given in (Buser & Karcher, 1981). Techniques are purely geometric but a complete proof is quite long and involved. Two good sketches of it are : (Sakai, 1996), pages 319 up and (Fukaya, 1990), § 8 and 9. Using Toponogov triangle control, one studies in detail the fundamental group as made up by isometries of the universal covering. The curvature being extremely small, the exponential map is a covering at very large distances and moreover not much different from an isometry. Then the elements of the fundamental group almost commute, as can be seen by controlling

also the parallel transport (remember the golden triangle : the curvature is related to the parallel transport around infinitesimal parallelograms). From the control on the commutators of the fundamental group one can boil down to nilpotent groups. Except for the best $\varepsilon(d)$ still to be found, this result is then optimal.

4. THE NEGATIVE PINCHING

Here the answer is even more no than for the zero case. First in (Gromov & Thurston, 1987c), then in (Farrell & Jones, 1989a), one finds many manifolds with curvature as close as wanted to -1 which never can have a metric of constant curvature. Do not hope either a general result around $1/4$ for the negative HypKP^n , as shown in (Farrell & Jones, 1994). The techniques used are very geometric. One uses smart gluings along totally geodesic hypersurfaces in one instance. In the other one makes connected sums with exotic spheres and then controls the curvature with the formulas for the curvature in Riemannian submersions. In the first type of construction the resulting topology can be quite sophisticated but in the second technique one gets only manifolds homeomorphic to space forms. Note that from section C.1 larger and larger diameters are forced. For real analytic constructions, which a harder game, see (Abresch & Schroeder, 1996a).

There is however the following optimal result. Assume M is with $-4 \leq K \leq -1$ and that $\pi_1(M)$ is group-isomorphic to the π_1 of a compact space form of negative curvature which is moreover not of constant curvature (i.e. not real hyperbolic but of rank 1). Then M is isometric to that space form. This was in (Ville, 1985a) for dimension 4, for the complex case independently in (Hernandez, 1991) and (Yau & Zheng, 1991). For the quaternionic and the Cayley case this follows from (Hernandez, 1991), (Corlette, 1992) and (Gromov, 1991c). Compare this rigidity result with the one in A. 1.

All the above was concerning various hypotheses to get space forms (specially of constant curvature) under various pinching conditions. These conditions were always first, at least on one side, involving almost always sectional curvature and the aim was to get space forms. They always started with $|K - \kappa| < \varepsilon$. Suppose we want to do the same for Ricci curvature. Then the first thing to do is to discover who are the manifolds with constant Ricci curvature. This will be the topic of the entire section III. C. Such manifolds are called Einstein. Roughly speaking the answer, starting dimension 5

, is almost completely open (the reader will agree or not with me after reading that section). But it is still a natural question to ask if a strong enough Ricci pinching on a given manifold implies for this manifold the existence of some Einstein metric. Moreover such kind of results could be a clue for the search of these wild animals.

There is a very good answer in (Anderson, 1990c) : there exists an universal ε depending only d , i , D , σ such that a d -manifold with diameter smaller than D , injectivity radius larger than i and Ricci pinching $|\text{Ricci} - \sigma| < \varepsilon$ always admits an Einstein metric. The ε is in the air since gotten by a contradiction proof using general compactness results (see C. 2).

B. Curvature of a given sign

We are going to see again how much different is the landscape for the positive and for the negative sign.

1. THE POSITIVE SIDE : SECTIONAL CURVATURE

Surveys are (Gromoll, 1990), (Abresch & Meyer, 1997), appendix A, and (Greene, 1997). We are still here in an almost completely mysterious situation. We describe now the only known examples of (compact) manifolds with **positive** sectional curvature : because of Myers's theorem, except for a finite classification problem (although not trivial), we can afford essentially to look only at the simply connected case. First come the spheres, the KP^n and few other homogeneous spaces of dimensions respectively 7 and 13 .

Among them the **Aloff-Wallach** manifolds $W_{p,q}$ are specially interesting : (Aloff & Wallach, 1975)). They are suitably defined homogeneous metrics on the quotient of the Lie group $SU(3)$ by the " (p,q) circle" : this means that the direction of the circle is given by the point of coordinates (p,q) in the integral lattice defining a maximal torus of $SU(3)$. Recall that all maximal tori of $SU(3)$ are of dimension 2 and conjugate. Moreover this "universal lattice" is the regular hexagonal one. In the semi-desertic area of manifolds of positive curvature Aloff-Wallach manifolds are in fact a fascinating family. We mention **three** of their properties. First according to the couples (p,q) they have an infinite number of homotopy types (and even for some couples one can get not diffeomorphic ones). Second they carry for every (p,q) a metric with a positive lower pinching not much different from $16/29.37$: see (Huang, 1981). Third they also admits homogeneous Einstein

metrics and if the volume is normalized to 1 the set of Einstein constants (which are all positive) is infinite : see (Wang, 1982) (Wang & Ziller, 1986). Note that these Einstein metrics first are not of Aloff-Wallach type, in particular they have sectional curvature of both signs. Another of their properties is that the set of those positive Einstein constants, when the volume is normalized to be one, does not converge to a smooth metric : this shows that the Palais-Smale "C- condition" (see (Jost, 1995)) is not valid for Riemannian metrics, at least in the today known possibilities : see **III. C** for this type of question.

There is a complete classification of positive curvature homogeneous spaces : there are no other homogenous space of positive curvature than the above ones : (Bérard Bergery 1976). It was discovered in (Wilking, 1997) that, if this result is true, there was some curious mistake in the proof, finally compensated by the fact that some space missing in the list of (Berger, 1961) turns out to be isometric to one of the Aloff-Wallach spaces.

Only four other types of examples (not homogeneous) are known. They are in dimension 6 and 7 : (Eschenburg, 1992), (Taimanov, 1997) and in dimension 13 : (Bazaikin, 1995). The last ones are in a family close to Aloff-Wallach examples. Pinching constants for these various spaces have funny coincidental properties which are explained in (Taimanov, 1997). Taimanov's study is pursued in (Püttmann, 1997) where the pinching constant $1/37$ appears and is explained via Taimanov's deformations of Aloff-Wallach metrics. The starting point is to embed Aloff-Wallach manifolds as totally geodesic submanifolds in the 13-dimensional examples. Moreover there is enough "room" (transversality) to deform the metric quite a lot and still keep the totally geodesic property. One even gets a series of metric which converge to a smooth one with a pinching constant equal to $1/37$. This supports the conjecture that the best pinching constant for Aloff-Wallach manifolds is $1/37$ and this is precisely proved in (Wilking, 1997).

No other manifolds are known of positive curvature, except of course small enough deformations of the preceeding ones.

If we turn to manifolds of nonnegative curvature, the situation is not really better. As examples we have all symmetric spaces of "positive" type (see **II**). As well as Riemannian products of the above known ones of positive and of nonnegative ones. Besides those to our knowledge the only examples are some exotic spheres

and the connected sum of two symmetric spaces of rank one : see (Cheeger, 1973) and (Gromoll & Meyer, 1974). Those examples are built by a clever mixture of large actions (low cohomogeneity) of isometry groups and Riemannian submersion technique to compute the curvature.

But one knows at least that not every manifold can carry a metric of nonnegative curvature (a fortiori of positive curvature). First came the basic theorem of (Gromov, 1981d) : manifolds of nonnegative curvature have a sum of Betti numbers (for every field the way the proof goes) universally bounded by the dimension. Hence we are left with only a finite number of (integral) **homology type** (see C. 1). The tools is somewhat, but not in reality, analogous to Morse theory. It requires subtle algebraic topology arguments, based on compressibility of balls and topological "content" of them. The basic lemma is that, by Toponogov theorem for triangles, a succession of critical points for the distance function (see A. 2) to a given point and whose distances are growing at least as a geometric progression, has to be **finite** (the cardinal being universal in the dimension). In short "critical points cannot be too far away". To prove this the idea is as follows : the initial speed vectors from the point p under consideration to two different critical points, when their distances are much different, has to be an angle large enough. This means putting on the unit sphere at p many points whose spherical all mutual distances are larger than a given number : such sets have to have a cardinal bounded from above. A rough (but not too bad by the way) estimate can be obtained just by a measure computing of spherical caps. For refined estimates, which is a central problem in pure and applied mathematics, that of spherical codes, see (Conway, 1995).

Then one has to play with distance functions to different points. It involves in particular the following idea : small fingers (fingers with a very small wedding ring) do not count in topology, but to discard them one has to look at what remains from the topology of a ball in the concentric ball of sufficiently larger radius. The reader can look at the various presentations, involving also simplifications of the original text : (Cheeger, 1991), (Meyer, 1989) and the text (Grove, 1993) which explains in a very short time what is happening. Today the bound for the Betti numbers is huge, the conjecture is that their sum is bounded by 2^d with equality only for tori (then necessarily flat, see below at least two reasons for that).

Gromov result was extended in (Abresch, 1985) to the non compact complete case. The play with concentric balls such the small one is contractible in the larger one is essential today, for example in Perelman above and for finiteness results in C. 1 .

Do not hope now to extend this to Ricci curvature. There are examples of nonnegative Ricci curvature with any Betti numbers : this was achieved simultaneously in (Anderson, 1990d) and (Sha & Yang, 1991). In (Sha & Yang, 1991) one builds up connected sums, using delicate metric surgery on double warped products. An extra difficulty here is that one needs to spread the modifications all over the manifolds under the gluing operation. In (Anderson, 1990d) one uses models from (Gibbon & Hawking, 1978). This construction was put in the more general framework of Schwarzschild metric in (Anderson, 1992c). More recently (Perelman, 1997) built up examples showing that Betti numbers can still be not bounded when the Ricci curvature is lower bounded even if both the volume is bounded from below and also the diameter from above (contrast this with the finiteness theorems in section C. 1).

The above still does not tell us the smallest thing about the question : is there any difference between positive and nonnegative sectional curvature ? A baffling remark in (Yau, 1982), page 670, is that one does not know any (compact) simply connected manifold with nonnegative curvature for which one can prove that it **does not** admit a metric of positive curvature. For example Gromov bound on Betti number does not make any difference between positive and nonnegative. Yau notice starts with Hopf's conjecture on $S^2 \times S^2$, see the last alinea of B. 5 . For the non-simply connected case, Rong's results below provide a partial answer.

Positive and nonnegative sectional curvature is a case where **non compact** manifolds enjoy optimal results, so that we make an exception to our rule. Moreover this is basic when studying compact manifolds with an infinite fundamental group. In (Gromoll & Meyer, 1969b) it is proved that a complete non-compact manifold of positive curvature is diffeomorphic to $\mathbf{R}^d$. The basic geometric notion in non compact manifolds is that of **ray** (half-line) : it is a geodesic defined on $[0, \infty[$ which a segment on every subinterval. There are in some sense at least as many rays as "ends" of the manifold. The basic fact is that to any ray is associated a strongly convex subset, namely the set of points which are at negative distance (suitably normalized) from the point at

infinity on that ray. We will come back to these objects called **Busemann functions** in section 2 below. But they were introduced and systematically studied in the general case in (Greene & Shiohama, 1981a) and (Greene & Shiohama, 1981b).

In (Cheeger & Gromoll, 1972) it is proved, with more technicalities, that a complete manifold of nonnegative curvature is vector bundle over a compact totally geodesic submanifold of positive curvature. Such a submanifold is called a **soul**, it need not to be unique (think of a flat cylinder). This theorem was recently precised in (Perelman, 1994b), see it for the references to intermediate results. Recent exposition in book form are in (Petersen, 1997b) and (Sakai, 1996).

A final note is in order. Topological spheres can be characterized (up to homeomorphism today for $d \geq 5$) by a suitable positive curvature condition, namely that of the **curvature operator** : (Micallef & Moore, 1988). The curvature operator is the symmetric endomorphism of the 2-exterior tangent bundle defined by the curvature tensor. The proof uses strong geometric analysis, namely existence of minimal harmonic maps of S^2 into our manifold, see **TOP. 7**. Then one analyses the second variation. This will finally yield the fact that the homotopy groups of the manifold are all vanishing in dimension smaller than that of the manifold and now use the solution of Poincaré's conjecture for $d \geq 5$. Using Bochner's technique for exterior forms (Meyer, 1971) was yielding only real homology spheres. For $d = 4$ one has even a diffeomorphism optimal result : (Hamilton, 1986). The tool this time is the deformation of Riemannian metrics, see comments in **O. F**.

An interesting byproduct of Micallef and Moore is the solution of the "pointwise" $(1/4)$ -pinching problem already mentioned in **A.2**: we assume that at every point m one has $1/4 < K(m) \leq 1$. Then the manifold is homeomorphic to the sphere. Note that such a result is completely out of the reach of **RCT** and/or Toponogov type results.

To go just at the zero level is only partially solved today : a latest reference is (Chen, 1990). The proof is interesting, using at the same time the Ricci flow and the holonomy classification (**TOP. 3. A**). The hope "of course" is to prove that besides spheres only the KP^n can appear.

Back to a "classification" of the manifolds of positive curvature, we are still lacking any general structure statement. We quote here **four** different approaches to attack the problem. The **first** and the most promising is that of (Rong, 1996a). Thanks to the finiteness theorem in C. 1, in a given even dimension, there are only finite diffeomorphism types of manifolds with $0 < \delta \leq 1$. If the dimension is odd, we no longer have Klingenberg's lower bound π for the injectivity radius. The first thing to do is to look at the odd-dimensional case, always here for given dimension and positive pinching δ . One of the results in (Rong, 1996a) is that, modulo a finite number of cases and in the simply connected case, one can modify the Riemannian metric into a new one, still of positive curvature, but admitting now a circle acting freely and by isometries. Since the quotient is also of positive curvature by the Riemannian submersion formulas, one has then reduced the classification to that of the even dimension case and the classification of some circle bundles over them (always modulo a finite number). For this kind of results see also (Tuschmann, 1996). The starting idea is to look at the conditions in Cheeger's finiteness theorem (C. 1) and to assume that things do not work, namely that we have a very small volume. But then we should be in the collapsing case (see C. 3). Thereafter one applies the structure results for the collapsing situation and the main point now is to change the metric in a new one which satisfy the condition above. This is achieved in particular using the Ricci flow, but note that the pinching is now smaller than δ .

For the non simply connected case, recall first that by Synge's theorem in O. C, this is again an odd-dimension story. The fundamental group π_1 is finite by Myers (O. C) and in (Gromov, 1978a) one finds a dimension-depending-only upper bound for its number of generators, using smart basis for π_1 as acting on the universal covering and Toponogov. But we would like a upper bound for the possible group structures, this in function of the pinching. An old conjecture of Chern was that every abelian subgroup of the π_1 of a manifold of positive sectional curvature is cyclic. In (Rong, 1996a), by the very technique used for the proof above, it is proved that this is true, but only up to an index which is bounded by an universal constant $w(d, \delta)$ in the dimension and the pinching δ . Rong's conjecture is today is that $w(d, \delta)$ can be chosen independent of δ . With a much deeper analysis of the collapsing case, (Rong, 1996b) is a partial result in that direction, namely there are $w(d)$ and $w'(d, \delta)$ such that π_1 either has a finite

cyclic subgroup of index less than $w(d)$ or has order less than $w'(d, \delta)$. In

(Rong, 1997) things are refined.

Of course, always looking for a complete classification, the basic question is : look at all manifolds M^d of a given dimension and admitting a positive curvature metric and call $\delta(M)$ its best pinching constant (see below if any doubt). Define $\delta(d)$ as the infimum of all these $\delta(M^d)$: is $\delta(d)$ positive ?

In another direction an interesting result was conjectured in (Gromov, 1988c) : nonnegative curvature manifolds "look like convex bodies, like ellipsoids". The precise notion is that of various **widths** for a Riemannian manifold : they are purely metric invariants. This conjecture was settled affirmatively in (Perelman, 1995a).

A third approach to nonnegative curvature is that of the so-called "elliptic rational homotopy" in (Grove & Halperin, 1983) which is linked to the study of manifolds all of whose geodesics are periodic, see V. B (this should not be too surprising, thinking of the KP^n). Rational homotopy is directly linked with the Sullivan's fine theory of de Rham cohomology, see TOP. 6. B. This approach leads to the so-called "double soul" problem, see more in (Petersen, 1996e), problem 21.

A last approach would be to look at the **best pinched** metric on a given manifold admitting a metric of positive curvature. There is an optimal pinching δ (of course $< 1/4$ to be of interest). At least in even dimension now the hypothesis $\delta_i \leq K \leq 1$ for a sequence of δ_i converging toward the upper bound δ satisfies the hypothesis of convergence in C.2 : the diameter is bounded above by Myers theorem and the volume from below because of Klingenberg's result on the injectivity radius. So that we have an optimal metric, only presently of class $C^{1,\alpha}$. It seems there is no text studying this approach. The main difficulty is that it seems very hard, at least today, to use the pinching condition. More precisely when one looks for extrema one can always first express that the differential is zero. This for differentiable objects and sectional curvature, for a given tangent plane, is of course differentiable for families of metrics (see the formulas and more in (Bourguignon, 1973)). But the curvature is such a weird animal that the points where the sectional curvature is extremal are still not understood even in

dimension 4 (unless the manifold is Einstein : (Singer & Thorpe, 1969)), recall the digression in 1. COMMENTS on I.

The last question concerns **moduli** : assume a manifold admits at least one metric of positive curvature. How looks the set of all such metrics ? We know only today that it can be non connected, an example being the two Aloff-Wallach manifolds $W_{-4638661, 582656}$ and $W_{-2594149, 5052965}$, which are diffeomorphic but cannot be connected by a path of positive scalar curvature between them : see (Kreck & Stolz, 1993) and section 3 below.

2. THE POSITIVE SIDE : RICCI CURVATURE

For the Ricci curvature (in the compact case), it is **still** a mystery today to know what are exactly the implications of its positivity, except the finiteness of π_1 and except in dimension 3, see below. But for the fundamental group one has very strong and optimal recent results for nonnegativity and just below. We saw already an application of Bochner technique in O. D. Precisely if the Ricci curvature is nonnegative, then the first real Betti number b_1 is bounded by the dimension, with equality only for flat tori. A homotopy result is that of (Milnor, 1968) to the effect that the fundamental group is of at most polynomial growth when the Ricci curvature is nonnegative : this is obtained from a counting balls argument by volume in the universal covering using Bishop's result in TOP. 1. A to control the volume of these balls. But sadly enough which groups are really possible is far to be known today. We know only from (Gromov, 1981c) that the groups looked for are discrete subgroups of nilpotent ones.

The challenge now is to relax the hypothesis and to go a little bit below zero. There is a heuristic hope for that. Recall now Bochner's result in O. C : since $\text{Ricci} \geq 0$ implies $b_1 \leq d$, it is reasonable to expect that b_1 , being an integer, will remain $\leq d$ when the Ricci curvature is not too negative (with some normalization). And moreover equality should be expected only for flat tori. Partial results appeared in (Gromov, Lafontaine, & Pansu, 1981a) and (Gallot, 1981). Using the basic Colding comparison theorem (see the digression in A. 2) one has today two very strong answers in (Colding, 1996b) and (Cheeger & Colding, 1996d). First there exists some $\varepsilon_1(d) > 0$ such that $\text{Ricci. Diameter}^2 > -\varepsilon_1(d)$ and $b_1 = d$ implies that the manifold is diffeomorphic to a torus ($d \neq 3$). Second there exists an $\varepsilon_2(d) > 0$ such that $\text{Ricci. Diameter}^2 > -\varepsilon_2(d)$ implies that the fundamental group is almost nilpotent. For

the moment both ε_1 and ε_2 not explicit, since obtained by a limit-contradiction argument.

The latter ε_2 -result is incredibly stronger than the "old" pinching around zero result in A. 3. In fact this result was conjectured by Gromov and proved in the case of a lower bounded sectional curvature in (Fukaya & Yamaguchi, 1992). The proof involves a deep study of collapsing under only a lower bound, which is done in (Yamaguchi, 1991) and the authors mentioned there that their technique could be extended to the Ricci case modulo some reasonable "splitting" conjectures. The proof of the Ricci case uses the frame of the proof of the sectional case, but also new techniques coming from a deep study of the geometry of non compact Riemannian with Ricci curvature lower bounded, in particular an extension to Gromov-Hausdorff limits (see below) the splitting theorem of Cheeger and Colding.

The ε_1 -result is proved as follows. One starts with Gromov's proof to the effect that $b_1 \leq d$: on the universal covering of a suitable finite covering of diameter D , one constructs a basis of the deck transformations such that the generators displace the points not too much but every element displace them enough (in order to avoid too squeezed triangles). Independently, on suitable balls, one constructs d harmonic functions which have L^2 -small Hessians and have L^2 -almost orthogonal gradients. One mixes this with predictability arguments and a smart basis as above to obtain, when $\inf(\text{Ricci} \cdot \text{Diameter}^2)$ goes to zero, a Gromov-Hausdorff limit which is a torus.

Last, not least. In dimension 3 at least the situation is completely understood. In (Hamilton, 1982) it is proved, by the deformation technique along the Ricci flow (see O. F), that any positive Ricci metric can be deformed into one of constant sectional curvature. Note that in dimension 3 Ricci and sectional curvature have the same number of parameters, namely 6. In (Sha & Yang, 1993) it is proved that a simply connected manifold of dimension 4 admits a metric of positive Ricci curvature if it is homeomorphic to one which admits a metric of positive scalar curvature.

The nonnegative versus the positive one is little mysterious: there is no known manifold having some metric with positive scalar curvature (see just below), simply connected and for which one can prove that there is not a single metric with positive Ricci curvature. This will be less surprising if you look here at a fact

from **III. C** : Ricci flat manifolds are very mysterious, one only knows some examples.

Manifolds of positive and nonnegative Ricci curvature have a very nice and useful behavior when non compact (it was heavily used already in the results above for the Ricci lower bounded case) , so that we make here again an exception to our rule. Here the basic notion is no longer that of ray but of a **line**, namely a geodesic defined on $]-\infty, \infty[$ which is a segment for any of its intervals. Assume there exists a line in a manifold with $\text{Ricci} \geq 0$. Define along this line two distance functions (one from $+\infty$ and the other from $-\infty$, normalized of course : they are called **Busemann functions**). Then use the basic formula for the Hessian of a distance function (see **0. F**), the Ricci condition finally implies that these two functions should coincide and that their gradient is a parallel vector field. This forces the manifold to split as a Riemannian product with $\mathbf{R}$. Looking now at the universal covering of a compact manifold with $\text{Ricci} \geq 0$, one can prove the **rigidity** result: the universal covering of M^d splits as a Riemannian product of $\mathbf{R}^k \times M^{d-k}$ with some compact manifold M^{d-k} . So that for example the fundamental group is extremely special, namely the situation is essentially reduced to the finite case . The above result is in (Cheeger & Gromoll, 1971), see also (Eschenburg & Heintze, 1984). Before a splitting was obtained in (Toponogov, 1964), asking more, namely the sectional curvature to be nonnegative. The proof consisted in looking at Toponogov's theorem" when one goes to the infinite" along a line on both sides. Namely one ultimately proves that equality has to occur in the comparison theorem and this is forcing all the transverse sectional curvatures to vanish.

A new technique for studying the complete nonnegative Ricci case was introduced in (Abresch & Gromoll, 1990a). One proves that under a diameter growth condition the topology is bounded. The result can also thought of as a weak quantitative generalization of the splitting theorem. The tool was the first appearance of "thin" triangles . Colding's L^2 -Toponogov theorem quoted in the digression in **A. 2** is a dramatic generalization. The strongest splitting results is the one for Gromov-Hausdorff limits, in (Cheeger & Colding, 1996d), §6 .

Moduli for positive Ricci curvature are not well understood. A recent result is in (Kreck & Stolz, 1993) : some Aloff-Wallach manifolds have a set of positive Ricci curvature metrics whose

number of connected components is infinite : in fact what is proved in those metrics cannot be connected even within positive scalar curvature, see 3 below.

3. THE POSITIVE SIDE : SCALAR CURVATURE

The story here is one of the most beautiful chapter of recent Riemannian geometry (even if still not completely finished). Today we know exactly which are the simply connected manifolds able to carry of metric of positive scalar curvature : the keystone is (Stolz, 1992), see the report (Rosenberg & Stolz, 1994) for the many intermediate results. Bring your simply connected compact manifold M to a topologist and he will for you : first check if that manifold admits a spinor structure (TOP. 6. B) or not, then if yes he will compute for you an invariant called the α -genus. Thereafter you, the geometer, will always be able to find a metric of positive scalar curvature in two cases : first, always if the manifold is not spin, second if its is spin, if and only is the α -genus is zero. This complete classification was achieved through successive efforts of both geometers and topologists. Among interesting examples which admit no metric of positive scalar curvature are some exotic spheres.

The main geometric tool is that of (Gromov & Lawson, 1980b). It consists in proving, using Riemannian submersions, that any topological surgery of codimension smaller than 3 can be done in order to obtain the positivity of the scalar curvature of the total manifold provided that both manifolds coming into the surgery have themselves already positive scalar curvature. We refer the reader to the survey (Rosenberg & Stolz, 1994). The first result, a necessary condition, goes back to (Lichnerowicz, 1963), see below for more. An other early partial result was (Lawson & Yau, 1974).

Another geometric tool was to use GMT (Geometric measure theory, see section O.F) to get stable minimal hypersurfaces : (Schoen & Yau, 1979). They proved that in a manifold with $\text{scal} \geq 0$ any codimension 1 cohomology class can be realized by a hypersurface which admits some metric with $\text{scal} \geq 0$. This metric is not the induced one but obtained after a conformal change. This change with finally positive scalar curvature is made possible if one uses the second variation formula for stable minimal hypersurfaces. But this works only in dimensions less than 8 , since it is needed in GMT to be sure to get submanifolds without any

singularity. See a nice exposition of the ideas in pages 91-95 of (Gromov, 1991b).

Comments on the advantages of both tools figure page 246 of (Rosenberg & Stolz, 1994) and section $5\frac{2}{3}$ of (Gromov, 1996).

According to Gromov a basic question is to find the geometric concept unifying both techniques, see more on this below.

The above results do not finish the story. **First** the non-simply connected case is well advanced but still not finished and in fact is connected in part with the famous Novikov conjecture, one of the driving force of recent topology, see (Gromov & Lawson, 1980c) . And also (Rosenberg & Stolz, 1994), (Gromov, 1996) and the references there for more on this.

Second the topologist description is not geometrical enough. Moreover, as we are going to see, the proof of the main result above is everything but geometry, except for the surgery control. A contrario the product of any manifold with the 2-sphere always carries a metric of positive scalar curvature (please just shrink enough the 2-sphere). Also minimal hypersurfaces of a positive scalar curved manifold carries a metric with the same property, at least "today" in low dimensions, this is the nice construction mentionned above. A geometric classification in that spirit is sketched in (Gromov, 1996).

So that an interesting question today is to look for a purely geometric study of manifolds with positive scalar curvature. There is in fact a pure geometric statement in (Llarull, 1996a) to the effect that a Riemannian metric g on S^d , such that $g \geq g_{can}$ (this means everywhere, i.e. all the lengths are larger or than equal), has necessarily one point m such that $scal(m) < d(d-1)$ (the value of the scalar curvature for g_{can} of S^d) unless g is the canonical metric. But today the proof is using all the "spinor game" (**TOP. 6. B**) of the proof sketched below. It would be great to have a geometric proof of Llarull's result, see also (Llarull, 1995).

Spinors do not really look metric objects but it is interesting to know that in (Connes, 1994) the Riemannian metric $d(.,.)$ is

computed with the Dirac operator. In (Connes, 1995-6) Riemannian geometry is put in a very general context.

Bishop's theorem on the growth of the volume of balls is false for scalar curvature, even locally. One has only an infinitesimal version of it, useless for global purposes. It would be basic, according to Gromov, to decide whether or not a C^0 -limit preserves the nonnegativeness of the scalar curvature, see **III. D**. Note that this is definitely false for the negative side by Lohkamp's result in **5** below. According again to Gromov, a starting point for proving this and understanding what is going on is the following : the nonnegativity of the sectional curvature is equivalent to the non divergence of geodesics as compared to the euclidean case. For scalar curvature one has, in a sense to be precised, a "non divergence result" but this only in at least one direction.

Third we have the **trichotomy** problem. We are interested to limit case of nonnegative (scalar) curvature. On the positive side, things are solved, on the negative side also. A long time ago in (Kazdan & Warner, 1975a) and (Kazdan & Warner, 1975b), it was proved that any manifold can carry a metric of negative scalar curvature. But between ? With a little more than Kazdan-Warner results (see (Besse, 1987), section 4. E for details), one knows that for the class of manifolds between there is only a third possible case : that of manifolds which cannot carry a metric of positive scalar curvature but can carry a metric of zero Ricci curvature. We will meet again this mostly open question in **III. C** and **TOP. 3**. The zero Ricci curvature class of Riemannian manifolds is still very mysterious today.

Fourth the total set of positive scalar curvature metric on a given compact manifold can be **non connected**. We saw above that this implies examples of non connectedness for both the total spaces of positive sectional and positive Ricci curvature. The first example was in (Hitchin, 1974) where the π_0 (the number of connected components) and the π_1 of the set of positive scalar metrics on a given manifold are shown to be non zero in many cases. In (Gromov & Lawson, 1983c) one shows that π_0 is infinite for the sphere S^7 . In (Kreck & Stolz, 1993) the subject goes on. They make use there of the Aloff-Wallach manifolds met before : we quoted this both in **2** and **3**. For more examples one can consult among others : the survey (Lohkamp, 1996) (for both signs by the way) and (Lawson & Michelsohn, 1989), page 329. Note that non connectedness is important but does not say much on the topological structure of

the components : does it have more or less trivial topology ? Strikingly different will be the case of negative Ricci curvature in 5 below.

At this point it is important both historically and for the future to look at the **proof** of the above results. An expository sketch can be found in pages 95-100 of (Gromov, 1991b). The tools are of three completely different types. The first is the above surgery, 100% geometrical. The second part is pure algebraic topology, namely cobordism theory. This theory will tell you which manifolds can be built up by surgery with simple building blocks (among them some of the $K P^n$). Precisely one has to show that any manifold satisfying the conditions above can be obtained by surgery with building blocks which do have positive scalar curvature. All this will insure the sufficiency of the condition.

The necessity is a completely different story and needs a new tool. It started with (Lichnerowicz, 1963). This was historically the first condition obtained from the positivity of the scalar curvature (and even in those times of the positivity of the sectional curvature !). Assume the manifold admits a spin structure (see **TOP. 6. B** and the general scheme in **O. D**). Then the generalization of Bochner technique is available through the theory of elliptic operator to yield **harmonic spinors**. The new "Bochner's formula" is surprisingly simple here and reads : $\delta^2 = D^*D + \frac{\text{scal}}{4}$. To the effect that $\text{scal} > 0$ implies the non-existence of any harmonic spinor. Then the Atiyah-Singer index theorem (see **TOP. 6**) implies that some invariant called the $\hat{A}$ -genus has to vanish. To go to the α -genus needed some extra work which was achieved in (Hitchin, 1974).

Last, not least, we cannot resist to quote the beautiful result : on a torus a metric of nonnegative scalar curvature has to be flat (Gromov & Lawson, 1983c). The technique of the proof is put in a general frame in (Lawson & Michelsohn, 1989), IV. §7.

4. THE NEGATIVE SIDE : SECTIONAL CURVATURE

This topic is an entire world in itself and is quite satisfactory for Riemannian geometers. In fact, since Hadamard in 1898 and 1901, results kept flowing on it. First by Cartan in the 20's, by E. Hopf in 39, Preissmann in the 40's and then the ones below.

The Cartan-Hadamard theorem of the 20's, clarified by Hopf-Rinow, implies that nonpositive sectional curvature manifolds have a universal covering which is diffeomorphic to $\mathbf{R}^d$. This follows from the stronger fact that any pair of points is joined by a unique geodesic which is automatically a segment. So the simply connected case "looks like hyperbolic or/and euclidean geometry". Then apparently for the naive observer all the problem consists in studying the fundamental group π_1 : algebraic nature, classification, differences between the negative and the nonpositive case, etc. But things turned out to be not so simple and in particular the characterization of possible groups is still far away. Also one can hope that the compact case looks close to space forms, see II. But the case of the negative pinching seen in A. 4 can rise suspicion.

In fact the subject enjoyed many strong results on one hand. But at the same time more and more difficult appears to get a real understanding of what is going on for the π_1 . In particular to remain in the compact case is definitively misleading so that we will make again an exception to our rule. This is one of the contributions of Gromov and a justification is this. We will see below many nice results on the π_1 of compact negatively curved manifolds. But as far as today there is no property known they satisfy and which is not satisfied by hyperbolic groups. This notion of hyperbolic group was introduced in the founding (Gromov, 1987a). A book exposition is (Ghys & de la Harpe, 1990). The force of this notion sits in its four equivalent definitions, ranging from group theory to Riemannian manifolds. Roughly speaking (and using the word metric for groups) those groups enjoy: the same asymptotic isoperimetric behavior and the same large triangles inequalities as in classical hyperbolic spaces. All this is a strong incentive to leave, at least for the moment, the compact (or even the finite volume case).

We will try now to ask questions and give results. Very informative is (Pansu, 1990/91) as well as the survey: (Eberlein, Hamenstädt, & Schroeder, 1990), but note that the authors admit not to be complete even in their 50 pages. In book form (Ballmann, Gromov, & Schroeder, 1985) was quite complete. The more recent book (Ballmann, 1995a) is a complete exposition of the "rank rigidity" for manifolds of nonpositive curvature and finite volume, see below. For various points of view on the topic see the book (Farrell, 1997). Let us just extend the above remark by

noticing that space forms appear in many instances in this survey. See II for completing our very partial and biased exposition. We follow now Gromov's philosophy and refer for much broader context to (Gromov, 1987a), (Gromov, 1993), see also the "beginner text" (Gromov, 1991b).

As remarked in the introduction of (Eberlein, Hamenstädt & Schroeder, 1993) the topic involves various techniques coming from different fields of mathematics. Here we note interferences with dynamical systems in V. B, space forms of II and harmonic maps of TOP. 7. In particular to simplify the exposition we have put as much as possible dynamical results in V and the rigidity results for space forms (of negative curvature) in II.

The simply connected story starts naïvely by asking to compare different metrics of negative curvature on $\mathbf{R}^d$. The good notion is that of **quasi-isometry**. Two metrics are said to be quasi-isometric if there exist two constants such that one metric is squeezed between the other multiplied by these two constants. It turns out that it is very difficult to decide when two given metrics (say of negative curvature) are quasi-isometric or are not. We will soon feel the difficulty. This notion itself arises simply from the compact context since any diffeomorphism between two compact Riemannian manifolds lifts up trivially to a quasi-isometry between their universal coverings. Now the universal covering of a compact negatively curved manifold satisfies $a \leq K \leq b < 0$. So we forget this now and start with our first question : classify up to quasi-isometries metrics on $\mathbf{R}^d$ such that $a \leq K \leq b < 0$ (this situation is a particular case of "bounded geometry"). The situation turns out to be extremely difficult as soon as $d \geq 3$. For $d=2$ just use conformal representation and compute the curvature. One gets a Laplacian for the conformity function and the bounds on curvature finally yields a bounded function. Hence when $d=2$ all our metrics are quasi-isometric. The only similar case is when both metrics under considerations admit a full orthogonal one point symmetry. This time the Jacobi field equation yields again a conformity function which is bounded.

But that is the end of the easy part. To realize it just try to prove that the first two space forms which come under consideration, namely Hyp^4 and Hyp^2 , are **not** quasi-isometric. This is hard to see, it is a corollary of Mostow rigidity theorem (see II. B). In that direction it seems very difficult to classify our spaces under our

condition when one has $-4 < K \leq -1$ or $-4 \leq K \leq -1$. For example it was already hard to get an example which can never be put in the $-4 \leq K \leq -1$ range (see (Mostow & Siu, 1980)). We meet here the difficulty to construct manifolds of negative curvature (see also A. 4). Even if one performs "after" some geometry they always come finally from "Number Theory". A contrario, as remarked in TOP. 9, singular objects of negative curvature represent in some sense most of the natural geometric objects (glue hepta or more -gons and mutatis mutandis for higher dimensions). One now has a quite large family (depending on real parameters) of two by two non quasi-isometric manifolds, this is in essence in (Pansu, 1989). Moreover those metrics are homogeneous.

We stay in the difficulty at the $\mathbf{R}^d$ level with some nonpositive metric but look at it from the other side : can any discrete finitely generated group act by isometries with non bounded orbits ? There is today no restriction known besides the trivial one obtained from the fact there is a quotient which is a $K(\pi;1)$: our group should have some non-zero homology in some dimension. No conjecture seems in view, see (Gromov, 1993). The above considerations play a role in Novikov's conjecture, which is still pervading geometry and topology.

Oppositely one has a good understanding of how act special groups. In (Kleiner & Leeb, 1997) it is shown that if the group of an irreducible symmetric group of rank ≥ 2 acts by quasi-isometries it has to be by isometries. This is false for the reducible case. See also (Schwartz, 1995).

Tools are what the reader can ask about. First we have the global triangle comparison with hyperbolic geometry. This is a trivial extension of Cartan-Peissmann super-euclidean inequality seen in 0. B. This inequality is not enough, it has to be refined. This is done by introducing the notion of convexity for functions in a Riemannian manifold. Negative curvature implies strict convexity for many functions, e.g. Busemann ones and the distance between sides of a quadrilateral

Basically the smarter tools all consist in compactify the situation, since working with the infinite is hard. We are still on $\mathbf{R}^d$ with a nonpositive metric, most often with a lower bound for K . We extract few things from the basic reference (Ballmann, Gromov & Schroeder, 1985) for detailed proofs and from the survey

(Eberlein, Hamenstädt & Schroeder, 1990) . One defines the sphere at infinity $S(\infty)$ of this situation. This was already with his language in (Hadamard, 1898) when $d=2$. The first thing one gets on $S(\infty)$ is a topology, the cone-topology defined in an obvious way through rays. It does not depend on the origin chosen and if one varies only by quasi-isometries one still gets homeomorphisms.

The next is to have measure(s) or even metric(s) on $S(\infty)$. The metric called "Tits metric" by Gromov is very special. It is always nowhere defined (as equal to $+\infty$) when the curvature is negative (more precisely $K \leq b < 0$) and this is linked with the visibility condition introduced in (Eberlein & O'Neill, 1973). The visibility between two points at infinity is that there exists a geodesic joining them, which is definitively not the case for example for two different directions in euclidean spaces. More subtle are some other families of metrics one can define on $S(\infty)$. They have the properties to undergo quasi-conformal properties under quasi-isometrical changes. When those objects depend on the base point they still are changed in a reasonable way.

When the curvature is only nonpositive then Tits metric become finite for some pairs of points. This finite parts are associated to flat parts which we are going to meet below. The totality is some kind of a jig-saw (railway) type metric.

We turn now to properties of compact manifolds of negative (resp. nonpositive) curvature and of their fundamental group . First to come are **volumes**. As it will be the case for space forms in II volumes are discrete and in particular isolated from 0 for space forms of any type. Basic is (Heintze, 1976) : when $-1 \leq K < 0$ then the volume is lower bounded by $c(d) > 0$ depending only on the dimension. This is often called Heintze-Margulis lemma, because it was obtained in Margulis in the special case of space forms. In both authors the proof is purely geometric, using a "collar" argument. The strong divergence of geodesics in negative curvature as well as the nature of hyperbolic isometries are forcing the tube around the smallest periodic geodesic to have not too small a diameter and then one is done with the comparison theorem as for the tube argument in I. C. 1 . So that one can have some hope here in the negative case to have finiteness results with milder conditions than the four ones needed in the general Cheeger finiteness theorem in I. C. 2 . This was achieved in (Gromov, 1978b), is detailed in (Ballmann, Hamenstädt & Schroeder, 1985) and improved in

(Fukaya, 1984). We mention here a finiteness theorem of Gromov which is true for **real analytic** manifold but false in the differentiable case. It is extremely rare in Riemannian geometry to have such a kind of situation. The result is that if $-1 \leq K < 0$ for (M^d, g) and if the metric g analytic then the sum of its Betti numbers is bounded by $c(d) \cdot \text{Volume}(g)$, where $c(d)$ is universal in d . This is optimal, as shown by examples : they are obtained by connected sums, a typical smooth but non analytic operation. See (Ballmann, Gromov & Schroeder, 1985) for details of the proof.

Compact space forms of negative curvature were already hard to construct, see II. For variable curvature we saw examples in A. 4 but real analytic are harder to construct, see (Abresch & Schroeder, 1996a).è

Next are **fundamental groups**. The first statement is in (Milnor, 1968) : for compact manifolds negative curvature implies that the fundamental group has exponential growth (see also (Svarc, 1955)). In (Gromov, 1978b) (see (Buser & Karcher, 1981) for details) a bound is obtained for the number of generators of this group. Milnor's tools are just **RCT** and Bishop's theorem (see **TOP. 1. A**). Gromov uses the triangle comparison theorem and then one has only to count points on the unit sphere whose mutual distances are bounded from below as in 1 above.

The most general statement is (Farrell & Jones, 1989b), which is a topological version of Mostow's rigidity theorem (see II) : let M be a hyperbolic (compact) manifold, i.e. a space form and of dimension larger than 5 and let N be any topological manifold which is homotopically equivalent to M . Then M and N are homeomorphic. In a book form see (Farrell, 1997).

Next is the product situation : if the manifold is compact with nonpositive curvature and with a fundamental group which is algebraically a group product (with moreover no center), then the manifold itself is a Riemannian product (Gromoll & Wolf, 1971). The result of Preissmann seen in **O. C** was mastered in (Lawson & Yau, 1972) : in compact manifold with $K \leq 0$, commuting in π_1 means flat parts, namely any free abelian subgroup of π_1 implies the existence of flat totally geodesic tori in the manifold. This can be seen as a converse of the Bieberbach theorem for the classification of flat space forms : II.

For negative curvature we saw that, at least today, hyperbolic groups are the "final" answer". The big problem now, in the compact manifold downstairs as well as for the fundamental groups, is to appreciate the difference between negative and nonpositive curvature. There are today extremely strong results which show in essence that the two cases can be dramatically separated, otherwise said : to go from one to the other you have to make quite of a jump. We give the latest results and refer to (Eberlein, Hamenstädt & Schroeder, 1993) for a survey and to (Ballmann, 1995a) for the proofs.

We recall briefly that symmetric spaces (see II. C) have a **rank** which is the common dimension of their totally geodesics flat submanifolds, which are moreover all conjugate by the isometry group. This is of course rich only when the rank is ≥ 2 otherwise we just have geodesics. In particular every direction is contained in at least one such flat. At the infinitesimal Jacobi field level, rank ≥ 2 implies this : given any unit vector, along the geodesic it generates there is a non trivial **parallel** Jacobi field (orthogonal to the geodesic of course and non zero). This never happens for generic manifolds. In a Riemannian manifold we define the **rank** of a unit vector to be the dimension of the space of parallel Jacobi fields along the geodesic it generates. The **rank** of a Riemannian manifold is the minimum of the rank of its unit vectors. The strongest result today is the following "tout ou rien" result for compact Riemannian manifolds of nonpositive curvature. We assume any reasonable irreducibility condition. The theorem (for a manifold of nonpositive curvature) is this : "either the rank of the space is ≥ 2 and then we are on a space form with its symmetric space metric. Either the rank is 1 and then the geodesic flow is ergodic on the subset of UM made up by the vectors of rank 1 ". Today it is still open to know if this subset of rank 1 vectors is of full measure or not.

Note that the rank, a geometric invariant, can be at least conceptually computed from the fundamental group : (Eberlein, Hamenstädt & Schroeder, 1993). The existence of flats and what they really imply is a major issue today, see for example (Hummel & Schroeder, 1997).

The above dichotomy admits a lot of extensions with weaker hypothesis, like : finite volume to replace compactness, or/and a group restriction so-called " duality condition" and also to metric spaces mimicking Riemannian manifolds of nonpositive curvature.

For all this the reader can look at the given references and **TOP. 9**

Finally we end with the question to relax just a little but from above the condition $K \leq 0$. It is important to realize that without extra hypothesis "everything is permitted". A construction of Gromov was published in (Buser & Gromoll, 1988), (Bavard, 1987b). There for any $\varepsilon > 0$ are built examples of metrics on spheres and other 3-dimensional manifolds whose curvature satisfies $K \leq \varepsilon$ and are of bounded diameters. See also (Fukaya & Yamaguchi, 1991) : the results here are that one can obtain restrictions on the fundamental group, namely that the universal covering is diffeomorphic to $\mathbb{R}^d$, but with an extra hypothesis, like $K > -1$ and an upper bound on the diameter.

5. THE NEGATIVE SIDE : RICCI CURVATURE

We have here the strongest possible answer : (Lohkamp, 1994) belongs to a series of papers where it is shown that , caricaturally speaking, negative Ricci curvature "means nothing" or, equivalently, "permits everything". Worst : **any** metric can be continuously approximated by a metric of negative Ricci curvature (of course not with too much differentiability). Tools are from Analysis, it has to do with the h-principle of (Gromov, 1986) and also looking at the formula giving the variation of Ricci curvature under deformation of metrics where one can see that it is quite easy to lower Ricci curvature.

Another very important point (compare e.g. with section **B. 2**) is the fact that, on any manifold, the total set of negatively Ricci-curved metrics is trivial. That is much more than only connected, namely contractible . We suggest the reader to consult the very informative (Lohkamp, 1996).

A final note . Two of the three favourite questions that Hopf was asking since the 30's are still completely open : does a given sign for the curvature implies a given sign for the Poincaré characteristic ? Does $S^2 \times S^2$ admit or not a metric with positive curvature? In **O. E** we saw that Hopf's conjecture is true for $d=2$ and 4. For $d \geq 6$ it is completely open. Both questions are leaving today experts with no solid guess. The latest news for $S^2 \times S^2$ are in (Kuranishi, 1990). Even more itching is Yau's question mentioned

in 1. This question is most irritating, since Synge's theorem (see O. C) trivially excludes $\mathbf{RP}^2 \times \mathbf{RP}^2$ (as well as many other products of manifolds). Rong's results quoted in 1 exclude even a lot more. A recent general list of problems is in (Petersen, 1996e).

C. Finiteness, compactness, collapsing and the space of all Riemannian structures

The logical mathematical order to present the results below in 1 and 2 is the inverse of the one we are going to follow. This is the way things are presented in the surveys (Abresch, 1990b), (Fukaya, 1990), in a book form in the Appendix 6 of (Sakai, 1996), in (Petersen, 1997b), in the second edition (Gromov, 1997) to appear of the pioneer book (Gromov, Lafontaine & Pansu, 1981a). Historically the order was reversed and is the one we are going to follow.

1. FINITENESS RESULTS

This topic started simultaneously in 1966 with (Cheeger, 1967) and (Weinstein, 1967). For expository reasons we present first the simplest case, which is the only one treated by Weinstein. In despair to classify positive sectional curvature manifolds, Weinstein proved that there are only a finite number $N(\delta)$ of possible different homotopy types of even-dimensional manifolds with $0 < \delta \leq K \leq 1$. Unhappily $N(\delta)$ goes to infinity (and exponentially) when δ goes to 0. Note that oppositely Gromov's theorem quoted above on Betti numbers of nonnegatively curved manifolds implies a bound but only for (integral) **homology** types. The proof of the homotopy finiteness was purely geometrical, covering the manifold with convex balls whose number can be estimated only in function of δ with Bishop's bound for the volume (needing only Ricci curvature and hence is valid a fortiori for sectional one, see TOP. 1. A). This is done by the standard, but still wonderful, purely metric trick : packing as many as possible balls of radius r implies that the corresponding balls of radius $2r$ are covering. Such packings are called **efficient**. They have also the extra advantage that one can universally control the number of balls which are meeting a given one. Then the homotopy type is that of a simplicial complex with as many vertices as the above balls. The evenness of the dimension is needed because here the convexity radius is

bounded from below by $\pi/2$ thanks to Kingenberg's theorem seen in A. 2.

We will after look at the two next levels : **homeomorphy** types then **diffeomorphy** types. Note also that in any such finiteness result one can also ask for explicit bounds for the cardinal of those finite sets. We will see that in some cases there is no such explicit bound because the proof is obtained by a contradiction argument based on convergence of some infinite set.

We very briefly recall here facts about the three ratios **homotopy/homology**, **homeomorphy/homotopy**, **diffeomorphy/homeomorphy**. The two first ratios are in general infinite. For the first we just mention that even the classification of homology spheres is not finished. For the second the theory of characteristic classes is an ideal tool, a lucid exposition is in (Milnor & Stasheff, 1974). The third ratio is always finite as soon as $d \geq 5$, basic texts are (Hirsch & Mazur, 1974), (Kirby & Siebenmann, 1977). These finite cardinals can be computed explicitly using various topological information on the manifold under consideration. Dimension 4 is an exception, counter-examples can even be taken among algebraic surfaces.

It is basic to realize that one cannot extend the above result to odd dimensions, for example we saw that there is an infinite set of all Alon-Wallach manifolds $W_{p,q}$ (see B. 1) made up of all homeomorphic together but not diffeomorphic two by two, see the theorem 3.9 of (Kreck & Stolz, 1993). But one can still get not only homotopy but even diffeomorphy finiteness, but under stronger hypothesis. The pioneer work of (Cheeger, 1967; Cheeger, 1970a), independent of Weinstein's one, yielded this : the set of simply connected possible types of diffeomorphism is finite within the set of Riemannian manifolds of a given dimension $d > 4$ satisfying the four conditions : $K \geq a$, $K \leq b$, $\text{volume} \geq v > 0$, $\text{diameter} \leq D$ (where a , b , v , D are four fixed real numbers such that v has to be positive the time being a and b can be of any sign). So that one has finiteness not only for homotopy type but also for homeomorphism and, from there, to diffeomorphism and this, without any assumption on the sign of the curvature

One remark is now in order : in some instances one can effectively get rid of the condition $\text{volume} > v > 0$. For example when one works within the class of manifolds with a non-zero characteristic number : as remarked in (Cheeger, 1970a) this comes immediately

from Chern's formulas seen in O. E. Another instance is that of negatively curved manifolds, see section B. 4 above.

The way the proof goes is important for the future. The first thing we need is to be able to cover the manifold by an efficient packing of convex coordinate balls, the cardinal of the packing being is universally bounded (universal means of course in a , b , v , D and course the dimension d). This is of course done with efficient packings. The estimation of the number of balls is done thanks to Bishop volume estimate with Ricci curvature (which comes here from the sectional one), as soon as one can get a lower positive bound for the injectivity radius. This is the first part of Cheeger's work : the injectivity radius has a universal explicit (positive) lower bound function of only a , b , v , D , d . The original proof was a nice geometric argument (called the "butterfly") using RCT and Toponogov. Today the best proof is that of (Heintze & Karcher, 1978) : there is an estimate of the volume of the tube of given radius around a periodic geodesic, using a refinement of RCT left side estimate. These four bounds are necessary to control the injectivity radius. The four needed counter-examples are : flat rectangles of sides $(\epsilon, \epsilon^{-1})$ or $(\epsilon, 1)$, very flat ellipsoids of revolution and very much strangled spheres.

The second part goes by contradiction and shows that the number of homeomorphism types is finite (and even if desired explicit bounds for the numbers could be obtained). Finally one will have to control the coordinate changes between Riemannian coordinate balls for two different metrics. This is exactly what RCT can do for you when you know that $K \geq a$ and $K \leq b$ but note for the sequel that this is only a C^0 -estimate. To go from homeomorphism to diffeomorphism types, Cheeger used the results from algebraic-differential topology available mentioned above : the number of possible different diffeomorphism types on a given topological and simply connected manifold is finite when $d > 4$. However later on Cheeger managed to prove diffeo-finiteness for any dimension (under these four bounds) without appeal to differential topology. References are (Cheeger & Ebin, 1975) (theorem 7. 37) to be completed by (Peters, 1984).

A digression : does the curvature determines the metric ? We insist on the fact that the curvature tensor does not determine the metric (except in very special cases) : this can shock the naive reader since

the number of parameters is so large, as soon as $d \geq 4$. Be careful also that Cartan's result in **O. B** needs to know the curvature but also the geodesics and well as the parallel transport along them. The subject started quite lately : they are many examples of non isometric Riemannian manifolds admitting diffeomorphisms preserving their respective curvature tensors. A trivial case is to move the metric of a surface along the level lines of Gauss curvature K . In higher dimension things are of course more subtle, since the curvature has not only many, many parameters but is a very weird algebraic object. For surfaces it was just a numerical function. The complete mastery of this is not achieved today, see below and the bibliographies of the references given

A contrario there is (Kulkarni, 1970) to the effect that, when the sectional curvature is not constant and the dimension larger than 3, diffeomorphisms preserving the **sectional** curvature are isometries. This is not in contradiction with the above examples, because the definition of the sectional curvature involves not only the curvature tensor but also the metric. For lower dimensions see (Ruh, 1985).

Another meaning can be attributed to "does the curvature determines the metric?" . It consists in looking for metrics whose curvature tensor satisfies at every point some imposed purely algebraic condition. A typical example is : the curvature tensor is the same as that of a complex projective space. One would expect then in some cases isometry (local) and in general at least local homogeneity, as well as a description of the moduli. Since the founding (Ambrose & Singer, 1958) and (Singer, 1960) the field today uses various definitions and enjoy many results. The books (Tricerri & Vanhecke, 1983), (Berndt, Tricerri, & Vanhecke, 1995), (Boeckx, Kowalski, & Vanhecke, 1996) can be used as surveys. See also (Prüfer, Tricerri, & Vanhecke, 1996). The result in (Tricerri & Vanhecke, 1986) is exemplary of simplicity : if the curvature tensor is at every point the same as the one of an irreducible symmetric space, then the manifold is locally symmetric and then isometric to this model. The result is local, it uses a formula in (Lichnerowicz, 1950) which expresses the Laplacian of the full square norm of the curvature tensor as $-(1/2) \Delta(\|R\|^2) = \|DR\|^2 + \text{Univ}(R, R, R)$, where $\text{Univ}(R, R, R)$ is an universal cubic form in the curvature tensor itself. Then the hypothesis implies immediately that $\|R\|^2$ is constant and also $\text{Univ}(R, R, R)$, because it is obvious in the symmetric case which is characterized by $DR = 0$. It seems to us that this basic formula is still not used as much as one would have expected.

For the curvature with the parallel transport see **O. B** and **TOP. 4** (Ambrose's problem).

Many people think that the curvature and its derivatives are the only Riemannian invariants. This is true and classical when looking at the strong condition : algebraic invariants which stem from the connection, see page 165 of (Schouten, 1954) and the references there. But things are dramatically different if one asks only for tensors which are invariant under isometries (called "natural"). Then there is even no hope to get any kind of classification as explained in (Epstein, 1975). For more see (Munoz & Valdés, 1996).

Back to finiteness there are two ways to improve Cheeger's result. The problem is to control coordinate changes more than C^0 with only the sectional curvature (without the metric) . One way is to use **compactness** theorems, see section 2 below. The other is to use the **center of mass** technique. The center of mass in Riemannian geometry can be obtained in convex balls by two equivalent definitions (hence its usefulness). The first is to minimize the sum of the square of the distances, the second is to ask for linear dependence between the speed vectors from the center of mass to the points into consideration. The first being purely metric and the second involving tangent vectors one gets automatically some C^1 behavior from a metric one via **RCT** . For the center of mass and some applications one can see : (Karcher, 1989), (Buser & Karcher, 1981) chapter 8, (Cheeger, Müller, & Schrader, 1984). But it should be remarked that the existence of a unique center of mass and in the large for nonpositive curvature was already proved and used by Elie Cartan back in the 20's (see **O. B**). And also that for the general case (but locally only of course, think of the standard sphere) by Calabi in his unpublished result quoted in **A. 2** to solve the differentiable pinching problem. The center of mass technique was the one used in (Peters, 1984) : one put suitably chosen $d+1$ points in very small balls.

Now back to the favourite game since the 80's : ask for results with weaker and weaker hypothesis. The first kind of results is in (Gromov, 1981d), where it is remarked that the technique used to bound Betti numbers for nonnegatively curved manifolds (see **B. 1**) extends without any modification to manifolds with $K \geq k$ (for any real k) and diameter $\leq D$. More precisely the Betti numbers (over

any field) of Riemannian manifolds of a given dimension are exponentially bounded with the real number $(\text{diameter})^2 \cdot (-\inf K)$ as exponent. Then we have finiteness for homology types (in the simple connected case of course) with only bounds on $\inf K$ and diameter. There is an improvement of this bound in (Abresch, 1985).

Very new was (Grove & Petersen, 1988). A priori it seems impossible in Cheeger type of results to get rid of some injectivity radius estimate in order to get contractible, or more, convex, balls. But remember Gromov's proof for the Betti number of nonnegatively curved manifolds (see B. 1) : one of the basic facts is that critical points for the distance function cannot "appear too far from the center". In (Grove & Petersen, 1988), using the center of mass technique and Toponogov's theorem, it is proved that the class of Riemannian manifolds of given dimension satisfying $K \geq k$, volume $\geq v > 0$ and diameter $\leq D$ has only finite possible **homotopy** types. The cardinals were bounded explicitly in d, D, k, v . Cheeger's result on the injectivity radius is replaced here by a crucial lemma using Toponogov and the fact already mentioned that unit vectors at some points cannot be too many when their mutual angles are not too small (see B. 4). Easy examples show that no one of these three bounds can be removed from the statement.

In (Grove, Petersen, & Wu, 1990-1991) under the same hypothesis the result was extended to finiteness for **diffeomorphism** types. Dimension restrictions are : when $d=3$ due to the status of the Poincaré's conjecture and when $d=4$ when one has to be happy (today at least) only with homeomorphisms. We remark here that it is also open today if the counter-examples (mentioned above) yielding infinitely many differentiable structures on some topological manifolds of dimension 4 can or not satisfy these bounds. It seems also that the cardinals cannot be universally bounded today with the kind of proof used. But if one has bounds at the homeomorphy level then would have them also for the diffeo-level. The proof definitively involves new ideas after 1988 result. It uses fundamentally convergence techniques from the section 2 below and a lot of algebraic topology for manifolds, in particular the technique called "controlled topology".

In the same article there is the astonishing result : finiteness of homeomorphism types (hence diffeomorphism if in addition $d \neq 4$) for the class of manifolds of a given dimension $d \neq 3$ under only **two** bounds : volume $\leq V$ and injectivity radius $\geq i > 0$. We will

comment on this in **TOP. 1. C** in the **embolic story**. Before homotopy finiteness was obtained under the same conditions in (Yamaguchi, 1988). This Grove, Petersen and Wu result used essentially the basic Croke's local embolic result mentioned in **TOP. 1. E** and the ingredients for the proof of the above result.

Back to Grove-Petersen-Wu finiteness result, (Greene, 1994) and (Greene & Petersen, 1992) throw interesting light to it, in particular expliciting the role of the volume of tubes around (periodic) geodesics.

Again there are results obtained by replacing pointwise bounds on the curvatures by various integral bounds. Precise statements are always quite elaborated, hence not stated here, see (Yang, 1992), (Petersen & Wei, 1996b), (Petersen, Wei, & Ye, 1997a) and in a book form there is (Petersen, 1997b). One geometrical point was to control the volume of a tube around a periodic geodesic, this is done in (Petersen, Shteingold, & Wei, 1996a). The proofs are not only geometrical but involve a lot of Analysis.

Now how about finiteness with only bounds on **Ricci** curvature ? We saw above that already Gromov's homology finiteness for $K \geq k$ and $\text{diameter} \leq D$ does not extend to $\text{Ricci} \geq r$. However this did not stop geometers to see what $\inf \text{Ricci}$ means, since Ricci curvature is hopefully a "reasonable" curvature control. Results started with (Abresch & Gromoll, 1990a) and are now very strong. We make a partial choice among them and refer to the very informative survey (Anderson, 1994) and the references there. Note that some of them are optimal with respect to the ingredients in view of the examples in (Perelman, 1997).

First there is finiteness on diffeomorphisms types in a given dimension under : $\text{Ricci} \geq \lambda$ (of any sign), $\text{vol} < V$ and $\text{inj} \geq i > 0$: (Anderson, 1990c). Since one succeeded for sectional curvature to go beyond the injectivity radius, the urge is here too. The story is not finished but we know this from (Anderson, 1990c) together with (Anderson & Cheeger, 1991a) : there are in a given **odd** dimension only finite diffeomorphism types under the conditions :

$|\text{Ricci}| \leq \lambda$, $\text{vol} \geq v > 0$, $\text{diam} \leq D$ and $\int_M \|R\|^{d/2} \leq \Lambda$. When the dimension is even it is compulsory (there are examples) to add

orbifolds (see **TOP. 9**) to manifolds. Moreover in dimension 4 the $L^{d/2}$ -bound on the full curvature tensor is not needed thanks to the Allendoerfer-Weil formula (in **O. E**). It is not clear if it is reasonable to conjecture that this extra condition is not really needed. Note finally that all these results being obtained by contradiction hence they do not yield today explicit bounds for the cardinals.

The idea for the above results is to prove $C^{1,\alpha}$ -compactness. Basic is the fact (see the digression in the **1. COMMENTS on I**) that Ricci curvature is in harmonic coordinates, up to first order terms, equal to the Laplacian of the metric. But one has to control the **harmonic radius** (i.e. the largest balls on which harmonic coordinates are well defined and linearly independent, see next

section for more on it). When using the bound on $\int_M \|R\|^{d/2}$ the

difficulty is to understand how the curvature concentrates, this is achieved by looking for singular points in sequences converging in the Gromov-Hausdorff space to an orbifold by (Anderson, 1990c). Then one rescales the metrics around the singular points and shows that iteration of this process ends after a finite number of times.

For the fundamental group one has in (Anderson, 1990c) finiteness of group structures with a lower positive volume, an upper diameter and a lower Ricci bounds.

From **B. 3** it is clear that no finiteness statement is to be expected from bounds on scalar curvature.

2. COMPACTNESS, CONVERGENCE RESULTS

Climbing Jacob's ladder, even in the fog, comes the natural question to look at any kind of convergence and/or compactness within the set of Riemannian manifolds or/and of various subsets of it. Moreover we met before many problems where a convergence existence theorem would have been useful.

Look for example at finiteness diffeomorphism theorems or even more simply to the differentiable pinching problem. It is intuitive that - of course in a sense to be precised - the set of differentiable structures is *discrete*. Then within a compact subset we have finiteness. Another example is the isolation of the standard sphere. Note that would be a very pleasant proof but "in the air", in particular not yielding any explicit pinching constant.

Another motivation is that of **III** : look at some functional on the set (or some subsets) of Riemannian metrics on a given manifold M . Does there exist some Riemannian manifold realizing the infimum of that functional (a best, an extremal Riemannian structure on M)? Sadly enough all the convergence results we are going to see are not able to give an answer in that domain, see e.g. the minimal volume and the embolic volume in **TOP. 1 . C** and **D** as well as Einstein manifolds in **III. C**.

It seems that the first appearance in that direction was in (Shikata, 1967), where an isolation result was obtained for differentiable structures. Thereafter considerations of that kind were implicit in (Cheeger, 1970a). Then appeared the theorem (no number on it) on page 74 of (Gromov, 1983b). The proof uses implicitly a convergence theorem that the author always took for granted. According to him since some people doubted it or at least asked for more details he published a proof in (Gromov, Lafontaine & Pansu, 1981a). The proof was still a little incomplete, then (Katsuda, 1985) offered a complete proof. In 1985 various proofs of the optimal result (see below) started circulating, all using centers of mass and harmonic coordinates. Printed references are (Greene & Wu, 1988), (Peters, 1987) and (Kasue, 1989). We now have to make things precise since the notion of convergence needs a topology.

We now present the notions and the related results in the order which is the good one now, moreover it is well suited to be generalized to more general geometric objects than Riemannian manifolds. This is in fact unavoidable for understanding what is going on and moreover simplifies considerably the proofs quoted above. We follow mainly (Fukaya, 1990) and refer to it for precise definitions. Surveys are (Petersen, 1990) and (Petersen, 1997c). And the second edition to come of (Gromov, Lafontaine & Pansu, 1981a) : (Gromov, 1997).

Things start with the **Gromov-Hausdorff distance** $d_{G-H}(X, Y)$ between isometry classes of abstract compact metric spaces. Its definition and properties appeared first in (Gromov, Lafontaine & Pansu, 1981a). The set of all compact metric spaces, denoted by **MET**, endowed with d_{G-H} , is Hausdorff and complete, this extends the well-known analogous statement for compact subsets of $\mathbf{R}^d$.

Already for MET we have the Riemannian geometry result from (Gromov, Lafontaine & Pansu, 1981a) : inside (MET, d_{G-H}) the subset of Riemannian manifolds of a given dimension d , with diameter $\leq D$ and $\text{Ricci} \geq -(d-1)$ is precompact. The argument is just an "efficient packing". The cardinal can be bounded from above precisely with the improvements due to Gromov to Bishop's theorem and explained in **TOP. 1 . A** . But of course the closure of such sets is not in the realm of (smooth) manifolds of dimension d , just think for example of flat tori.

The next topology to come in our story is the Lipschitz's one. In MET we call it d_L . Its value $d_L(X,Y)$ is the best Lipschitz constant for the Lipschitz homeomorphisms $f : X \rightarrow Y$ together with the constant of f^{-1} . Back in (Shikata, 1967), the following is proved , which we called a discreteness result in the motivations above. We work within the set $\text{RM}(d,a,b,D,v)$ of Riemannian manifolds of given dimension d , with $a \leq K \leq b$, diameter $\leq D$ and volume $> v > 0$. Then if N is the d_L -limit of a sequence $\{M_i\}$ then for i large enough M_i is diffeomorphic to N . This is how Shikata solved the differentiable pinching problem.

But this is still not a convergence statement for **any** sequence (or some subsequence) in the set $\text{RM}(d,a,b,D,v)$. This is achieved in two steps. The first is in (Gromov, Lafontaine & Pansu, 1981a), to be completed with (Katsuda, 1985). It says that if $\lim_{i \rightarrow \infty} d_{G-H}(N, M_i) = 0$ then $\lim_{i \rightarrow \infty} d_L(N, M_i) = 0$. At this stage we do not know if the limit metric is a Riemannian metric. This needs more work and is in Gromov, Greene-Wu and Peters references mentioned above. We saw above in the digression in 1 that the difficulty lies in the fact that the control on curvature gives a **metric** control but not a differentiable one in the metric itself. This is because **RCT** shows only a C^0 behavior. The basic trick is to replace normal (geodesic) coordinates by apparently less geometric ones but in fact more adapted finally. Those are the **harmonic coordinates** (see **O. F** and **1. COMMENTS on I**) : one chooses d harmonic functions which are linearly independent and satisfy ad hoc boundary conditions. It works only within the **harmonic radius** which was first systematically introduced and controlled in (Anderson, 1990c). Such a control is basic in many results and we refer for this to the various references which we will meet. With sectional curvature bounds it is more easy than with Ricci ones.

The exact statement is the convergence of some subsequence in any infinite sequence $\{M_i\}$ in $\text{RM}(d,a,b,D,v)$ toward a Riemannian manifold (N,g) which is diffeomorphic to M_i for i large enough

but the Riemann tensor g is known in general to be only of class $C^{1,\alpha}$ (for any α in $(0,1)$). The proof uses coverings with suitable balls whose number is controlled as explained in 1 above and using harmonic coordinates on them. One can show that this is optimal, one cannot in general hope C^2 .

If one wants to get smoother limits one uses **smoothing techniques**, which are important in other instances and were obtained first, using the Ricci flow (see 1. COMMENTS on III), in (Bemelmans, Min-Oo, & Ruh, 1984). Thereafter in (Abresch, 1988) and in (Shi, 1989). See other statements in (Fukaya, 1990). Smoothness means that the absolute value of the covariant derivatives of the curvature tensor of any order, specially the first one which is technically very useful, can be made as small as wanted. One of the main interest of smoothing is that it respects most hypothesis and that one can for example add smoothness in $RM(d,a,b,D,v)$, etc. See also the norms in (Petersen, Wei & Ye, 1997a).

As in instances met above one is tempted to obtain results of compactness (convergence) with less hypothesis, in particular with **only Ricci curvature control**. Convergence results where the curvature control is mainly $\text{Ricci} \geq -(d-1)$ started in (Gao, 1990a), and where followed by (Anderson, 1990c). The best result today is (Anderson & Cheeger, 1992a) : one has precompactness in the $C^{0,\alpha}$ topology (for any α in $(0,1)$) under the conditions : $\text{Ricci} \geq r$, injectivity radius $\geq i > 0$ and volume $\leq V$. The link between finiteness and compactness is explained in (Anderson & Cheeger, 1992a) and is of quite general nature. If moreover the k -covariant derivatives of Ricci are bounded in absolute value by suitable constants one has precompactness in the $C^{k+1,\alpha}$ topology. See a extremely brief exposition in (Hebey & Herzlich, 1995b). Note that these extra conditions can be achieved by smoothing as seen above. The proof uses essentially (with a lot of Analysis, e.g. Sobolev inequalities) harmonic coordinates, study of the harmonic radius which can be bounded with the metric injectivity radius (this is not too surprising) together with Ricci bound : (Anderson, 1990c). See also (Cheeger, Colding, & Tian, 1997b).

For applications of convergence theorems we refer the reader to the various surveys above. We just note that in most cases, like pinching theorems or "just below", the convergence theorem yields an ε "in the air". This still gives job to people to get "direct" proofs and explicit constants.

A final remark is in order. Most of the above results work for (complete) non compact manifolds provided one works in the **pointed** category. These results are essential for example when one studies the fundamental group as in B .

3. THE SET OF ALL RIEMANNIAN METRICS : COLLAPSING

The desire here is very natural : if a sequence of Riemannian manifolds does not converge nicely (as we have seen this is mainly because the injectivity radius goes to zero) then what is really happening ? Do we have some limit space of some kind ? E.g. manifolds but of smaller dimension, or a reasonable generalization of Riemannian manifolds (see **TOP. 9**), or just some metric space ? In some sense one is looking for a **compactification** of the set of Riemannian metrics (of a given dimension) and an understanding of what goes on when going to the boundary. This of course does make sense only within suitable subsets, namely when one imposes some curvature bound(s) and some metric invariant(s). When there is no convergence towards a Riemannian manifold (of same dimension even permitting singularities) the situation is called **collapsing**. This is vague, we now study things more precisely. Again the reference we follow here is (Fukaya, 1990) , at least up to 1990. An informative text is (Pansu, 1983-1984). The core is built on : (Cheeger & Gromov, 1986-1990), (Fukaya, 1987a), (Fukaya, 1989) and by the whole team : (Cheeger, Fukaya, & Gromov, 1992), whose introduction is very informative.

The first situation we study is when one drops in $RM(d,a,b,D,v)$ the last condition $\text{volume} \geq v > 0$ and work in $RM(d,-1,1,D)$ (obvious notation) after the normalization from (a,b) to $(-1,1)$ which does not hurt when looking only for general statements. Then we have to study the situation where the volume goes to zero, which by Croke's local embolic theorem in **TOP. 1. C** implies that the injectivity radius goes to zero uniformly. The first thing to do is to have examples. We saw in the pinching around zero (see **A. 3**) that nilmanifolds are in our class. Note then that the **limit space** is the smallest possible one, namely reduced to a **point** .

The other basic general example appeared in (Gromov, 1983b) : any manifold admitting a circle action without fixed point, or say fibered by S^1 over some other manifold N , collapses to N . This is

just an application of Riemannian submersions formulas : take first a fixed Riemannian metric invariant by S^1 and make the fibres smaller and smaller keeping the "horizontal" metric component fixed. One get the nilmanifolds as a particular case applying this trick to the successive S^1 fibrations which comes from the nilpotent structure. From the Hopf fibration $S^1 \rightarrow S^{2n+1} \rightarrow \mathbb{C}P^n$ one gets a collapsing of the sphere S^{2n+1} to $\mathbb{C}P^n$. A fortiori of course this extends to manifolds admitting a torus simple action.

A more elaborated type of example is as follows. Take two 2-dimensional manifolds N and N' of same dimension with a boundary which is a circle and take the products $M = S^1 \times N$ and $M' = S^1 \times N'$. They are manifolds with boundary, in both case the boundary has the topological type of a torus T^2 . Then glue M and M' along that T^2 , not in the trivial way but **interchange** the two circles in T^2 (exchange the parallels and the meridians). The resulting manifold does not admit in general an global S^1 action, only locally on both parts, since on the common part they do not coincide but locally they agree as belonging both to the same 2-torus action. Then it is still possible to define a collapsing structure in $RM(3,-1,1,D)$ for this manifold. This example has been put in a very general context in (Cheeger & Gromov, 1986-1990), where the notions of **F- and T-structures** are defined. Extra important notions are that of **polarized** and **pure-polarized** F-structures. Detailed definitions are also to be found at the end of the survey (Fukaya, 1990). Existence of various structures of those types is linked with the topology of the manifold but how exactly is still partially open, see the various references for the precised statements. What is difficult is to decide when one is or not within manifolds with bounded diameter.

For example for a general F-structure one can build up only Riemannian metrics with $-1 \leq K \leq 1$ and injectivity radius going to zero. If the structure is moreover polarized then one can get the volume going to zero, and if moreover it is pure-polarized then one has really a collapsing in $RM(d,-1,1,D)$. In (Cheeger & Gromov, 1986-1990) many results are obtained for the study of what is happening during collapsing to the various Riemannian invariants. The above theory is used also to study integral formulas for and the existence of characteristic numbers when the manifold is no longer compact but still of finite volume : see (Cheeger & Gromov, 1985a). For what happen with the spectrum (chpater IV) see (Fukaya, 1987b).

A byproduct of (Cheeger & Gromov, 1986-1990) is a **structure** statement for any Riemannian manifold. Namely there exists an universal number $\varepsilon(d)$ depending only of the dimension d such that, given any complete Riemannian manifold M of dimension d and with $-1 \leq K \leq 1$, then there exists an open set U (the **thin** part) of M with the following properties : on U there is an F -structure (of positive dimension) and at points in $M \setminus U$ (the **thick** part) the injectivity radius is larger than $\varepsilon(d)$. The results on F -structures are used in (Anderson, 1996a).

We look now at the structure of the collapsing itself when the limit set of $\{M_i\}$ is a (compact) manifold N . The answer appeared in (Fukaya, 1987a), (Fukaya, 1989). We are in $RM(d, -1, 1, D)$ and assume $\lim_{i \rightarrow \infty} d_{G-H}(M_i, N) = 0$. then, at least for large i , there exists maps $f_i : M_i \rightarrow N$ which are fibrations. The fibers are infranil manifolds and f_i is almost a Riemannian submersion. In (Cheeger, Fukaya & Gromov, 1992) the two points of view are united to get very strong statement for collapsing. See this text for detailed statements and open problems in the topic. Finally the bold question of the nature of the d_{G-H} - closure of $RM(d, -1, 1, D)$ was addressed in (Fukaya, 1988). When one is interested to various closure on a **given** manifold things are much easier, see **III . D**.

We look now at to boundary structure for subsets involving weaker curvature bounds. The first thing to do is to discard $K \leq 1$ as we did in section 1 for finiteness results but still keeping $K \geq -1$. It is intuitive that limits can now have singularities, think e.g. of small fingers, bubbles, etc. The major discovery was that there exists a reasonable generalization of Riemannian geometry which is stable by metric limit under the condition $K \geq -1$. This is the notion of Alexandrov's space, see **TOP. 9** for more on this. In case of collapsing under only $K \geq -1$ with a smooth limit one has a strong structure result in (Yamaguchi, 1991).

The urge is now for control only with the **Ricci** curvature plus of course various metric invariants. This is an extremely active topic today and hence hard to present concisely. Besides references of the preceeding two sections, we mention (Anderson, 1992c) ; (Cheeger & Colding, 1997a) is the beginning of a series taking in account previous results. Recall that Colding's L^2 -Toponogov

theorem (see the digression in A. 2) plays the role of Toponogov's one when one has only a lower Ricci bound. For the insight we mention that those results are part of a programme of Anderson and Cheeger, including the conjecture : " d_{G-H} convergence in presence of a lower Ricci curvature bound implies volume convergence". This is now proved in (Colding, 1996b).

Next in weakening the assumptions is to use only **integral** bounds, sometimes mixed with various other ones. We just mention (some of them already met) : (Gallot, 1986), (Gallot, 1987), (Yang, 1992), (Gao, 1990a), (Gao, 1990b), (Petersen, 1997b) and the expository (Anderson, 1990e).

II : THE GEOMETRY HIERARCHY OF RIEMANNIAN MANIFOLDS : SPACE FORMS

Defining, constructing, classifying space forms can be very hard and is still not finished. We explain at the same time what we mean by this hierarchy. Here again the situation will be completely different according to the signs under consideration. There exist books and partial surveys : (Buser, 1992), (Wolf, 1972), (Gromov & Pansu, 1990a), (Vinberg, 1993a). For the hyperbolic case excellent references are (Benedetti & Petronio, 1992) and (Ratcliffe, 1994). So that we will give below only few references, mainly to locate the dates. Do not believe that the geometry of space forms is completely mastered today. Amazing relations between geometry (volumes of polytopes, cross-ratio and its generalizations) and Number Theory are in full study and enjoy beautiful results. We just mention the chapter 14 by Kellerhals of the book (Lewin, 1991) and the informative (Oesterlé, 1992/93)

Again we work only in the compact case but most natural and important cases are those of finite volume, see **TOP. 5**. On rigidity, surveys are (Gromov & Pansu, 1990a) and (Pansu, 1993-94), and add the book (Farrell, 1997).

A. The constant curvature case

A fundamental property of euclidean spaces is the congruence axiom : two triangles with equal respective sides are always deduced one from another by an isometry of the space. Let us say that they cannot be distinguished : "they are the same" . Another way to formulate this is to say that they are metric spaces which are **3-point** transitive. So the beginning of the hierarchy is to ask for all metric spaces having this property, globally first then locally only. The global answer was known in the 20's and follows from Hopf-Rinow and Cartan philosophy, see for example the book (Cartan, 1928a), but the second edition (Cartan, 1951) is also good of course. Look at infinitesimal triangles. Their infinitesimal shape will yield the sectional curvature and the spaces we are looking for are those of constant sectional curvature. The global answer is the simply connected ones, namely the spheres (with the canonical

metric if with curvature 1), euclidean spaces and hyperbolic spaces Hyp^d (most often normalized with the curvature -1).

Local (non simply connected but compact) answers are then the quotients of one of those three spaces by discrete groups of their isometry group. What is the state of affairs today ? Things were quite clear by the end of the 30's for **surfaces**. We discard first the trivial case of positive and zero sign. On the negative side it is easy to geometrically build up examples. One way is to take suitable triangles or polygons in the hyperbolic plane and make tessellations with them. But then the group yielding the quotient is hard to visualize. Another way is to glue together pantaloon pieces along their boundaries, provided they are closed geodesics. The pantaloon pieces are obtained by gluing to identical hyperbolic hexagons all of whose angles are equal to $\pi/2$. This latter way enables us to build up all examples, as seen by working back by dissection (this is perfectly exposed in (Buser, 1992)). The other way is to use conformal representation for compact topological surfaces of any genus. Finally the complete classification was worked out by Teichmüller in the late 30's . On an orientable surface of genus γ , the constant curvature structures build up a space with $6\gamma-6$ parameters. Building those space forms by group theory is harder and is basically Number Theory, see for example (Vigneras, 1980a). The link between the algebra and the genus of the surface is already subtle.

Starting dimension three difficulties appear formidable. We do things very sketchy. For all dimensions the positive and the zero case were understood almost completely in the 60's and are detailed in the book (Wolf 1972) (try to get the second edition). Contrary to what most people think the classification is not completely finished but this is more an algorithmic problem. Moreover the rigidity problem is almost not studied in the positive case, in contrast with the negative case to come now. To get some feeling about the mysterious and hard component in the positive case see (Milnor, 1966). The flat case is based on the famous Bieberbach theorem (1911-12) : the flat compact manifolds are tori or finite quotient of them.

The negative case is already difficult starting dimension three. It was only in 1931 that examples were built in (Löbell, 1931). This was geometric, using tessellation with polyhedra in three-dimensional hyperbolic space. In the 60's existence of negative curvature space forms were in the air, specially with Selberg. More surprising : many papers appeared which were studying them

without knowing their existence. The general result has to wait (Borel, 1963). Important to know is that Borel construction is entirely Number Theory, finding suitable discrete subgroups (called "arithmetic") of the isometry group of Hyp^d .

There is a good reason for that. In (Vinberg, 1984) it was realized that geometric tessellation constructions **cannot** exist in large dimensions. Left today is the exact value of the limit dimension, see (Vinberg, 1993a).

How about now some kind of Teichmüller theory (deformations, number of parameters) ? The big event was (Mostow, 1968) : space forms of constant negative curvature ($d > 3$) are **rigid**. As soon as the fundamental group is given as a group, one has Riemannian uniqueness. So important and conceptual is that rigidity that successive simpler and simpler proofs appeared. The last one is nothing but a direct corollary of (Besson, Courtois, & Gallot, 1995a) and (Besson, Courtois, & Gallot, 1995b) to be seen also in V. A and B. See also the survey (Besson, 1996b), the expository (Pansu, 1996-97), (Pansu, 1993-94) and the book (Farrell, 1997).

This rigidity does not imply the classification. Today one still does not know everything. On one side, we have the Borel type arithmetic examples (a precise definition of **arithmeticity** is a little long). On the other hand, first in (Gromov & Piatetski-Shapiro, 1988a), then in (Farrell & Jones, 1993), are built up examples which are definitively **not** arithmetic. If arithmetic ones or non-arithmetic ones are more numerous is the principal open question.

An important issue for negative space forms are **volumes**. When $d=2$ Blaschke-Gauss-Bonnet formula implies immediately that the set of volumes is an arithmetic progression. The existence of an universal positive lower bound for any dimension is in (Wang, 1972) where it is proved that starting dimension 4 and for space forms of any rank the set of volumes is discrete. In (Prasad, 1989) the volumes of all arithmetic space forms are explicitly computed. The case of dimension 3 was solved in (Jorgensen, 1977) and (Thurston, 1978) and the result is fascinating : volumes are isolated from zero (however the best value is still unknown today) but they have accumulation points located on a discrete scale. For more see (Gromov, 1981b) and (Ratcliffe, 1994), page 501. See the recent (Goncharov, 1997) to realize the depth of the problem.

B. Space forms of rank one

Pursuing our hierarchy we look now for metric spaces which are "only" **2-point** transitive i.e. there is always an isometry carrying one pair of points into another pairs (provided of course that their respective distances are equal). Another formulation is : looking for spaces which are **isotropic**, namely all unit tangent directions are metrically equivalent. If simply connected they are all known. Discarding the constant curvature preceeding ones, one is left with the KP^n and their negative curvature analogues, denoted here by $HypKP^n$. Their curvature range is from $1/4$ to 1 or from -1 to $-1/4$ after normalization. The CP^n together with the $HypCP^n$ are the complex geometries corresponding to the spheres and the hyperbolic spaces. This classifications is in (Wang, 1952) for the compact case and in (Tits, 1955), for the noncompact one. It is still a long story to do it in detail, the best reference today to our knowledge is that of (Karcher, 1988) (see page 120). The difficult part of the proof is to show that one is in a symmetric space, see (Szabo, 1991) for a short proof. Then it is forced obviously to be of rank one, see C below.

We now look for compact quotients of those simply connected ones. In the positive case (spheres excepted) there are no such quotient (except up to Z_2 because dimensions are even here and we have then Synge's theorem, see 0. C). We turn to the negative curvature case. The complex case is particularly interesting because of the relations with complex analysis and algebraic geometry. General existence is again in (Borel, 1963), of course of arithmetic type. Non-arithmetic examples appeared in (Mostow, 1980), built up by subtle tessellating polyhedra in $HypCP^2$ and in $HypCP^3$. Also later on in (Deligne & Mostow, 1986) by different and expensive Number Theory techniques. The question of deciding if there exists non arithmetic examples for the $HypCP^n$ is still open today for higher n . But for the quaternionic and the Cayley case, $HypHP^n$ and $HypCaP^2$ it is proved in (Gromov & Schoen, 1992d) that all compact quotients of these spaces are necessarily arithmetic. The proof uses hard Analysis, precisely harmonic maps into manifolds with singularities, see TOP. 7.

There is a rigidity theorem as well for the compact quotients of the $HypKP^n$ (Mostow), but today it can be gotten by the general result of (Besson, Courtois & Gallot, 1995a) already mentioned.

The search for spaces which are only **measure-isotropic** (so called **harmonic**) started in the early 40's. There are many equivalent definitions, the first says that at every point the solid angle (the infinitesimal measure along a geodesic starting from this point) depends only on the distance. The words harmonic was chosen because this condition is equivalent, at least locally, to the fact that the value at every point p of any harmonic function f ($\Delta f=0$) is equal to its mean value on every distance sphere centered at p . Lichnerowicz conjectured in 1944 that such spaces are, at least locally, isometric to a space form of rank one, so that measure-isotropy will force metric isotropy. After many intermediate results the conjecture was settled positively for the compact case by Szabo in 1990 and then negatively in the non compact case by Damek and Ricci. All the references can be found in (Berndt, Tricerri, & Vanhecke, 1995) and its bibliography.

C. Space form of symmetric spaces of rank larger than one

Next to 2-point transitive spaces come 1-point ones. But this is nothing more than **homogeneous** spaces. In some sense this category is too general, even if met it many times below when constructing examples of Riemannian manifolds.

But there is an basic intermediate category, the **symmetric spaces**, discovered by Elie Cartan in the 30's. They are the manifolds for which the geodesic symmetry around any point is (at least locally) an isometry. This implies immediately (local) homogeneity by composing those symmetries at pairs of points. But another equivalent definition (this has to be proven and is due to Elie Cartan) is that the curvature is invariant by parallel transport. It is the most natural homogeneity condition (think of the golden triangle in **O. A** and of Ambrose's problem at the very end of **TOP. 4**). Elie Cartan showed that these (local) geometries can in fact completely classified, a job done first at the level of Lie algebra. The result for irreducible ones is a very restricted list, containing infinite series and a finite number of exceptions in low dimensions. All members of the infinite series are spaces known in geometry: the simple Lie groups themselves, the Grassmann manifolds over the fields $\mathbf{R}$, $\mathbf{C}$, $\mathbf{H}$, the set of complex structures, of Lagrangian subspaces, and few others.

To a symmetric space is attached an integer, called its **rank**. It is the common dimension of all maximal totally geodesic flat submanifolds. They are all conjugate under the group action. The rank one spaces are those of section **B** above. These flat

submanifolds are, in the simply connected case : tori for the compact case and euclidean spaces for the noncompact case. This completely determines the geodesic flow (see V. B). In particular for the spheres and the KP^n all their geodesics are simple, closed (periodic) and with a common length.

These irreducible symmetric geometries split again in two classes : those of nonnegative curvature and those of nonpositive one. Finding all the quotients in the positive case is a finite job, which has been done back in (Cartan, 1927). For the negative case it is again in (Borel, 1963) that compact quotients were proved to exist for the first time, always by arithmetic considerations. But here, if there is a classification problem, it is for number theoreticians. Because in (Margulis, 1977) a **superrigidity** result was proven, which implies not only topological rigidity but also arithmeticity for the rank ≥ 2 case. Margulis tools were expensive, today there are other ways to recover Margulis result : see for example (Jost & Yau, 1990). Results like (Besson, Courtois & Gallot, 1995a) still do not yield the superrigidity.

Now one can look at generalizing euclidean geometry with spaces more general than Riemannian manifolds. In fact one is forced more or less to do so when working in Riemannian geometry, this was seen in I. C. 3 : see TOP. 9.

III : THE SET OF RIEMANNIAN STRUCTURES ON A GIVEN COMPACT MANIFOLD : IS THERE A BEST METRIC ?

A. The problem

We still stick to the compact case. Very natural is the following question : given a compact differentiable manifold, does it carry a **best**, or a family of best, Riemannian structure(s)? We saw in II. A that in the case of surfaces things go wonderfully. Any compact surface admits a metric of constant curvature. Moreover on a given one they are completely classified (Teichmüller theory), building up a finite-dimensional "moduli" space. How about higher dimensions ? This question was asked to the author by Thom circa 1960 and is the very first problem in the list (Yau, 1990). In the low dimensions 3 and 4 the question is of primary importance since Riemannian

geometry seems to be a basic tool to solve the various still open topological conjectures (see TOP. 8).

Today there is not a single general satisfying statement in dimension larger than 4, only very partial results. The lower dimensions are better understood. One of the reasons which makes the problem difficult is the following one. To get a best metric it is reasonable to work within the condition $-1 \leq K \leq 1$. But we saw in I. C. 2 that in order to get a limit we need moreover volume $\geq v > 0$ and diameter $\leq D$. Even when the topology of the manifold implies the volume condition there is still no way to control the diameter.

The scheme of attack is to look at a suitable numerical functional $F(g)$ defined on the space of metrics g and then search for those for which $F(g)$ is minimum (or at least critical, that is to say the derivative of $F(g)$ at g is zero for any variation of the metric). If the functional is not scale independent, just normalize it. If criticality says us something, we can stop here. If we need more, one can then look for an extremum. For the general context see the chapter 4 in (Besse, 1987), (Anderson, 1990e) and (Sarnak, 1997a).

Sarnak considers in particular as functional the determinant of the Laplacian, see IV.

The first thing to do is to study the space of all Riemannian metrics on a given compact manifold, but also its quotient by the set of all diffeomorphisms since we are interested mainly not in metrics but in structures (metrics up to isometry). There is no reason for this (infinite dimensional) structure space to be smooth, but there is a nice slice result which gives a local structure and its "tangent space", see 12. C of (Besse, 1987). See also (Gil-Medrano & Michor, 1991). For the topology of the set of all Riemannian structures on a given manifold we know only of (Bourguignon, 1975).

B. The minimal volume and the $\text{Min} \|R\|^{d/2}$

It seems to us that the most natural definition of an optimal metric is : "it is the less curved one". So that the obvious functional is

$\int_M \|R\|^{d/2}$ and we immediately attach to a compact manifold M the

invariant $\text{Min} \|R\|^{d/2}(M)$ which is the infimum of all $\int_M \|R\|^{d/2}$

when considering all Riemannian metrics on M . Note that the Blaschke-Gauss-Bonnet theorem shows that this functional is certainly not good in dimension 2. But precisely we are interested

in dimensions higher. It turns out that, at least in dimensions larger than 4, the condition for g to be critical is not understood today : it is a partial differential equation for g and its curvature which actually is undecipherable. There are only results telling us that in some cases this invariant is zero and non zero in some other. But the exact delimitation is still a mystery. As well as to know if its vanishing is or not equivalent to that of the minimal volume, see

TOP. 1. D. The functional $\int_M \|R\|^{d/2}$ is also encountered in

finiteness theorems with Ricci curvature : see **I. C. 1.**

But the lower dimensions 3 and 4 are exceptions. For 4 the generalization of Blaschke-Gauss-Bonnnet theorem (section **0.E**) reads :

$$8 \pi^2 \chi(M) = \int_M (\|R\|^2 - \|Ricci - \frac{scal}{4} \cdot g\|^2) dm$$

Then, on a given M , our functional is an absolute minimum when the Ricci curvature of g is proportional to it : $Ricci(g) = \frac{scal}{4} g$.

Passing by this proves that the characteristic of an Einstein 4-manifold is always nonnegative, see **C**. But the converse works only when $\chi(M)$ is nonnegative. The metrics of whose Ricci curvature is proportional to the metric itself are nothing but what we are going below to call **Einstein**. And we will study them quite in detail in **C**, because it is the sole notion for a best metric which seems really systematically workable today in general dimensions.

In dimension 3 partial (but deep) results are obtained in (Anderson, 1996a) and (Anderson, 1996b). and are paving the way to the Thurston geometrization conjecture.

Three other functional are more geometric (in fact two of them are curvature-free) and look interesting : they are the minimal volume, the systolic volume for some very special manifolds (see **TOP. 1. E**) and the embolic volume. The **minimal volume** of a compact differentiable manifold is the infimum of the volume of all Riemannian metrics on this manifold which have a sectional curvature between -1 and 1. It is equivalent to look at the infimum of $\sup |K|$ under the condition volume = 1. This shows more the "less curved" definition. The **embolic volume** is the

infimum of the volume under the condition that injectivity radius is equal to π .

Despite such simple definitions results on them are so few that we discard them here and postpone things to TOP. 1. D for the MinVol. For the systolic story, which is quite interesting in itself, see TOP. 1. E. Finally for the embolic story recent references are (Croke, 1988b) and (Grove, Petersen, & Wu, 1990-1991) (see theorem C), see also TOP. 1. C. The difficulty of studying these invariants can be seen by just telling that it is already not known when they are zero or not. This is a place Chern's integral formulas are useful but not enough (see O. E and I. C. 1).

However we mention here the main result of (Besson, Courtois & Gallot, 1995a) in V. B. One of its many byproducts is that the best metric is known on compact space forms of negative curvature (rank 1): it is unique and is precisely this locally symmetric metric. This is true for more than the minimal volume only, see for this result V. B and TOP. 1. D.

In the case a manifold admits a metric of a given sign, a best metric would naturally be one with the best pinching. Except the cases met above in I. A. 2 this question has today no answer and we already explained why it is hard at the end of I. B. 1.

C. The case of Einstein manifolds

We turn now to Einstein manifolds. The story started with (Hilbert, 1915). There, for theoretical physics motivations, he computed the directional derivative in the space of all Riemannian metrics of the total scalar curvature

$$F(g) = \int_M \text{scal}(g) \, dg.$$

Without normalization, critical implies for g that $\text{Ricci}(g) = 0$, which is too restrictive, see below. So we normalize the volume to be some given constant (which does not matter) and get the condition " $\text{Ricci}(g)$ is proportional to g " : $\text{Ricci}(g) = \frac{\text{scal}(g)}{d} g$ (this is nothing but the Lagrange multiplier classical technique). Using Bianchi identity the proportionality function is necessarily constant as soon as $d \geq 3$. Such a metric will be called **Einstein**. In

dimension 3 this condition is too strong, it implies constant sectional curvature, so that we discard this dimension from now on. In dimension 4, we saw that this is equivalent to minimizing the total square norm of the full curvature tensor. Note that Einstein manifolds come automatically into the picture when one looks for Ricci pinching, see then end of I. A. 4.

The fact that the equation defining Einstein metrics is a PDE (because the number of parameters are $d(d+1)/2$ on both sides) is rising great hopes. As well as the fact to have a smooth functional. However today there is no way to use any available kind of Morse theory to get some existence (of critical points). In fact experts are divided, some think it cannot by essence. Some other think a tool like Floer homology (see (Hofer, Taubes, Weinstein, & Zehnder, 1995)) could help. See the end of I. B. 1, Besse's book and below for more comments.

This Hilbert functional received a new consideration in (Connes, 1995-6) where it appears as the $(d-2)$ -volume of the manifold, therefore an "area" in dimension 4.

Digression : the Yamabe problem. We make here an historical interlude together with a topic completely omitted elsewhere in this survey, except here and indirectly in TOP. 8. This is the Yamabe problem. In (Yamabe, 1960) the author tried, starting with any Riemannian metric g , to deform it conformally into one $f.g$ of constant scalar curvature, where f is a positive numerical function. One can think that the secret hope of the author was in dimension 3 to attack the Poincaré conjecture (compare with Hamilton result in I. B. 2). To go further in dimension 3 is the content of (Anderson, 1997) and references there.

The problem to solve was a non-linear elliptic equation in the unknown conformity function f . This question in Analysis is extremely hard and was extremely studied. This because the equation is of the non-linear type $\Delta f + \frac{d-2}{4(d-1)} \text{scal} = f^{(d+2)/(d-2)}$ where the unknown conformity function f comes with the "limit" exponent for Sobolev embedding. For larger or smaller exponent standard results are available but are of no use here. Today the Yamabe problem is almost completely understood after basic contributions of Aubin and Schoen. We refer to the surveys (Besse,

1987), 4.D, (Hebey, 1993), (Hebey, 1995a) and the references there. We just note that the set of metrics of constant scalar curvature, as soon as $d > 2$, is always an infinite dimensional space and is definitively not what we want only when looking for a "best" metric, see also (Anderson, 1997).

We come back now to Einstein manifolds. How are things today with existence, uniqueness or/and moduli of Einstein metrics ? This is a whole subject in itself and the book (Besse 1987) is entirely devoted to it and was up to date at that time. The introduction (chapter 0) is a good resumé. Past to 1987 few papers appeared on the subject, very recent ones are (Eschenburg & Wang, 1995), (Boyer, Galicki, & Mann, 1994a), (Boyer, Galicki, Mann, & Rees, 1997) and the survey (Anderson, 1994). We now give very briefly the main results and the main open questions.

In dimension 4 there are topological obstructions. The first type uses Allendoerfer-Weil formula, which yields among other conditions the nonnegativity of the Euler characteristic seen above in B. With a more refined analysis of the integrand one can obtain the inequality $\chi(M) \geq \frac{3}{2} |\tau(M)|$ between the characteristic and the signature τ : (Thorpe, 1969) and see the chapter 6 of (Besse, 1987) for more. The second type uses the minimal volume of Gromov (TOP. 1. D). Mixing both one can see that in some sense "many 4-dimensional manifolds do not admit any Einstein metric" : see (Sambusetti, 1996), which uses techniques from (Besson, Courtois & Gallot, 1995a), and (LeBrun, 1996).

Oppositely the most baffling open question is that, starting dimension 5, there is no obstruction known to the existence of an Einstein metric on any compact manifold. This means that it is not excluded that any compact differentiable manifold (with this dimension condition) can be put on an Einstein metric. It seems that experts have only one guess today, namely that most manifolds ($d > 5$) will admit Einstein metrics but with reasonable singularities : (Anderson, 1994).

So that we turn now to the existing known examples and results. First are the locally irreducible symmetric spaces (in particular generalized space forms). The reason is that the isotropy group is irreducible and cannot leave, up to a scalar, more than one invariant quadratic form : use diagonalization. For homogenous but

non symmetric spaces, even if in theory curvature is algebraically computable, to find Einstein homogeneous metrics can be very subtle, and even can be proved to be impossible in some cases. For this, see a series of papers by Wang and Ziller quoted in (Besse, 1987) and (Wang, 1992). Needless to say that we are far from a classification of homogeneous Einstein manifolds. A recent text is (Kowalski & Vlasek, 1993) which in particular gives a complete classification of the Einstein and homogeneous metrics on the Aloff-Wallach spaces $W_{p,q}$; we met those examples in I. B. 1.

Then come a quite a large variety of non-homogenous examples, obtained by different techniques : spaces with a group action and "more or less almost homogeneous", fibre bundles of various types, twistor theory (see TOP. 6), 3-Sasakian manifolds (TOP. 3. A). Today examples are getting more and more numerous. We only give some references posterior to Besse's book. Note that an up to date second edition of it will come soon. The very recent (Böhm, 1996) is important in two respects : first it gives among others non standard Einstein metrics on even-dimensional spheres. The technique uses cohomogeneity one manifolds and the equations obtained in (Bérard Bergery, 1982) for obtaining an Einstein metric. Second it presents examples of infinite sequences of Einstein metrics on given compact manifolds with fixed volume which is not converging to any metric, and this shows that the Palais-Smale "condition C" (Palais & Smale, 1964) is not satisfied in general for Riemannian metrics. For this condition see (Jost, 1995) and see other types of such "counterexamples in I. B. 1. Before, in (Jensen, 1973), were built up non standard Einstein metrics on the S^{4n+3} spheres.

Uniqueness and classification is in its infancy. The above examples show the complexity of the topic. We meet here our favourite paradox : one knows uniqueness for the compact space form which are quotients of Hyp^4 and $HypCP^2$. For the latter it is in (LeBrun, 1995), where the tool are the new Witten invariants (TOP. 8). For Hyp^4 it is a byproduct of the strong result of (Besson, Courtois & Gallot, 1995a) which is quoted in V. B and in TOP. 1. D. An exceptional case is that of $K3$ surfaces. Using ingredients of various horizons and the Kähler facts just below one has a complete description of all the Ricci flat metrics on them : see details in (Besse, 1987), 12. K.

A puzzling question is that of the signs of Einstein manifolds. This is the sign of the proportionality factor (i.e. of the scalar curvature).

For example it is plus (resp. zero, minus) in the case of space forms of positive (resp. zero, negative) curvature. It is still open today if some manifold will bear Einstein metrics with at least two different signs.

Noncompact Einstein manifolds are also studied, see among others (Heber, 1997) and (Hitchin, 1995).

We turn now to the only domain where things are very satisfactory, namely for **Kähler** manifolds (see **TOP. 3. B**). Moreover this topic turned out to be important in mathematical physics. The basic remark is the following : using the complex structure one can transform the symmetric differential form which is the Ricci curvature in a exterior form (of degree 2) which is automatically closed . It was easy folklore in the 50's that, by Chern formulas and de Rham theorem, this form belongs to the first Chern class of the Kähler manifold under consideration. So that we have an immediate necessary condition for Kähler manifold to admit any Einstein metric : the first Chern class should have some de Rham representative 2-form which (via the complex structure) is either : zero, positive definite or negative definite. This condition is not that strong. Many algebraic manifolds satisfy it and this can be decided by using today standard techniques (see e.g. (Hirzebruch, 1966), (Griffiths & Harris, 1978)) to compute their Chern classes (see **TOP. 3. B**).

We briefly explain here why things are "workable" for Kähler manifolds. In fact the Kähler structure g yields an exterior closed 2-form ω of complex type (1,1) and Kähler variations in the same cohomology class are easily seen to be necessarily of the form $\omega + \sqrt{-1} \partial \bar{\partial} f$ where f is a numerical function on the manifold. The partial differential equation needed for f to yield an Einstein metric is of Monge-Ampère type and then hopefully solvable.

This lead Calabi in the pioneer paper (Calabi, 1954) to conjecture that as soon as the zero or the definiteness of the Chern class was insured, then one can change the initial Kähler metric into a new one which is finally Einstein. More : in the negative case one has uniqueness. It was only in (Aubin, 1970) that any progress was obtained and in (Aubin, 1976) and (Yau, 1978) that the following was proven. Aubin proved the existence in the negative case, Yau proved it in the zero and the negative case. Then in (Futaki, 1983) an obstruction was found for the positive case. In (Tian, 1990) one has a definite answer for complex surfaces ($d=4$) : necessary and

sufficient conditions to get Einstein Kähler metrics of the positive type. For larger dimensions, the latest result is (Real, 1996). A conjecture is formulated in (Tian, 1996). See also (Bando & Mabuchi, 1987). For the zero case and a proof following the Ricci flow see (Cao, 1985).

Calabi-Yau existence results are extremely useful. First the uniqueness of the Kähler structure on $\mathbb{CP}^n$ (for any n) : (Yau, 1977) . Besides Yau existence result one needs the deep (Hirzebruch & Kodaira, 1957). For $\mathbb{CP}^2$ one even has uniqueness of the complex structure. Another application is the inequality $c_1^2 - 3 c_2 \leq 0$ between the first two Chern classes of any Kähler manifold. For this and more see 11. B in (Besse, 1987). A still mysterious applications is the fact that, for a Kähler manifold, the vanishing of the two first Chern classes c_1 and c_2 implies that the manifold can be made flat and in particular is a finite quotient of a torus : see references and comments in page 67 of (Bourguignon, 1993). The mystery lies in the fact that nobody knows a direct proof.

To stay in the Kähler Einstein domain, we mention (Hulin, 1995). The author uses the very interesting notion of **diastasis** introduced by Calabi in 1953, see the expository (Berger, 1993b) and **TOP. 3. B** . And then proved very strong results for the complex submanifolds of $\mathbb{CP}^n$ which are Einstein. The diastasis was rarely used up to now but it might still have a rich future.

Moduli of Einstein metrics are not bad, in the sense that we know they are always finite-dimensional and real analytic stratified spaces. But they are explicitly known in extremely few cases, the most spectacular is that of K3 - surfaces seen just above. For all this see the chapter 12 of (Besse, 1987). In (Anderson, 1992b) there is a deep study of the moduli space of Einstein metrics for any compact manifold of dimension 4 and this for the three possible signs. Special attention is given to the closure, here orbifolds (**TOP. 9**) appear and in particular the above results for K3- surfaces are precised and (algebraic) surfaces with singularities come into the picture.

Of special interest and completely mysterious is the case of Ricci flat metrics (the zero sign case). Today no example are known for which the holonomy group is not a special one, namely $SU(n)$ and $Sp(n)$ in the Kähler case, then G_2 and $Spin(7)$ which are very special, see **TOP. 3** .

Some final facts on Einstein manifolds. First they are, in a sense to be precised, always real analytic. This is a good use of harmonic coordinates, see section 5. E of (Besse, 1987) and the references there. Second, and posterior to 1987, there exist many results on pinching, convergence for Einstein manifolds. They are mainly all contained in the texts mentionned in the corresponding sections : end of I. A. 2 and I. C. 1,2. Non compact Ricci flat manifolds are important and extremely studied today, see TOP. 5 and the recent (Cheeger & Tian, 1994).

In dimension 4 there is notion weaker than that of Einstein, namely **anti-self-duality**. See the entire chapter 13 of (Besse, 1987) and complete it with (Donaldson & Kronheimer, 1996b), (Donaldson, 1996a) and the expository (Gauduchon, 1992/93) on (Taubes, 1992). For the need of the proof of Taubes's result special norms are introduced, which are close to the various global norms we mentionned in different places in I.

D. Some topological closures

We come back to a strict interpretation of the title of the present section. In the space of all Riemannian metrics on a given M in the spirit of I. C. 2 one looks for the closure of the various subsets defined by some curvature condition. This is surveyed systematically in (Lohkamp, 1992). We just mention few cases. The notations are obvious for those spaces, e.g. C^α -closure($\text{Ricci} \geq \kappa(M)$). Following (Gromov, Lafontaine, & Pansu, 1981a) the **RCT** theorems imply the strongest possible closure : d_{G-H} -closure($K \geq k(M)$) = $K \geq k(M)$ and the same is true for $\leq k$. This implies any desired C^α smoothness. For a lower Ricci it is an open problem for d_{G-H} but true for C^0 . For a lower scalar curvature bound things are a complete mystery, see I. B. 3. For an upper bound, Lohkamp's result in I. B. 5 can be written d_{G-H} -closure($\text{Ricci} \leq \alpha(M)$) = {all metrics on M } . See also TOP. 9 .

IV : THE SPECTRUM, THE EIGENFUNCTIONS

Recall that on a Riemannian manifold we have a canonical differential operator Δ of second order operating on functions, namely just take the trace (with respect to the metric) of the Hessian (the second covariant derivative) : $\Delta f = - \text{trace}_g(\text{Hess } f)$. At the center m of normal coordinates this is written

$$\Delta f(m) = - \sum_{i=1, \dots, d} \frac{\partial^2 f}{\partial x_i^2}(m) . \text{ The Laplacian of Riemannian}$$

manifolds controls for them first the *heat* equation $\Delta f = - \frac{\partial f}{\partial t}$ where $f(m,t)$ is the temperature at the point m and at time t .

Also the *wave* equation $\Delta f = - \frac{\partial^2 f}{\partial t^2}$: the wave equation for the height $f(m,t)$ of the "water" after time t at the point m if you consider more or less abstractly (M,g) as a thin sheet of water. To the acoustic and/or tidal language one can prefer the equivalent language of geometrical optics and talk about propagation of **light**. The wave equation can also be considered as describing (M,g) as an abstract vibrating object . Finally the Schrödinger equation uses **complex** valued functions. It is written as $\hbar^2 \Delta f = i \hbar \frac{\partial f}{\partial t}$ (where

$i = \sqrt{-1}$ and $\hbar$ is the Planck constant). Very close in fact to the heat equation is **Brownian motion** on Riemannian manifolds, whose "propagation speed is the Ricci curvature". We can quote : (Stroock, 1996), the text by Elworthy in (Diaconis, Elworthy, Nelson, Papanicolaou, & Varadhan, 1985-87) and (Elworthy, 1988), (Pinsky, 1989), (Pinsky, 1990).

Despite of its importance, the spectrum of Riemannian manifolds ("spectrum" is a short cut for the study of the eigenvalues and the eigenfunctions of the operator Δ) does not seem to have much interested interesting mathematicians before (Minakshisundaram & Pleijel, 1949), (Avakumovich, 1956). Exceptions are the spherical harmonics and the computation of the spectrum of the complex projective space in (Cartan, 1931). Add (Lichnerowicz, 1958) result on λ_1 which will be commented at the very end of this section.

There are books devoted to the topic. The first was (Berger, Gauduchon, & Mazet, 1971) ; some people still like it because it is a fast introduction to the topic, (Buser, 1992) is extremely complete for the case of Riemann surfaces (namely the curvature is a negative constant), (Chavel, 1984) and (Chavel, 1993) cover a good part of "classical" material, then (Bérard, 1986) which, besides many things, contains a systematic bibliography up to date but only up to 1982. See also (Colin de Verdière, 1992). The books (Gilkey, 1995) and (Berline, Getzler, & Vergne, 1992) concern

the case of general elliptic operators, specially in connection with **TOP. 6.** (Hörmander, 1985) is encompassing many things.

The **isospectral** problem started to be in the air in the 60's, when the question of isospectrality was raised by Leon Green and soon answered negatively in (Milnor, 1964). This paper launched a lot of activity to find examples of isospectral (non isometric) manifolds, see below. But the real triggering paper was (McKean & Singer, 1967). The problem was to find what is really possible to extract as informations from the asymptotic expansion of the heat kernel obtained in (Minakshisundaram & Pleijel, 1949), see below. Apparently not much in a naive sense. But with the help of (Patodi, 1971), and the sitting next by paper in the same journal, people managed to link this with the already existing Atiyah-Singer index theorem. Then one got a complete set of results. Although quite special, was the discovery of the η -invariant, see **TOP. 6 . A**, but Δ was needed not only on functions but as well on all exterior forms). This, in turn, triggered the heat equation technique for various bundles over manifolds, as well as spinor bundles. We will not insist on this, for which we refer the reader to the books (Gilkey, 1995), (Berline, Getzler & Vergne, 1992) and (Lawson & Michelsohn, 1989). Spinors were met in section **I. B. 3** when positive scalar curvature was in question, see **TOP. 6** for more. For the above, techniques are mixing Analysis and topology.

We now come back down to the manifold itself and to the most natural question : describe the behavior of the spectrum, as a subset of $\mathbf{R}$, under various conditions. First, in general, then under geometric conditions (as curvature, volume, diameter), finally as linked with the geodesic flow, in particular with ergodic theory for the geodesic flow and periodic geodesics. Here is the state of affairs today.

Already in (Minakshisundaram & Pleijel, 1949), who used a parametrix and the ζ -function of the spectrum $\zeta(s) = \sum_{\lambda \neq 0} \lambda^{-s}$, one

knew that the spectrum has an asymptotic behavior whose first term is given by the dimension and the total volume $\text{Vol}(g)$ of the manifold :

$N(\lambda)$ = number of eigenvalues smaller than λ

$$\approx \lambda \rightarrow \infty \quad \frac{\beta(d)}{(2\pi)^d} \text{Vol}(g) \lambda^{d/2} + o(\lambda^{d/2})$$

(where $\beta(d)$ is the volume of the ball of radius 1 in $\mathbf{R}^d$)

This is the so-called Weyl's estimate. The name comes from (Weyl, 1911) (or Complete Works) where this estimate is obtained for plane domains with boundary by cutting them in smaller and smaller pieces and using the minimax principle, see an exposition in III. C of (Bérard, 1986)). But the second term of the asymptotic expansion is a completely different story which we will address below.

In (Minakshisundaram, 1953) the so-called **heat kernel**

$K(x,y,t) = \sum_i \exp(-\lambda_i t) \phi_i(x) \phi_i(y)$ (where the λ_i are the eigenvalues and ϕ_i are corresponding orthonormalized eigenfunctions) was constructed in a way which yields immediately, when t goes to zero, an asymptotic expansion of the following form :

$$K(x,x,t) \approx \frac{1}{(4\pi t)^{d/2}} \left(\sum_{k=0}^{\infty} u_k(x) t^k \right)$$

The numerical functions $u_k(x)$ are given by universal (but "in the air") universal formulas involving, thanks to Elie Cartan philosophy of normal coordinates (O. B), only the curvature tensor of (M,g) and its covariant derivatives at x . In particular

$$\sum_i \exp(-\lambda_i t) \approx_{t \rightarrow \infty} \frac{1}{(4\pi t)^{d/2}} \left(\text{Vol}(g) + U_1 t + U_2 t^2 + \dots \right)$$

$$\text{where } U_k = \int_M u_k(x)$$

This arose great hopes since the knowledge of the spectrum is equivalent to that of the (always converging for $t > 0$ series

$\sum_i \exp(-\lambda_i t)$. Sadly enough these invariants U_k are of almost no direct use except the first one which is the volume and the second one only in dimension 2. Indeed $u_1(x) = \frac{1}{6} \text{scal}(x)$ then the Blaschke-Gauss-Bonnet (O. E) formula yields the Euler-Poincaré characteristic in function of the spectrum. To see the weakness of

the U_k in general, think of manifolds of constant curvature (space forms, e.g. flat manifolds). They will have by their universality all the same U_k for every k as soon as the volume is known. And the volume is certainly not enough (except in dimension 1 and already not when 2) to recover the manifold up to isometry. However, although not too much of direct use, what is happening in higher dimension was in fact a tremendous incentive. For example a query of the middle 60's was if U_k is a topological invariant for every manifold of dimension $d = 2k$. This turned out to be false but see in **TOP. 6** what came out of it. It is fair to say that the asymptotic expansion, and its universal curvature form, is of basic use in the finiteness and compactness results to be seen soon below.

Finally we remark that it is hard from the spectrum (on functions) to extract (except when the dimension is 2) any topological information. To our knowledge there is no connection between the spectrum (only on functions) and the topology. But the story is different when using the spectrum of objects more general than functions, in particular all exterior forms, see **TOP. 6**. One reason for the case of dimension 2 to be special is that the spectrum of functions determines the spectrum of all differential forms by duality (just use Hodge theory, **O. D**) and add the first Betti number for harmonic 1-forms).

Coming back now to the asymptotic value of $N(\lambda)$ a classical theorem in Analysis yields the Weyl asymptotic from the first

term in that of $\sum_i \exp(-\lambda_i t)$, but only with a "small o". Note here

$\sum_i \exp(-\lambda_i t)$ is a "practically weaker" (when using only its asymptotic expansion in t near zero) as we are going to see that the

classical ζ -function $\zeta(s) = \sum_{\lambda \neq 0} \lambda^{-s}$. In (Avakumovich, 1956) (only

for dimension three) and (Hörmander, 1968) in general (in fact for any elliptic operator with the obvious modifications in the powers involved) showed that the next order term is in fact a "capital o": $O(\lambda^{(d-1)/2})$. This is the best possible as seen on standard spheres. This last result will control the **gaps** in the spectrum: they cannot be too large, there is some regularity. But Hörmander result was yielding a $O(\lambda^{(d-1)/2})$ completely "in the air". It is only in (Gromov, 1996) that was first obtained an explicit control on the gaps with

the geometry. Gromov needs upper and lower bounds for the curvature and a lower bound for the injectivity radius. This not surprising in view of **I. C. 2** . The proof is extremely involved and indirect. This is contrasted with Hörmander proof, based on the wave equation. The question to obtain the desired control with the wave equation is open today. Even in the case of flat tori a fine study of the gaps is extremely subtle, see (Sarnak, 1997b).

It is important to realize that a control of the gaps **from the beginning** needs some geometric conditions. Because (Colin de Verdière, 1987) proved this basic fact : given any manifold of dimension three or more, one can always find on it a metric whose beginning of the spectrum is any finite given subset of $\mathbf{R}^+$ (including any desired multiplicities) .

In the spirit of control theory as in the digression in **I. A. 2**, one can expect to be able to get both upper and lower bounds for the eigenvalues from various conditions on the geometric invariants of the metric. There is a very satisfying answer. It started with (Gromov, 1980a) and was finished in (Bérard, Besson, & Gallot, 1985). More references are in the book (Bérard, 1986). Upper bounds are obtained with only the volume and a lower bound for the Ricci curvature. Lower bounds need a lower bound for the Ricci curvature and the diameter. Examples show that this is optimal as far as the ingredients are concerned, only the explicit constants might be improved. Both bounds are in $\lambda^{d/2}$, in agreement with Weyl's asymptotic. For the first eigenvalue λ_1 this sharpens considerably the preceedings "classical" inequalities of Lichnerowicz and Cheeger, see the very end of this chapter.

The proof is a function-symmetrization of the heat kernel in order to compare it with the heat kernel of the comparison space of constant curvature. On a simple function, this technique was introduced in (Faber, 1923) and (Krahn, 1924). So that in fact the result of (Bérard, Besson & Gallot, 1985) yields much more, since it controls $K(x,x,t)$ at every point and for all t . Another basic tool in the proof is the control of isoperimetric inequalities in a Riemannian manifold, which was achieved also in (Bérard, Besson & Gallot, 1985), controlling the so-called "isoperimetric profile" with the help only of a lower bound for the Ricci curvature and the diameter. These ingredients are optimal, only the best constants are still to be found. For more on the isoperimetric profile, see **TOP. 1 . B** .

In the spirit of chapter I the work (Gallot, 1987) improved the preceeding results on the eigenvalues by replacing uniform lower bound on Ricci curvature by integral norms. The basic tool is still the isoperimetric profile and the work consists in extending the control of it with only integral norms on Ricci, see TOP. 1 .

The link with periodic geodesics came first in (Colin de Verdière, 1973). One uses here the heat equation with complex time, which is Schrödinger equation in disguise. But the more simple and essential wave equation explanation came just after in (Chazarain, 1974). Immediately after came (Duistermaat & Guillemin, 1975). The first essential link is : the distribution

$$\sum_{k \geq 0} \exp(+_{-} i \sqrt{\lambda_k} t) \text{ (this is happily never a converging series,}$$

unlike $\sum_k \exp(-\lambda_k t)$) has a **singular support** contained in the

union of the lengths of the periodic geodesics (including 0). A very caricatural heuristic explanation is that the waves emanating from a periodic geodesic and with a good (bad) frequency in $2\pi N/\text{length}$ will give birth to some kind of resonance (a tidal wave).

This is a one direction result. In the special case of space forms, one gets much more with Selberg's trace formula : see (Iwaniec, 1975) and (Hehjal, 1976) . But in general it is only in dimension 2 that results are very strong. The reason is that one needs to know not only the length of the periodic geodesics, but what is happening when making one turn around it. This involves two things : the Poincaré return map and the holonomy (parallel transport). Recently progress in higher dimensions was done in (Guillemin, 1993), see also (Guillemin, 1996). There it is proved that the spectrum completely determines the symplectic invariants of the Birkhoff canonical form for the Poincaré map.

There are many links between the spectrum (for functions) and the length spectrum (the set of lengths of all periodic geodesics). The belief, guessed in (Balian & Bloch, 1972), is that the oscillations in the eigenvalue density in the spectrum is related in some sense to the length spectrum. The jumps can be understood, for example, roughly as deviations from the Weyl's asymptotic law.

An interesting link between the spectrum and the periodic geodesics is that of the **quasi-modes**. The story started in (Babich & Lazutkin, 1967) and is far from being finished today and remains quite mysterious, see (Colin de Verdière, 1977). Briefly speaking, to one given periodic geodesic (satisfying certain conditions) one can associate a series of numbers which are approaching quite a lot of eigenvalues. The idea of the proof is to build up approximative solutions of the wave equation which will propagate along the geodesic. A natural question is : are there many cases for which one can in that way obtain the whole spectrum ? The answer is that is exceptional and happens only when the geodesic flow is integrable. In general the hyperbolic zones between the KAM tori will yield a contradiction. The entire book (Lazutkin, 1993) is devoted to this topic.

Those were **direct** problems : I know the manifolds and some of its invariants. What can I say about the spectrum ? **Inverse** problems are : I know various things about the spectrum, what can I recover of the metric ? This can also be seen as a "recognition" problem. The book (Andersson & Lapidus, 1997) can be used as a survey.

The first question is the **uniqueness** : are two isospectral manifolds necessarily isometric ? The first counter-example came in (Milnor, 1964) : they are two flat tori of dimension 16, quotients of $\mathbf{R}^{16}$ by two lattices famous in Number Theory : E_{16} and $E_8 + E_8$. Another topic : is the set of isospectral metrics compact ? Are generic Riemannian manifolds determined by their spectrum ? For an eventual compactness of isospectral sets there are today only partial results in dimensions 2, 3, 4 : (Osgood, Phillips, & Sarnak, 1989), (Anderson, 1991b) and (Brooks, Perry, & Petersen, 1994). For topological finiteness of isospectral sets, see (Brooks, Perry, & Petersen, 1990) and (Brooks, Perry, & Petersen, 1992). One

introduces the ζ - function : $\zeta(s) = \sum_{\lambda \neq 0} \lambda^{-s}$ and one can manage by

analysis to define its derivative $\zeta'(0)$ at the origine. This value is linked to a suitably defined **determinant** of the spectrum which it

is formally defined as the infinite product $\prod_i \lambda_i$. Note that the

determinant is studied as a functional for trying to obtain extremal metrics in (Sarnak, 1997a), see **III. A.**

The curvature nature of the asymptotic expansion of the heat kernel is also of basic use.

The generic situation is completely open today, except within the very restricted class of Riemann surfaces. In dimension 2 the conjecture is the finiteness of isospectral sets : (Sarnak, 1997c). Uniqueness is specially irritating, one can prove it today only for $\mathbb{RP}^2$, the 2-dimensional flat tori and the spheres up to dimension 6 : (Tanno, 1980), the proof uses only Minakshisundaram's asymptotic expansion of the heat kernel.

For isospectrality the results extend from 1964 to still strongly going on research, mostly looking to get more and more examples of less and less geometrically special isospectral manifolds, and even one-parameter isospectral deformations. A survey is (Bérard, 1989). Important intermediate texts were (Vignéras, 1980b), (Sunada, 1985). See (Gordon & Mao, 1994b), (Gordon, 1994c), (Pesce, 1992) and the references there. A major question remains : are generic Riemannian manifolds spectrally isolated (solitary) ?

For special cases one has finiteness or non deformability results : for Riemann surfaces see the chapter 13 of (Buser, 1992), (Buser, 1996) for "solitary surfaces" and for negatively curved surfaces (Guillemin & Kazhdan, 1980a). This last text puts together the fact seen above that the spectrum determines the length of periodic geodesics and the fact that negative curvature implies a lot of well distributed periodic geodesics. Then one uses a Fourier analysis of the derivative of the metric as a function on the unit tangent bundle. This beautiful technique seems not to have been much used since.

This is the place to remark that the wave equation technique used in these results is mixing both **microlocal analysis** (which is essentially Analysis but in the tangent (phase) bundle) and **symplectic geometry**. The symplectic structure properly said is on the cotangent bundle T^*M , on UM one has only what is called a contact structure and more precisely a **Sasakian** structure, see page 56 of (Sakai, 1996).

Another important and natural question is to know if the ergodicity (V. B) of the geodesic flow implies or not a very good regularity for both the spectrum and the eigenfunctions. First : is the spectrum very well distributed ? For simplicity sake let us look us only in dimension 2 because then Weyl's asymptotic is linear : here $d/2 = 2/2 = 1$. One then searches for a probabilistic result on the

variance of the quantities $|N(\lambda+L) - N(\lambda-L)|$, suitably averaged in λ and L .

It was conjectured in (Pandey, Bohigas, & Giannoni, 1984) that the spectrum is that of a GOE (i.e. Gaussian random symmetric matrix) when the geodesic flow is ergodic. Today there is no theoretical general answer. There are many numerical experiments by theoretical physicists, see also (Lazutkin, 1993). At the other end stands the integrable case, see (Sarnak, 1997b).

From some numerical experiments it seems that the distribution could be not GOE for some arithmetic Riemann surfaces. This was mathematically proved in (Luo & Sarnak, 1996). However the geodesic flow is ergodic. So what is today belief is that still the distribution will be GOE for generic Riemann surfaces. To explain heuristically why arithmetic forms are exceptional, one remembers what was said above, namely that the jumps in the spectrum are linked with the structure of the length spectrum. But precisely the length spectrum of arithmetic forms is very "degenerate" in the sense that the lengths are given (after normalization) by integers. The asymptotic exponential behaviour (see V. A) forces then all these periodic geodesics to have very large multiplicities, hence huge jumps in the length spectrum . References are (Sarnak, 1995), (Luo & Sarnak , 1995a), (Luo & Sarnak, 1996), (Rudnick & Sarnak, 1996) and references there.

Another question : are the nodal sets (the set of points where the eigenfunction vanish) evenly distributed ? even more important : are the eigenfunctions evenly distributed ? This is proved today only for a subset of the eigenfunctions (but however of full density) : (Colin de Verdière, 1985) which follows ideas of (Schnirelman, 1974). See also (Zelditch, 1987) and (Zelditch, 1992) for the non compact case.

This is linked with the question of "scars" : in some numerical experiment people found that the nodal lines of some surfaces where, in some sense, accumulating along some periodic geodesics. But in (Sarnak, 1995) it is proved that this can never happen for arithmetic space forms (for some definition of what is a scar). This is an amusing paradox : the arithmetic case implies more regularity at the time it is a less common case (in the realm of space forms). What is the general state of affairs is still dividing experts, since

scarring is only today almost purely experimental and because the definition for scars varies according to authors . See (Rudnick & Sarnak, 1996), (Shimizu & Shudo, 1995) and the references there.

For the volume of the nodal sets (i.e. the set of zeroes of an eigenfunction) a conjecture of Yau in 1972 on the asymptotic behavior of their volume is partially solved in (Donnelly &

Fefferman, 1988). One has $c(g) \sqrt{\lambda} \leq \text{Vol}(\phi_\lambda^{-1}(0)) \leq c'(g) \sqrt{\lambda}$ for every eigenfunction ϕ_λ attached to the eigenvalue λ . The story is not finished because today one first needs analyticity and second the constants $c(g)$ and $c'(g)$ are "in the air", the time being one would like to have for them an explicit estimation based on Riemannian invariants. See also (Donnelly & Fefferman, 1990).

Another paradox is this : first, as one will see in the next chapter, the sole existence of periodic geodesics is an almost completely open question, at the time we have very good general results for the spectrum. But in the case of ergodic manifolds, specially for manifolds of negative curvature and even more specially for space forms, one has oppositely a very good control of the periodic geodesics : asymptotic expansion of the length counting function, regular distribution in space and even more in phase. In exchange for these space forms we just saw that both the eigenvalues and the eigenfunctions are not still completely understood. An open problem is the question of the multiplicity of these various lengths.

One can ask : why to be so much interested in the asymptotic properties and not to the people in the beginning ? There are more than one reason for that. The first is that, most often, only asymptotic properties are accessible, recall Colin de Verdière result above : any finite part of the reals can be realized as the beginning of the spectrum of some Riemannian manifold. In physics asymptotics are very significative in quantum theory. A basic question is to understand to which extent classical and quantum mechanics are linked. The hope is that the classical one is the limit of the quantum one when the Planck constant h goes to zero in the Schrödinger equation $h^2 \Delta f = i h \frac{\partial f}{\partial t}$. One sees that this amounts to study what is happening for large eigenvalues. The various studies connected with this type of questions are called **semi-classical analysis**.

A contrario it is also extremely important to have a good lower control on the first non zero eigenvalue λ_1 depending on the geometry. This for example in order to exclude some possible "resonances". The first inequality came with (Lichnerowicz, 1958) (see page 135) : from a positive lower bound on Ricci one deduces a lower bound for λ_1 which is optimal and with equality only for the standard sphere (Obata, 1962). For more references see (Croke, 1982) which has more, namely a pinching type theorem. Lichnerowicz proof was just to use the Bochner's formula given in O. F., but his bound appears in many contexts. A typical example for the geometer when studying distance functions, as in Colding's triangle theorem seen in section I. A. 1. Thereafter came (Cheeger, 1970b), which is looking for the isoperimetric profile (TOP. 1. B) at its middle value $h(\frac{1}{2})$: $\lambda_1 < \frac{1}{4} h^2(\frac{1}{2})$ and had a lot of impact. Now Bérard et alia inequality seen above is much stronger and this is not surprising in view of Cheeger's inequality since the author estimation is obtained with their bound for the whole isoperimetric profile $h(t)$.

The nature of the first eigenvalue for Riemann surfaces and their locations with respect to $\frac{1}{4}$ is fascinating and deep, see a description in (Buser, 1992).

V : Periodic geodesics, the geodesic flow

A. Periodic geodesics

Here comes one, if not the, of the most irritating unsolved question for Riemannian manifolds (except in the negative curvature case). A survey is (Bangert, 1985) and there is the book (Klingenberg, 1978a) (the lemma 4.3.4 contains gaps as well as other statements, but there is a Russian translation with critics). See also (Klingenberg, 1982). Note that many authors use the word "closed" instead of periodic, but this can be misleading when one thinks of geodesic loops.

First let us state what seems reasonable to ask. One would expect first the existence of infinitely many periodic geodesics on any compact manifold. This means "geometrically distinct", one does

not consider as geometrically different two periodic geodesics which are both an iteration of a common periodic one, hence having the same geometric support (for the physicist they will be different, as having different lengths then different energy levels). Second, assuming such an existence, what is the growth of the counting length function ? The **counting function** $CF(L)$ is the number of periodic geodesics of length smaller than L . One can expect different results depending on the geometry of the manifold. Note that if the counting function has exponential growth the iterates will not appear because giving only linear contributions in the counting function.

Before giving the positive existing results, the fact is that today one does not know that there is more than one periodic geodesic for every manifold of dimension larger than 2 (this starts with S^3). So that the counting function question does not even make sense. Also one thing is sure since (Weinstein, 1970) : in the general case the periodic geodesics are not dense in the phase space. But they still could always be dense in the manifold itself, this is open.

Two points to help the reader realizing the difficulty of the problem and that one cannot work purely with topology, i.e. the metric enters essentially in the game. First, the following surprising theorem in (Morse, 1934) shows how weird can be the set of periodic geodesics in an apparently very nice manifold. Given any length L (think of it as very large) there exists an $\epsilon(L) > 0$ with the following property. Assume an ellipsoid in $\mathbf{R}^d$ has all its principal axis $a_1 < a_2 < \dots < a_d$ satisfying $|1 - a_i| < \epsilon(L)$, then the only periodic geodesics of length smaller than L are the $d(d-1)/2$ intersections with the 2-planes of coordinates. The proof given in (Klingenberg, 1982), lemma 3. 4. 7, is just a subtle play with the Sturm-Liouville type equations for the coordinates of a geodesic. In some sense, except a finite number of them, all the other geodesics "disappear" when the ellipsoid approaches more and more the sphere : they are "ghosts". It implies that it can be expensive to find them. Note also the paradox for the limit case, the standard sphere.

Second, an example, due to Katok (see it in (Ziller, 1982) with many generalizations), exhibits a Finsler (today still non symmetric) metric on S^2 with only two periodic geodesics. But the free loop space structure is basically the same for Riemannian and for Finsler metrics. For Finsler metrics see **TOP. 9**.

The initial positive fact is that, as soon as the manifold is compact, in any non zero free homotopy class (they are the conjugacy classes of the fundamental group) there exists always at least one periodic geodesic : just take only the shortest curve of the class (see O.F). So that there is (again) a radical difference between the simply connected case and at the extreme opposite side the **negative curvature** one. Then we have immediately a huge number of periodic geodesics since π_1 is huge. And in the case of space forms of negative curvature (those of rank 1) results are beautiful. Back in (Katok, 1988) and thanks to conformal representation, it was proved that **any** metric on such a negative curvature surface has as many or more periodic geodesics than the constant curvature one. More precisely the counting function is **exponential**, with an exponent which is larger than for the constant curvature case, unless we are precisely in this case. One can paraphrase this result by saying that these spaces are very susceptible animals, reacting very strongly : any time you touch them, even as slightly as possible, they develop exponentially more periodic geodesics. Behind these statements is inevitably "hidden" the various notions of **entropy**, see the digression in **B** below.

In (Besson, Courtois & Gallot 1995a) this result is generalized to any dimension. But today negativity of the curvature is still required. Things work for rank 1 symmetric spaces and their products but not today for higher ranks. This means as much susceptibility as described above. See in **B** where the proof is coming from.

The way the periodic geodesics are distributed in those manifolds of negative curvature is subtle but more or less understood, see (Eberlein, Hamenstädt, & Schroeder, 1993) and the section 3 of (Besse, 1994). One fact is that even distribution for the Liouville measure characterizes space forms. See also for this and the next section the expository (Pansu, 1990/91), also chapter 20 of the quite complete (Katok & Hasselblatt, 1995).

As soon as one leaves the negative curvature realm, things are not understood (except what is said in **I. B. 4**), for example the case of various metrics on tori seems intractable at the moment (see however (Bangert, 1988)). The fundamental group via its conjugacy classes will easily yield a counting function which is polynomial of degree equal to the dimension, but it might well be that the

counting function is always exponential (at least for dimensions larger than 2).

A digression : geodesics joining two points. We comment here in an indirect way on the difficulty to obtain periodic geodesics. Morse's theorem above on ellipsoids was one remark. To feel the other difficulties let us look first before at the collection of geodesics joining two given points p and q . Morse theory is basic for that. In (Serre, 1951) algebraic topology and Morse theory yielded an infinite number of geodesics between any pair of points of compact manifold. They might be all covering a periodic one like for the standard sphere, but this situation is exceptional (if it is true for any pair of points, it probably implies that we are in a manifold all of whose geodesics are periodic, see below for this). In (Morse, 1934) this conclusion was obtained only for topological spheres.

Briefly : between two given points Morse theory provides at least as many geodesics of index equal to k as the k^{th} -Betti number $b_k(\Omega)$ of the space Ω of all curves in the manifold joining these two points. The **index** is the number of conjugate points between the two extremities but it does not matter here. Now Serre's contribution was that for any compact manifold there is an infinite number of non zero $b_k(\Omega)$. A minor technical point here is that Morse theory needs the two points to be not conjugate (this along any geodesic joining them). Two points are said to be conjugate along a geodesic joining them when there is a non zero Jacobi field vanishing at both ends. This does not matter at the end since, given any point, the set of its conjugate points is of measure zero. But in the case of periodic geodesics the difficulty is inevitable and a major one besides the other ones to be explained below.

Passing by we quote a very nice recent result which says that the double integral of the counting function is exponential with an exponent equal to the (topological) entropy of the geodesic flow : (Mané, 1996) and (Paternain & Paternain, 1994). Entropies will be defined below in section B . Define the **geodesic entropy** h_{geod}

$$\text{as} \quad h_{\text{geod}}(M,g) = \lim_{L \rightarrow \infty} \frac{1}{L} \int_{M \times M} \log(CF(p,q;L)) \, dp \, dq$$

where the counting function $CF(p,q,L)$ is the number of geodesics joining p to q and of length smaller than L . Then both Mané and the Paternain proved that $h_{\text{geod}} \leq h_{\text{top}}$. The other inequality $h_{\text{geod}} \geq h_{\text{top}}$ is a particular case of a more general inequality for smooth flows in (Przytycki, 1980). The proof of the first inequality is

very Riemannian, working on the product manifold $M \times M$. There the set of geodesics joining two points in M becomes the set of geodesics initiating orthogonally from the diagonal. To obtain this : start from the middle point of the geodesic and go equal length on both directions. This "naive" idea was already used in (Grove & Petersen, 1988). To get the entropy at the end one uses a hard theorem of (Yomdin, 1987)(see also (Gromov, 1987b)) where the computation of the entropy uses differentiability. Note that h_{geod} is for the Riemannian geometer a definition of the entropy of an absolute simplicity. It is not cleared today if, possibly in some cases like negative curvature or genericity, one can just take the same definition but with only two fixed points (no mean value to perform). Or at least for almost all pair of points. This is certainly not true in general because in (Burns & Paternain, 1996b) one finds on S^2 a large set of metrics for which the strict inequality

$$\limsup_{L \rightarrow \infty} \frac{1}{L} \log(\text{CF}(p,q;L)) < h_{\text{top}}$$

holds for an open set of pairs (p,q) .

For obtaining periodic geodesics "à la Morse" in following the scheme above one has to replace the space of curves joining two points by the space of all closed curved in the manifold. Serre's theory still formally applies but there are two difficulties. The first concerns the "non-degeneracy" of a periodic geodesic which here plays the role of conjugacy for pairs of points. Even if overcome there is the fact that this infinite number of periodic geodesics so obtained might be only the coverings of a finite number of them! The second difficulty is that the space of maps from the circle S^1 to a manifold is much more subtle that of the maps from $[0,1]$. The job to do in algebraic topology is already treacherous. Many mathematicians wrote wrong statements concerning the topology of the space of closed curves in a manifold. Moreover, even a perfect knowledge of this topology will not be enough, in view of the Morse and Katok examples, since the fact to be "only" in a Finsler manifold does not make any difference at the level of the topology of the space of closed curves.

So that the job is now to look more carefully **simultaneously** at Morse theory and at the geometry of the manifold. Indices here matter a lot. One should **decouple** indices and homology classes, looking at the behavior of the indices of the various iterates of

periodic geodesics. The basic tool here is from (Bott, 1956) : the index of the iterates of a given periodic geodesic grows linearly plus or minus a fixed constant. Here is now what one can be done today.

The case of surfaces is not too bad, the problem is for the sphere, the only simply connected compact surface and was raised in (Poincaré, 1905). Despite his efforts he could not show the existence of at least one periodic geodesic. Birkhoff got it in 1917 and this made him famous overnight. He achieved that by a minimax principle applied to "tapestries" of the sphere and reducing this infinite dimensional problem by approximating curves by broken geodesics (using the injectivity radius). This is now a finite-dimensional story and results can be obtained with the help of classical compactness results. This was the birth of Morse theory. Note that the Birkhoff geodesic is not necessarily **simple** (i.e. without self-intersection), see (Calabi & Cao, 1992a).

In (Lusternik & Schnirelman, 1929) was claimed the existence of at least three simple ones. The topic is very treacherous. Things seem to be OK now after many incomplete proofs, see (Taimanov, 1993) and the analysis there. It is fair to say that the "intermediate" results were bringing true and important facts. Note in recent works the use of a deformation technique involving Analysis (see also O.F). The idea was first used in the plane in (Gage & Hamilton, 1986) . Deforming a plane curve by normal variations which are proportional to the curvature makes the curve becoming more and more close to a circle ("the" curve of constant curvature). In a Riemannian manifold the curve will (hopefully, this has to be proved) have as a limit a curve of zero geodesic curvature, i.e. a geodesic. The equation is a parabolic PDE for which the existence of solutions for small time solutions is quite easy, most of the job is to show first the existence for any time and second a good geometric behavior of the limit curve. The best proof today is (Grayson, 1989).

The big event was the conjunction of (Bangert, 1993) with (Franks, 1992) yielding at least an infinite number of periodic geodesics. This made the joyce of popular writers : "a million rubber bands around a potato". Precised in (Hingston, 1993) one knows even that the counting function grows at least as a constant times $L / \log L$. Still there is not a single explicit surface for which the asymptotic order of the counting function is exactly known. No expert seems to even have any guess, like : is it always polynomial, exponential, or exponential generically ? Note that you discard the question when a

manifold, e.g. surfaces of revolution and ellipsoids, possesses a one-parameter continuous family of periodic geodesics (then necessarily locally of same length).

For higher dimension, one has to wait (Lusternik & Fet, 1951) just to get one periodic geodesic. The proof is in Birkhoff spirit. There is no higher dimension generalization of Lusternik-Schnirelman result today, only with the strong $\frac{1}{4}$ -pinching extra hypothesis, see (Ballmann, Thorbersson, & Ziller, 1983) and do not forget Morse's theorem above. Thereafter came (Gromoll & Meyer, 1969a) : one is sure to get infinitely many periodic geodesics as soon as the Betti numbers of the space of closed curves of our manifold are not bounded. The beautiful idea is to use the fact that the index of the iterates of periodic geodesic grows only very close to linearly when iterating it. By contradiction a finite number only of periodic geodesics will then forces the Betti numbers to be bounded. But, as explained above, one has to handle the case of degenerate geodesics. The authors succeeded in doing it by a technical "tour de force".

This leaves us with the topology problem to find when this Betti number condition is true. The story is still not completely finished today but it is almost true that the only exceptions are the positive curvature space forms, the spheres and the KP^n . What a paradox : these manifolds have all their geodesics periodic for their canonical metric! So we are left with the first example : on the three dimensional sphere one still does not know that there is, for any metric, always more than one periodic geodesic. For periodic geodesics of manifolds having the rational homotopy type of spheres or the KP^n , see (Rademacher, 1994b).

In fact a real understanding of the situation needs more technicalities. One knows that, as soon as a periodic geodesic is of the so-called "twist type", in any tubular neighborhood of it there is an infinity of periodic geodesics, this is due already to Birkhoff. So one of the main questions is perhaps not about an infinite number of geodesics but : does there exist on S^2 a metric all of whose geodesics are hyperbolic ? The same question in fact for every simply connected manifold in any dimension, and one can also demand this for only an open set of such metrics. Hyperbolicity for a periodic geodesic means that the Poincaré return map (met also in IV) has no eigenvalues of modulus equal to one.

In despair of a general statement we look for the **generic** situation (on any manifold). Recently it was proved in (Rademacher, 1994a) that one can decouple the indices of periodic geodesics by small perturbations of the metric. Then one knows that, at least in the **generic** case, there exist always infinitely many periodic geodesics. This result uses intermediate ones on generic metrics (called "bumpy") : (Klingenberg & Takens, 1972) and (Anosov, 1983).

Moreover the proof yields more. If one adds the validity of the conjecture above on the haunting KP^n and puts together Gromoll-Meyer, Hingston, Rademacher and Moser results then the counting function grows at least as a constant times $L / \log L$ in the generic case. This $L/\log L$, as well as the one above for surfaces, is not mysterious. It comes from the fact that one has to look finally at prime numbers in a given arithmetic progression. Dirichlet's theorem gives precisely such an order of magnitude. But nobody seems to have any guess if always (or in the generic case) things grows more, say like a polynomial of a degree possibly related to the dimension, or even exponentially, etc.

B. The geodesic flow (geometry and dynamics)

The **geodesic flow** is the set made up by the collection of the G^t (where t is any real number) which act on the unit tangent bundle UM by walking along geodesics a length equal to t . Explicitly : $G^t(v)$ is the speed vector $\gamma_v'(t)$ at the point $\gamma_v(t)$ of the geodesic γ_v whose initial speed vector at 0 is $v = \gamma_v'(0)$. In all this study basic is the fact (Liouville theorem) that the flow respects the canonical contact structure of UM , hence the attached measure. For the importance of contact structure in Mechanics, see (Arnold, 1988), (Arnold, 1978). The symplectic structure is on the cotangent bundle T^*M (isomorphic to TM for Riemannian manifolds). So that we have a natural and geometrical dynamical system in Riemannian geometry. The inevitable invariant is the **entropy**. Two things are difficult with it. First the definition is never very short, second they are more than on entropy, see the digression at the end of this section and the general references given there.

Problems are direct or inverse (recognition). Direct : knowing things for the manifold, what can we say about its geodesic flow ? Inverse problems : assume the geodesic flow is known, what can be said about the geometry of the manifold ? In particular there is the uniqueness problem : if two manifolds M and N have the same

geodesic flow (the precise definition leads rather to the "conjugacy of geodesic flows") are they then isometric ? A partial survey is section 3 of (Besse, 1994) and very detailed is (Eberlein, Hamenstädt & Schroeder, 1993). **Conjugacy** means the existence of a map $UM \rightarrow UN$ commuting for every t with the two geodesic flows $G_M^t: UM \rightarrow UM$ and $G_N^t: UN \rightarrow UN$. Commutation with the projections $UM \rightarrow M$ and $UN \rightarrow N$ is not demanded and would be too strong to be of interest. This means that you are interested in your plane trajectory in itself but not asked to look by the window above which point of the earth you are.

(Hopf, 1939) was the first direct result : the geodesic flow of a compact surface of negative curvature is **ergodic** (this is Eberhard Hopf, not Heinz Hopf). Ergodicity means for a dynamical system that almost every trajectory is everywhere dense. It is equivalent (thanks to Birkhoff main theorem) to the extremely practical and useful definition : "space and time averages coincide". For entropies see definitions at the end of this section and the references there. Much weaker is topological transitivity : there exists at least one trajectory which is everywhere dense.

In fact Hopf's result had a precursor in (Morse, 1921a), (Morse, 1921b) where first topological transitivity was obtained and better: a coding for the trajectories. **Coding** is extremely important, it enables people to study the geodesic flow by looking only at the model called the "discrete "shift". We mention only some recent references : (Parry & Pollicott, 1990), (Katok & Hasselblatt, 1995), (Katok, 1996). H. Hopf's result was extended in any dimension in (Anosov, 1967) where it is proved moreover that the geodesic flow is stable (one can trace this type of question about "structural stability" to (Hadamard, 1901)). Anosov's proof is quite involved even today. The best exposition is in the appendix by Brin in (Ballmann, 1995a).

In (Katok, 1982) for surfaces and in (Besson, Courtois & Gallot, 1995a) for any dimension there is a very strong statement : when one looks at various metrics on a space form locally symmetric of negative curvature (i.e. of rank 1), the topological entropy of the geodesic flow is a minimum for and only for the locally symmetric metric. This strong result has many corollaries, we saw in **III** one of them in the case of Einstein metrics. Another one is for the inverse problem of the counting function, see **A** above. These manifolds can also be recognized by counting their periodic geodesics (at least in the negative curvature case). As seen above in

A one can present this as a "susceptibility" result.

We will come back in detail below to that result. We first address two questions stemming out from Hopf and Anosov results. First : is really negative curvature (or nonpositive one) needed to get ergodicity ? Does one really need a manifold with a large fundamental group ? Recent results seem to say : no. We know now that there exists ergodic metrics on S^2 (Donnay, 1988) but it is still open if one can do it with positive curvature (a convex surface). They are real analytic convex surfaces (close to ellipsoids) with positive topological entropy : (Paternain, 1993b). More generally there are C^2 -deformations of ellipsoids in any dimension with positive topological entropy and exponential growth for the counting functions : (Petroll, 1996) , see also (Knieper & Weiss, 1994). And more : any compact manifold has some ergodic metric according to Lohkamp (unpublished yet). For what means "chaos" see the end of the section. Things are open for the measure (metric, Liouville) entropy. It seems that the main question (see also A above) is to know if S^2 can or not carry a metric all of whose geodesics are hyperbolic. This never happen for a $\frac{1}{4}$ -pinching : (Ballmann, Thorbergsson, & Ziller, 1982). This implies that one will have to go quite far to find such a metric.

The second question is an inverse (recognition) problem in the spirit of IV . Does the length spectrum (the set of the length of all periodic geodesics) determines the Riemannian manifold ? In all generality certainly not . Because we have counterexamples of isospectral surfaces and for surfaces the knowledge of the length spectrum and the knowledge of the ordinary spectrum are equivalent. The situation is radically different for the **marked length** spectrum. This means one "remembers" for lengths which free homotopy class they are coming from. And for manifolds on nonpositive curvature the equal marked length spectrum condition is equivalent of conjugacy of geodesic flow. Then many partial but strong results are available, in particular (Otal, 1990a). We refer to section (9.2) of (Eberlein, Hamenstädt & Schroeder, 1993) for more details.

Let us be now more precise on (Besson, 1996b) and (Besson, Courtois & Gallot, 1995a). In the various statements we will always assume the volume is normalized. The authors address the more general problem of maps of non zero degree but we simplify

things by working on a given manifold (namely we pick up the identity map). Now let (M, g_0) be a compact space form of negative curvature (hence locally symmetric of rank 1 by definition). The theorem is that for any other metric g on M the volume entropy $h_{\text{vol}}(g)$ satisfies $h_{\text{vol}}(g) \geq h_{\text{vol}}(g_0)$ and equality only if $g = g_0$. Since always $h_{\text{top}} \geq h_{\text{vol}}$ with equality for nonpositive curvature, the same result is valid for h_{top} .

The very definition of the volume entropy and Bishop's theorem **TOP. 1. A** imply immediately the minimal volume result (see **TOP. 1. 4**) in the case of constant sectional curvature. The result on counting periodic geodesics comes from the estimation of their counting function with the topological entropy and the fact that $h_{\text{vol}} = h_{\text{top}}$ in the nonpositive curvature case. The result on Einstein 4-manifolds (see **II. B**) comes from the formula in **O. E** for the characteristic and see the original text for other corollaries.

The more natural and conceptual proof would be to prove that the entropy is a strictly convex functional on the space of all Riemannian metrics and that the locally symmetric metric is a (the) critical point. Unhappily this program does not work as such except in a conformal class, see (Robert, 1994). The original proof was quite involved but using nice techniques, like the center of mass for the structure of the sphere at infinity (see **I. B. 4**) and the technique of calibration (see **TOP. 3. B**). In (Besson, Courtois & Gallot, 1995b) the proof is greatly simplified by introducing the Patterson-Sullivan measures on the sphere at infinity : with them it is easier to work on center of mass. Finally the result yields an extremely simple proof of Mostow theorem (see **II.A** and **B**) for the case of rank 1 . For an expository text, see (Pansu, 1996/97).

The theorem also shows that conjugacy of the geodesic flow to that of a space form implies isometry : one can recognize the compact space forms of rank one (and of negative type) by their geodesic flow. This is now the place to mention another recognition procedure for those forms among negatively curved manifolds.

In Anosov main result basic in the proof are the stable and unstable foliations. For the Riemannian geometer they are the vector fields normal to a family of spheres centered at the same point at infinity. Equivalently one can prefer to look at Jacobi fields which are 0 at infinity when given an initial condition : the stable one when 0 is at $+\infty$ and unstable when 0 is at $-\infty$. Those foliations are always defined for negative curvature and are of course smooth for space forms. Historically they appeared first in

(Hadamard, 1901). But smoothness when one looks from infinity is known to be a difficult game in Analysis. And this is dramatically seen now : if the curvature is negative then smoothness of the Anosov foliations implies that we are metrically on a space form. This is a very strong recognition result, just by smoothness. After intermediate results (see (Eberlein, Hamenstädt & Schroeder, 1993)) this is now obtained by putting together (Besson, Courtois & Gallot, 1995a) and (Benoist, Foulon, & Labourie, 1992).

Back to the geodesic flow, in particular for tori, there were partial results, see e.g. (Croke, 1992). We mention the recent proof of a long standing conjecture stemming from the result in (Hopf, 1948) in dimension 2, to the effect that a metric without conjugate points on a torus (of any dimension) has to be flat : (Burago & Ivanov, 1994). We also recall here the strong results of I. B. 4 which study the hard problem to distinguish between manifolds of negative curvature and those of nonpositive one, see also Finsler spaces in TOP. 9.

In all the above results the proofs mix more or less classical results of ergodic theory with various geometric techniques. The notion of Busemann function was already mentioned. Variations of them can be caricaturally described as follows : one looks at the universal covering of the manifold of nonpositive curvature under consideration. It looks topologically like $\mathbf{R}^d$. One defines on it various kinds of structures "at infinity" and look at the metric "from infinity". We already met this technique in section I. B. 4.

The extreme opposite case is that of space forms of positive curvature, the spheres and the $\mathbf{K}P^n$. They have a beautiful geodesic flow, since all the geodesics are periodic, simple and of the same length (this is understood for their canonical metric): see II. C. But the inverse problem is almost completely open. The spheres and the $\mathbf{K}P^n$ behave very differently. Back in (Zoll, 1903) one finds on the two-dimensional sphere many examples of metrics all of whose geodesics are periodic (understand : non isometric to the standard metric). They are completely classified only : for surfaces of revolution (Besse, 1978), theorem 4.13 and for metrics closed enough to the standard one (Besse, 1978), section 4.H or (Guillemin, 1976). For higher dimensional spheres a very partial technical result is (Kiyohara, 1984). But the classification is untouched.

For the reader who might underestimate the difficulty of the subject, we mention few facts. For more and references see Besse's

book. First, if it is trivial to show that the lengths are all multiple of a smallest one, it is true but hard to show that the lengths are finally bounded. Worse : if they are plenty of examples for which the lengths are different (not of the iterates of course), e.g. lens spaces and more generally any space form of curvature 1. But is open to know if in the simply connected case all the lengths are equal. There is also the question of simple or self-intersecting periodic geodesics. The only result we know of is on S^2 : (Gromoll & Grove, 1981) shows that then in fact all the lengths are equal and the geodesics all simple. But the proof is very two-dimensional and uses Lusternik-Schnirelman three geodesic result above.

Oppositely there are no known metric on any KP^n all of whose geodesics are periodic (except of course the standard one, see II. C). The case $K = \mathbb{R}$ is solved, only standard metrics have that property : it is a direct corollary (look at the universal covering) of the result on spheres all of whose cut-loci are reduced to a point to be seen in TOP. 4. For the other K , we have only a partial result, i.e. this is true for metrics close enough to the standard one : (Tsukamoto, 1981). This is used in (Kiyohara, 1987) to prove the spectral isolation (solitaryness) of the KP^n : the link with the spectrum is insured by the results quoted above between periodic geodesics and the spectrum. Other results of this type "infinitesimal rigidity" are in (Gasqui & Goldschmidt, 1994) and various texts by the same authors, past and future.

In the field "geometry and dynamics" a special mention should be given to (d'Ambra & Gromov, 1991), see also TOP. 9. Some geodesic flows are very special, they are called **integrable**. This is the case of surface of revolutions as well as of ellipsoids. For more on them, see (Spatzier, 1990).

Digression : the entropies in Riemannian geometry. Entropy is a difficult concept . Heuristically not : it is nothing but the measurement of the exponential factor of anything you want to compute in a dynamical system : the loss of information when the time goes on, the dispersion (divergence) of trajectories, etc. Strict definitions are less simple. Moreover there is more than one entropy. Two are standard : the metric (measure theoretic) and the topological ones. Up to quite recently the labeling was ridiculous : metric entropy needs only a measure to be defined and is really a measure notion, topological entropy needs a metric (even if finally

is independent of it) and is really a topological notion. The third entropy (volume entropy) makes sense only for manifolds whose universal covering is "huge". Metric (measure theoretic) entropy was historically the first. General references are : up to 1982 (Walters, 1982) was the standard and efficient book, now there is (Katok & Hasselblatt, 1995). See (Sinai, 1976) for its very informative style. Besides (Katok & Hasselblatt, 1995), and (Mané, 1987) specially for the smooth case. For result references we refer mainly to the bibliographies there.

Entropy is defined for **discrete** dynamical systems $f : X \rightarrow X$ where X is a compact space and f some homeomorphism. In case of the geodesic flow one takes $X = UM$ to be the unit tangent bundle and f to be $f = G^1$ (the geodesic flow for the length equal to 1). It does not matter if the "length" 1 looks special because in fact one is interested only to what is happening for large iterates f^k and this means essentially that $t \rightarrow \infty$.

The simplest one to define is the **volume entropy** in compact Riemannian manifolds (M, g) . Let M^* be its universal covering. We look at the metric balls $B(p, R)$ in M^* and set :

$$h_{\text{vol}}(g) = \lim_{R \rightarrow \infty} \frac{1}{R} \log (\text{Vol}(B(p, R)))$$

which is easily seen to exist and to be independent of the base point $p \in M^*$. Unhappily as such it is of no interest unless M^* (that is $\pi_1(M)$) is "huge". For "smaller" fundamental groups and a polynomial renormalization, see (Babenko, 1992).

For a general dynamical system $f : X \rightarrow X$ put any metric on X and look at the iterates f^k of f . One gets new metrics d_n for every integer n by setting $d_n(x, y) = \sup_{0 \leq k \leq n} d(f^k(x), f^k(y))$. Let now $N_n(\epsilon)$ be the minimum number of ϵ -balls in the metric d_n needed to cover X and define the **topological entropy** to be :

$$h_{\text{top}}(X; f) = \lim_{\epsilon \rightarrow 0} \lim_{n \rightarrow \infty} \frac{1}{n} \log (N_n(\epsilon))$$

It is easy to check that it exists and does not depend on the metric chosen on the compact manifold under consideration.

Finally the **metric entropy** (more and more called now **measure theoretic** or just **measure**) was the first to be defined historically and by a mixing definition but quite lengthy and moreover almost

intractable to compute explicitly in given situations. We still use an auxiliary metric which can be forgotten at the end. This time we define $N_n(\epsilon, \delta)$ to be the minimum number of ϵ -balls (still for the metric d_n) needed to cover some subset of X whose complement has measure less than δ . Then one checks that the following definition makes sense :

$$h_{\text{met}}(X;f) = \lim_{\delta \rightarrow 0} \lim_{\epsilon \rightarrow 0} \lim_{n \rightarrow \infty} \frac{1}{n} \log(N_n(\epsilon, \delta))$$

These entropies enjoy many inequalities and properties in various instances. First : always $h_{\text{vol}} \leq h_{\text{top}}$ and equality as soon as the curvature is nonpositive : (Manning, 1979) and references there for intermediate results. The link with periodic geodesics is (see (Margulis, 1969)) : when the curvature is negative then

$$h_{\text{top}} = \lim_{L \rightarrow \infty} \frac{1}{L} \log(\text{CF}(L))$$

Do not think that positive entropy implies ergodicity, even locally, it only implies "some kind of local chaos". Metric (measure) entropy is the strongest invariant. For example positive metric entropy in a Riemannian manifold implies the existence of a set of positive measure where a geodesic is everywhere dense. This is definitively not the case for the topological entropy. In Riemannian geometry for the geodesic flow one also calls the metric (measure) entropy the **Liouville entropy** : $h_{\text{Liouville}}$.

We end with a formula giving explicitly a Riemannian quantitative information on divergence. This is the formula of (Ballmann & Wojtkowski, 1989) for the geodesic flow of compact manifolds of nonpositive curvature :

$$h_{\text{Liouville}} \geq \int_{\text{UM}} \text{trace}(\sqrt{-R_v}) \, dv$$

where the integral is on unit vectors v and R_v denotes the linear map $u \rightarrow R(v,u)v$ from the curvature tensor. The trace is taken with respect to the Riemannian metric. Moreover equality holds only for locally symmetric spaces. This formula has been preceded by weaker ones, see references in (7.3) of (Eberlein, Hamenstädt & Schroeder, 1993). In (Foulon, 1997) there is a generalization of the above formula for Finsler spaces (TOP. 9).

TOP. SOME OTHER RIEMANNIAN GEOMETRY TOPICS OF INTEREST

1. Volumes

A. BISHOP'S THEOREM

In (Bishop, 1963) one studies manifolds M with $\text{Ricci} \geq (d-1)k$ where k can be of any sign or zero. Then it is proved that the volume of any ball $B(p, R)$ of radius R is bounded by the volume of "the" ball $B^*(R, k)$ in the simply connected space form $M^*(k)$ of constant curvature equal to k and of same dimension. This was stated for balls which are within a coordinate ball. More strong was the statement : the ratio $\text{Vol}(B(p, R))/\text{Vol}(B^*(R, k))$ is nonincreasing with R . The original proof was in the spirit of RCT. Modern proofs are more simple, using the distance function "à la Gromov", see (Eschenburg, 1994) for a very detailed and nice proof. Despite its simplicity, the importance of Bishop's theorem should not be underestimated. We saw above many applications of it, specially amplified with Gromov's next double remark.

In (Gromov, Lafontaine, & Pansu, 1981a) it is proved (even if it is not difficult once Bishop's is granted) that the statement remains valid for any metric ball, that is to say one can go "over the injectivity radius". Then a Ricci lower bound yields first an upper bound for the volume of balls. But the non increasing property gives now a lower bound once the diameter or the volume of the manifold are known, since always trivially $\text{Vol}(B(p, \text{diameter})) = \text{Vol}(M)$. This is the key of many results, the most spectacular being the precompactness of I. C. 2. Such a two-sided bound on the volume of balls with only a lower bound on Ricci curvature should be compared with Colding's formula seen in the digression of I. A. 2.

B. THE ISOPERIMETRIC PROFILE

We used in IV the fact that there is for general Riemannian manifolds an isoperimetric inequality involving only the diameter and the lower bound of the Ricci curvature. The **isoperimetric profile** $h(t)$ of a compact Riemannian manifold (M^d, g) is defined as the map $h : [0, 1] \rightarrow \mathbb{R}_+$ where $h(t)$ is the infimum of the measure of the boundary $\partial(\Omega)$ of all domains Ω in the manifold

whose volume is $\text{volume}(\Omega) = t \cdot \text{volume}(g)$. Note that $h(1-t) = h(t)$ (look at complements). The domains have to be reasonable and then the measure of their boundary is any classical $(d-1)$ -dimensional measure. What are the explicit domains realizing the infimum are in explicit (standard) manifolds is in general impossible to know. Typically they are not metric balls, so that Bishop's inequality is not of any help. What is important, and very often enough for applications, is to have a lower bound for $h(t)$. A Ricci curvature lower bound (together with the diameter) is still enough as was discovered in (Gromov, 1980a). This was made more precise in (Bérard, Besson, & Gallot, 1985), the case of a negative Ricci lower bound being more tricky because there is no optimal comparison manifold. In the positive case one of course compare things with spheres. The proof goes as follows and is very important to know as it is a very new technique. Moreover Gromov's result (called Lévy-Gromov in the literature), apparently pure Riemannian geometry, triggered a lot of applications in Banach spaces as well as in probability theory : see the report (Ledoux, 1992/93).

The basic tool here is **GMT** (see **TOP. 10. C**). It insures first us of the existence - under reasonable extra conditions - of a (smooth or with reasonable singularities) submanifold realizing the minimal volume in its homology or homotopy class. Here for the isoperimetric profile it insures us with existence of extremal domains. Gromov's idea is to fill up the "inside" of the domain Ω by the geodesics normal to its boundary $\partial(\Omega)$. One then gets an inequality between $\text{volume}(\Omega)$, $\text{volume}(\partial(\Omega))$ and the mean curvature η of $\partial(\Omega)$. This inequality is in (Heintze & Karcher, 1978) and generalizes Bishop's one and applies to the sheaf of geodesics orthogonal to a given hypersurface, involving only the mean curvature of the hypersurface and the lower Ricci bound. Here η is constant because **GMT** insures us that almost everywhere the boundary is smooth and then its mean curvature is constant by the first variation formula for hypersurfaces. But this is apparently dramatic since we have no information on this constant η . Gromov's amazing trick is to look both at Ω and its complement $M \setminus \Omega$ and one is done because the effect is just to change η in $-\eta$.

For the amateur of problems naive to state, seeming obvious or at least simple but still open, we mention here the question of computing the exact value (and also if possible the shape of the optimal domains) for our preferred manifolds : the spheres S^d and the KP^n . One can add the cubic flat tori. For the sphere the

isoperimetric profile (called then "the isoperimetric inequality") is known since (Schmidt, 1948-1949). Who by the way solved also the problem for hyperbolic geometry (the simply connected space forms of constant negative curvature). The optimal domains as expected are metric balls. The tool is symmetrization and today the best reference is (Burago & Zalgaller, 1988).

But symmetrization does not work any longer, globally for the real projective spaces $\mathbf{RP}^d$ and even locally for the other $\mathbf{KP}^n$. The problem today is solved only for $\mathbf{RP}^3$ in (Ritoré & Ros, 1992). The solution is as follows : for small t optimal domains are metric balls but, starting at a number t trivial to evaluate, the tubes around a periodic geodesic (a projective line) gives a better answer. The proof in $\mathbf{RP}^3$ is finished because one is able there to completely classify the constant mean curvature surfaces in $\mathbf{R}^3$ (which moreover have a nonnegative second variation and then called "stable"). A reasonable conjecture is : in all the $\mathbf{KP}^n$ the isoperimetric profile is given by the succession of the tubular neighborhoods around : a point, a projective line, a projective plane, etc.

Another important control is that of λ_1 with the Cheeger's "h-constant" (which is the value $h(\frac{1}{2})$ of the isoperimetric profile for the value equal to just one-half of the total volume) : (Cheeger, 1970b). His inequality was $\lambda_1 > \frac{1}{4} h^2(\frac{1}{2})$ and is optimal by (Buser, 1978) but never attained by a smooth object, see the very end of chapter III.

The beautiful formula in (Savo, 1996a), or (Savo, 1997), studies the volume of various tubes and then encompasses many results of the above type with a very nice proof : the second derivative with respect of the radius of the tube is computed and linked with the Laplacian.

Recently, in the spirit of many results in Riemannian geometry (some of them were met above), the article (Gallot, 1987) succeeded in controlling the isoperimetric profile with only integral bounds on the Ricci curvature, see this text for precise statements. Corollaries are many : finiteness theorems (see I. C. 1), control on the spectrum (see IV). Moreover Gallot also controls the volume of

tubes around hypersurfaces. A key tool as we saw in I. C would be to control the volume of tubes around geodesics. With only an integral Ricci bound this was achieved in (Petersen, Shteingold, & Wei, 1996a).

A special mention should be attributed to the case of **nonpositive** curvature case. The conjecture is : in a simply connected manifold of nonpositive curvature the isoperimetric profile is **super-euclidean**. This means that for any compact domain Ω the ratio $\text{Vol}^d(\partial(\Omega))/\text{Vol}^{d-1}(\Omega)$ is larger than equal to its value for balls in $\mathbb{R}^d$. Today things are mysterious. For surfaces this was probably conjectured by Paul Lévy in a talk at Hadamard seminar in 1926, then immediately proved by André Weil using conformal representation : (Weil, 1926). In (Croke, 1984b) it is also proved when $d=4$ using integral geometry and no hope to make it working in any other dimensions. For $d=3$ this is the result of (Kleiner, 1992). Then in (Cao & Escobar, 1995) it is proved for piecewise linear (PL) manifolds of $\text{CAT}(0)$ (see TOP. 9 for those manifolds) and of any dimension. However is not clear today is any (smooth) Riemannian manifolds of nonpositive curvature can be approximated by PL-manifolds of $\text{CAT}(0)$. The main difficulty is that **GMT** is of no longer of use in a straightforward way : because of the lack of compactness optimal domains can "escape at infinity".

A basic tool in Analysis are the Sobolev inequalities, they are used in many results quoted in this text. For a numerical function $f : M \rightarrow \mathbb{R}$ on the Riemannian manifold (M^d, g) and the integrers (p, q) with $\frac{1}{p} + \frac{1}{q} = \frac{1}{d}$ they say that between the function and its gradient one has always for the global L^p and L^q norms :

$$\|f\|_p \leq A \|df\|_q + B \|f\|_p$$

In most cases the Riemannian geometer is interested not to have only some A and B but by a control on them with the curvature, etc. The problems of the optimal A and B are much different. For B it is basically the work of Gallot, see the various references to this author above. For A it is the work of Aubin, see for example (Aubin, 1982). See the recent reference (Hebey, 1996b) and its bibliography. We just mention that in both works the control of the isoperimetric profile is basic.

C. THE EMBOLIC VOLUME

This was alluded to in III. B . The simplest numerical and curvature free functional invariant on the set of compact differentiable manifolds M is the embolic volume, namely the infimum $\text{MinEmb}(M)$ of the ratio $\text{Vol}(g)/\text{Inj}^d(g)$ when g runs through all possible Riemannian metric on M . Today one does not know if there is some kind of convergence result for MinEmb which will insure the existence of a minimal metric on M , even if one admits metrics with (reasonable) singularities. However the situation is not completely desperate thanks to the following three results. First the very strong finiteness theorem of (Grove, Petersen, & Wu, 1990-1991) says that (at least when $d \neq 3, 4$) the possible diffeomorphism types with $\text{MinEmb} < A$ are always finite, so that the various real numbers MinEmb can be considered as a nice scale using Riemannian geometry to evaluate the degree of complexity of a manifold. The second result is (Berger, 1980) : $\text{MinEmb}(M)$ is always larger than or equal to that of the standard sphere, namely $\beta(d)/\pi^d$ (where $\beta(d)$ denotes the volume of the standard sphere of dimension d) and equality for $\text{Vol}(g)/\text{Inj}^d(M)$ is attained only for the case of constant curvature. In particular any embolic volume is always positive. The third result is the isolation in (Croke, 1988b) : there is some $c > 0$ such that for any manifold M which is not a topological sphere one has $\text{MinEmb}(M) > \beta(d)/\pi^d + c$. It seems an interesting question to relate in various ways $\text{MinEmb}(M)$ and the different topological invariants of M . Finally we cannot resist to tell the reader that the exact value of the embolic volume for the KP^n is completely unknown. If it were the value obtained for the standard metric, this will solve (or at least help a lot) the problem of characterizing the KP^n by their geodesic flow.

Of basic importance in various domain of Riemannian geometry is the local embolic result of (Croke, 1980) which tells us that, when the injectivity radius of the manifold is given, then every ball of radius half of the injectivity radius (or smaller) has a volume bounded from below by a universal constant in the dimension (and the radius). This is used above in many finiteness and convergence theorems of I. C. Open is to know if the value of this universal constant is always or not nothing but the value it takes on the standard sphere. This of course would be the optimal value.

D. THE MINIMAL VOLUME

One natural functional to evaluate when one wants to get the "less curved" metric a Riemannian metric g is the **minimal volume** of Gromov. $\text{MinVol}(M)$ is the minimum of the volumes of all Riemannian metrics on M such that $-1 \leq K \leq 1$. Here this functional can be zero, think of flat tori or even nilmanifolds. More generally it is zero any time the manifold M admits a sequence of collapsing metrics, see I. C. 3. Thinking back of the finiteness results in I. C. 1 for manifolds with $a < K < b$, diameter $< D$ and volume $> v > 0$, the importance of deciding if MinVol is positive or zero cannot be exaggerated. This because $a \leq K \leq b$ can always be written $-1 \leq K \leq 1$ after normalization when one is not interested in precise values. Also when looking for a **best metric** on a given manifold (see III). Then finally a positive minimal volume is an insurance against collapsing. There is no general survey on the minimal volume, but (Fukaya, 1990) devotes one of its sections to it.

Apart the case of surfaces where Blaschke-Gauss-Bonnet formula (O. E) tells us everything, the minimal volume remained quite mysterious up to 1996. Let us see briefly the state of affairs today. We saw that the main point is to know when the minimal volume is zero or is positive. We first remark (following Cheeger) that the minimal volume is positive as soon as the manifold has some non zero characteristic number : this is thanks to Chern integral formulas for characteristic classes seen in O. E. Note that this works only in even dimension and mainly in dimensions multiple of 4. In (Gromov, 1983b) it is proved that the minimal volume is non zero for space forms (see II) of negative type. And hence for many other manifolds using various functorial properties of the minimal volume. But note that those are always highly non simply connected. Today one still does not know any **odd-dimensional simply connected** manifold for which one can prove that its minimal volume is positive.

On the other hand (Gromov, 1983b) exhibits many manifolds with $\text{MinVol} = 0$. It started with manifolds where there is a free action of the circle, the first example (besides the obvious flat space forms) being of course the odd-dimensional sphere where the circle action is that of their Hopf fibration). Then this was extended to various manifolds which were laying the foundations for the study of collapsing, see I. C. 3. So that we are left with the basic question : are there more manifolds with $\text{MinVol} > 0$ or more manifolds with $\text{MinVol} = 0$? The fact that $\text{MinVol} = 0$ implies the

existence of an F-structure (see I. 3 . B) is not enough. We will see now that the recent work (Cheeger & Rong, 1996b) seems to show that "most" manifolds have a zero minimal volume.

Let us finish the MinVol story by addressing three questions. The first is : how looks the subset of the reals made up by the minimal volume of all manifolds of a given dimension d ? Is it a discrete set ? For surfaces Blaschke-Gauss-Bonnet formula tells us that our subset is the arithmetic progression $2\pi N$. The discreteness at 0 is the isolation of 0 (existence of a gap). This is true in dimension 3 and is implicit in (Cheeger & Gromov, 1986-1990). Oppositely the proof of the isolation of 0 in dimension 4 obtained in (Rong, 1993) is extremely involved and not completely geometric. It uses the η -invariant (see TOP. 6) in a basic way. But for $d \geq 5$ to know if there is or not a gap (one of Gromov's conjectures) is open. In (Cheeger & Rong, 1996b) the existence of a gap is proved when one restricts the problem to manifolds whose diameter is bounded. Namely, for any d and D , there exists $\epsilon(d,D)$ such that $\text{Vol} < \epsilon(d,D)$, $|K| \leq 1$ and $\text{Diameter} < D$ imply $\text{MinVol} = 0$. By Cheeger's finiteness theorem (I. C. 1), this implies also the discreteness of the set of all MinVol when the diameter is bounded. The problem is to know if, when D goes to infinity, a gap and/or discreteness keep preserved. The proof uses collapsing theory (I. C. 3) and consists in constructing, when $\text{Vol} < \epsilon(d,D)$, a polarized F-structure on the manifold. When the manifold is simply connected the result is an application of the collapsing previous results of Cheeger-Fukaya-Gromov where one can obtain a pure-polarized structure.

The second question is to compute the minimal volume of "standard" manifolds, e.g. space forms of different types. Things are again dramatically different according the sign of the curvature. For positive curvature or nonnegative one not a single minimal volume (when non zero of course) is known. It starts with S^4 . The only known fact is a local result for the even-dimensional spheres in (Ville, 1987). Oppositely for negative curvature space forms (hyperbolic) one has an unbelievably strong result. As a corollary of the main theorem in (Besson, Courtois, & Gallot, 1995a) (we already saw more than one application of it in III, the result is detailed in V. B) : under the sole condition $\text{Ricci} \geq -(d-1)$ any metric on a compact hyperbolic space form has a volume larger than equal to that of the hyperbolic metric. Moreover it is equal to it if and only if the metric is the hyperbolic one. So that once again we know that the hyperbolic metric on a negative space forms is "the best one".

The last and **third** question is the relation of MinVol with the functional $g \rightarrow \int_M \|R\|^{d/2}$ (see III. B). The minimum of it for various metrics on a given compact M is called $\text{Min}(\|R\|^{d/2})$ and is of interest, except for $d=2$ because to the Blaschke-Gauss-Bonnet formula. It plays a role in I. C. Trivially it is universally bounded by MinVol so that $\text{MinVol} = 0$ implies $\text{Min}(\|R\|^{d/2}) = 0$. The converse should be obvious with the convergence theorem in $\text{RM}(d, -1, 1, v, D)$ since we can always normalize our sequence within $-1 \leq K \leq 1$ and argue by contradiction. But we forgot about the diameter. Today the question is open to know if this converse is true or not, despite many attempts.

E. THE SYSTOLIC STORY

In A above the game was to control volumes of different kinds with control from the metric, e.g. curvatures of various types. The systolic story is entirely different : one wants to have inequalities for volumes independent from the metric, depending on the toological structure of the manifolds. Here the results are basically complete, all major questions have been solved. Only remains open few points (which might be very hard however). The story is as follows and is mainly Gromov's work. An almost complete survey is (Berger, 1993a) and add (Gromov, 1992a) which contains a lot of information. We now add few recent references.

Things started with Loewner's theorem (1949, unpublished). Call the **systole** of a two-dimensional Riemannian (T^2, g) torus the infimum of the length of closed curves which are not contractible. Let $\text{Sys}(g)$ be this positive real number. Then the area $\text{Area}(g)$ of

our torus satisfies the inequality $\text{Area}(g) \geq \frac{\sqrt{3}}{2} \text{Sys}^2(g)$ and equality occurs only for the flat regular hexagonal torus. The proof is not too hard and rests essentially on conformal representation : when averaging under the translations of the flat underlying structure the systole can only goes up by its very definition and the area goes down by Schwarz inequality.

The challenge is obvious : generalize this to surfaces of higher genus, then to (non simply connected) manifolds of higher dimension. And also to higher dimensional systoles. Namely call

$\text{Sys}_k(g)$ the infimum of the volume of all homologically non trivial k -dimensional submanifolds (for some Riemannian metric g on M) of M . In this broader context René Thom question in the 60's to the present author was to find when there is some universal inequality between the different systoles, the total volume of M^d being nothing but $\text{Sys}_d(g)$. In short we know today that we have almost a complete positive answer in the case of the 1-systoles and a negative answer for higher dimensional systoles. Let be more precise.

For the 1-systole and surfaces of higher genus in (Accola, 1960) and independently in (Blatter, 1961) came an answer but the lower constant for Area/Sys^2 was very poor, say anti-intuitive since going to zero extremely fast with higher genus the time being one oppositely expects some growing up with the genus. For the special case of the real projective plane the expected optimal answer was obtained "à la Loewner" in (Pu, 1952). Nothing more came on the subject before (Gromov, 1983a) which gave first an optimal answer first for surfaces (at least asymptotically with the genus γ) with a lower bound in $c \gamma / \log^2 \gamma$ (with c universal). The result is essentially the same for the homotopy systole and for the homology systole.

Second for the homotopy 1-systole and for any dimension d Gromov proved the existence of a positive lower bound for Vol/Sys^d for any reasonable manifold, namely for a class containing all acyclic ones and some more, like $\mathbb{R}P^d$. The technical definition is that of an **essential** manifold.

Gromov's proofs are very involved. The one for higher dimensional manifolds introduces many new interesting geometric invariants for Riemannian manifolds : the **filling radius**, the filling volume. Like the minimal and the embolic volumes their explicit value for standard manifolds are very difficult to compute explicitly. The latest state of affairs for the important filling radius is in (Katz, 1983) and (Katz, 1991) : this radius is known for the spheres, the real projective spaces and the complex projective plane, but still unknown for the remaining $\mathbb{K}P^n$. These invariants are promised to some future in Riemannian geometry. For example the filling radius is used in (Greene & Petersen, 1992) to improve the recent finiteness theorems mentioned in I. C. 1.

Back to Gromov's proof the game is to produce various inequalities between the above invariants (even with a non optimal constant)

and relate them finally to the systole. The proof uses an important trick : embed a Riemannian manifold in a Banach space by the distance functions to various points (see **TOP. 9**) (even if after during the proof one makes a systematic use of finite dimensional approximations). The proof for optimal (in the ingredients) systolic inequalities for surfaces of high genus is even harder, it uses the notion of **simplicial volume** and a diffusion process for simplicial cycles. These two ingredients are basic for the results we mentioned for the minimal volume in (Gromov, 1983b), see **D** above.

We are left with three questions. The first is to find the optimal value and if "extremal metrics" exist and who they are. For this, besides the "Filling paper" (Gromov, 1983a), see : (Calabi, 1992b) and (Gromov, 1992a). The second is to characterize manifolds for which we have such an universal inequality. This is done in (Babenko, 1992) for the **homotopical** $\text{Sys}_{1,\pi}$ associated to the fundamental group. Such a result can be seen as a "systolic" characterization of a topological property. This is also valid for the **homological** systole $\text{Sys}_{1,H}$ (but does not figure in the text), just be careful about the non-orientable case.

For higher dimensional systoles , negative results started to appear in (Gromov, 1992a). They were intermediate results, the Gromov 's example of (1,3)-softness on $S^1 \times S^3$ is basic and consists simply in gluing on $[0,1] \times S^3$ after twisting enough with the Hopf fibration. We have now a extremely large category of negative examples in (Babenko & Katz, 1997) and (Babenko,Katz, & Suciu, 1998), see also Katz appendix in (Gromov, 1997)). One says that a manifold M^d is **systolically (k,d-k)-soft** (or **free**) is the infimum of the quotients $\text{Vol}(g)/\text{Sys}_k(g)\text{Sys}_{d-k}(g)$ for all metrics g on M^d is zero. The above authors proved softness in the following cases : first, for any orientable (k-1)-connected M^d with $d \geq 3$, $k < d/2$ and k not multiple of 4 , there is (k,d-k)- softness. Second for simply connected $M^{d=2n}$ with $d \geq 6$, provided $H_n(M)$ is torsion free. Startling examples are the $S^k \times S^k$ for any $k \geq 3$ and $\mathbf{HP}^2$. The (4,4)-softness fo $\mathbf{HP}^2$ is most surprising, since the projective lines really fill up the whole space in the most geometric possible fashion and should have been sniffing around enough to prevent softness. The "freedom" can be very large, for example on $S^3 \times S^3$ one can even use only homogeneous metrics. It is not clear today if the various topological restrictions are really necessary as well as the dimension ones. In fact there is **no** known example of a **hard** inequality as soon as the involved systoles are 2 or more

dimensional. However the dimension 4 might be exceptional. In (Babenko & Katz, 1997) it is shown that the (2,2)-softness of $S^2 \times S^2$ and CP^2 are equivalent

An important parallel is to mentionned with (Besicowitch, 1952) (see chapter 7 of (Gromov, 1983a)) and concerns manifolds with boundary (see TOP. 9). The author proves that the volume of a cube with any Riemannian metric has a volume larger than equal to the product of the infimum of the lengths of the curves joinging two opposite pair of faces. But for a cylinder, there are metrics with smaller and smaller volume even when both the infima of the length of the curves joinging the two boundary disks and the area of disks whose boundary belong to the cylinder itself are larger than one.

The techniques for the above results are completely geometric. One exhibits submanifolds which are "roomy" as in Besicowitch cylinder, but construced like the Gromov $S^1 \times S^3$ counterexample above. Then one can find associated exterior forms which are calibrating (see TOP. 6. A) to insure that the systoles are lower bounded. All the above softness results can be interpreted as instability results for the homology, see (Babenko & Katz, 1997) for more on this.

2. Isometric embedding

Since (Whitney, 1936) one knows that they are no more abstract manifolds that topological submanifolds of euclidean spaces. Since (Nash, 1956) one knows now that abstract Riemannian manifolds are no more that submanifolds of euclidean spaces endowed with the Riemannian induced metric. The game of finding the optimal codimension is hard. Questions here can be local, global or adressing real analyticity. For the general case a good part of the book (Gromov, 1986) is devoted to it. We refer to it and just mention few points.

A spectacular case is : any abstract metric on S^2 of positive curvature can be realized by a convex surface. This was achieved through successive efforts : in (Lewy, 1938) for the analytic case, in (Nirenberg, 1953) for the twice differentiable case. There is a related result for the case of polyhedra (and going to the limit) back in 1941-42 by A. D. Alexandrov. For more precisions and the most general statements in connection with singular Riemannian manifolds, see the book (Pogorelov, 1973).

It is important to note that the differentiability assumptions are basic in this game and in the general result of existence above. For example any compact Riemannian manifold M^d admits a local C^1 -isometric embedding in $\mathbb{R}^{d+1}$ and a global one into $\mathbb{R}^{2d}$ (a dimension which is already needed topologically). This was achieved in (Kuiper, 1955) and (Nash, 1954). The idea is like Lebesgue handkerchief for developable surfaces : clever crumpling, creasing. This result can be compared with Lohkamp's one on negative Ricci curvature in I. B. 5. Very interesting to visualize is the result of (Bleecker, 1995). This text first extends Kuiper's result to one-parameter families, then uses it to obtain C^1 -isometric deformations of, for example prolate enough ellipsoids in $\mathbb{R}^3$, which are paradoxically **increasing** the volume. The wording of the author is "finely corrugated wrinkling". This of course does not exist with a C^2 condition since Gauss theorem would implies convexity, hence rigidity, by the theorem quoted in TOP. 10. B.

3. Holonomy groups and special metrics : another (very restricted) Riemannian hierarchy, Kähler manifolds

Since the parallel transport is the most natural geometric operation in Riemannian manifolds, we ask the following : are they manifolds for which the parallel transport, besides of course the euclidean metric of the tangent spaces, preserve some "extra" structure ? Manifolds could then, in some sense, be classified by those structures. A typical case is the Kähler one. It was made clear in (Lichnerowicz, 1955) that to be Kählerian is equivalent to the complex structure to be preserved by parallel transport.

A. HOLONOMY GROUPS

Such a classification was Elie Cartan hope in (Cartan, 1926a) where he introduced the notion of **holonomy** group. The holonomy group of a Riemannian manifold is simply the subgroup of the full orthogonal group made up by the parallel transport along all possible loops based at a given point. The change of point does not change the structure of the group, the only possible difference is if the manifold is simply connected or not. Then one has to distinguish between the full and the restricted holonomy group (use only contractible loops). Surveys of holonomy groups are : the chapter 10 of (Besse, 1987) and the book (Salamon, 1989).

Holonomy groups were completely forgotten (or might be at least found too hard to be studied) between Elie Cartan and the 50's. They came back to study in (Borel & Lichnerowicz, 1952). Things here are in some sense completely local, holonomy groups make sense even for non complete manifold. The only problem left is this still unbelievably open question : is the holonomy group of a compact manifold compact ? The question arises of course only in the non simply connected case. The answer is yes for many cases of the classification below, but indirect ones : one first uses the fact that most groups in the classification have a **finite** normalizer in the full orthogonal group. Second, for the Ricci flat cases, one uses the splitting theorem : see theorem 6 in (Cheeger & Gromoll, 1971).

Next to look at is the reducibility of the group. Things were finished to be cleared up in (de Rham, 1952). The reducibility of the group implies quite easily that the metric is locally a product. Then simply connectedness implies moreover a global product but it is harder to prove.

Then in (Lichnerowicz, 1955) (pages 258-261) it is proved that the Kähler condition is equivalent to the holonomy group to be a subgroup of the unitary group $U(n)$ for M^{2n} . Moreover the special unitary group $SU(n)$ characterizes Ricci flat Kähler metrics. Before Yau's solution of Calabi's conjecture in 1978, no (compact) manifold with holonomy $SU(n)$ was known (see the end of III. C).

How about other irreducible subgroups of the orthogonal group ? The result of (Berger, 1953) (see (Simons, 1962) for a much better proof or the indicated above surveys) says this. Note first that the result is completely local, no completeness is needed. Caricaturally speaking the holonomy group is almost always the full orthogonal group in general or the unitary group for the Kähler case. So that there exists a two-step hierarchy : the general case and the Kähler one. There are however few exceptions in this hierarchy. First the space forms of the general type, namely of any locally symmetric space, so that our new hierarchy is finally not too far from the one in II.

Then come two other exceptions in general dimensions : the first is the case of manifolds called **quaternionic** (even if they do not really admit a quaternionic structure) : their holonomy groups is that of HP^n or $HypHP^n$ which is the group $S(Sp(n \times Sp(1))) \subset SO(4n)$. It is open today if there exists many quaternionic manifolds besides the locally symmetric ones, no compact one is known today.

Recent references are (Le Brun & Salamon, 1994), (Boyer, Galicki, & Mann, 1994) as well the whole Proceedings (Gentili, Marchiafava, & Pontecorvo, 1994). The recent notion of **3-Sasakian** manifolds is very useful for constructing examples.

The second is the case of the group $\mathrm{Sp}(n) \subset \mathrm{SO}(4n)$. Those manifolds are called **hyperkähler**. Paradoxically there are quite a few such manifolds. The first complete (non compact) example appeared in (Calabi, 1979). Compact ones can be built using the solution of Calabi's conjecture and some algebraic geometry : (Beauville, 1983). Practically they are Kähler manifolds with more than one Kähler structure, in fact really admit a quaternionic structure. They are today encountered in mathematical physics, in particular in the theory of "mirror manifolds" : (Voisin, 1996). Recent references for them are (Salamon, 1996), the report (Hitchin, 1991/92), (Besse, 1987) and (Biquard & Gauduchon, 1996).

The naive reader should be warned against thinking that hyperkähler manifolds are the quaternionic analogue of complex manifolds and enjoy coordinates like complex ones : holomorphy ("quaternionomorphy"), quaternionic derivatives, etc. This can, sadly enough, happen only in the flat case (then holonomy is zero and not $\mathrm{Sp}(n)$). This, due to Fueter before the 40's, can be found in modern language in (Salamon, 1982), see also (Sommese, 1975) and (Batson, 1992). However in (Joyce, 1997) one manages to develop a theory of "quaternionic functions" on hyperkähler manifolds.

Then the list in (Berger, 1953) ended with three exceptions in dimensions 7, 8 and 16. The case of dimension 16 was that of the subgroup $\mathrm{Spin}(9) \subset \mathrm{SO}(16)$. But this group has to disappear from the list, because of (Alekseevskii, 1968) where it is proved that this group is forcing so strong curvature relations that in fact it implies local isometry with the panda CaP^8 or its negative analogue, which are symmetric spaces, see (Brown & Gray, 1972) for a more detailed proof. One is left with two possible groups which are respectively $G_2 \subset \mathrm{SO}(7)$ and $\mathrm{Spin}(7) \subset \mathrm{SO}(8)$. It is only in (Bryant, 1987) that the first examples appeared, sadly enough local only. Complete ones appeared in (Bryant & Salamon, 1989). Compact ones exist now by (Joyce, 1996a), (Joyce, 1996b). It is not clear if manifolds with these holonomy groups are numerous or scarce. Joyce's construction is extremely expensive but very nice. It uses deformation of metrics, singular manifolds and the Atiyah-Singer theorem. For example for G_2 the idea is to start with a

manifold with a reasonable G_2 -structure on its frame bundle. Such a structure has a **torsion** which measures the defect for it to be parallel transported. Then one use a one-parameter deformation technique to get finally a torsion vanishing metric.

Note that, as well as $SU(n)$ and $Sp(n)$, holonomy G_2 and $Spin(7)$ forces the manifold to be Ricci flat, see the end of section III. There is also a strong relation between these special holonomy groups and parallel spinors : see TOP. 6. B. This is due to the fact that both G_2 and $Spin(7)$ are, oppositely to $SO(d)$, simply connected hence induce on the manifold a canonical spin structure. Using parallel objects one can define the G_2 (resp. $Spin(7)$) case very simply by just asking the manifold to have an invariant by parallel transport (i.e. with zero covariant derivative) exterior form of degree 3 (resp. 4). The picture below is taken from (Salamon, 1989)

Recently holonomy groups became important in mathematical physics : see (Fröhlich, Granjean, & Recknagel, 1997)) for a systematic theoretical physics presentation of this hierarchy and (Andersen, Dupont, Pedersen, & Swann, 1997). The holonomy group classification is also useful when studying nonpositive curvature manifolds, see section I. B. 4, and, on the nonnegative side for the characterization of symmetric spaces among compact Kähler manifolds by their bisectional curvature : (Mok, 1988) and I. B. 1.

We mention here an important heuristic meaning of the holonomy classification, whose consequences are still not exhausted : **objects (tensors) invariant by parallel transport do not**

exist in general in a Riemannian manifold. A contrario when someone exists it implies extremely strong restrictions for the Riemannian structure. The reason is that the full orthogonal group has "no invariant" besides trivial ones. Oppositely as soon as one object is invariant, it has to be very special, e.g. the Kähler form, etc.

B. KÄHLER MANIFOLDS

This is an entire topic in itself and we will have to be extremely brief. We know of no recent survey. A classic is the book (Weil, 1958), thereafter there is (Wells, 1980). Although motivated by algebraic geometry (Griffiths & Harris, 1978) is invaluable for Kähler geometry. See also the chapter 2 of (Besse, 1987), moreover Besse's book contains many facts on Kähler manifolds, in particular a detailed study of the homogeneous ones. The book (Amoros, Burger, Corlette, Kotschik, & Toledo, 1996) (see also (Toledo, 1997)) is entirely concerned with the fundamental group of Kähler manifolds but can be used indirectly, with its references, as a survey or a partial survey of the "Riemann-Kähler" domain. For more general complex geometry one can consult the three books (Bedford, d'Angelo, Greene, & Krantz, 1991).

An analysis of the historical contribution of Kähler himself can be found in (Bourguignon, 1993). Of course part of the interest of Riemannian geometers was to add the Kähler hypothesis to attack many problems too difficult to be solved in the general Riemannian context. However this is not purely pride, since Kähler manifolds appear most naturally in algebraic geometry. More : Kähler manifolds became recently extremely important in mathematical physics : see (Voisin, 1996) and the references there. Just be careful that Kähler manifolds are almost never to be considered as "complexifications" of "real" manifolds. However they are of course wonderful to work on since we have for them the holomorphic calculus.

By definition a Kählerian manifold is one with a complex structure (this means in particular that the coordinates changes are holomorphic for the complex coordinates) together with a Riemannian metric which has with this complex structure the best possible link, namely that the complex structure is first compatible with the metric but moreover invariant by parallelism. This is equivalent to the condition for the holonomy group to be included into the unitary group, hence equivalent also to ask the existence of

a 2-form of maximal rank of zero covariant derivative. An equivalent definition is this (but the proof is a little tricky) : we have a complex structure J with a Riemannian metric g which are compatible : $g(Jx, Jy) = g(x, y)$ for every x, y . Then one gets an exterior 2-form ω , called the **Kähler form**, defined as $\omega(x, y) = g(x, Jy)$. Then the manifold will be Kählerian if and only if ω is closed : $d\omega = 0$.

For many people Kähler manifolds look like algebraic ones. There is a good reason for that : a celebrated theorem of Kodaira asserts that this is the case under the sole condition that, via de Rham, the form ω belongs to the rational cohomology (equivalently has rational periods). For a proof in book form see (Griffiths & Harris, 1978) or (Wells, 1980).

Beware also that Riemann-Cartan normal coordinates can never be complex except in the flat case. But basic (even if trivial) is heredity: a complex submanifold of a Kählerian one is also Kählerian. The parallelism of the complex structure J implies immediately the curvature relation : $R(Jx, Jy) = R(x, y)$ as endomorphisms. Note that a Kähler manifold is then canonically a **symplectic** manifold, so that all symplectic geometry becomes available. Recent books on the topic are : (McDuff & Salamon, 1995), (Hofer & Zehnder, 1994) and see also (Audin & Lafontaine, 1994). Another strong point is that the Kähler notion is hereditary, not only for (complex) submanifolds but also for many algebraic geometry operations, in particular blowing-up. Note also that ω and its exterior powers ω^k are "calibrating", see **TOP. 6. A**.

Gromov's complaint is that the above definition is not too much geometric : (Gromov, 1992b). A geometrical tool, rarely used, is the **diastasis**. Invented in (Calabi, 1953), it is some kind of adapted metric (in fact it should be viewed more as a potential) but in general defined only locally. Everything concerning it can be found in (Hulin, 1995) and its bibliography. In this text the diastasis is used to study Einstein manifolds (see **III. C**).

Concerning curvature, besides the sectional one, two notions are natural for Kählerian manifolds : holomorphic curvature, namely sectional curvature of the real 2-planes which are complex lines. Explicitly they are the $K(x, Jx)$ (where $\|x\|=1$). Metrics of constant holomorphic curvature are locally isometric to $\mathbb{C}^n$, $\mathbb{C}P^n$, $\text{Hyp}\mathbb{C}P^n$ (please normalize) : this is standard now and is one very special case among the general type of question raised in the digression of **I. C. 1**. It was stated in (Bochner, 1947), see (Igusa, 1954) or

(Hawley, 1953) for a detailed proof. A weaker notion is that of bisectional curvature : $B(x,y) = R(x,Jx,y,Jy) = K(x,y) + K(x,Jy)$ for x,y orthonormal. A very strong result is that of (Siu & Yau, 1980) and independently (Mori, 1979) : positive bisectional curvature implies that the underlying manifold is CP^n . The nonnegative case is settled in (Mok, 1988) : one can have only biholomorphy with CP^n and isometry with Hermitian symmetric space, plus of course products and coverings.

For complex manifolds the theory of exterior forms is richer, they enjoy (at least the pure ones) a **type** (p,q) which refers to their writing in complex coordinates. But one has more, namely that the

exterior differentiation can also be split into two ones : ∂ and $\bar{\partial}$. The holomorphic forms are those killed by $\bar{\partial}$. If moreover the manifold is Kähler then (total) Laplacian $\frac{1}{2}\Delta$ coincides with both

the partial Laplacians coming from ∂ and $\bar{\partial}$. Via Hodge-de Ram theorem and the exterior product with ω (which is of course harmonic) this yields a lot of information. Among them let us first mention the growing of Betti numbers up to the middle dimension : $b_{p+2} \geq b_p$. Another simple example is that positive Ricci curvature forbids the existence of holomorphic form of any degree.

Second, here the Sullivan theory mentionned in **TOP. 6. A** is much stronger. It is proved in (Deligne, Griffiths, Morgan, & Sullivan, 1975) that the homotopy of a compact Kähler manifold, over the real numbers, is a formal consequence of the real homology ring (compare with **I. C. 1**). It is also easy to imagine now that in the Kähler case the Bochner vanishing technique will yield a lot of strong results under various assertions on curvature. This is indeed the case and is used heavily for various bundles over a Kähler manifold, especially (complex) line-bundles. There is an immense literature, see the classic (Hirzebruch, 1966). And thereafter, among others, (Siu, 1980) and (Sampson, 1986).

On the spectrum of a Riemannian manifold can one read if it is a Kähler one ? We are far from it. Using the asymptotic expansion there are some special results in (Gilkey, 1973a). Then (Gromov, 1992b) is more ambitious. There are good results for λ_1 :

(Lichnerowicz, 1958) where there is lower bound for λ_1 depending only on a positive Ricci lower bound and in (Bourguignon, Li, & Yau, 1994) an upper bound with the volume for algebraic manifolds

4. Cut-loci

A natural question on a Riemannian manifold is to ask when two points p, q are joined by more than one segment (a segment is a shortest path, hence supported by a geodesic). There is a technical problem here, which partly explains the difficulty of the topic.

Assume compactness for simplicity. Starting from a given point p a geodesic will be a segment (realizing the distance) up to a certain value, called its **cut-value** (certainly smaller than or equal to the diameter). The **cut-locus of p** is by definition the set of these cut-points. But at such a point two things can happen (simultaneity is possible) : either a second segment is appearing, either the point is conjugate to p , i.e. the exponential map is not of maximal rank. This is the "dichotomy" for the cut-locus. The cut-locus of a point p is always the topological closure of the set of points joined to p by more than one segment, see page 589 of (Warner, 1965). The injectivity radius at p is the (positive) distance between p and its cut-locus. On a compact manifold the injectivity radius, introduced in I. A. 2, is the infimum of the various injectivity radius at p when p runs through the manifold. It is a positive number and we saw above that is often a basic tool I. C and an important invariant in III. B. In book form the cut-locus is quite detailed in the chapter III of (Sakai, 1996), in the last chapter of (do Carmo, 1992) and in (Kobayashi, 1989).

We address now the question of the structure of the cut-locus. Oppositely to many topics we have considered in Riemannian geometry, the cut-locus had a continuous coming of results. But very few and quite spaced in time. It was first considered for surfaces in (Poincaré, 1905) who called it "ligne de partage". Then Elie Cartan used it for the topology of Lie groups in (Cartan, 1936) and studied it also in (Cartan, 1951) (it seems that he was unaware of Poincaré and Myers results). For surfaces it was well studied in (Myers, 1935b), (Myers, 1936) and (Whitehead, 1935), who introduced the words cut-locus and conjugate locus (and by the way proved the general existence of local convex balls). Then it appeared explicitly in the proof of the sphere theorem in (Klingenberg, 1959). One then has to wait (Wall, 1977), (Buchner, 1978) to get structure results for the generic and the real analytic

case. At the same time (Gluck & Singer, 1978) showed that in general the cut-locus can be very bad, typically non-triangulable for an open set of points. Very recently the cut-locus made a strong comeback. Essentially "cut-loci are not that bad", more precisely their Hausdorff dimension is in direct relation with the smoothness of the metric, see (Itoh & Tanaka, 1997b) and the references there.

Still the cut-locus remains mysterious and basically unknown on a given manifolds. The sole exception is the case of symmetric spaces: (Sakai, 1978) and (Takeuchi, 1979). See for space forms (Hebda, 1995). Oppositely the cut-locus on a two-dimensional (even of revolution) ellipsoid is still not known. For an umbilic it is reduced to its antipodal umbilic. For all other points it is conjectured to be a topological segment. But this latter assertion depends on the scandalously unproved Jacobi "statement" : the conjugate locus of a non-umbilical point m of an ellipsoid has exactly four cusps. Geometrically this conjugate locus is the **envelope** of the geodesics emanating from m .

Inverse results for the cut-locus are completely unknown up to one exception : the cut-locus of every point is always reduced to a single point (Wiedersehenmannigfaltigkeiten) only for the standard sphere, see in book form the chapter VI of (Sakai, 1996). This was obtained for surfaces in (Green, 1963) and was a conjecture of Blaschke in the 20's. For higher dimensions it is the content of the appendix D of (Besse, 1978). An inverse problem could be : who are the manifolds all of whose conjugate loci are submanifolds, in particular of codimension larger than 1 ? The problem is untouched today, even with stronger conditions, see in (Besse, 1978) what concerns the Blaschke conjecture. A very itching question, which is a very particular case of the above, is : who are the Riemannian manifolds for whom the diameter is equal to the injectivity radius ? This happens for the (standard) spheres and all the KP^n (with the standard metric). Are they other ones ? The question is raised at large in (Besse, 1978) where it is called Blaschke conjecture. It is not too hard to prove that the cut-locus of any point is a submanifold of constant dimension (that of a K -hyperplane) and also true, but more subtle, by algebraic topology that almost only the KP^n can appear as topological manifolds. We just saw that for the sphere the question is solved. Taking the sphere covering an RP^d the question is also solved for them : no other ones. But for $K = C, H, Ca$ the question is scandalously open.

We now come back to the question raised in 0. B about Cartan local statement : parallel transport and the curvature determine the

metric. In (Ambrose, 1956) is raised the global problem when one knows the parallel transport and the curvature for all geodesics segments starting from a given point. He conjectures that the answer was a complete determination of the metric. This is linked with the structure of the cut-locus and is still an open problem. It was solved recently for surfaces in (Itoh, 1997a) using (Hebda, 1994).

A final remark is in order : the cut-locus is essentially a Riemannian notion if one expects a reasonable behavior. As soon as one goes to more general metric spaces things can become very very wild. See for example (cf. TOP. 9) : (El Alaoui, Gauthier, & Kupka, 1996) for Carnot-Caratheodory spaces, (Shiohama & Tanaka, 1992) for Alexandrov surfaces and (Zamfirescu, 1996) (as well as the references there) for the boundary of convex bodies. For Riemannian manifolds with boundary see the references in TOP. 9

5. Non compact manifolds

In various instances we already mentioned extensions of results for compact manifolds to complete ones. It is clear that one should restrict oneself to some kind of non compact Riemannian manifolds to hope some results. Let us mention in particular : replace compactness by **finite volume**, ask for an **asymptotic behavior at infinity** of some given type, e. g. quadratic decay, quadratic curvature decay, volume behavior, etc.

A typical example is that of manifolds with nonnegative Ricci curvature. In the case of nonnegative sectional curvature we saw in I. B. 1 a perfect structure theorem (splitting and soul) of Cheeger-Gromoll and the bounded topology result of Gromov-Abresch. Note that these results are valid without any extra condition on the geometry (bounded or behavior at infinity). But if one asks the same question for nonnegative Ricci curvature, the splitting theorem is still valid but there is no structure theorem and there is no bounded topology result as seen on examples. This implies that results on nonnegative Ricci curvature on compact manifolds should use extra hypothesis, typically some kind of growth. The results one expect could be of different nature. Corresponding in the case of negative curvature case to various structures on the sphere at infinity (I. B. 4), it seems that the right structure to infinity to be defined (even if not unique in general) is the following. The idea is to look at the structure at infinity by considering, for a given (M, g) , the sequence of the (pointed)

manifolds $(M, p, r_i^{-1}g)$ when the "radii" r_i go to infinity. The convergence theorems in C. 2 extend easily to the category of **pointed** manifolds, in book form see (Petersen, 1997b). In particular one can extract a subsequence converging toward "some" limit "cone" (a cone at infinity, denoted by abuse M_∞) ; this cone need not be unique and also might depend on p . In various instances, one can prove that point M_∞ is a volume cone, or better a metric cone, and finally sometimes the euclidean space itself. Here is a small choice of results among many recent ones : §7 of (Cheeger & Colding, 1996d), (Cheeger & Colding, 1997a) and the followers. Exemplary is the conjecture of Anderson and Cheeger to the effect that if a cone at infinity is isometric to $\mathbf{R}^d$ and the Ricci curvature nonnegative, then the manifold itself is isometric to $\mathbf{R}^d$; this conjecture is proved now in (Colding, 1996b).

But we insist that non compact manifolds are, to some respect, more important than compact ones. This is why our partial survey should absolutely be completed by the reader in some way or another. The reader could discover that in the references given. Non compact manifolds appear with surfaces, specially space forms, where the eigenvalue behavior with respect to the value $1/4$ is even more delicate than for compact ones, see for example (Luo, Rudnick, & Sarnak, 1995b).

We saw in I. B. 2 and I. B. 4 non compact manifolds appearing naturally in the case of compact manifolds bounded Ricci curvature, of negative or of nonpositive curvature, when one looks at their universal coverings. But they are also used heavily in different instances, like in collapsing with **not bounded** diameters or, more generally, when one drops the diameter bound in various situations (see I. C and references there). See also the very geometric (Babenko, 1992).

A natural case is that of **finite volume**. Here many results we met above extend, with few or much pain, depending of the question. To give even a short list of references is impossible. What follows is very biased. Note just, to start with the naturality of the topic, that the famous modular domain $SL(2, \mathbf{R})/SL(2, \mathbf{Z})$ is of finite volume for its hyperbolic canonical metric (but also has singularities) : see (Luo, Rudnick & Sarnak, 1995b). We also mention the problem of extending the integral Chern formulas to the finite volume case. This turned out surprisingly to be an extremely hard subject. Even for surfaces, where it started in (Cohn-Vossen, 1935) and (Cohn-Vossen, 1936) for the problem of extending Blaschke-Gauss-Bonnet formula. However a lot of work remained to be done for surfaces.

We refer to (Shioya, 1992) and the intermediate references there, and just mention that many topics introduced in I. B. 4 (for negatively curved manifolds) play a role here. For higher dimensional formulas, involving then the characteristic χ but also the characteristic (Pontryagin) numbers, the job was started in (Cheeger & Gromov, 1985a) : in the non compact case one can get **irrational** characteristic numbers. More results are to be found recently in (Rong, 1995), see also the survey (Lück, 1996) .

Finite volume is specially a strong condition in the negative curvature realm, where most results valid for compact manifolds extend with often not too much pain to finite volume. For space forms and Mostow's rigidity see (Farrell & Jones, 1989b), for the general case see (Ballmann, Gromov, & Schroeder, 1985) and of course (Eberlein, Hamenstädt, & Schroeder, 1993).

Another condition is that of **bounded geometry**. It comes naturally for coverings of compact manifolds and homogeneous spaces. The question is studied in (Semmes, 1996) in relation with the quite recent notion of **complexity**.

An interesting question is to know when a complete Riemannian manifold can admit a non constant harmonic function ($\Delta f = 0$) or even more a positive spectrum. This possibility is directly linked with various curvature properties. The founding text is (Yau, 1975) where positive Ricci curvature was shown to be enough to forbid non constant harmonic functions (this is a "Liouville type" result). Then results did not stop to appear. We refer only to recent ones and there bibliography : (Colding & Minicozzi II, 1997) and (Yu, 1997). See also (Benjamini & Cao, 1996) for relations with sectional curvature.

In the non compact realm various results address the question : obtain results when the structure at infinity satisfies some condition, like to be asymptotically euclidean (with an asymptotic order to be precised). See among others (Shen, 1996) and references there. There are also **gap** results, which forbid various compact metrics to be glued with completely flat ones outside, etc. see the introduction of (Lohkamp, 1996) and (Greene & Wu, 1990), as well as what was seen in I. B. 4 . Many results have to be looked upon in (Eberlein, Hamenstädt & Schroeder, 1993), like for example asymptotically harmonic manifolds. A deep result is (Cheeger & Tian, 1994), it is used in the results of Cheeger- Colding in I. B. 2.

Techniques in non compact Riemannian geometry are many. Besides using the sphere at infinity in various ways, we mention the very geometrical **chopping** technique of (Cheeger & Gromov, 1989). This is an exhaustion technique where, when one goes to infinity, the boundaries of the successive compact pieces keep a bounded second fundamental form and are also volume-controlled.

Very interesting also is the technique for proving the "mass conjecture" in (Schoen & Yau, 1981) (see also the expository (Kazdan, 1991/82)) : using GMT one look at minimal hypersurfaces in the manifolds and study what they become when they go to infinity. For other viewpoints see the references in (Cao, 1996).

On compact manifolds essentially all homology and cohomology theories coincide. In the non compact case things are much more subtle. A good notion for Riemannian geometry is that of L^2 -Betti numbers. There are surveys (Pansu, 1996) and (Lück, 1996). We just mention that the "only-compact" motivated Riemannian geometer should consider this with respect. A striking example is (Gromov, 1991a), where L^2 -Betti numbers are used to solve the H. Hopf's conjecture mentioned above (at the very end of **I. B**) for the Kähler case : a compact Kähler manifold M^{2n} of negative curvature has a characteristic of sign equal to $(-1)^n$.

6. Bundles over Riemannian manifolds

We introduce bundles in the order of their Riemannian geometric character. First canonical or almost canonical ones (spinors), then those obtained by twisting canonical ones, then Yang-Mills fields. The most natural ones are those of **exterior differentials forms** met in **0.D**. As reference in book form there is (Gilkey, 1995), but add also (Berline, Getzler, & Vergne, 1992) for the index theorem in a very general context and for the spinors (Lawson & Michelsohn, 1989). Of basic importance are characteristic classes for various bundles over general differentiable manifolds, so one should look at their relations with Riemannian geometry.

A. EXTERIOR DIFFERENTIAL FORMS (AND SOME OTHERS)

Exterior differential forms exist on any differentiable manifold, their degree extend from 0 to the dimension d of the manifold. The corresponding vector spaces are denoted by $\Omega^p M$. And the basic link is the exterior differential $d : \Omega^p \rightarrow \Omega^{p+1}$ with $d \circ d = 0$ (this collection is called the **differential complex of M**). The **closed** forms are the ω with $d\omega = 0$. This is leading, as seen in **O. D**, to de Rham theorems via Stokes formula. Although not a Riemannian story, one should know that if this was only yielding for the topologist the real Betti numbers, more can now be extracted from the sole exterior differential complex, as it was discovered in (Sullivan, 1977), which yields for example some topological finiteness type results (see **I. C. 1**), see also **TOP. 3. B**. What more can now ask and do a Riemannian geometer? Of course for interesting relations between the metric and exterior forms.

A basic fact is the existence of the $*$ -operator : $\Omega^p \rightarrow \Omega^{d-p}$. It is involutive or antiinvolutive : $* \circ * = (-1)^{d(p+1)+1}$ and needs an orientation. The form $*\omega$ is defined by $*\omega(x_{p+1}, \dots, x_d) = \omega(x_1, \dots, x_p)$ for any positive orthonormal basis $\{x_1, \dots, x_d\}$. For example on an oriented manifold the $*$ of the constant function 1 is the **volume form**, namely the d -form which takes the value +1 on any positive orthonormal set (of tangent vectors). Its existence is equivalent to orientability.

Natural is to look at the canonical norms $\|\omega\|$ and thinking of de Rham theorem to look (if it exists) at the minimum of the L^2 -norm

$\int_M \|\omega(m)\|^2 dm$ for ω running into a given cohomology class.

Hodge- de Rham basic theorem (met in **O. D**) is that this minimum is attained and this by a unique form in the class which is characterized by $\Delta\omega = 0$ where the Laplacian Δ on forms is $d \circ d^* + d^* \circ d$, with the adjoint $d^* : \Omega^p \rightarrow \Omega^{p-1}$ of d is defined as

$(-1)^{d(p+1)}(* \circ d \circ *)$. There is no need of an orientation since one uses $*$ twice. **Harmonic** forms ω are those with $\Delta\omega = 0$. Stokes formula implies that $\Delta\omega = 0$ is equivalent to $d\omega = 0$ (closed form) and $d^*\omega = 0$ (a co-closed form). The second condition is exactly where the Riemannian structure enters.

Concerning harmonic forms one can wonder about the "inverse problem" : if harmonic forms are special or not among closed differential forms. This is a tricky question, attacked first for 1-

forms and solved in (Calabi, 1969). For higher degrees see (Farber, Katz, & Levine, 1996).

One can also wonder about the fact that, except for 1-forms, exterior forms are not good in general for computing volumes when one restricts them to submanifolds. But there are exceptions, this was first noticed in (Wirtinger, 1936). In a Kählerian manifold the Kähler form ω enjoys the following property : for any orthonormal pair x, y one has $\omega(x, y) \leq 1$ with equality only for complex lines : $y = Jx$ and this works also mutatis mutandis for the exterior powers $\wedge^p \omega$. Because ω is closed, and by Stokes formula, this implies immediately that any complex submanifold of a Kählerian manifold has an absolute minimal volume in its homology class (this is much stronger than being only a minimal submanifold and is called stability). Wirtinger used his result for various applications, for example to compute explicitly the volume of any algebraic submanifold of given degree in $\mathbb{C}P^n$.

The above inequality can hold for some forms of degree p on some Riemannian manifolds : $\omega(x_1, \dots, x_p) \leq 1$ for every orthonormal k -uple. This property is called **calibration** (calibrating form, calibrated manifold) in (Harvey & Lawson, 1982). The straightforward property of a calibrating form ω is that, if a

submanifold N^p has a volume equal to $\int_M \omega$, then it is an absolute

minimum in its homology class, in particular a stable minimal submanifold. For this it is necessary and sufficient that ω takes the value 1 on each tangent space to N^p . Calibration was used in (Berger, 1972) to prove the systolic inequality for the standard metric of KP^n for H and Ca (for C this is Wirtinger). In (Harvey & Lawson, 1982) a general theory of calibration is made. In particular some generalizations of Wirtinger are presented, in the sense that calibration is also linked with some PDE the same way complex submanifolds can be defined as holomorphic, namely by $\bar{\partial} = 0$. Recently calibration was used for systolic softness (freedom) in (Babenko & Katz, 1997) (see 1. E) and in (Besson, Courtois & Gallot, 1995a) in a Hilbert framework.

We now present the two outcomes of spectral considerations for exterior forms which have a Riemannian geometry flavor. The Kähler case is more rich as seen above in I. B. 1 . First with (McKean & Singer, 1967) a firework started and then yielded deep understanding and many applications. We describe it briefly, a complete reference is (Gilkey, 1995) (this second edition is very up to date). Roughly speaking what happens is the following. We look

back at the asymptotic expansion in IV for $\sum_i \exp(-\lambda_i t)$ with the

$U_k(x)$ integrals, which are universal in the curvature tensor (this will mean always including its covariant derivatives). People were in despair of the fact that $U_{d/2}$ was not a topological invariant as soon as $d > 2$. Since they have also a canonical Laplacian we can do the same (it is not too much more expensive and appeared first in (Gaffney, 1958)) for p -exterior forms and get the pointwise invariants, denoted by $u_{p;k}(x)$, coming in the asymptotic expansions of the corresponding heat kernels for the term in t^k . They are still universal in the curvature but differ in general with various p . Their integrals over M will be denoted by capitals : $U_{p;k}$ and the eigenvalues of the p -spectrum by $\{\lambda_{p;i}\}$. Now let us perform the alternate double sum

$$\sum_p (-1)^p \sum_i \exp(-\lambda_{p;i} t) .$$

Because both the operators d and d^*

commute with the Laplacian Δ they transform eigenfunctions into eigenfunctions. Now the Hodge decomposition theorem of any form into a harmonic part, a closed part and a coclosed part shows that into this alternate summation everything will disappear except at the harmonic level : there the zero eigenvalue $\lambda_{p;0}$ has a multiplicity equal to the p^{th} -Betti number b_p . So in the alternate sum of the corresponding asymptotic expansions everything should also disappears for any k except when $k = d/2$. Hence the

alternate pointwise sums $\sum_s (-1)^p u_{p;k}(x)$, when integrated on M and adding after multiplication by t^k , will yield **identically** the constant $\sum_s (-1)^p b_p = \chi(M)$. This explains McKean and Singer

dream : a fantastic pointwise cancellation might well take place in the pointwise $u_{p;k}$ to yield the forced integrated cancellation. This was indeed proved in (Patodi, 1971).

The rebound was taken first in (Gilkey, 1973b) then in (Atiyah, Bott, & Patodi, 1973). One studies Patodi cancellation result but put it in successive more and more general bundles equipped with suitable elliptic operators, including the Dirac one on spinors and use Gilkey's results. It then turns out that those structures are many enough to yield all elliptic operators, giving a new proof of the index theorem (see C below). Using the theory of invariants "à la Gilkey" and functorial behavior is important. The (final) result is to use this technique for more general statements. The harvest is larger : Hirzebruch's signature theorem (see 0. E) can be obtained this way and of course this new insight yields many results in differential topology . This domain is still in full activity, see the two books already mentionned. One point in this philosophy is that "pointwise cancellation" show that **local type index** theorems can exist. But Riemannian geometry is quite far away. However here comes the second byproduct of the rebound : the η -invariant.

The main trick in the founding papers (Atiyah, Patodi, & Singer, 1975-1976) is to obtain the characteristic $\chi(M)$, not as the alternating sum of the zero-eigenvalues of the various Laplacians on the exterior forms of a given degree on (M, g) , but in one shot as the index of the first order operator $B = d - d^*$ as acting on the total set of exterior forms on M (one just has to be careful to put right signs in front of B). The eigenvalues of Δ are of the form λ^2 where λ is eigenvalue of B but different signs are possible here. Hence the function

$$\eta(s) = \sum_{\lambda \neq 0} (\text{sign}(\lambda)) |\lambda|^{-s}$$

makes sense for suitable s . In an strict sense (as usual for this kind of function) $\eta(0)$ is not defined but with some extra work one can still make sense out of it . It is then called the **η -invariant** of (M, g) and measure the "spectral asymmetry". This invariant is specially interesting for manifolds with boundary. For a $4k$ -manifold M' with a $(4k-1)$ boundary M (and provided that locally at the boundary the metric is a product) one can express the

signature $\text{sign}(M')$ by the integral formula $\text{sign}(M') = \int_{M'} L(R) -$

$\eta(M)$ <here L is the universal curvature integrand for the signature of Hirzebruch seen in 0. E . This invariant has many applications

when looking at the subtle problem of the non-existence of pointwise invariant integration formulas for the "signatures". Besides the original papers we refer the reader to (Atiyah, Donnelly, & Singer, 1983) and (Gilkey, 1995). There are also relations with the secondary characteristic classes below, also with $\hat{A}$ -genus when spinors are in view. An application of the η -invariant for 3-manifolds is the isolation result of (Rong, 1993) for the minimal volume in dimension 4 seen in TOP. 1. D. The η -invariant is also used in Number Theory : (Atiyah, Donnelly & Singer, 1983).

Another invariant based on the spectral analysis of differential forms is to be found in (Ray & Singer, 1971). The result is that, from a suitable combination of the $\zeta_p'(0)$ value of the ζ_p -functions associated to the spectrum of the p -differential forms of all degree p , one can recover a topological invariant. They conjectured that their invariant was coinciding with the topological invariant called the Reidemeister torsion and gave some evidence for that. The conjecture was proved independently in (Müller, 1978) and (Cheeger, 1979), the proof is very involved and was one of Cheeger's motivation for the study of the spectrum on some singular manifolds, see TOP. 9.

There is no reason to stick with exterior forms and not to look at other tensors. Lichnerowicz introduced the theory of canonical and special Laplacian in (Lichnerowicz, 1961). Particularly important is his Laplacian for symmetric bilinear differential forms, as they can be interpreted as infinitesimal variations of a Riemannian metric. For Hodge-de Rham type theorems concerning it, see section 12. C of (Besse, 1987) where it is used to study deformations of Riemannian structures, i.e. metrics up to isometries (diffeomorphisms). Lichnerowicz' Laplacians are **natural**, in the sense explained in 0. D

B. SPINORS

When it exists, the **spinor bundle** is "almost canonical". For all what follows and more on "spin geometry" see (Lawson & Michelsohn, 1989), and add for some parts (Gilkey, 1995), (Berline, Getzler & Vergne, 1992). We are going to present things very caricaturally. The basic idea is : Riemannian geometry has a euclidean structure on every tangent space, so that, if moreover the space is oriented, the special orthogonal group $SO(d)$ is "the basic

object". Besides the fundamental representation on $\mathbf{R}^d$ the special orthogonal group has a basic one on the exterior algebra and, by tensor products, one can build all the other ones. So that, apparently, the Riemannian geometer should not look further than for those. But the Lie group $SO(d)$ is not simply connected and its (nontrivial twosheeted when $d > 2$) universal covering group $Spin(d)$ has an extra representation in a "canonical" complex vector space, the space of spinors of the initial euclidean space.

First let us make some purely algebraic facts. The canonical quadratic form on $\mathbf{R}^d$ gives birth to the **Clifford algebra** $Cliff(d)$, of dimension 2^d . This algebra, only as a vector space, is canonically isomorphic to the exterior algebra $\wedge(\mathbf{R}^d)$. Now there is a basic link between $Spin(d)$ and $Cliff(d)$ to the effect that one finally obtains two things in a canonical way. First, a complex vector space $\Sigma(d)$ of dimension $2^{[d/2]}$ (where $[n]$ denotes the integral part of n). This space shows up when one tries to write $Cliff(d)$ as an endomorphism algebra $End(?)$ and is called the (complex) spinor space of $\mathbf{R}^d$. Second, because $Spin(d)$ can be realized as a subgroup of the group made up by the units elements of the algebra $Cliff(d)$, one finally gets a representation of $Spin(d)$ on the space $\Sigma(d)$. The theory is complicated by the fact that this representation is irreducible only when d is odd but splits into non isomorphic ones denoted $\Sigma^+(d)$ and $\Sigma^-(d)$ of half the dimension when d is even.

There also exists real spinors, but the situation is more complicated with them and is to be considered mod 8. It is a basic fact that this game cannot be played without having fixed a quadratic form, because the universal cover of the special linear group $SL(d, \mathbf{R})$ does not admit any faithful finite dimensional representation except the ones in the exterior algebra.

Since all the above construction depends only of oriented euclidean geometry, namely a (the) positive definite quadratic form on $\mathbf{R}^d$, the game is to do this two step construction for any oriented Riemannian manifold (M^d, g) . Since the (principal) bundle of oriented frames $PSO(d)M$ has $SO(d)$ as structure group one first tries to define a universal double covering of this bundle, asking moreover of course compatibility with the group covering $Spin(d) \rightarrow SO(d)$. This is not always possible and a necessary and sufficient condition for this is the vanishing of $w_2(M)$, the second Stiefel-Whitney class of M : (Haefliger, 1956). Up to isomorphism, the

spinor structures so obtained are classified by $H^1(M, \mathbb{Z}_2)$. They are for example unique in the simply connected case. However one in general keeps a unique notation, namely $P_{\text{Spin}(d)}M$ to denote any spin-bundle covering $P_{\text{SO}(d)}M$. Now, attached to any spin structure on the manifold, (see C below) the representation of $\text{Spin}(d)$ on $\Sigma(d)$ automatically produces a (complex) vector bundle denoted by $\Sigma(M, g)$ and called the ("a") spinor bundle on (M, g) . A section of it is what is called a **spinor field** on M . Still using general facts on bundles (C below) the above construction and the Levi-Civita connection yield a canonical connection D on $\Sigma(M)$.

The third point, a major one, is that using D one can define on $\Sigma(M)$ a canonical differential operator of degree one, called the **Dirac operator** and denoted by δ . In even dimensions it exchanges the spinors $\Gamma(\Sigma^+)$ and $\Gamma(\Sigma^-)$. Now for δ^2 there exists a Bochner-Weitzenböck type formula which is $\delta^2 = D^*D + \frac{1}{4} \text{scal}$ (the Lichnerowicz formula) and was already used heavily in I. B. 3. In comparison with formulas for Δ involving the collection of the Curv_p (see O. D) this looks a priori a disaster. But it is just the opposite : with much less information on the curvature, one still gets information on the topology with the Hodge-type theorem for δ , namely for harmonic spinors σ , i.e. those satisfying $\delta\sigma = 0$.

The history of spinors is fascinating. It started with Elie Cartan when, in (Cartan, 1913), he completely classified complex irreducible representations of simple Lie groups : besides the expected orthogonal and exterior representations he found an extra one bringing a new space to life. Moreover he prophetically indicates how this representation can generate all others. Then, completely independently, Dirac invented spinors on physical grounds (hence their name) in 1928 and defined the Dirac operator, this not for manifolds but just for Minkowski space $\mathbb{R}^{3,1}$. The link was built by Cartan in 1937, but he concluded his book (Cartan, 1937) by noting the "impossibility" to build up a satisfying theory : this is so because coordinate changes with general Riemannian metrics on manifolds are only linear (not orthogonal) and we saw above that the universal cover of the special linear group has no finite dimensional representation besides those coming from the genuine linear group. The construction of spinor bundles and of the Dirac operator on them are hard for us to precisely date, it seems that they were more or less folklore. But historical is the year 1963 when at the same time appeared the index theorem, which among others yielded the fact that the index for the Dirac

operator is the Hirzebruch $\hat{A}$ -genus (hence an integer) and Lichnerowicz scalar curvature formula above. For this history and more on spinors see the introductions of (Lawson & Michelsohn, 1989) and of (Berline, Getzler & Vergne, 1992), as well as the postface "Spinors in 1995" by Bourguignon in "The algebraic theory of spinors and Clifford algebras" in (Chevalley, 1997).

We saw "the" basic applications of this in **I. B. 3**. For more on that see the books above. But the basic idea is that, when one defines much more general spin-type-bundles, one can often prove that $\frac{1}{4}$ scal is the dominant term. Other applications are : some proofs of the index theorem, construction of special holonomy groups, see **TOP. 3**.

A strong warning : as opposed to the case of the Laplacian and harmonic forms, the kernel of the Dirac operator δ depends on the metric chosen on a given manifold. The dimension of this kernel (the space harmonic spinors) can change, as it was first discovered in (Hitchin, 1974). Worst : any spin-manifold can carry a metric with some non trivial harmonic spinor (at least today in dimensions $4n+3$) : (Bär, 1995). However, harmonic spinors are conformal invariants : (Hitchin, 1974). The dependence on the metric is not easy to control because, when one varies the metric, the bundle also changes. In (Bourguignon & Gauduchon, 1992) is presented a detailed computation for everything in this context, in particular the first variation of the Dirac operator and of its eigenvalues (see references there in for intermediate results).

Finally, for the very geometrically minded reader, we mention that in (Connes, 1994) one finds a formula giving the distance of two points using the Dirac operator, which has the advantage to carry over the non commutative setting.

Not much more complicated to define are the Spin_c structures which exists on many more manifolds than spin-ones. In particular they exist on any 4-dimensional manifolds, see **TOP. 8** below for references to recent very significant applications. Beware of the notation : $\text{Spin}_c(d)$ is not the complex group $\text{Spin}(n, \mathbb{C})$ but "only" $(\text{Spin}(d) \times S^1)/\mathbb{Z}_2$.

C. VARIOUS OTHER BUNDLES

The theory of bundles can be found in most books of differential geometry and in detail in (Kobayashi & Nomizu, 1963-1969). From the topology viewpoint, the classics are (Steenrod, 1951), (Husemoller, 1975). Bundles E are differentiable manifolds equipped with a map $E \rightarrow B$ onto the basis B and a typical fibre F , such that the base B is covered with charts domains such that the counter-images in E are products with F . On their intersections, these bundles charts should be smooth on the fibers and most often preserve some given structure. For principal bundles the fiber is acted simply and transitively by a group, and charts changes have to be group automorphisms. For vector bundles they should of course be linear maps. Important is the possibility, when given a principal bundle together with an action of the group on some object (e.g. being a representation in a vector space), of canonically deducing from it an associated bundle. Such a construction was basic in order to construct canonically the spinor bundle above.

For bundles one has a notion of connection, then of curvature, of parallel transport and of holonomy. The idea of connection is the same as the one explained in O. A, namely one wants to be able to compare (infinitesimally) two fibers and to develop some kind of differential calculus. Of course, the connection one uses should preserve in a reasonable sense the various structures the bundle carries. In particular, Riemannian vector bundles appear when there is given a positive quadratic form on every fiber (with smooth dependence on the base point). For vector bundles one has characteristic classes and Chern's type formulas are still valid. Books are (Hirzebruch, 1966), (Berline, Getzler & Vergne, 1992), (Gilkey, 1995).

Secondary characteristic classes : we mention in O. E that there are no other integral formulas involving curvature besides Chern's ones. In (Chern & Simons, 1974) new subtle invariants were introduced. They take place in various bundles, they are not "downstairs" Riemannian objects. Although typically topologically trivial their connections are not and the parallel transport is yielding those new invariants. In (Cheeger & Simons, 1985b) they were put downstairs on the manifold where they live as "differential characters. These new invariants became more and more important, in particular recently in mathematical physics and in Number Theory : (Gillet & Soulé, 1994). In book form one can

look at (Berline, Getzler & Vergne, 1992) and (Gilkey, 1995), section 3. 11.

Yang-Mills fields "started to exist" in the late seventies and by a demand of theoretical physicists. A book appeared as soon as (Atiyah, 1979). But since the topic completely blew up and is now a topic in itself, with many subtopics. Yang-Mills question can be stated roughly as follows. On a given compact Riemannian manifold, most often a "standard" one like the sphere, one picks up some interesting vector bundle over it, some "bundle-Riemannian" metric on it and some connection preserving this metric. Then one looks for the "less twisted" one, typically in the L^2 -sense for the curvature of this connection. This is a problem of calculus of variations and leads for the curvature of the bundle to a condition looking like the harmonicity of the curvature. There is a rich harvest on Yang-Mills because what one looks for is like looking for harmonic forms, instead that the harmonicity condition is on the curvature. There are results like Hodge-de Rham. Besides its natural differential geometry setting is basic in theoretical physics (**gauge theory**). Surveys and/or recent references are : (Donaldson & Kronheimer, 1996b), (Donaldson, 1996a) and (Andersen, Dupont, Pedersen, & Swann, 1997). Intermediate references were (Freed & Uhlenbeck, 1984), (Bourguignon & Lawson, 1982).

We just mention here that there is a drastic difference between Yang-Mills structures on various bundles and the condition for this manifold to be Einstein (see III). In the Yang-Mills game one keeps fixed the metric downstairs and varies the connection in the bundle, while the Einstein condition is **coupling** the Riemannian structure of the basis with the tangent bundle connection. To do both at the same time is today very difficult as we saw.

Twistor theory belong more to Riemannian geometry than Yang-Mills ones, even if there are strong links between them. They are mostly bundles with compact fibers. They are constructed from a Riemannian manifold using its metric and some given additional structure e.g. a complex structure, special holonomy, etc. In dimension 4 every Riemannian manifold has a canonical twistor space above it, with fiber S^2 . The main point is that twistor space have a richer structure than their basis, e.g. Kählerian. In many cases they have pure geometric applications. One is the classification of minimal surfaces in standard spheres of high

dimension done in (Calabi, 1967). They are also of basic use for finding holonomy groups of some types (TOP. 3), but also to construct Einstein manifold, see for example (Hitchin, 1995). Besides a brief appearance in (Lawson & Michelsohn, 1989), one can consult in book form (Besse, 1987), and its second edition to come. An intermediate reference is (Atiyah, Hitchin, & Singer, 1978). Twistor-like considerations appear also in 3-Sasakian manifolds in connection with quaternionic holonomy : 3. A.

General vector bundles are of course only differential geometry at the beginning. The fact the basis is Riemannian gives in particular a measure and also a canonical connection on the tangent bundle. The interest of vector bundles lies in the fact that the so-called **K-theory** is an algebra made of all complex vector bundles over some given manifold. So one has succeeded bringing in an algebraic tool to study differentiable manifolds, the maps between them, etc. For example one of the important facts about the Dirac operator is that it represents the fundamental class in K-theory. For real vector bundles one had to use the more subtle **KO-theory**.

For general Riemannian real vector bundles, the idea of Atiyah and Singer was to extend to any such bundle the spinor construction of B (which was sketched for the complex case). The topological construction is the same as for the tangent bundle. One can consider both the complex and the real case. Moreover associated to Riemannian connections there is a canonical Dirac operator and a generalized Lichnerowicz formula which is again surprisingly simple, even for twisted bundle : see II. §8 of (Lawson & Michelsohn, 1989). For various vanishing theorems as used in I. B. 3 an important fact is that the scalar curvature of the base manifold appears separately of the curvature term of the vector bundle under consideration. Finally these generalized Dirac operators became essential in understanding more the index theorem below. This was through various contributions of Atiyah-Bott, McKean-Singer, Patodi and Gilkey : see the introduction of (Berline, Getzler & Vergne, 1992). This book is precisely devoted to the index approach with generalized spinors and Dirac operators.

If one now very broadly generalizes the Laplacian acting on functions and on exterior forms, by considering any elliptic operator on some vector bundle, there exists a very deep and universal result, the **Atiyah-Singer index theorem** which was a great event of 1963. This has been a query of Gelfand who

conjectured the topological invariance of the index. See (Atiyah & Singer, 1963) for historical references and intermediate results. The index of an elliptic operator is the dimension of its kernel minus the dimension of the kernel of its adjoint (cokernel). The theorem is that this integer, on a compact manifold M , is equal to a number computed only with two things : first the Todd class of M which is expressible with the set of its Pontryagin classes, second with an invariant computed with the symbol of the elliptic operator considered as acting on the various Chern classes of the vector bundles on which the operator is defined. In the case of differential forms one just recover the characteristic via Hodge-de Rham theorem. Applications of the index theorem are numerous (and far to be finished), e.g. yielding integrality of various invariants which a priori only real numbers. In the "pure Riemannian case" we saw above in I. B. 3 the case of the scalar curvature. References are (Gilkey, 1995), (Berline, Getzler & Vergne, 1992) and the references there.

Starting in 1982 with Witten, in 1985 with Quillen and Bismut in 1986, most of the above various notions for bundles were raised at a **super** level, e.g. superspace, superconnection, supersymmetry, etc. In particular Bismut Levi-Civita superconnection is basic for bringing the "super" concepts into Riemannian geometry. Since, things did not stop and they drastically enlarge the panorama and changed the vision, including connections, Laplacians and Dirac operators, asymptotic expansions of the heat kernel, the index theorem. The book (Berline, Getzler & Vergne, 1992) is the ideal reference for all this. For the holomorphic side, see (Bismut, Gillet, & Soulé, 1988).

7. Harmonic maps between Riemannian manifolds

It was realized a long time ago that geodesics are not only

minimizing the length $\int_0^1 \|f'(t)\| dt$ but also the **e n e r g y**

$\int_0^1 \|f'(t)\|^2 dt$. The quadratic character, as opposed to the absolute

value, makes the computation of the variational derivatives much easier. Moreover this energy can be interpreted as follows : it concerns a map f from the interval into the Riemannian manifold and is defined only with the Riemannian structure of it. More

generally to a map f between two Riemannian manifolds (M, g) and (N, h) one can attach its **energy** $E(f)$ which is the integral over M (for its canonical measure) of the trace of the quadratic form obtained from the derivative $f'(m)$ using the induced by the (N, h) Riemannian structure (the trace is taken with respect to the Riemannian structure of (M, g)). A map f is said to be **harmonic** when its energy is critical among all maps from M to N .

This very general notion was introduced in the founding (Eells & Sampson, 1964). The case of surfaces is quite special because harmonic maps are directly related to minimal surfaces thanks to the conformal representation. But in higher dimensions there is in general no direct connection between minimal submanifolds and harmonic maps, only in some special cases.

Since, it had a rich history, for the natural questions as well as for applications. The reader will find references, in particular of surveys, in the book (Eells & Rato, 1993). At its date, the book size (Eells & Lemaire, 1988) was a very systematic, informative and complete survey at that date. Important are various regularity results. In some cases harmonic maps are defined in a more general context that purely Riemannian (see also TOP. 9), see the books (Jost, 1995), (Helein, 1996) and for applications to Riemann surfaces (Jost, 1996).

Harmonic maps are a basic tool in today Riemannian geometry as very well illustrated in Eells-Lemaire report. For example see : their use for space forms in II. B, where the notion had to be extended to manifolds with singularities, in I. B. 1 for manifolds with positive curvature operator, in (Jost & Yau, 1990) for nonpositive curvature manifolds (I. B. 4). Harmonic maps are basic for studying Kähler manifolds, in particular their fundamental groups, see (Amoros, Burger, Corlette, Kotschik & Toledo, 1996) and the references in 3. B.

8. Low dimensional Riemannian geometry

If dimension 2 is wonderful we already saw above that the dimensions 3 and 4 have to be treated with care. For examples many finiteness theorems mentioned in I. C. 1 start to hold only in dimension 5 or more. A contrario in some instances one has very strong results in dimensions 3 and 4.

One of the interest of Riemannian geometry in low dimension is the hope to use Riemannian tools to solve topology and differential topology problems which are still haunting topologists. Recent reports are (Donaldson, 1996a), the books (Donaldson & Kronheimer, 1996b) and (Morgan, 1996). They present in particular the basic tools : Yang-Mills theory, twistors, anti-self-duality in dimension 4 and from (Seiberg & Witten, 1994) : the Spin_C theory, see the end of 6. B.

For the "best metric" approach based on functionals and their critical points, and the link with the geometrization program of Thurston, see the various references to Anderson we gave in chapter III, in particular (Anderson, 1997).

Finally we recall here the two Hamilton results, using the Ricci flow, for the Ricci curvature in dimension 3 (I. B. 2) and for the curvature operator in dimension 4 (I. B. 1).

9. Some generalizations of Riemannian geometry

A first generalization is that of **Riemannian manifolds with boundary**. Except for the η -invariant above we never mentioned them. For the general case one can see (Alexander, Berg, & Bishop, 1990) and the references there. For relation with positive scalar curvature see (Lawson & Michelsohn, 1984). For the cut-locus see (Alexander, Berg, & Bishop, 1993) and (Alexander & Bishop, 1997).

We mention one interesting and practical problem in this area. Assuming some kind of convexity on the boundary $\partial(M)$ of a Riemannian manifold (M, g) , one can define a distance $d(p, q)$ between points of $\partial(M)$. To which extent does this distance d as a map $\partial(M) \times \partial(M) \rightarrow \mathbf{R}^+$ determines the "inside" metric g (up to isometry-diffeomorphism of the inside). Thinking a while of earthquakes, tomography, X-rays and scanners one sees the practical importance of such a kind of problems. In an indirect way they were also met in I. B. 4 and in the "gap" results in TOP.5. References are (Otal, 1990b), (Michel, 1994), (Croke, 1991), (Arconstanzo, 1994) and its references, (Bourdon, 1996). See also the Besicovitch results at the end of TOP. 1. E.

Back to generalizations, the subject blew up recently and results are flowing now so that we will be very concise. Some cases appeared

before sporadically, today strong incentives are coming from understanding Riemannian geometry more deeply in various instances, e.g. that of I. C., namely to look at limits of sequence of Riemannian manifolds. Interesting generalizations of Riemannian geometry can be of different types since of course to look at general metric spaces would be too general to obtain reasonably serious results, for example to define objects playing the role of tangent vectors, angles, curvature. We will try to guide the reader through this new realm by suitable recent references and of course survey type ones when they exist. Of general type we know only of (Berestovskii & Nikolaev, 1993).

Orbifolds were introduced in (Thurston, 1978). Essentially they are quotients of Riemannian manifolds by isometries when one permits fixed points with moreover the isotropy group having to be finite. This is typical in the case of space forms. But they came in more subtle situations, for example as limit spaces of some subsets of Riemannian manifolds with Ricci curvature bounded from below as well as when working on Einstein manifolds, see (Anderson & Cheeger, 1992a), (Anderson, 1992b), (Anderson, 1994) and see also (Ballmann & Brin, 1995b). For orbifolds and quaternionic holonomy see (Galicki & Laswon, 1988), for orbifolds in the Kähler realm see (Tian, 1991).

Spaces with an isolated **conical singularity** look quite simple, but for applications it is important to see what is happening to the spectrum (see IV). More (and less) generally one can think to study Riemannian manifolds by approximating them by **PL** (piecewise linear) ones, metrically the pieces are all flat and "the curvature is concentrated (distributionally)" at the vertices, the edges, etc. The PL-manifolds are the very natural locally euclidean version of general Riemannian manifolds but with singularities. What is happening to the curvature and to various formulas is the object of (Cheeger, Müller, & Schrader, 1984; Cheeger, Müller, & Schrader, 1986), see (Lafontaine, 1985/86) for an expository text. This is in fact a very subtle subject. We saw one case in **TOP. 1. B** for the isoperimetric profile of nonpositive curved manifolds.

An extensive study of the spectral behavior of singular spaces was done by Cheeger in a long series of papers: we refer to (Cheeger, 1983) and its bibliography. This work came out of the proof in (Cheeger, 1979) of Ray-Singer conjecture for the Reidemeister torsion (see **TOP. 6. A**). One of its byproducts was in Analysis,

namely results on the diffraction of waves meeting some kind of obstacles, another one was an explicit solution of the heat equation on standard spheres ; for both see (Cheeger & Taylor, 1982).

Alexandrov spaces is a wonderful class because they enjoy two simultaneous properties. First they come naturally in the boundary of the space of all Riemannian manifolds whose curvature is bounded (only) from below. Second they share many properties with (smooth) Riemannian manifolds. There are many texts of more or less of survey type : (Perelman, 1994c), (Berestovskii & Nikolaev, 1993), (Reshetnyak, 1993), (Burago, Gromov, & Perelman, 1992b) , (Otsu & Shioya, 1994) and (Yamaguchi, 1992). We now give a extremely brief account.

These spaces appeared first in 1948 in the work of A. D. Alexandrov's on convex surfaces, but in 1957 Alexandrov made a more systematic study. Then came the founding paper (Burago, Gromov & Perelman, 1992b) and between was (Gromov, Lafontaine & Pansu, 1981a).

In I. C we had the desire to look at the boundary of suitable subsets of the set of all Riemannian manifolds (compactness and convergence results). The description was possible (to some extent) for bounded curvature, say $-1 \leq K \leq 1$. But for $K \geq k$ we had only finiteness theorem and no description of collapsing. The solution lies in the theory of Alexandrov spaces with curvature bounded below.

On the one hand Alexandrov spaces of curvature bounded below are defined very simply. They are first locally compact metric spaces which are **length spaces**, namely the metric coincides with the infimum of the length of curves joining the two given points. Note here a question of terminology. In (Gromov, Lafontaine & Pansu, 1981a) the wording was "length space", in (Gromov, 1987a) "geodesic space" if moreover any pair of points can be joined by at least one shortest path. Busemann in (Busemann, 1955) was speaking of "intrinsic" metrics. Here such a length space is called Alexandrov with curvature $\geq k$ if it enjoys (besides completeness of course) everywhere locally a Toponogov comparison theorem (see I. A. 2) with the standard space form of constant curvature equal to k . One can then prove a global Toponogov theorem but also much more. Only with such a mild condition one has a notion of angles, a notion of a tangent cone which is itself an Alexandrov

space of curvature ≥ 0 and can be defined in many equivalent ways. The points where this cone is not euclidean are the singular points and their set is of Hausdorff codimension at least 2 so that our spaces are almost everywhere locally Riemannian manifolds. Note that the singular set can be everywhere dense, just think of a suitable limit of convex polyhedra. Important for generalizations of Riemannian results in the spirit of I. A and B is that the notion of critical point for distance functions is well defined and that regions without critical points can be deformed one another.

On the other hand, the definition of curvature $\geq k$ being purely metric within the set (MET, d_{G-H}) introduced in I. B, then limit points of Riemannian manifolds with $K \geq k$ are automatically Alexandrov spaces with curvature $\geq k$. Hence the results above constitute the answer to our main query and any further results on those spaces will be a progress in the study of the boundary of the set of Riemannian manifolds with $K \geq k$. The following recent references show how much the topic is active. (Grove, 1992) extends some results of I. A, (Shiohama & Tanaka, 1992) studies cut-loci and distance spheres, (Yamaguchi, 1992) generalizes the collapsing structure result of I. C, various type of results generalizing those in I can be found in (Petersen, 1996e), for example the bound for Betti numbers (see I. B. 2) of nonnegatively curved manifolds (see the references there).

In (Cheeger & Colding, 1997a), (Cheeger & Colding, 1997c) and (Cheeger & Colding, 1998) many of the above results are generalized to "limit spaces with Ricci curvature bounded below". These spaces occur of course as limit spaces, in the Gromov-Hausdorff space, of Riemannian manifolds with the same Ricci condition.

In the opposite direction to the above one can look for a large class, but not too, of spaces generalizing Riemannian manifolds of negative curvature. A good notion is that of **hyperbolic spaces**, which were introduced and studied in (Gromov, 1987a) (not to be confounded with hyperbolic space forms which are only a very very special case). This is a game where the interesting case is when the "curvature" now is bounded above. For the spirit see I. B. 4. After the founding paper (Gromov, 1987a) then the topic became an entire world. We just mention references : (Gromov,

1993), (Ghys & de la Harpe, 1990) and the book (Bridson & Haefliger, 1997).

Here the analogous condition of Alexandrov spaces of curvature bounded below for an upper bound of the curvature is the $CAT(k)$ condition. Say that we work in simply connected spaces. The $CAT(0)$ condition is that one has, for every triple of points, the triangle inequality discovered by Cartan and rediscovered by Preissmann (see O. B and C). For $CAT(k)$ it is the same condition but the comparison is no longer with euclidean spaces but with hyperbolic spaces of curvature $-k$. The CAT initials stand for Cartan, Alexandrov, Toponogov. One of the difficulty in hyperbolic space is that they can have branch points since they can even be graphs. Think also of a one sheet hyperboloid converging toward the total cone and look at the origin. A contrario one sheet of the two sheet hyperboloid associated with the same asymptotic cone will converge with a conical point (of positive curvature), a nice Alexandrov space.

Examples of them are to be taken among the PL-manifolds (piecewise linear i.e. euclidean). For PL-surfaces the $CAT(0)$ condition is what you expect : at every vertex the sum of the angles of the triangles meeting there has to be larger than or equal to 2π . For higher dimensions is not too easy to explicit this condition. The general $CAT(k)$ condition goes the reverse way from the Alexandrov case. What is happening to the curvature and to various formulas was mentioned above, see also (Bourdon, 1996).

Other recent newcomers in generalizing Riemannian geometry are the **Carnot-Caratheodory metric spaces**. For them one can say in a caricatural sense that singularities are at every point and very strong. They are the subject of the book (Bellaïche & Risler, 1996). Precisely one is given in some manifold M^d at every point m some vector subspace $V(m)$ of the tangent space $T_m M$ of a constant dimension $k < d$. On those subspaces one puts an euclidean structure. The metric is then defined as the distance $d(m,n)$ between two points being the infimum of the length of the curves whose speed vector at every of its points q should always belong to $V(q)$. Hence the other name for the topic : **sub-riemannian geometry**. This distance can be infinite but it is always finite when the distribution $V(\cdot)$ is wild enough, that is to say "completely non integrable". Carnot-Caratheodory spaces are basic in control theory since in this setting only special ways are

permitted due to various restrictions imposed by practical situations. In street traffic the geodesics are what you have to do to park and unpark in a narrow slot. See the recent (El Alaoui, Gauthier & Kupka, 1996) and the introductory text (Pelletier & Valère Bouche, 1992). For the naive reader or a naive PDE person (Bryant & Hsu, 1993) is extremely informative.

Finsler geometry I have been keeping for the middle of this section. This is not historically correct. As very well explained in (Spivak, 1970) where he analyses Riemann dissertation, Riemann in fact generalized euclidean geometry by introducing Finsler manifolds. Those are differentiable manifolds M where at every point m is given some Banach structure $g(m)$ (i.e. a convex body symmetric around the origin of $T_m(M)$). With this lengths and hence a metric can be as well defined. Then Riemann has that prophetic sentence : "we will now stick to the case of ellipsoids (quadratic forms) because if not the computations would become very complicated". One can also relax the symmetry condition, e.g. in the Katok type examples mentioned in V. A.

There are few geometrical objects as natural as Finsler manifolds, e.g. in any kind of "Mechanics" as soon as the energy is no longer quadratic. However Finsler geometry destiny was not that of Riemannian geometry. It is clear I am biased with the subject but I think that, before very recently, there were very few deep typically Finsler results and not already Riemannian. Or, when of course it makes sense, Riemannian and not true for Finsler. Up to very recently we knew two of them : Leon Green and Burago-Ivanov in V. B . One counterexample was in (Busemann, 1955) where are built Finsler tori without conjugate points and non flat. The other was in (Skorniakov, 1955), where it is proved that any system of curves on RP^2 satisfying the axioms of projective geometry (and some obviously needed reasonable smoothness) is that of the geodesics of some Finsler (non Riemannian in general thanks to Desargues theorem) structure, Green's theorem figures in V. B and in TOP.4. Interesting is (Vérovic, 1996) where it is proved that the result of (Besson, Courtois & Gallot, 1995a) (quoted in I. B. 4, III and V) can be true or false for Finsler manifolds. "Yes or no" is because Finsler manifolds can be given different notions of measure in their unit bundle. Oppositely most of the results in the "filling paper" (Gromov, 1983a) are in fact valid for the Finsler case.

Here is a personal comment. Finsler spaces look more general than Riemannian metrics. They are in some sense. But in another one they are more special. In an affine (finite dimensional) real vector space a (symmetric) convex body has privileged directions (points) e.g. those where the affine curvature is critical, etc. Oppositely an ellipsoid is **infinitesimally isotropic**, all the directions (the points) are equivalent. This is for us one reason for the importance of Riemannian geometry.

There are of course many places to find information on Finsler manifolds. A recent reference is (Bao, 1995). The Note (Foulon, 1997) is a perfect blitz introduction to the modern point of view, see the formula at the end of V. We mention the fact that Finsler manifolds have a curvature, but what implies constancy of this curvature is not clear today. The result of the preceeding Note is again a rigidity result for compact negative curvature manifold. The positive case remains mysterious, but (Bryant, 1996) shows some light on it. (Alvarez, Gelfand, & Smirnov, 1997) characterizes Finsler metrics for which the geodesics are the straight lines (Hilbert's fourth problem).

Riemannian foliations is also a topic in itself as well as totally geodesic foliations of Riemannian manifolds : books are (Tondeur, 1988) and (Godbillon, 1991). For relation with scalar curvature see $1\frac{7}{8}$ of (Gromov, 1996) and see also in (Connes, 1994) a new viewpoint on foliations.

Lorentzian manifolds are completely similar to Riemannian ones. They are 4-dimensional manifolds with everywhere a definite quadratic form but of signature $(+, +, +, -)$. More generally **pseudo-riemannian** geometry is any dimensional and with definite quadratic forms of any signature. At the beginning they have many things in common. This is definitively misleading at the long range as was remarked by the physicist C. N. Yang, with the help of a beautiful picture : (Yang, 1980). The picture can also be seen on page 11 of (Besse, 1987). The golden triangle is mainly what is common to both objects. Then questions and results diverge. Lorentzian "geometry" is deriving its inspiration mainly from General Relativity. Books on Lorentzian geometry are : (Beem & Ehrlich, 1981), (Hawking & Ellis, 1973), (Misner, Thorne, & Wheeler, 1973), (O'Neill, 1983), (Sachs & Wu, 1977). A special mention

should be given to (d'Ambra, 1988) (see also (d'Ambra & Gromov, 1991)) where it is proved that the isometry group of a compact and simply connected Lorentzian manifold is itself compact (presently analyticity is needed but this looks only technical). For surfaces (Weinstein, 1996) studies a Lorentzian conformal concept which is analogous to that of Riemann surfaces as opposed to that Riemannian surfaces. Of Riemannian flavor is and (Christodoulou & Kleinerman, 1993) , see the expository (Bourguignon, 1990/91). Recently pseudo-riemannian geometry appeared in various contexts, see for example (Benoist, Foulon, & Labourie, 1992) and (Kühnel & Rademacher, 1995).

Infinite-dimensional Riemannian geometry is still very young. The book (Lang, 1972) was the first to treat Riemannian geometry systematically in an infinite dimensional setting. Then appeared (Klingenberg, 1982) and now the second edition (Lang, 1996) and we suggest strongly the reader to look at it. See also (Bourbaki, 1969) and (Gil-Medrano & Michor, 1991). We mention only few directions.

The **first** is to use suitable embeddings of Riemannian manifolds in the standard Hilbert space. This can be done for example by using suitably normalized eigenfunctions of the Laplacian , see (Besson, Courtois, & Gallot, 1991) and (Bérard, Besson, & Gallot, 1997) for the setting and applications to finiteness results.

One can also embed any Riemannian manifold M in the Banach space of the continuous functions $C^0(M)$ by the "trivial" map made of the various distance functions, namely $m \rightarrow d(m, \cdot)$. This childish map turned out to be basic in the systolic inequality of (Gromov, 1983a), see **TOP. 1. E** .

The **second** most natural desire is to define a notion of infinite-dimensional Riemannian manifolds, starting with some infinite-dimensional differentiable manifold M (whatever it means) and by endowing every tangent space $T_m M$ with some Hilbertian structure. This seems easy. As we see things, it is in fact difficult, at least today, for the following fact : for a simple Hilbertian structure one gets in general non-complete metric spaces. With more complicated ones (typically in the Sobolev range) one has completeness but then the geodesics seem to have no good and/or useful geometrical properties, their equation is too complicated. In (Ebin, 1970), (Ebin, 1972) one uses such a structure to solve some fluid mechanics equation, see also (Ebin & Marsden, 1970). The text

(Gil-Medrano & Michor, 1991) addresses the question to make a manifold with the set of all Riemannian metrics on a given manifold.

In the Appendix 2 of (Arnold, 1978) the group of diffeomorphisms of a manifold is given an (infinite-dimensional) Riemannian structure which is proved to be always of negative curvature.

For **path-space** Riemannian geometry see (Stroock, 1996).

Finally special attention should be given to (Connes, 1995-6) : this is the beginning of a complete program to put most (if not all) geometries into a very general frame. The frame is that of algebras of operators and is called "noncommutative geometry". In this frame, according to various extra suitable axioms one can recover almost any kind of geometry, including of course Riemannian ones. The next in the program will be to generalize every concept one could wish, e.g. curvature. For the metric itself this is done and we saw in **TOP. 6. B** that one can compute the distance with spinors. See also **III. C** for the Hilbert functional in this context.

10. Submanifolds

Among other motivations, Riemannian geometry came out when trying to generalize the geometry of surfaces in $\mathbf{R}^3$. There is no reason not to consider the geometry of submanifolds inside a given Riemannian manifold. Because euclidean geometry is hereditary for subspaces, it follows that Riemannian geometry is also **hereditary** for submanifolds (but of course the metric is not the induced one in the rough metric sense), this is one reason more why Riemannian geometry is important. We do not know of any systematic survey for this topic, even in the special and much studied case of submanifolds of euclidean spaces. Few books give the general equations for a submanifold of any codimension in a Riemannian manifold (the so-called Gauss and Codazzi-Mainardi equations). Among them : chapter VII of (Kobayashi & Nomizu, 1963-1969), chapter XX (see section 20. 14. 8) of (Dieudonné, 1972) and chapter 7 of (Spivak, 1970) present the foundations of the theory of submanifolds in a Riemannian manifold. But there is a huge number of local and of global results. Recent surveys are : (Terng, 1990), (Terng & Thorbergsson, 1995), (Dillen & Vestraelen, 1992), (Palais & Terng, 1988), see also in (Willmore, 1993). In particular **isoparametric hypersurfaces** deserve a special mention as

relating different topics. The founding text is (Cartan, 1939), for isoparametric submanifolds in an infinite dimensional setting, see (Heintze & Liu, 1997).

The naive geometer will ask what in Riemannian geometry is generalizing affine (euclidean) and/or projective subspaces. The basic remark is that a general Riemannian manifold admits **no** submanifold which is stable under geodesy (i.e. local geodesics in it are geodesics in the ambient one and this is certainly not the case in general). Such submanifolds are called **totally geodesic**. Exceptions are extremely few and among them are the symmetric spaces of rank ≥ 2 , see II. C. We saw in I. B. 4 how the fact of having "flats" is very strong in some instances. An extreme case occupied people in the early century : the axiom of free mobility. Things were completely clarified by Elie Cartan in (Cartan, 1928a)(one can also see the second edition (Cartan, 1951)) where he proved that only space forms of constant curvature satisfy it.

A most naive problem is this : what is the convex enveloppe of k points in a Riemannian manifold of dimension $d \geq 3$? Even for three points and $d \geq 3$ the question is completely open (except when the curvature is constant). A natural example to look at would be CP^2 , because it is symmetric but not of constant curvature. The only text we know of addressing the question is (Bowditch, 1994).

A.THE CASE OF SURFACES IN R^3

They were the main object of differential geometers in the preceeding century, the bible was (Darboux, 1887, 1889, 1894, 1896) or (Darboux, 1972). Many things were put in a modern way in (Spivak, 1970), a survey is (Burago and Zalgaller 1992a). Even for compact surfaces the subject enjoyed many results in our half century. A famous recently open problem is Willmore's conjecture, see (Willmore, 1993).

In 1955 Alexandrov proved that constant mean curvature embedded surfaces have to be round spheres, and this for any topological type. For the simply connected cases (topological spheres) this was Hopf "old" result (Hopf, 1951). Besides Burago-Zalgaller book, one could see (Hopf, 1983), (do Carmo, 1976). These surfaces are physically the **soap-bubbles** and Hopf proved that they are really round spheres. The left open problem was about the possibility of immersions, which are impossible for the sphere

topological type. An event was (Wente, 1986) who constructs immersed tori of constant mean curvature. This started a field still not finished, see (Pinkall & Sterling, 1989). A recent point of view is (Kamberov, 1997) where spinors come into the picture.

In higher dimensions, Alexandrov's theorem is still valid for embedded hypersurfaces : only rounds spheres have constant mean curvature. But for immersions things are radically different : there are many non spherical immersed topological spheres S^{d-1} of constant mean curvature in $\mathbf{R}^d$. See (Eells & Rato, 1993) for references, the topic is directly related tharmonic maps.

An old result of Hilbert says that there is no immersed (embedding is not even needed for the conclusion) surface isometric to the hyperbolic plane. This means constant Gauss curvature and completeness. This explains why hyperbolic geometry could not be discovered in $\mathbf{R}^3$. This result was generalized in the 50's by Efimov in a deep paper, see (Klotz Milnor, 1972) and (Burago & Zalgaller, 1992a) : there is no immersion in $\mathbf{R}^3$ of a complete surface of curvature $K \leq a < 0$. But there is still a deep (naive) problem : is there any complete and negative curvature surface staying in a bounded region of $\mathbf{R}^3$? Some people call the preceeding question "Hadamard's conjecture" because Hadamard took for granted the non existence. An analogous conjecture, Calabi-Yau's one, was that it cannot exist a complete minimal surface staying in a bounded region. In (Nadirashvili, 1996) both conjectures are killed, one constructs a surface living in a bounded region, which is at the same minimal and of negative curvature. The tool is Weierstrass formulas for minimal surfaces.

Questions which are still not finished concern the isometric non trivial deformations of surfaces in $\mathbf{R}^3$. First globally : all non deformation results are for convex surfaces and non congruence examples are C^∞ surfaces. Locally it looks obvious that any piece of surface can be deformed (and with a lot of parameters : take a tennis balls with some piece removed). In fact the question is unbelievably hard, we refer to the survey in (Burago & Zalgaller, 1992a). We mention that they are examples of pointed surfaces which cannot be deformed no matter how small is the neighborhood of the point. See also Bleecker's result at the end of TOP. 2.

A special mention should be given to the case of zero mean curvature namely the minimal surfaces (Plateau's problem). This old topic underwent tremendous developments. There are many good survey and entire books on the topic. The latest one is (Dierkes, Hildebrandt, Küster, & Wohlrab, 1992), others are (Nitsche, 1996), (Jost, 1991), (Struwe, 1988) ; see also GMT in C below and in O. F.

B. HIGHER DIMENSIONS

An important remark is in order, which is folk and which we owe to Lafontaine. The topic A is important not only for historical reason but because of this. When $d \geq 4$ take in $\mathbf{R}^d$ some hypersurface which is generic. Then it is trivial that the sectional curvature (hence the isometry type) determines the second fundamental form. Classically this implies congruence (this is a pure local statement). The proof is : in a basis which diagonalizes the second fundamental form, the Gauss equations imply that, for the sectional curvature, one has $K(e_i, e_j) = \eta_i \eta_j$ where the η_i are the principal curvatures (this is exactly the generalization of the theorem egregium in O. A). And now, if I know the three products ab , bc and ca , then I know a , b , c as soon as none is 0. So that in the generic case the metric of an hypersurface determines automatically also its second fundamental form. The generalized Gauss-Codazzi equations show that together the metric and the second fundamental form determine completely the embedding (see any of the books quoted at the begining of this section). So that, in some sense, there is no "congruence versus isometry problem" left. Of course the above philosophy this applies also to hypersurfaces in general Riemannian manifolds M^d as soon as $d \geq 4$.

Things are completely different when $d=3$, the Gauss curvature (hence the isometry type) yields only the product of the two principal curvature. Then there is a huge place for a fascinating game, not finished as just seen above. Let us look for example at rigidity : two compact strictly convex hypersurfaces which are abstractly isometric are congruent (deduced by a global isometry of the whole $\mathbf{R}^d$). For $d > 3$ it follows from the above folk remark, but, when $d=3$, the result remains but the proof is much more sophisticated and is due to Cohn-Vossen and Herglotz : for references and more see (Berger & Gostiaux, 1988), section 11.4 and/or (Klingenberg, 1978b), 6.2.8 .

This being realized, there remain many topics in the field, specially global questions and we already mention some of them above. **Integral geometry** is very interesting, but works only (with one exception to come) in euclidean spaces and space forms. See the survey (Schneider & Wieacker, 1993) and add the classic (Santalo, 1976), in particular Chern's kinematic formula. In Riemannian manifold integral geometry works for hypersurfaces and geodesics starting from them, it is basic for Croke's local isoperimetric inequality of **TOP. 1. C** and the isoperimetric inequality for nonpositive curvature manifolds in **TOP. 1. A** : see (Croke, 1980) and (Croke, 1984b).

C. GEOMETRIC MEASURE THEORY AND PSEUDO-HOLOMORPHIC CURVES

Submanifolds enjoying various strong geometric properties are basic in some instances. Two seem to be important today. First the dramatic appearance of **GMT** (geometric measure theory, see **O. F**) with its harvest of results provided Riemannian geometers with almost all the existence theorems they could dream of forgetting **minimal** (or constant mean curvature) objects. This was used above, in Schoen-Yau approach for positive scalar curvature in **I. B. 3** and for the isoperimetric profile in **TOP. 1. B**. **GMT** is an extremely hard subject, in the founding book (Federer, 1969) as well as in many other articles on the subject, it is very hard to find one's way through in a reasonable time. We suggest to read first (Morgan 1988), then (Simon, 1983).

The second tool appeared in the pioneer (Gromov, 1985). A recent survey book is (Audin & Lafontaine, 1994). Although those pseudo-holomorphic curves are mainly used in symplectic geometry, one meet them in Riemannian geometry, see for example the article of Labourie in the preceeding book and (Gromov, 1992a).

Bibliography

- Abrahamov (1951). On the topological invariants of Riemannian spaces obtained by integration of tensor fields. Doklady Akad. Nauk. SSSR, 81, 125-128.
- Abresch, U. (1985). Lower curvature bounds, Toponogov's theorem, and bounded topology. Ann. scientif. Ecole norm. sup., 18, 651-670.
- Abresch, U. (1988) Über das glatten Riemannscher Metriken. thesis, Bonn.
- Abresch, U. (1990b). Endlichkeitssätze in der Riemannschen Geometrie. Jber. d. Dt. Math-Verein., Jubiläumstagung, 152-176.
- Abresch, U., & Gromoll, D. (1990a). On complete manifolds with nonnegative Ricci curvature. J. of the Amer. math. Soc., 3, 355-374.
- Abresch, U., & Meyer, W. (1994b). Pinching below $1/4$, injectivity radius, and conjugate radius. J. differential geometry, 40, 643-691.
- Abresch, U., & Meyer, W. (1996b). A sphere theorem with a pinching constant below $1/4$. J. of differential geometry, 44, 214-261.
- Abresch, U., & Meyer, W. (1997). Injectivity radius estimates and sphere theorems. In Comparison geometry, MSRI publications, Cambridge University Press.
- Abresch, U., & Schroeder, V. (1996a). Analytic manifolds of nonpositive curvature.
- Accola, R. (1960). Differential and extremal lengths on Riemann surfaces. Proc. Nat. Acad. Sci. USA, 46, 540-543.
- Alekseevskii, D. (1968). Riemannian spaces with exceptional holonomy groups. Functional analysis and applic., 2, 97-105.
- Alexander, S., & Bishop, R. (1997). Thin Riemannian manifolds with boundary. Math. Annalen.
- Alexander, S., Berg, D., & Bishop, R. (1990). The distance geometry of Riemannian manifolds with boundary. In R. Greene & S.-T. Yau (Ed.), Differential geometry, 54, prt 3 (pp. 31-36). Los Angeles: American math. soc. 1993
- Alexander, S., Berg, D., & Bishop, R. (1993). Cut-loci, minimizers and wave fronts in Riemannian manifolds with boundary. Michigan math. j., 40, 229-237.

- Allendoerfer, C., & Weil, A. (1943). The Gauss-Bonnet theorem for Riemannian polyhedra. Trans. Americ. math. Soc., 53, 101-129.
- Aloff, S., & Wallach, N. (1975). An infinite family of distinct 7-manifolds admitting positively curved Riemannian structures. Bull. of Americ. math. Soc., 81, 93-97.
- Alvarez, J., Gelfand, I., & Smirnov, M. (1997). Crofton densities, symplectic geometry and Hilbert's fourth problem No. Univ. cath. Louvain, Belgique.
- Ambrose, W. (1956). Parallel translation of Riemannian curvature. Annals of math., 64, 337-363.
- Ambrose, W., & Singer, I. (1958). On homogenous Riemannian manifolds. Duke math. j., 25, 647-669.
- Amoros, J., Burger, M., Corlette, K., Kotschik, D., & Toledo, D. (1996). Fundamental groups of compact Kähler manifolds. American math. soc.
- Andersen, J., Dupont, J., Pedersen, H., & Swann, A. (Ed.). (1997). Geometry and physics. Marcel Dekker.
- Anderson, M. (1990a). Metrics of positive Ricci curvature with large diameter. Manuscripta math., 68, 405-415.
- Anderson, M. (1990c). Convergence and rigidity of metrics under Ricci curvature bounds. Inventiones math., 102, 429-445.
- Anderson, M. (1990d). Short geodesics and gravitational instantons. J. of differential geometry, 31, 265-275.
- Anderson, M. (1990e). Degeneration of metrics with bounded curvature and applications to critical metrics of Riemannian functionals. In R. Greene & S.-T. Yau (Ed.), Differential geometry, part 3 (pp. 53-79). Los Angeles: American math. soc. 1993
- Anderson, M. (1991b). Remarks on the compactness of isospectral sets in low dimensions. Duke math. j., 63, 699-711.
- Anderson, M. (1992b). The L^2 structure of moduli spaces of Einstein metrics on 4-manifolds. Geometric and functional analysis, 2, 29-89.
- Anderson, M. (1992c). Hausdorff perturbations of Ricci-flat manifolds and the splitting theorem. Duke math. j., 68, 67-82.
- Anderson, M. (1994). Einstein metrics and metrics with bounds on Ricci curvature. In International congress of mathematicians, 1 (pp. 443-452). Zürich: Birkhäuser.
- Anderson, M. (1996a). Scalar curvature and geometrization conjectures for 3-manifolds. In Comparison geometry (pp. 47-80). MSRI publications, Cambridge university press.
- Anderson, M. (1996b). Extrema of curvature functionals on the space of metrics in 3-Manifolds. In Calculus of variations and PDE, .

Anderson, M. (1997). The static vacuum Einstein equations and degenerations of Yamabe metrics on 3-manifolds. .

Anderson, M., & Cheeger, J. (1991a). Diffeomorphism finiteness for manifolds with Ricci curvature and $L^{n/2}$ curvature bounded. Geometric and functional Analysis, 1, 231-251.

Anderson, M., & Cheeger, J. (1992a). C^α compactness for manifolds with Ricci curvature and injectivity radius bounded below. J. differential geometry, 35, 265-281.

Anderson, M., & Cheeger, J. (1992a). C^α compactness for manifolds with Ricci curvature and injectivity radius bounded below. J. differential geometry, 35, 265-281.

Andersson, S., & Lapidus, M. (Ed.). (1997). Recent progress in inverse spectral geometry. Birkhäuser.

Anosov, D. (1967). Geodesic flows on closed Riemannian manifolds with negative curvature. Proc. Steklov Inst., 90.

Anosov, D. (1983). On generic properties of closed geodesics. Math. USSR Izv., 21, 1-29.

Arconstanzo, M. (1994). Des métriques finslériennes sur le disque à partir d'une fonction distance entre les points du bord. Commentarii math. helvetici, 69, 229-248.

Arnold, V. (1978). Mathematical methods of classical mechanics. Springer.

Arnold, V. (1988). Geometrical methods in the theory of ordinary differential equations, second edition. Springer.

Atiyah, M. (1979). Geometry of Yang-Mills fields. Pisa: Academia naz. dei Lincei and Scuola norm. superiore.

Atiyah, M., & Singer, I. (1963). The index of elliptic operators on compact manifolds. Bulletin of the A.M.S., 69, 422-433.

Atiyah, M., Bott, R., & Patodi, V. (1973). On the heat equation and the index theorem. Inventiones math., 19, 279-330.

Atiyah, M., Donnelly, H., & Singer, M. (1983). Eta-invariants, signature defects of cusps, and the value of L-functions. Annals of math., 118, 131-177.

Atiyah, M., Hitchin, N., & Singer, I. (1978). Self-duality in four-dimensional geometry. Proc. royal soc. London, 362, 425-461.

Atiyah, M., Patodi, V., & Singer, I. (1975-1976). Spectral asymmetry and Riemannian geometry I, II, III. Math. Proc. Cambridge phil. soc., 77-78-79, 43-69, 405-432, 71-99.

Aubin, T. (1970). Métriques riemaniennes et courbure. J. of differential geometry, 4, 383-424.

- Aubin, T. (1976). Equations du type de Monge-Ampère sur les variétés kählériennes compactes. C. R. Acad. Sc. Paris, 283, 119-121.
- Aubin, T. (1982). Nonlinear analysis on manifolds. Monge Ampère equations. Springer.
- Audin, M., & Lafontaine, J. (Ed.). (1994). Pseudo-holomorphic curves in symplectic geometry. Birkhäuser.
- Avakumovich, V. (1956). Über die Eigenfunktionen auf geschlossenen Mannigfaltigkeiten. Mathematische Zeitschrift, 65, 327-344.
- Babenko, I. (1992). Asymptotic invariants of smooth manifolds. Russian acad. sc. Izv. math., 41, 1-38.
- Babenko, I., & Katz, M. (1997). Systolic softness of orientable manifolds. Annales scientifiques Ecol. norm. sup.
- Babenko, I., Katz, M., & Suciu, A. (1998).
- Babich, & Lazutkin, V. (1967). On eigenfunctions concentrated at a closed geodesic. In Problems in math. phys. N2 Leningrad University.
- Balian, R., & Bloch, C. (1972). Distribution of eigenfrequencies for the wave equation in a finite domain. Annals of Physics, 69, 401-447.
- Ballmann, W. (1995a). Lectures on spaces of nonpositive curvature. Birkhäuser.
- Ballmann, W., & Brin, M. (1995b). Orbihedra of nonpositive curvature. Publications math. IHES, 82, 169-209.
- Ballmann, W., & Wojtkowski, M. (1989). An estimate for the measure theoretic entropy of geodesic flows. Ergodic theory and dynamical systems, 9, 271-279.
- Ballmann, W., Gromov, M., & Schroeder, V. (1985). Manifolds of nonpositive curvature. Birkhäuser.
- Ballmann, W., Thorbergsson, G., & Ziller, W. (1982). Closed geodesics on positively curved manifolds. Annals of math., 116, 213-247.
- Ballmann, W., Thorbergsson, G., & Ziller, W. (1983). Existence of closed geodesics on positively curved manifolds. J. of differential geometry, 18, 221-252.
- Bando, S., & Mabuchi, T. (1987). Uniqueness of Einstein Kähler metrics modulo connected group actions. Advanced studies in pure math., 10, 11-40.
- Bangert, V. (1985). Geodätische Linien auf Riemannschen Mannigfaltigkeiten. Jahrbuch der deutsche math., 87, 39-66.

- Bangert, V. (1988). Mather sets for twist maps and geodesics on tori. In U. Kirchgraber & H. Walther (Eds.), Dynamics reported (pp. 1-56). Teubner, Wiley.
- Bangert, V. (1993). On the existence of closed geodesics on the two-sphere. International J. math., 4, 1-10.
- Bao, D. (1995). Finsler geometry. In D. Bao, S.-S. Chern, & Z. Shen (Ed.), Joint summer research conference on Finsler geometry, 196. Seattle, Washington: American math. society. 1996
- Bär, C. (1995). Metrics with harmonic spinors No. 15/1995). Math.Fakultät Freiburg.
- Batson (1992). Quaternionic complexes. J. geometry and physics, 8, 29-52.
- Bavard, C. (1987b). Courbure presque négative en dimension 3. Compositio math., 63, 223-236.
- Bazaikin, J. (1995).
- Beauville, A. (1983). Variétés Kähleriennes dont la première classe de Chern est nulle. J. differential geometry, 18, 755-782.
- Bedford, E., d'Angelo, J., Greene, R., & Krantz, S. (Ed.). (1991). Several complex variables and complex geometry. Santa Cruz 1989. Proc. symposia in pure math. n°52, American math. Soc.
- Beem, J., & Ehrlich, P. (1981). Global Lorentzian geometry. Dekker.
- Bellaïche, A., & Risler, J.-J. (1996). Sub-Riemannian geometry. Birkhäuser.
- Bemelmans, J., Min-Oo, & Ruh, E. (1984). Smoothing Riemannian metrics. Math. Zeitschrift, 188, 69-74
- Benedetti, R., & Petronio, C. (1992). Lectures on hyperbolic geometry. Springer.
- Benjamini, I., & Cao, J. (1996). Examples of simply- connected Liouville manifolds with positive spectrum. Differential geometry and its applications, 6, 31-50.
- Benoist, Y., Foulon, P., & Labourie, F. (1992). Flots d'Anosov à distributions stables et instables différentiables. J. american math. soc., 5, 33-74.
- Bérard, P. (1986). Spectral geometry : direct and inverse problems. Springer.
- Bérard, P. (1988). From vanishing theorems to estimating theorems ; the Bochner technique revisited. Bull. Amer. Math. Soc., 19, 371-406.
- Bérard, P. (1989). Variétés riemanniennes isospectrales non isométriques. In Séminaire Bourbaki, 177-178 (pp. 127-154). Paris: Soc. math. France.
- Bérard, P., Besson, G., & Gallot, S. (1985). Sur une inégalité isopérimétrique qui généralise celle de Paul Lévy. Inventiones math., 80, 295-308.

- Bérard, P., Besson, G., & Gallot, S. (1997). Embedding Riemannian manifolds by their heat kernel. .
- Bérard Bergery, L. (1976). Les variétés riemanniennes homogènes simplement connexes à courbure strictement positive. J. math. pures et appl., 55, 47-68.
- Bérard Bergery, L. (1982). Sur de nouvelles variétés riemanniennes d'Einstein. Publications de l'Institut Elie Cartan (Nancy), 4, 1-60.
- Berestovskii, V., & Nikolaev, I. (1993). Multidimensional generalized geometry. In Encyclopedia of math. sciences Geometry IV, 70, Springer.
- Berger, M. (1953). Sur les groupes d'holonomie des variétés riemanniennes. Bulletin Soc. math. France, 83, 279-330.
- Berger, M. (1960). Les variétés riemanniennes $(1/4)$ -pincées. Ann. Scuola. norm. sup. Pisa, 14, 161-170.
- Berger, M. (1961). Les variétés riemanniennes homogènes normales simplement connexes à courbure positive. Ann. scuola norm. sup. Pisa, 15, 179-2546.
- Berger, M. (1972). Du côté de chez Pu. Ann. Scient. Ec. Norm. Sup., 5, 1-44.
- Berger, M. (1980). Une borne inférieure pour le volume d'une variété riemannienne en fonction du rayon d'injectivité. Ann. Inst. Fourier Grenoble, 30, 259-265.
- Berger, M. (1983). Sur les variétés riemanniennes pincées juste au-dessous de $1/4$. Ann. Institut Fourier, 33, 135-150.
- Berger, M. (1993a). Systoles et applications selon Gromov. In Séminaire Bourbaki (pp. 279-230), Astérisque 216, Société Mathématique de France.
- Berger, M. (1993b). Encounter with a geometer : Eugenio Calabi. In P. de Bartolomeis, F. Tricerri, & E. Vesentini (Ed.), Conference in honour of Eugenio Calabi, Manifolds and geometry (pp. 20-60). Pisa: Cambridge University Press.
- Berger, M., & Gostiaux, B. (1988). Differential geometry : manifolds, curves and surfaces. Springer.
- Berger, M., Gauduchon, P., & Mazet, E. (1971). Le spectre d'une variété riemannienne. Springer.
- Berline, N., Getzler, E., & Vergne, M. (1992). Heat kernels and Dirac operators. Springer.
- Berndt, J., Tricerri, F., & Vanhecke, L. (1995). Generalized Heisenberg groups and Damek-Ricci harmonic spaces. LNM 1598, Springer.

- Besicowitch, A. (1952). On two problems of Loewner. J. London math. soc., 27, 141-144.
- Besse, A. (1978). Manifolds all of whose geodesics are closed. Springer.
- Besse, A. (1987). Einstein manifolds. Springer.
- Besse, A. (1994). Some trends in Riemannian geometry. In M. Artin, H.-P. Kraft, & R. Remmert (Eds.), Duration and change : fifty years at Oberwolfach (pp. 71-105). Springer.
- Besson, G. (1996b). Volumes and entropies. In M. Lovric, M. Min-Oo, & M. Wang (Eds.), Riemannian geometry (pp. 1-22). American math. soc.
- Besson, G., Courtois, G., & Gallot, S. (1995a). Entropies et rigidités des espaces localement symétriques de courbure strictement négative. Geometric and funct. Analysis, 5, 731-799.
- Besson, G., Courtois, G., & Gallot, S. (1995b). A simple proof of the rigidity and the minimal entropy theorem No. 314). Institut Fourier, Grenoble.
- Biquard, O., & Gauduchon, P. (1996). La métrique hyperkählérienne des orbites coadjointes de type symétrique d'un groupe de Lie complexe semi-simple. C. R. acad. sciences Paris, 323, 1259-1264.
- Bishop, R. (1963). A relation between volume, mean curvature and diameter. Notices of the Amer. math. Soc., 10, 364.
- Bismut, J.-M., Gillet, H., & Soulé, C. (1988). Analytic torsion and holomorphic determinant I, II, III. Communic. in math. physics, 115, 49-78, 79-126, 301-351.
- Blaschke, W. (1921). Vorlesungen über Differentialgeometrie. I (first edition). Springer.
- Blaschke, W. (1930). Vorlesungen über Differentialgeometrie. I (third edition). Springer.
- Blatter, C. (1961). Über Extremallänge auf geschlossenen Flächen. Comment. Math. Helvetici, 35, 153-168.
- Bleecker, D. (1995). Isometric deformations of compact hypersurfaces. Geometriae dedicata.
- Bochner, S. (1946). Vector fields and Ricci curvature. Bulletin of Amer. math. Soc., 52, 776-797.
- Bochner, S. (1947). Curvature in Hermitian metrics. Bull. Amer. Math. Soc., 53, 179-195.
- Bochner, S. (1948). Curvature and Betti numbers. Annals of math., 49, 379-390.
- Bochner, S., & Yano, K. (1953). Curvature and Betti numbers. Princeton University press.

Boeckx, E., Kowalski, O., & Vanhecke, L. (1996). Riemannian manifolds of conullity two. World scientific

Böhm, C. (1996). Sequences of Einstein metrics with bounded volumes. .

Bonnet, O. (1855). Sur quelques propriétés des lignes géodésiques. C. R. Acad. Sc. Paris, 40, 1311-1313.

Borel, A. (1963). Compact Clifford-Klein forms of symmetric spaces. Topology, 2, 111-122.

Borel, A., & Lichnerowicz, A. (1952). Groupes d'holonomie des variétés riemanniennes. C. R. Acad. Sciences Paris, 234, 1835-1837.

Bott, R. (1956). On the iteration of closed geodesics and the Sturm intersection theory. Communications in pure and applied math., 9, 171-206.

Bourbaki, N. (1969). Fascicule de résultats des variétés. Hermann.

Bourdon, M. (1996). Sur le birapport au bord des CAT(-1)-espaces. Publications math. IHES, 83, 95-104.

Bourguignon, J.-P. (1973). Some constructions related to Hopf's conjecture on product manifolds. In S.-S. Chern & R. Osserman (Ed.), Summer Institute on Differential geometry, 25 (pp. 33-37). Stanford: American math. soc.

Bourguignon, J.-P. (1975). Une stratification de l'espace des structures riemanniennes. Compositio math., 30, 1-41.

Bourguignon, J.-P. (1988). The "magic" of Weitzenböck formulas. In C. Berestycki Ekeleland (Ed.), Variational methods, (pp. 251-271). Paris: Birkhäuser.

Bourguignon, J.-P. (1990/91). Stabilité par déformation non-linéaire de la métrique de Minkowski(d'après D. Christodoulou et S. Kleinerman). In Séminaire Bourbaki, Astérisque 201-202-203 (pp. 321-358). Paris: Société math. France.

Bourguignon, J.-P. (1993). Eugenio Calabi and Kähler metrics. In P. de Bartolomeis, F. Tricerri, & E. Vesentini (Ed.), Manifolds and geometry, (pp. 61-85). Pisa: Cambridge University Press.

Bourguignon, J.-P., & Gauduchon, P. (1992). Spineurs, opérateurs de Dirac et variations de métriques. Communications in math. phys., 144, 581-599.

Bourguignon, J.-P., & Lawson, B. (1982). Yang-Mills theory : its physical origins and differential geometry aspects. In Seminar on differential geometry Princeton university press.

Bourguignon, J.-P., & Polombo, A. (1981). Intégrands des nombres caractéristiques et courbure : rien ne va plus dès la dimension 6. J. of differential geometry, 16, 537-550.

- Bourguignon, J.-P., Li, P., & Yau, S.-T. (1994). Upper bounds of the first eigenvalue for algebraic submanifolds. Commentarii math. Helvetici, 69, 199-207.
- Bowditch, B. (1994). Some results on the geometry of convex hulls in manifolds of pinched negative curvature. Commentarii math. helvetici, 69, 49-81.
- Boyer, C., Galicki, K., & Mann, B. (1994a). The geometry and topology of 3-Sasakian manifolds. J. für reine und angew. math., 455, 183-220.
- Boyer, C., Galicki, K., & Mann, B. (1994b). Quaternionic geometry and 3-Sasakian manifolds. In G. Gentili, S. Marchiafava, & M. Pontecorvo (Ed.), Quaternionic structures in mathematics and physics, . Trieste: SISSA Trieste.
- Boyer, C., Galicki, K., Mann, B., & Rees, E. (1997). Compact 3-Sasakian 7-manifolds with arbitrary second Betti number
- Bridson, M., & Haefliger, A. (1997). Metric spaces of non-positive curvature. Springer.
- Brooks, R., Perry, P., & Petersen, P. (1990). Finiteness of diffeomorphisms types of isospectral manifolds. In R. Greene & S.-T. Yau (Ed.), Differential geometry, 54, part 3 (pp. 89-94). Los Angeles: American math. soc.
- Brooks, R., Perry, P., & Petersen, P. (1992). Compactness and finiteness theorems for isospectral manifolds. J. reine and ang. math., 426, 67-89.
- Brooks, R., Perry, P., & Petersen, P. (1994). Spectral geometry in dimension 3. Acta mathematica, 173, 283-305.
- Brown, R., & Gray, A. (1972). Riemannian manifolds with holonomy group Spin(9). In Differential geometry, in honour of K. Yano, (pp. 41-59). Tokyo: Kinokuniya.
- Bryant, R. (1987). Metrics with exceptional holonomy. Annals of math., 126, 525-576.
- Bryant, R. (1996). Finsler structures on the 2-sphere satisfying $K = 1$. Contemporary math., 196, 27-41.
- Bryant, R., & Hsu, L. (1993). Rigidity of integral curves of rank 2 distributions. Inventiones math., 114, 435-461.
- Bryant, R., & Salamon, S. (1989). On the construction of some complete metrics with exceptional holonomy. Duke math. j., 58, 829-850.
- Buchner, M. (1978). The structure of the cut-loci in dimension less than or equal to six. Compositio math., 37, 103-119.
- Burago, Y., & Ivanov, S. (1994). Riemannian tori without conjugate points are flat. Geometric and functional analysis, 5, 801-808.
- Burago, Y., & Zalgaller, V. (1988). Geometric inequalities. Springer.

- Burago, Y., & Zalgaller, V. (Ed.). (1992a). Geometry III. Encyclopedia of math. sc. 48, Springer.
- Burago, Y., Gromov, M., & Perelman, G. (1992b). Alexandrov's spaces with curvatures bounded from below I. Russian mathematical surveys, 47, 1-58.
- Burns, K., & Paternain, G. (1996b). Counting geodesics on a Riemannian manifold and topological entropy of geodesic flows. .
- Busemann, H. (1955). The geometry of geodesics. Academic Press.
- Buser, P. (1978). Über eine Ungleichung von Cheeger. Math. Zeitschrift, 158, 245-252.
- Buser, P. (1992). Geometry and spectra of compact Riemann surfaces. Birkhäuser.
- Buser, P. (1996). Inverse spectral geometry on Riemann surfaces. In S. Andersson & M. Lapidus (Eds.), Recent progress in inverse spectral geometry Birkhäuser.
- Buser, P., & Gromoll, D. (1988). On the almost negatively curved 3-sphere. In T. Sunada (Eds.), Geometry and Analysis on manifolds , LNM 1339, Springer.
- Buser, P., & Karcher, H. (1981). Gromov's almost flat manifolds. Société math. France.
- Calabi, E. (1953). Isometric embedding of complex manifolds. Annals of Math., 58, 1-23.
- Calabi, E. (1954). The space of Kähler metrics. In Proceedings of the International Congress of Mathematicians (pp. 206-207). Amsterdam
- Calabi, E. (1958). An extension of E. Hopf's maximum principle with an application to Riemannian geometry. Duke math. j., 25, 45-56.
- Calabi, E. (1967). Minimal immersions of surfaces in Euclidean spheres. J. Differential Geometry, 1, 111-125.
- Calabi, E. (1967). Minimal immersions of surfaces in Euclidean spheres. J. Differential Geometry, 1, 111-125.
- Calabi, E. (1969). An intrinsic characterization of harmonic one-forms. In D. C. S. a. S. Iyanaga (Eds.), Papers in Honor of K. Kodaira (pp. 101-117). Princeton University Press.
- Calabi, E. (1979). Métriques kählériennes et fibrés holomorphes. Annales scientifiques Ecol. norm. sup., 12, 269-294.
- Calabi, E. (1992b). Extremal isosystolic metrics for compact surfaces. In A. Besse (Ed.), Table ronde de géométrie différentielle, Séminaires et congrès 1 (pp. 167-204). Luminy: Soc. Math. France.
- Calabi, E., & Cao, J. (1992a). Simple closed geodesics on convex surfaces. J. Differential Geometry, 36, 517-549.

Cao, H.-D. (1985). Deformations of Kähler metrics to Kähler Einstein metrics on compact Kähler manifolds. Inventiones math., 81, 359-372.

Cao, J. (1996). The Martin boundary and isoperimetric inequalities for Gromov-hyperbolic manifolds ,No. 251, Univ. Notre-Dame, dept. of math.

Cao, J., & Escobar, J. (1995). An isoperimetric theorem for PL-manifolds of non-positive curvature.

Cartan, E. (1913). Les groupes projectifs qui ne laissent invariante aucune multiplicité plane. Bull. soc. math. France, 41, 1-44.

Cartan, E. (1926a). Les groupes d'holonomie des espaces généralisés. Acta math., 48, 1-42.

Cartan, E. (1926b). Sur une classe remarquable d'espaces de Riemann. Bull. Soc. math. France, 214-264.

Cartan, E. (1927). Sur certaines formes riemanniennes remarquables de géométries à groupe fondamental simple. Annales scientif. Ecole norm. sup., 44, 345-367.

Cartan, E. (1928a). Leçons sur la géométrie des espaces de Riemann. Gauthier-Villars.

Cartan, E. (1928b). Sur les nombres de Betti des espaces de groupes clos. C. R. Acad. sciences Paris, 187, 196-198.

Cartan, E. (1929a). Sur les invariants intégraux de certains espaces homogènes clos et les propriétés topologiques de ces espaces. Annales Soc. math. polonaise, 8, 217-252.

Cartan, E. (1929b). Groupes simples clos et ouverts et géométrie riemannienne. J. math. pures et appl., 8, 1-33.

Cartan, E. (1931). Leçons sur la géométrie projective complexe. Gauthier-Villars.

Cartan, E. (1936). La topologie des espaces représentatifs des groupes de Lie. L'Enseignement math., 35, 177-200.

Cartan, E. (1937). La théorie des spineurs. Hermann.

Cartan, E. (1939). Sur quelques familles remarquables d'hypersurfaces. Math. Zeitschrift, 45, 335-367.

Cartan, E. (1946). Leçons sur la géométrie des espaces de Riemann, seconde édition, Gauthier-Villars.

Chavel, I. (1984). Eigenvalues in Riemannian geometry. Academic Press.

Chavel, I. (1993). Riemannian Geometry : a modern introduction. Cambridge University Press.

Chazarain, J. (1974). Formule de Poisson pour les varietes riemanniennes. Inventiones math., 24, 65-82.

Cheeger, J. (1967) Comparison and finiteness theorems for Riemannian manifolds. PhD, Princeton University.

Cheeger, J. (1970a). Finiteness theorems for Riemannian manifolds. American J. of math., 92, 61-74.

Cheeger, J. (1970b). A lower bound for the smallest eigenvalue of the Laplacian. In Symposium in honour of S. Bochner, (pp. 195-199). Princeton University Press.

Cheeger, J. (1973). Some examples of manifolds of nonnegative curvature. J. differential geom., 8, 623-628.

Cheeger, J. (1979). Analytic torsion and the heat equation. Annals of math., 109, 259-322.

Cheeger, J. (1983). Spectral geometry of singular Riemannian spaces. J. of differential geometry, 18, 575-667.

Cheeger, J. (1991). Critical points of distance functions and applications of geometry. In J. Cheeger, M. Gromov, C. Okonek, & P. Pansu (Eds.), Geometric topology : recent developments (pp. 1-38). LNM 1504, Springer.

Cheeger, J., & Colding, T. (1996d). Lower bounds on Ricci curvature and the almost rigidity of warped products. Annals of math., 144, 189-237.

Cheeger, J., & Colding, T. (1997a). On the structure of spaces with Ricci curvature bounded below : I. Annals of math.

Cheeger, J., & Colding, T. (1997c). On the structure of spaces with Ricci curvature bounded below, II. .

Cheeger, J., & Colding, T. (1998). On the structure of spaces with Ricci curvature bounded below, III. .

Cheeger, J., & Ebin, D. (1975). Comparison theorems in Riemannian Geometry. North-Holland.

Cheeger, J., & Gromoll, D. (1971). The splitting theorem for manifolds of nonnegative Ricci curvature. J. Differential Geometry, 6, 119-128.

Cheeger, J., & Gromoll, D. (1972). On the structure of complete manifolds of nonnegative curvature. Annals of math., 96, 413-443.

Cheeger, J., & Gromoll, D. (1980). On the lower bound for the injectivity radius of $1/4$ - pinched Riemannian manifolds. J. differential geometry, 26, 437-442.

Cheeger, J., & Gromov, M. (1985a). On the characteristic numbers of complete manifolds of bounded curvature and finite volume. In I. Chavel & H. Farkas (Eds.), Differential geometry and complex analysis (pp. 115-154). Springer.

Cheeger, J., & Gromov, M. (1986-1990). Collapsing Riemannian manifolds while keeping their curvature bounded. J. differential geometry, 23-32, 309-346, 269-298.

Cheeger, J., & Gromov, M. (1989). Chopping Riemannian manifolds. In B. Lawson & K. Teneblat (Ed.), Differential geometry : a symposium in honour of Manfredo do Carmo, (pp. 85-94). Rio de Janeiro: Longman scientific.

Cheeger, J., & Rong, X. (1996b). Existence of polarized F-structures on collapsed manifolds with bounded curvature and diameter. Geometric and functional analysis, 6, 411-429.

Cheeger, J., & Simons, J. (1985b). Differential characters and geometric invariants. In Geometry and topology, LNM 1167, Springer.

Cheeger, J., & Taylor, M. (1982). On the diffraction of waves by conical singularities, I and II. Communications in pure and appl. math., 35, 275-231 and 487-529.

Cheeger, J., & Tian, G. (1994). On the cone structure at infinity of Ricci flat manifolds with Euclidean volume growth and quadratic curvature decay. Inventiones math., 118, 493-571.

Cheeger, J., Colding, T., & Tian, G. (1997b). Constraints on singularities under Ricci curvature bounds

Cheeger, J., Fukaya, K., & Gromov, M. (1992). Nilpotent structures and invariant metrics on collapsed manifolds. J. of the American math. society, 5, 327-372.

Cheeger, J., Müller, W., & Schrader, R. (1984). On the curvature of piecewise flat spaces. Communic. Math. Phys., 92, 405-454.

Cheeger, J., Müller, W., & Schrader, R. (1986). Kinematic and tube formulas for piecewise linear spaces. Indiana Univ. math. J., 35, 737-753.

Chen, H. (1990). Manifolds with 2-nonnegative curvature operator. In R. Greene & S.-T. Yau (Ed.), Differential geometry, 54, part 3 (pp. 129-133). Los Angeles: American math. soc. 1993.

Cheng, S. (1975). Eigenvalue comparison theorems and its geometric applications. Math. Zeitschrift, 143, 289-297.

Chern, S.-S. (1944). A simple intrinsic proof of the Gauss-Bonnet formula for closed Riemannian manifolds. Annals of math., 45, 747-752.

Chern, S.-S. (1946). Characteristic classes of Hermitian manifolds. Annals of math., 47, 87-121.

Chern, S.-S. (1955). On curvature and characteristic classes of a Riemannian manifold. Abh. math. sem. Hamburg, 20, 117-126.

Chern, S.-S., & Simons, J. (1974). Characteristic forms and geometric invariants. Annals of math., 99, 48-69.

Chevalley, C. (1997). Collected work, volume 2. Springer.

Christodoulou, D., & Kleiner, S. (1993). The global linear stability of Minkowski space. Princeton University Press.

Cohn-Vossen, S. (1935). Kürzeste Wege und Totalkrümmung auf Flächen. Compositio math., 2, 63-133.

Cohn-Vossen, S. (1936). Totalkrümmung und geodätische Linien auf einfach zusammenhängenden vollständigen Flächenstücke. Recueil math. Moscou, 43, 139-163.

Colding, T. (1996a). Large manifolds with positive Ricci curvature. Inventiones math., 124, 193-214.

Colding, T. (1996b). Ricci curvature and volume convergence. Inventiones math.

Colding, T. (1996c). Shape of manifolds with positive Ricci curvature. Inventiones math., 124, 175-191.

Colding, T., & Minicozzi II, W. (1997). Weyl type bounds for harmonic functions.

Colin de Verdière, Y. (1973). Spectre du laplacien et longueurs des géodésiques périodiques I et II. Compositio math., 27, 83-106 et 159-184.

Colin de Verdière, Y. (1977). Quasi-modes sur les variétés Riemanniennes. Inventiones Math., 3, 15-52.

Colin de Verdière, Y. (1985). Ergodicité et fonctions propres du laplacien. Communic. math. physics, 102, 497-502.

Colin de Verdière, Y. (1987). Construction de laplaciens dont une partie finie du spectre est donnée. Annales scient. Écol. norm. sup., 20, 599-615.

Colin de Verdière, Y. (1992). Spectre du laplacien : survol partiel depuis le BGM et problèmes. In A. Besse (Ed.), Table ronde de géométrie différentielle, Congrès et séminaires 1 (pp. 233-252). Luminy: Soc. math. France 1996.

Connes, A. (1995-6). Brisure de symétrie spontanée et géométrie du point de vue spectral. In Séminaire Bourbaki, Paris, Astérisque, Société math. France.

Connes, A. (1994). Noncommutative geometry. Academic Press.

Conway, J. (1995). Sphere packings, lattices, codes and greed. In Proceedings of the International Congress of Mathematicians Birkhäuser.

Conway, J. H., & Sloane, N. J. A. (1993). Sphere Packings, Lattices and Groups Springer

Corlette, K. (1992). Archimedean superrigidity and hyperbolic geometry. Annals of math., 135, 165-182.

- Croke, C. (1980). Some isoperimetric inequalities and eigenvalue estimates. Ann. sci. Ec. norm. sup. Paris, 13, 419-435.
- Croke, C. (1982). An eigenvalue pinching theorem. Inventiones math., 68, 253-256.
- Croke, C. (1991). Rigidity and the distance between boundary points. J. differential geometry, 33, 445-464.
- Croke, C. (1984b). A sharp four dimensional isoperimetric inequality. Commentarii math. helvetici, 59, 187-192.
- Croke, C. (1988b). An isoembolic pinching theorem. Inventiones math., 92, 385-387.
- Croke, C. (1991). Rigidity and the distance between boundary points. J. differential geometry, 33, 445-464.
- Croke, C. (1992). Volume of balls in manifolds without conjugate points. International journal of math., 3, 455-467.
- d'Ambra, G. (1988). Isometry groups of Lorentz manifolds. Inventiones math., 92, 555-565.
- d'Ambra, G., & Gromov, M. (1991). Lectures on transformations groups : geometry and dynamics. Surveys in differential geometry : supplement to the J. of differential geometry, 1, 19-111.
- Darboux, G. (1887, 1889, 1894, 1896). Leçons sur la théorie générale des surfaces (4 vol.). Gauthier-Villars.
- Darboux, G. (1972). Leçons sur la théorie générale des surfaces. Chelsea.
- de Rham, G. (1931). Sur l'analysis situs des variétés à n dimensions. J. math. pures et appliquées, 10, 115-200.
- de Rham, G. (1952). Sur la réductibilité d'un espace de Riemann. Commentarii math. helvetici, 26, 328-344.
- de Rham, G. (1954). Variétés différentiables. Hermann.
- Deligne, P., & Mostow, G. (1986). Monodromy of hypergeometric functions and no-lattice integral monodromy. Publications math. IHES, 63, 5-89.
- Deligne, P., Griffiths, P., Morgan, J., & Sullivan, D. (1975). Real homotopy of Kähler manifolds. Inventiones math., 29, 245-274.
- Diaconis, P., Elworthy, D., Nelson, E., Papanicolaou, G., & Varadhan, S. (Ed.). (1985-87). Ecole d'été de probabilités de Saint-Flour. LNM 1362, Springer.
- Dierkes, U., Hildebrandt, S., Küster, A., & Wohlrab, O. (1992). Minimal surfaces I and II. Springer.
- Dieudonné, J. (1972). Treaty of Analysis. Academic Press.
- Dillen, F., & Vestraalen, L. (Ed.). (1992). Geometry and topology of submanifolds. World scientific.

- do Carmo, M. (1976). Differential geometry of curves and surfaces. Prentice-Hall.
- do Carmo, M. (1992). Riemannian geometry. Birkhäuser.
- Dombrowski, P. (1979). 150 years after Gauss. Astérisque 62, Société mathématique de France.
- Donaldson, S. (1996a). The Seiberg-Witten equations and 4-manifolds topology. Bull. American math. soc., 33, 45-70.
- Donaldson, S., & Kronheimer, P. (1996b). The geometry of four-manifolds. Oxford University Press.
- Donnay, V. (1988). Geodesics flow on the two-sphere II : ergodicity. In Lecture notes in math. n° 1342 : Dynamical systems (pp. 112-153). Springer.
- Donnelly, H., & Fefferman, C. (1988). Nodal sets of eigenfunctions on Riemannian manifolds. Inventiones math., 93, 161-183.
- Donnelly, H., & Fefferman, C. (1990). Growth and geometry of eigenfunctions of the Laplacian. In C. Sadosky (Eds.), (pp. 635-655). Dekker.
- Duistermaat, J., & Guillemin, V. (1975). The spectrum of positive elliptic operators and periodic bicharacteristics. Inventiones math., 29, 39-79.
- Eberlein, P., & O'Neill, B. (1973). Visibility manifolds. Pacific J. of math., 46, 45-109.
- Eberlein, P., Hamenstädt, U., & Schroeder, V. (1990). Manifolds of nonpositive curvature. In R. Greene & S.-T. Yau (Eds.), Differential geometry, 54, part 3 (pp. 179-227). Los Angeles: American math. society 1993.
- Ebin, D. (1970). The space of Riemannian metrics. In Global analysis, 15 (pp. 11-40). Berkeley: American math. soc.
- Ebin, D. (1972). Espace des métriques riemanniennes et mouvement des fluides via les variétés d'applications. Ecole polytechnique.
- Ebin, D., & Marsden, J. (1970). Groups of diffeomorphisms and the motion of an incompressible fluid. Annals of math., 92, 102-163.
- Eells, J., & Lemaire, L. (1988). Another report on harmonic maps. Bull. London Math. Soc., 20, 385-524.
- Eells, J., & Rato, A. (1993). Harmonic maps and minimal immersions with symmetries. Princeton University Press.
- Eells, J., & Sampson, J. (1964). Harmonic mappings of Riemannian manifolds. American J. of math., 86, 109-160.
- Einstein, A. (1916). Näherungsweise Integration der Feldgleichungen der Gravitation. S. B. Preuss. Akad. Wiss., 688-696.

- El Alaoui, E.-L., Gauthier, J.-P., & Kupka, I. (1996). Small sub-Riemannian balls on $\mathbb{R}^3$, J. of dynamical control systems, 2, 359-421
- Elworthy, K. (1988). Geometric aspects of diffusions on manifolds. LNM 1362, Springer.
- Epstein, D. (1975). Natural tensors on Riemannian manifolds. J. of differential geometry, 10, 631-645.
- Eschenburg, J.-H. (1982). New examples of Riemannian manifolds of strictly positive curvature. Inventiones math., 66, 469-480.
- Eschenburg, J.-H. (1986). Local convexity and nonnegative curvature : Gromov's proof of the sphere theorem. Inventiones math., 84, 507-522.
- Eschenburg, J.-H. (1992). Inhomogeneous spaces of positive curvature. Differential geometry and its applications, 2, 123-132.
- Eschenburg, J.-H. (1994). Comparison theorems in Riemannian geometry. Trento: Universita di Trento.
- Eschenburg, J.-H., & Heintze, E. (1984). An elementary proof of Cheeger-Gromoll splitting theorem. Annals of global analysis and geometry, 2, 141-151.
- Eschenburg, J.-H., & Wang, M. (1995). The initial value problem for cohomogeneity one Einstein metrics No. 341). Mathematik, Univ. of Augsburg.
- Faber, G. (1923). Beweis, dass unter allen homogenen Membranen von gleicher Fläche und gleicher Spannung die kreisförmige den tiefsten Grundton gibt. Sitzungsberichte der Bayrischen Akad. Wissen., 169-172.
- Farber, M., Katz, G., & Levine, J. (1996). Morse theory for harmonic forms No. M/96/34). I H E S.
- Farrell, F. (1997). Surgical methods in rigidity. Tata Institute lectures, Springer.
- Farrell, F., & Jones, L. (1989a). Negatively curved manifolds with exotic smooth structures. J. of Amer. math. Soc., 2, 899-908.
- Farrell, T., & Jones, L. (1989b). A topological analogue of Mostow's rigidity theorem. J. American math. soc., 2, 257-370.
- Farrell, F., & Jones, L. (1993). Non-uniform lattices and exotic smooth structures. J. of differential geom., 38, 235-261.
- Farrell, F., & Jones, L. (1994). Complex hyperbolic manifolds and exotic smooth structures. Inventiones math., 117, 57-74.
- Federer, H. (1969). Geometric measure theory. Springer.
- Federer, H., & Fleming, W. (1960). Normal and integral currents. Annals of math., 72, 458-520.

- Flach, N. (1994). Diamètre des variétés à courbure sectionnelle positive. C. R. Acad. sciences Paris, 318, 827-830.
- Foulon, P. (1997). Les espaces de Finsler localement symétriques à courbure négative. C. R. Acad. sciences Paris.
- Franks, J. (1992). Geodesics on S^2 and periodic points of annulus diffeomorphisms. Inventiones math., 108, 403-418.
- Freed, D., & Uhlenbeck, K. (1984). Instantons and four-manifolds. Springer.
- Fröhlich, J., Granjean, O., & Recknagel, A. (1997). Supersymmetric quantum theory No. ETH-Höngerberg, Zürich.
- Fubini, G. (1903). Sulle metriche definite da una forma Hermitiana. Atti Inst. Veneto, 6, 501-513.
- Fukaya, K. (1984). A finiteness theorem for negatively curved manifolds. J. of differential geometry, 20, 497-521.
- Fukaya, K. (1987a). Collapsing Riemannian manifolds to ones of lower dimension. J. of differential geometry, 25, 139-156.
- Fukaya, K. (1987b). Collapsing Riemannian manifolds and eigenvalues of the Laplace operator. Inventiones math., 87, 517-547.
- Fukaya, K. (1988). A boundary of the set of the Riemannian manifolds with bounded curvature and diameters. J. of differential geometry, 28, 1-21.
- Fukaya, K. (1989). Collapsing Riemannian manifolds to ones of lower dimension II. J. math. society of Japan, 41, 333-356.
- Fukaya, K. (1990). Hausdorff convergence of Riemannian manifolds and its applications. In Recent topics in differential and analytic geometry (pp. 143-238). Tokyo: North Holland and Kinokuniya.
- Fukaya, K., & Yamaguchi, T. (1991). Almost nonpositively curved manifolds. J. differential geometry, 33, 67-90.
- Fukaya, K., & Yamaguchi, T. (1992). The fundamental group of almost nonnegatively curved manifolds. Annals of math., 136, 253-333.
- Futaki, A. (1983). An obstruction to the existence of Einstein-Kähler metrics. Inventiones math., 73, 437-443.
- Gaffney, M. (1958). Asymptotic distributions associated with the Laplacian for forms. Communic. pure and appl. math., XI, 535-545.
- Gage, M., & Hamilton, R. (1986). The heat equation shrinking plane curves. J. of differential geometry, 23, 69-96.
- Galicki, K., & Laswon, B. (1988). Quaternionic reduction and quaternionic orbifolds. Math. annalen, 282, 1-21.

- Gallot, S. (1981). A Sobolev inequality and some geometric applications. In M. Berger, S. Murakami, & T. Ochiai (Ed.), Spectra of Riemannian manifolds, (pp. 45-55). Kyoto: Kaigai.
- Gallot, S. (1986). Inégalités isopérimétriques et analytiques sur les variétés riemanniennes. In Sulle geometria delle varietà differenziabili, 163-164 (pp. 31-92). Rome. Astérisque 163-164, Société mathématique de France.
- Gallot, S. (1987). Isoperimetric inequalities based on integral norms of Ricci curvature. In Colloque Paul Lévy sur les processus stochastiques (pp. 191-216). Ecole polytechnique. Astérisque 157-158, Soc. math. France.
- Gallot, S., Hulin, D., & Lafontaine, J. (1990). Riemannian geometry. Springer.
- Gao, L. (1990a). Convergence of Riemannian manifolds ; Ricci and $L^{n/2}$ curvature pinching. J. differential geometry, 32, 349-381.
- Gao, L. (1990b). $L^{n/2}$ curvature pinching. J. differential geometry, 32, 713-774.
- Gauduchon, P. (1992/93). Variétés riemanniennes autoduales. In Séminaire Bourbaki, Astérisque 216 (pp. 151-186). Paris: Société math. de France.
- Gasqui, J., & Goldschmidt, H. (1994). The infinitesimal rigidity of the complex quadrics of dimension four. American J. of Math., 116, 501-539.
- Gentili, G., Marchiafava, S., & Pontecorvo, M. (Ed.). (1994). Quaternionic structures in mathematics and physics. Trieste: SISSA, Trieste.
- Ghys, E., & de la Harpe, P. (Ed.). (1990). Sur les groupes hyperboliques d'après Mikhael Gromov. Birkhäuser.
- Gibbon, G., & Hawking, S. (1978). Gravitational multi-instantons. Physics letters B, 78, 430-432.
- Gil-Medrano, O., & Michor, P. (1991). The Riemannian manifold of all Riemannian metrics. Quarterly j. of math. Oxford, 42, 183-202.
- Gilkey, P. (1973a). The spectral geometry of real and complex manifolds. In Differential geometry, (pp. 265-280). Stanford: American math. soc. 1975
- Gilkey, P. (1973b). Curvature and the eigenvalues of the Laplacian for elliptic complexes. Adv. in math., 10, 344-382.
- Gilkey, P. (1974). The index theorem and the heat equation. Publish or Perish, Boston.
- Gilkey, P. (1995). Invariance theory, the heat equation, and the Atiyah-Singer index theorem (second edition). Boca Raton: CRC Press.
- Gillet, H., & Soulé, C. (1992). An arithmetic Riemann-Roch theorem. Inventiones math., 110, 473-543.
- Gluck, H., & Singer, D. (1978). Scattering of geodesic fields, II. Annals of math., 110, 205-225.

- Godbillon, C. (1991). Feuilletages. Etudes géométriques. Birkhäuser.
- Goncharov, A. (1997). Volumes of hyperbolic manifolds, and mixed Tate motives.
- Gordon, C. (1994c). Isospectral closed Riemannian manifolds which are not locally isometric, part II. Contemporary math., 121-131.
- Gordon, C., & Mao, Y. (1994b). Comparisons of Laplace spectra, length spectra and geodesic flows of some Riemannian manifolds. Mathematical research letters, 1, 677-688.
- Grayson, M. (1989). Shortening imbedded curves. Annals of math., 129, 71-111.
- Green, L. (1963). Auf Wiedersehenflächen. Annals of math., 78, 289-299.
- Greene, R. (1990). Some concepts and methods in Riemannian geometry. In R. Greene & S.-T. Yau (Ed.), Differential geometry, 3 (pp. 1-22). Los Angeles: American math. soc. 1993
- Greene, R. (1994). Homotopy finiteness theorems and Helly's theorem. The journal of Geometric Analysis, 4, 317-325.
- Greene, R. (1997). Manifolds of nonnegative curvature. .
- Greene, R., & Petersen, P. (1992). Little topology, big volume. Duke math. J., 67, 173-290.
- Greene, R., & Shiohama, K. (1981a). Convex functions on complete noncompact manifolds : differentiable structure. Annales scientifiques Ecol. norm. sup., 14, 357-367.
- Greene, R., & Shiohama, K. (1981b). Convex functions on complete noncompact Riemannian manifolds. Inventiones math., 63, 129-157.
- Greene, R., & Wu, H. (1988). Lipschitz convergence of Riemannian manifolds. Pacific J. of math., 131-1, 119-141.
- Greene, R., & Wu, H. (1990). Nonnegatively curved manifolds which are flat outside a compact set. In R. Greene & S.-T. Yau (Ed.), Differential geometry, 54, 3 (pp. 327-335). Los Angeles: American math. soc. 1993
- Greene, R., & Yau, S.-T. (Ed.). (1993). Differential geometry, Los Angeles 1990, Proceedings of symposia in pure math. . American math. soc.
- Griffiths, P., & Harris, J. (1978). Principle of algebraic geometry. John Wiley.
- Gromoll, D. (1966). Differenzierbare Strukturen und Metriken positiver Krümmung auf Sphären. Math. annalen, 164, 353-371.
- Gromoll, D. (1990). Spaces of nonnegative curvature. In R. Greene & S.-T. Yau (Ed.), Differential geometry, 54, part 3 (pp. 337-356). Los Angeles: American math. society. 1993

Gromoll, D., & Grove, K. (1981). On metrics on S^2 all of whose geodesics are closed. Invent. math., 65, 175-177.

Gromoll, D., & Grove, K. (1987). A generalization of Berger's rigidity theorem for positively curved manifolds. Ann. sci. Ecole normale sup. Paris, 20, 227-239.

Gromoll, D., & Meyer, W. (1969a). Periodic geodesics on compact Riemannian manifolds. J. differential geom., 3, 493-510.

Gromoll, D., & Meyer, W. (1969b). On complete manifolds of positive curvature. Annals of math., 90, 75-90.

Gromoll, D., & Meyer, W. (1974). An exotic sphere with nonnegative curvature. Annals of math., 100, 401-406.

Gromoll, D., & Wolf, J. (1971). Some relations between the metric structure and the algebraic structure of the fundamental group in manifolds of nonpositive curvature. Bull. American math. soc., 77, 545-552.

Gromov, M. (1978a). Almost flat manifolds. J. of differential geometry, 13, 231-241.

Gromov, M. (1978b). Manifolds of negative curvature. J. of differential geometry, 13, 223-230.

Gromov, M. (1980a). Paul Levy's isoperimetric inequality No. M/80/320). Institut des Hautes Etudes Scientifiques.

Gromov, M. (1981b). Hyperbolic manifolds according to Thurston and Jorgensen. In Lecture Notes in Math. 842 (pp. 40-53). Springer.

Gromov, M. (1981c). Groups of polynomial growth and expanding maps. Publications math. I.H.E.S., 53, 53-73.

Gromov, M. (1981d). Curvature, diameter and Betti numbers. Commentarii math. helvetici, 56, 179-195.

Gromov, M. (1983a). Filling Riemannian manifolds. J. of Diff. Geometry, 18, 1-147.

Gromov, M. (1983b). Volume and bounded cohomology. Public. math. IHES, 56, 213-37.

Gromov, M. (1985). Pseudo-holomorphic curves in symplectic manifolds. Inventiones math., 82, 307-347.

Gromov, M. (1986). Partial Differential Relations. Springer.

Gromov, M. (Ed.). (1987a). Hyperbolic groups. in Essays in group theory. S. Gersten, ed. MSRI publications 8, Springer.

Gromov, M. (1987b). Entropy, homology and semi-algebraic geometry. In Séminaire Bourbaki, Astérisque 145-146 (pp. 225-240), Paris: Société mathématique de France.

Gromov, M. (1988c). Widths and related invariants of Riemannian manifolds. Astérisque, 163-164, 93-109.

Gromov, M. (1990b). Stability and pinching. In Giornate di Topologica e geometria delle varietà, . Bologna: Università degli studi di Bologna.

Gromov, M. (1991a). Kähler hyperbolicity and L^2 Hodge theory. Journal of Differ. geom., 33, 263-292.

Gromov, M. (1991b). Sign and geometric meaning of curvature. Rendiconti Seminario mat. e Fisic. di Milano, 56, 10-123.

Gromov, M. (1991c). Foliated Plateau problem I et II. Geometric and functional analysis, 1, 14-79 and 253-320.

Gromov, M. (1992a). Systoles and intersystolic inequalities. In A. Besse (Ed.), Table ronde de géométrie différentielle, 1 (pp. 291-362). Luminy, Séminaires et congrès 1, Soc. math. France 1996.

Gromov, M. (1992b). Metric invariants of Kähler manifolds. In R. Caddeo & F. Tricerri (Ed.), Differential geometry and topology, (pp. 90-116). Alghero: World scientific.

Gromov, M. (1993). Asymptotic invariants of infinite groups. In G. Noble & M. Roller (Eds.), Geometric group theory (pp. 75-263). London math. soc.

Gromov, M. (1996). Positive curvature, macroscopic dimension, spectral gaps and higher signatures. In S. Gindikin, J. Lepowski, & R. Wilson (Eds.), Functional analysis in the eve of the 21st century (pp. 1-213). Birkhäuser.

Gromov, M. (1997). Structures métriques pour les variétés riemanniennes, second edition. Birkhäuser.

Gromov, M., & Lawson, B. (1980b). The classification of simply connected manifolds of positive curvature. Annals of math., 111, 423-434.

Gromov, M., & Lawson, B. (1980c). Spin and scalar curvature in the presence of a fundamental group. I. Annals of math., 111, 209-230.

Gromov, M., & Lawson, B. (1983c). Positive scalar curvature and the Dirac operator on complete Riemannian manifolds. Publications math. I H E S, 58, 83-196.

Gromov, M., & Pansu, P. (1990a). Rigidity of lattices : an introduction. In P. de Bartolomeis & F. Tricerri (Ed.), Geometric topology: recent developments, Lecture Notes, n° 1504 . Montecatini: Springer.

Gromov, M., & Piatetski-Shapiro, I. (1988a). Non-arithmetic groups in Lobachevskij spaces. Public. Math. Inst. Hautes Etud. Sci., 66, 93-103.

Gromov, M., & Schoen, R. (1992d). Harmonic maps into singular spaces and p-adic superrigidity for lattices in groups of rank one. Publications math. IHES, 76, 165-246.

Gromov, M., & Thurston, W. (1987c). Pinching constants for hyperbolic manifolds. Inventiones math., 89, 1-12.

Gromov, M., Lafontaine, J., & Pansu, P. (1981a). Structures métriques pour les variétés riemanniennes. Cedric-Nathan.

Grove, K. (1985). Metric differential geometry. In V. L. Hausen (Ed.), Nordic summer school, 1263 (pp. 171-227). Lingby 1985: Springer.

Grove, K. (1990). Critical point theory for distance functions. In R. Greene & S.-T. Yau (Ed.), Differential geometry, 3 (pp. 357-385). Los Angeles: Amer. math. soc. 1993

Grove, K. (1992). Ramifications of the classical sphere theorem. In A.-L. Besse (Ed.), Table ronde de géométrie différentielle, 1 (pp. 363-376). Luminy, Séminaires et congrès 1, Société math. France 1996.

Grove, K., & Halperin, S. (1983). Contributions of rational homotopy theory to global problems in geometry. Publications math. IHES, 56, 379-385.

Grove, K., & Petersen, P. (1988). Bounding homotopy types by geometry. Annals of math., 128, 195-206.

Grove, K., & Petersen, P. (1993). A radius sphere theorem. Inventiones math., 112, 577-583.

Grove, K., & Shiohama, K. (1977). A generalized sphere theorem. Annals of math., 106, 201-211.

Grove, K., & Wilhelm, F. (1995). Hard and soft packing radius theorems. Annals of math., 142, 213-237.

Grove, K., Petersen, P., & Wu, J.-Y. (1990-1991). Geometric finiteness theorems via controlled topology. Inventiones math., 99 and 104, 205-213 and 221-222.

Guillemin, V. (1976). The Radon transform on Zoll surfaces. Advances in math., 22, 85-119.

Guillemin, V. (1993). Wave-trace invariants and a theorem of Zelditch. International math. research notices, 12, 303-308.

Guillemin, V. (1996). Wave trace invariants. Duke math. j., 83, 287-352.

Guillemin, V., & Kazhdan, D. (1980a). Some inverse spectral results for negatively-curved 2-manifolds. Topology, 19, 301-312.

Hadamard, J. (1898). Les surfaces à courbure opposées et leurs lignes géodésiques. J. Math. pures et appl., 4, 27-73.

Hadamard, J. (1901). Sur l'itération et les solutions asymptotiques des équations différentielles. Bulletin de la Soc. math. France, 24, 224-228.

- Haefliger, A. (1956). Sur l'extension du groupe structural d'un espace fibré. Comptes rendus Acad. sciences Paris, 253, 558-560.
- Hamilton, R. (1982). Three-manifolds with positive Ricci curvature. J. of differential geometry, 17, 255-306.
- Hamilton, R. (1986). Four-manifolds with positive curvature operator. J. of differential geometry, 37, 225-243.
- Harvey, R., & Lawson, B. (1982). Calibrated geometries. Acta mathematica, 148, 48-157.
- Hawking, S., & Ellis, G. (1973). The large structure of space time. Cambridge University Press.
- Hawley, N. (1953). Constant holomorphic sectional curvature. Canadian j. of math., 5, 53-56.
- Hebda, J. (1994). Metric structure of cut-loci in surfaces and Ambrose's problem. J. differential geometry, 40, 621-642.
- Hebda, J. (1995). Cut Loci of submanifolds in space forms and in the geometry of Möbius and Lie. Geometriae Dedicata, 55, 75-93.
- Heber, J. (1997) Geometric and algebraic structure of noncompact homogeneous Einstein spaces. Augsburg.
- Hebey, E. (1993). Courbure scalaire et géométrie conforme. J. of geometry and physics, 10, 345-380.
- Hebey, E. (1995a). Optimal Sobolev inequalities on complete Riemannian manifolds with Ricci curvature bounded below and positive injectivity radius. American Journal of Math.
- Hebey, E., & Herzlich, M. (1995b). Convergence of Riemannian manifolds : a status report No. 1096. Ecole Polytechnique, Centre de mathématiques.
- Hebey, E. (1996b). Sobolev spaces on Riemannian manifolds. LNM 1635, Springer.
- Hebey, E. (1997). Introduction à l'analyse non linéaire sur les variétés. Diderot.
- Hejhal, D. (1976). The Selberg trace formula and the Riemann zeta-function. Duke math. J., 43, 441-481.
- Heintze, E. (1976) Mannigfaltigkeiten negativer Krümmung.
Habilitationsschrift, Bonn.
- Heintze, E., & Liu, X. (1997). Homogeneity of infinite dimensional isoparametric submanifolds.

- Heintze, E., & Karcher, H. (1978). A general comparison theorem with applications to volume estimates for submanifolds. Ann. Sci. Ec. Norm. Sup. Paris, 11, 451-470.
- Helein, F. (1996). Applications harmoniques, lois de conservation et repères mobiles. Diderot.
- Hernandez, L. (1991). Kähler manifolds and 1/4-pinching. Duke math. J., 62, 601-611.
- Hilbert, D. (1915). Die Grundlagen der Physik. Nach. Ges. Wiss. Göttingen, 461-472.
- Hingston, N. (1993). On the growth of the number of closed geodesics on the two-sphere. International math. research notices, 9, 253-262.
- Hirsch, M., & Mazur, B. (1974). Smoothing of piecewise linear manifolds. Princeton university press.
- Hirzebruch, F., & Kodaira, K. (1957). On the complex projective spaces. J. de math. pures et appl., 36, 201-216.
- Hirzebruch, F. (1966). Topological methods in Algebraic geometry. Springer.
- Hitchin, N. (1974). Harmonic spinors. Advances in math., 14, 1-55.
- Hitchin, N. (1991/92). The hyperkähler manifolds. In Séminaire Bourbaki, (pp. 137-166). Paris, Astérisque 206, Société math. de France.
- Hitchin, N. (1995). Twistor spaces, Einstein metrics and isomonodromic deformations. J. of differential geometry, 42, 30-112.
- Hodge, W. (1941). The theory and applications of harmonic integrals (second edition 1952). Cambridge University Press.
- Hofer, H., & Zehnder, E. (1994). Symplectic capacities. Birkhäuser.
- Hofer, H., Taubes, C., Weinstein, A., & Zehnder, E. (Ed.). (1995). The Floer memorial volume. Birkhäuser.
- Hopf, E. (1939). Statistik des geodätischen Linien in Mannigfaltigkeiten negativer Krümmung. Ber. Vehr. Sächs. Akad. Wiss. Leipzig, 91, 261-304.
- Hopf, E. (1948). Closed surfaces without conjugate points. Proc. Nat. Acad. Sciences, 34, 47-51.
- Hopf, H. (1925a). Die Curvatura integra Clifford-Kleinischer Raumformen. Nach. Ges. d. Wiss. Göttingen, Math. Physik Klasse.
- Hopf, H. (1932). Differential Geometrie and topologische Gestalt. Jahrbuch der DMV, 41, 209-229.
- Hopf, H. (1951). Über Flächen mit einer Relation zwischen den Hauptkrümmungen. Math. Nachrichten, 4, 232-249.

- Hopf, H. (1983). Differential geometry in the large. LNM 1000, Springer.
- Hopf, H., & Rinow, W. (1931). Über den Begriff der vollständigen differentialgeometrischen Flächen. Commentarii math. Helv., 3, 209-225.
- Hörmander, L. (1968). The spectral function of an elliptic operator. Acta mathematica, 121, 193-218.
- Hörmander, L. (1985). The analysis of linear partial differential operators : I, II, III, IV. Springer.
- Huang, H.-M. (1981). Some remarks on the pinching problem. Bull. Inst. math. Sinica, 9, 321-340.
- Hulin, D. (1995). Sous-variétés complexes d'Einstein de l'espace projectif. Bull. soc. math. France, 124, 277-298.
- Hummel, C., & Schroeder, V. (1997). Tits geometry associated with closed real-analytic manifolds of non positive curvature and dimension 4 No. Univ. Pennsylvania and Zürich Univ.
- Husemoller, D. (1975). Fibre bundles. Springer.
- Igusa, J. (1954). On the structure of a certain class of Kähler manifolds. American j. of math., 76, 669-678.
- Im Hof, H.-C., & Ruh, E. (1975). An equivariant pinching theorem. Commentarii math. helvetici, 50, 389-401.
- Itoh, J. (1997a). The length of the cut-loci on a surface and Ambrose's problem. J. of differential geometry.
- Itoh, J.-I., & Tanaka, M. (1997b). On the Hausdorff dimension of a cut-locus on a smooth Riemannian manifold. .
- Iwaniec (1975). Introduction to the spectral theory of automorphic forms. Biblioteca de la revista matematica iberoamericana.
- Jensen, G. (1973). Einstein metrics on principal fibre bundles. J. of diff. geometry, 8, 599-614.
- Jorgensen, T. (1977). Compact 3-manifolds of constant negative curvature fibering over the circle. Annals of math., 106, 61-72.
- Jost, J. (1991). Two-dimensional geometric variational problems. Wiley-Interscience.
- Jost, J. (1995). Riemannian geometry and geometric analysis. Springer.
- Jost, J. (1996). Compact Riemann surfaces. Springer.
- Jost, J., & Karcher, H. (1982). Geometrischen Methoden zur Gewinnung von a-priori Schranken für harmonische Abbildungen. Manuscripta mathematica, 40, 27-77.

- Jost, J., & Yau, S.-T. (1990). Harmonic maps and superrigidity. In R. Greene & S.-T. Yau (Ed.), Differential geometry, 54 (pp. 245-280). Los Angeles: American math. Soc.
- Joyce, D. (1996a). Compact 8-manifolds with holonomy $\text{Spin}(7)$. Inventiones math., 123, 507-552.
- Joyce, D. (1996b). Compact manifolds with holonomy G_2 : I. II J. of differential geometry, 43, 291-328, 329-375.
- Joyce, D. (1997). A theory of quaternionic algebra, with applications to hypercomplex geometry. .
- Kamberov, G. (1997). Prescribing mean curvature : existence and uniqueness problems
- Karcher, H. (1988). A geometric classification of positively curved symmetric spaces and the isoparametric construction of the Cayley plane. In On the geometry of differentiable manifolds (pp. 111-136).Astérisque 163-164, Soc. Math. France.
- Karcher, H. (1989). Riemann comparison constructions. In S. S. Chern (Eds.), Global differential geometry (pp. 170-222). The mathematical association of America.
- Kasue, A. (1989). A convergence theorem for Riemannian manifolds and some applications. Nagoya math. j., 114, 21-51.
- Katok, A. (1982). Entropy and closed geodesics. Ergodic theory and dynamical systems, 2, 339-365.
- Katok, A. (1988). Four applications of conformal equivalence to geometry and dynamics. Ergodic Th. and Dynamical Syst., 8, 139-152.
- Katok, A., & Hasselblatt, B. (1995). Introduction to the modern theory of dynamical systems. Cambridge Univeristy Press.
- Katok, S. (1996). Coding of closed geodesics after Gauss and Morse. Geometriae dedicata, 63, 123-145.
- Katsuda, A. (1985). Gromov's convergence theorem and its applications. Nagoya math. J., 100, 11-48.
- Katz, M. (1983). The filling radius of two-point homogeneous spaces. J. of differential geom., 18, 503-511.
- Katz, M. (1991). The rational filling radius of complex projective space. Topology and its applications, 42, 201-215.
- Kazdan, J. (1981/82). Positive energy in general relativity. In Séminaire Bourbaki, Astérisque 92-93 (pp. 315-330). Paris: Société math. France.

- Kazdan, J., & Warner, F. (1975a). Existence and conformal deformation of metrics with prescribed Gaussian and scalar curvature. Ann. of math., 101, 317-331.
- Kazdan, J., & Warner, F. (1975b). Scalar curvature and conformal deformation of Riemannian structure. J. of differential geom., 10, 113-134.
- Kirby, R., & Siebenmann, L. (1977). Foundational essays on topological manifolds, smoothings, and triangulations. Annals of math. studies 88, Princeton University Press.
- Kiyohara, K. (1984). On infinitesimal $C_2\pi$ -deformations of standard metrics on spheres. Hokkaido math. Journal, 13, 151-231.
- Kiyohara, K. (1987). Riemannian metrics with periodic geodesic flow on projective spaces. Japan J. math., 13, 209-234.
- Kleiner, B. (1992). An isoperimetric comparison theorem. Inventiones math., 108, 37-47.
- Kleiner, B., & Leeb, B. (1997). Rigidity of quasi-isometries for symmetric spaces and euclidean buildings
- Klingenberg, W. (1959). Contributions to Riemannian geometry in the large. Annals of math., 69, 654-666.
- Klingenberg, W. (1961). Über Riemannsche Mannigfaltigkeiten mit positiver Krümmung. Commentarii math. Helvetici, 35, 47-54.
- Klingenberg, W. (1978a). Lectures on closed geodesics. Springer.
- Klingenberg, W. (1978b). A course in differential geometry. Springer.
- Klingenberg, W. (1982). Riemannian geometry Second edition in 1995. de Gruyter.
- Klingenberg, W., & Pinl, M. (1981). Die Bedeutung von Christoffel für die Geometrie, E. B . Christoffel Weg zum absoluten Differentialkalkül und sein Betrag zur Theorie des Krümmungstensors. In P. Butzer & F. Fehér (Eds.), E. B. Christoffel (pp. 462-473)
- Klingenberg, W., & Sakai, T. (1980). Injectivity radius estimate for (1/4)-pinched manifolds. Arch. math., 34, 371-376.
- Klingenberg, W., & Takens, F. (1972). Generic properties of geodesic flow. Math. annalen, 197, 323-334.
- Klotz Milnor, T. (1972). Efimov's theorem about complete immersed surfaces of negative curvature. Advances in math., 8, 474-543.
- Knieper, G., & Weiss, H. (1994). A surface with positive curvature and positive topological entropy. J. Differential geometry, 39, 229-249.
- Kobayashi, S. (1989). On conjugate and cut-loci. In S. S. Chern (Eds.), Global differential geometry (pp. 140-169). The math. Assoc. of America.

- Kobayashi, S., & Nomizu, K. (1963-1969). Foundations of differential geometry I and II. Interscience.
- Koszul, J.-L. (1960). Lectures on fibre bundles and differential geometry. Bombay: Tata institute of fundamental research.
- Kowalski, O., & Vlasek, Z. (1993). Homogeneous Einstein metrics on Aloff-Wallach spaces. Differential geometry and its applic., 3, 157-167.
- Krahn, E. (1924). Über eine von Rayleigh formulierte Minimaleigenschaft des Kreises. Math. Annalen, 94, 97-100.
- Kreck, M., & Stolz, S. (1993). Nonconnected moduli spaces of positive sectional curvature metrics. Journal of the Amer. math. Soc., 6, 825-849.
- Kühnel, W., & Rademacher, H.-B. (1995). Essential conformal fields in pseudo-riemannian geometry. J. math. pures et appl., 74, 453-481.
- Kuiper, N. (1955). On C^1 -isometric imbeddings I and II. Indagationes math., 17, 545-566 and 683-689.
- Kulkarni, R. (1970). Curvature and metric. Annals of math., 91, 311-331.
- Kuranishi, M. (1990). On some metric on $S^2 \times S^2$. In R. Greene & S.-T. Yau (Ed.), Differential geometry, 54 part 3 (pp. 439-450). Los Angeles: American math. soc. 1993
- Lafontaine, J. (1985/86). Mesures de courbure des variétés riemanniennes et des polyèdres (d'après Cheeger, Müller et Schrader). In Séminaire Bourbaki, 145-146 (pp. 241-256). Paris: Société mathématique de France.
- Lanczos, C. (1922). Ein vereinfachendes Koordinatensystem für die Einsteinschen Gravitationsgleichungen. Phys. Z., 23, 537-539.
- Lang, S. (1972). Differentiable manifolds. Addison-Wesley.
- Lang, S. (1996). Differentiable manifolds and Riemannian geometry. Springer
- Lawson, B., & Michelsohn, M.-L. (1984). Embedding and surrounding with positive mean curvature. Inventiones math., 77, 399-419.
- Lawson, B., & Michelsohn, L. (1989). Spin geometry. Princeton University Press.
- Lawson, B., & Yau, S.-T. (1972). Compact manifolds of nonpositive curvature. J. differential geometry, 7, 211-228.
- Lawson, B., & Yau, S.-T. (1974). Scalar curvature, non-abelian group actions, and the degree of symmetry of exotic spheres. Commentarii math. helvetici, 49, 232-244.
- Lazutkin, V. (1993). KAM theory and semiclassical approximations to eigenfunctions. Springer.

- LeBrun, C. (1995). Einstein metrics and Mostow rigidity. Mathematical research letters(2), 1-8.
- LeBrun, C. (1996). Four-manifolds without Einstein metrics. Math. research letters, 3, 1-15.
- LeBrun, C., & Salamon, S. (1994). Strong rigidity of positive quaternionic manifolds. Inventiones math., 118, 109-132.
- Ledoux, M. (1992/93). Inégalités isopérimétriques en analyse et probabilités. In Séminaire Bourbaki, Astérisque 216 (pp. 343-376). Paris: Société math. de France.
- Levi-Civita, T. (1917). Nozione di parallelismo in una varietà qualunque e conseguente specificazione geometrica della curvatura Riemanniana. Rendiconti di cir. Palermo, 42, 73-205.
- Levi-Civita, T., & Ricci, G. (1901). Méthodes de calcul différentiel absolu et leurs applications. Math. Annalen, 54, 125-201.
- Lewin, L. (Ed.). (1991). Structural properties of polylogarithms. American math. soc.
- Lewy, H. (1938). On differential geometry in the large I. Transactions of the A.M.S., 43, 258-270.
- Lichnerowicz, A. (1950). Courbure, nombres de Betti, et espaces symétriques. In International congress of math., (pp. 216-223). Harvard:
- Lichnerowicz, A. (1955). Théorie globale des connexions et des groupes d'holonomie. Ed. Cremonese, Rome.
- Lichnerowicz, A. (1958). Géométrie des groupes de transformations. Dunod.
- Lichnerowicz, A. (1961). Propagateurs et commutateurs en relativité générale. Publications math. IHES, 10, 293-344.
- Lichnerowicz, A. (1963). Spineurs harmoniques. C. R. Acad. Sciences Paris, 257, 7-9.
- Llarull, M. (1996a). Sharp estimates and the Dirac operator. .
- Llarull, M. (1995). Scalar curvature and volume of a Riemannian manifold. Geometriae dedicata, 56, 1-4.
- Löbell, F. (1931). Beispiele geschlossener dreidimensionaler Clifford-Kleinscher Räume negativer Krümmung. Ber. Verh. Sächs. Akad. Wiss. Leipzig. Math-Phys. Kl., 83, 167-174.
- Lohkamp, J. (1992). Ricci curvature modulo homotopy. In A. Besse (Ed.), Table ronde de géométrie différentielle, (pp. 437-451). Luminy, Séminaires et congrès 1, Soc. math. France.

Lohkamp, J. (1994). Manifolds of negative Ricci curvature, Annals of math., 140, 655-683.

Lohkamp, J. (1996). Global and local curvatures. In M. Lovric, M. Min-Oo, & M. Wang (Eds.), Riemannian geometry (pp. 23-52). American math. society.

Lück, W. (1996). L^2 invariants of regular coverings of compact manifolds and CW-complexes (survey) No. Münster Universität.

Luo, W., & Sarnak, P. (1995a). Quantum ergodicity of eigenfunctions on $PSL_2(\mathbb{Z}) \backslash \mathbb{H}^2$. Publications math. I. H. E. S., 81, 207-237.

Luo, W., & Sarnak, P. (1996). Number variance for arithmetic hyperbolic surfaces. .

Luo, W., Rudnick, Z., & Sarnak, P. (1995b). On Selberg's eigenvalue conjecture. Geometric and functional analysis, 5, 387-401.

Lusternik, L., & Fet, A. (1951). Variational problems in closed manifolds. Doklady Akad. Nauk SSR, 81, 17-18.

Lusternik, L., & Schnirelman, L. (1929). Sur le problème de trois géodésiques fermées sur les surfaces de genre 0. C. R. Acad. Sciences Paris, 189, 269-271

Mané, R. (1987). Ergodic theory and differentiable dynamics. Springer.

Mané, R. (1996). On the topological entropy of geodesic flows. J. of differential geometry.

Manning, A. (1979). Topological entropy for geodesic flows. Annals of math., 110, 567-573.

Margerin, C. (1991, 1193, 1994). Théorie de la déformation en géométrie métrique No. M1005.0890, M1100.1293, M1073.0593). Ecole polytechnique.

Margulis, G. (1969). Applications of ergodic theory to the investigation of manifolds of negative curvature. Functional analysis and applic., 3, 335-336.

Margulis, G. (1977). Discrete groups of motions of manifolds of non-positive curvature. A M S Translations, 39, 33-45.

McDuff, D., & Salamon, D. (1995). Introduction to symplectic topology. Oxford: Clarendon Press.

McKean, H., & Singer, I. (1967). Curvature and the eigenvalues of the Laplacian. J. Differential Geometry, 1, 43-69.

Meyer, D. (1971). Sur les variétés riemanniennes à opérateur de courbure positif. C. R. Acad. Sciences Paris, 272, 482-485.

Meyer, W. (1989). Toponogov's theorem and its applications. In College on differential geometry, . Trieste: ITP Trieste.

- Micallef, M., & Moore, J. (1988). Minimal two-spheres and the topology of manifolds with positive curvature on totally isotropic two-planes. Annals of math., 127, 199-227.
- Michel, R. (1994). Restriction de la distance géodésique à un arc et rigidité. Bull. soc. math. France, 122, 435-442.
- Milnor, J. (1963). Morse theory. Princeton Univ. Press.
- Milnor, J. (1964). Eigenvalues of the Laplace operator on certain manifolds. Proc. nat. Acad. Sci. USA, 51, 542.
- Milnor, J. (1966). Whitehead torsion. Bulletin of the A.M.S., 72, 351-426.
- Milnor, J. (1968). A note on curvature and the fundamental group. J. differential geometry, 2, 1-7.
- Milnor, J., & Stasheff, J. (1974). Characteristic classes. Annals of math. studies 76, Princeton University Press.
- Min-Oo, & Ruh, E. (1979). Comparison theorems for compact symmetric spaces. Annales scientif. Ecol. norm. sup., 12, 335-353.
- Min-Oo, & Ruh, E. (1981). Vanishing theorems and almost symmetric spaces of noncompact type. Math. Annalen, 257, 419-433.
- Minakshisundaram, S. (1953). Eigenfunctions on Riemannian manifolds. J. indian math. Soc., 17, 118-165.
- Minakshisundaram, S., & Pleijel, A. (1949). Some properties of the eigenfunctions of the Laplace operator on Riemannian manifolds. Canadian J. of math., 1, 242-256.
- Misner, C., Thorne, K., & Wheeler, J. (1973). Gravitation. Freeman.
- Mok, N. (1988). The uniformization theorem for compact Kähler manifolds of nonnegative holomorphic bisectional curvature. J. of differential geometry, 27, 179-214.
- Morgan, F. (1988). Geometric measure theory : a beginner's guide. Academic Press.
- Morgan, J. (1996). The Seiberg-Witten equations and applications to the topology of smooth 4-manifolds. Math. notes 44, Princeton university press.
- Mori, S. (1979). Projective manifolds with ample tangent bundles. Annals of math., 110, 593-606.
- Morse, M. (1921a). A one-to-one representation of geodesics on a surface of negative curvature. American J. of math., 43, 33-51.
- Morse, M. (1921b). Recurrent geodesics on a surface of negative curvature. Transactions of the A. M. S., 22, 84-100.

- Morse, M. (1934). Calculus of variations in the large. Providence: American math. soc.
- Mostow, G. (1968). Quasi-conformal mappings in n -space and the rigidity of hyperbolic space forms. Public. math. I H E S, 34, 53-104.
- Mostow, G. D. (1980). A remarkable class of polyhedra in complex hyperbolic space. Pacific J. of Math., 86, 171-276.
- Mostow, G., & Siu, Y.-T. (1980). A compact Kähler surface of negative curvature non covered by the ball. Annals of math., 112, 321-360.
- Müller, W. (1978). Analytic torsion and R-torsion of Riemannian manifolds. Advances in math., 28, 233-305.
- Munoz, M., J., & Valdés, A. (1996). Génération d'anneaux d'invariants différentiels des métriques riemanniennes. C. R. Acad. Sc. Paris, 323, 643-646.
- Myers, S. (1935a). Riemannian manifolds in the large. Duke math. J., 1, 39-49.
- Myers, S. (1935b). Connections between differential geometry and topology, I. Duke math. J., 35, 376-391.
- Myers, S. (1936). Connections between differential geometry and topology, II. Duke math. J., 2, 95-102.
- Myers, S. (1941). Riemannian manifolds with positive mean curvature. Duke math. J., 8, 401-404.
- Nadirashvili, N. (1996). Hadamard's and Calabi-Yau's conjectures on negatively curved and minimal surfaces. Inventiones math., 126, 457-466.
- Nash, J. (1954). C^1 -isometric imbeddings. Annals of math., 60, 383-396.
- Nash, J. (1956). The embedding problem for Riemannian manifolds. Annals of math., 63, 20-63.
- Nirenberg, L. (1953). The Weyl and Minkowski problems in differential geometry in the large. Communications in pure and applied math., VI, 337-394.
- Nitsche, J. (1996). Minimal surfaces. Springer.
- Nomizu, K. (1954). Affine connections on homogeneous spaces. American j. of math., 76, 33-65.
- O'Neill, B. (1966b). The fundamental equations of a submersion. Michigan math. journal, 13, 459-469.
- O'Neill, B. (1983). Semi-Riemannian geometry. Academic Press.
- Obata, M. (1962). Certain conditions for a Riemannian manifold to be isometric with a sphere. J. math. Soc. Japan, 14, 333-340.

- Oesterlé, J. (1992/93). Polylogarithmes. In Séminaire Bourbaki, Astérisque 216 (pp. 49-68). Paris: Société math. de France.
- Osgood, O., Phillips, R., & Sarnak, P. (1989). Moduli space, heights and isospectral sets of plane domains. Annals of Math., 129, 293-362.
- Otal, J.-P. (1990a). Le spectre marqué des surfaces à courbure négative. Annals of math., 131, 151-162.
- Otal, J.-P. (1990b). Sur les longueurs des géodésiques d'une métrique à courbure négative dans le disque. Commentarii math. helvetici, 65, 334-347.
- Otsu, Y. (1991). On manifolds of positive Ricci curvature and large diameter. Math. Zeitschrift, 2, 255-264.
- Otsu, Y. (1993). On manifolds of small excess. American J. of math., 115, 1229-1280.
- Otsu, Y., & Shioya, T. (1994). The Riemannian structure of Alexandrov spaces , Journal of differential geometry 39, 629-658.
- Otsu, Y., Shiohama, K., & Yamaguchi, T. (1989). A new version of the differentiable sphere theorem. Inventiones math., 98, 219-228.
- Page, D. (1979). A compact gravitational instanton. Phys. letters, 79 B, 235-238.
- Palais, R., & Smale, S. (1964). A generalized Morse theory. Bulletin of the Amer. math. soc., 70, 165-172.
- Palais, R., & Terng, C.-L. (1988) Critical point theory and submanifold geometry. LNM 1353, Springer
- Pandey, A., Bohigas, O., & Giannoni, M. (1984). Characterization of chaotic quantum spectra and universality of level fluctuations laws. Physical review letters, 52, 1, 1-4.
- Pansu, P. (1983-1984). Effondrement des variétés riemanniennes. In Séminaire Bourbaki, 618 (pp. 63-82). Paris: Société math. France.
- Pansu, P. (1989). Métriques de Carnot-Carathéodory et quasi-isométries des espaces symétriques de rang 1. Annals of math., 129, 1-60.
- Pansu, P. (1990/91). Le flot géodésique des variétés riemanniennes à courbure négative. In Séminaire Bourbaki, Astérisque 201-202-203 (pp. 269-298). Paris: Société math. France.
- Pansu, P. (1993-94). Sous-groupes discrets des groupes de Lie : rigidité, arithméticité. In Séminaire Bourbaki, 227 (pp. 69-105). Paris, Astérisque 618, Société math. France.
- Pansu, P. (1996). Introduction to L^2 Betti numbers. In M. Lovric, M. Min-Oo, & M. Wang (Eds.), Riemannian geometry (pp. 53-86). Fields Institute monographs, American math. soc.

Pansu, P. (1996/97). Volume, courbure et entropie (d'après G. Besson, G. Courtois et S. Gallot). In Séminaire Bourbaki, . Paris, Astérisque , Société math. France.

Parry, W., & Pollicott, M. (1990). Zeta functions and the periodic orbit structure of hyperbolic dynamics. Astérisque 187-188, Société math. France.

Paternain, G. (1993b). Real analytic convex surfaces with positive topological entropy and rigid body dynamics. Manuscripta math., **78**, 397-402.

Paternain, G. (1993b). Real analytic convex surfaces with positive topological entropy and rigid body dynamics. Manuscripta math., **78**, 397-402.

Paternain, G., & Paternain, M. (1994). Topological entropy versus geodesic entropy. International J. of mathematics.

Patodi, V. (1971). Curvature and the eigenforms of the Laplace operator. J. of differential geom., **5**, 233-249.

Pelletier, F., & Valère Bouche, L. (1992). The problem of geodesics, intrinsic derivation and the use of control theory in singular sub-riemannian geometry. In A. Besse (Ed.), Table ronde de géométrie différentielle, 1 (pp. 453-512). Luminy, Séminaires et congrès 1, Société math. France.

Perelman, G. (1994a). Manifolds of positive Ricci curvature with almost maximal volume. Journal of the AMS, **7**, 299-305.

Perelman, G. (1994b). Proof of the soul conjecture of Cheeger and Gromoll. J. of differential geom., **40**, 209-212.

Perelman, G. (1994c). Spaces with curvature bounded below. In International Congress of mathematicians, (pp. 517-525). Zürich: Birkhäuser.

Perelman, G. (1995a). Widths of nonnegatively curved spaces. Geometric and functional analysis, **5**, 445-463.

Perelman, G. (1995b). A diameter sphere theorem for manifolds of positive Ricci curvature. Math. Zeitschrift, **218**, 595-595.

Perelman, G. (1997). Constructions of manifolds of positive Ricci curvature with big volume and large Betti numbers. .

Pesce, H. (1992). Nilvariétés isospectrales. In A.-L. Besse (Ed.), Table ronde de géométrie différentielle, 1 (pp. 513-530). Luminy, Séminaires et congrès 1, Société math. France 1996.

Peters, S. (1984). Cheeger's finiteness theorem for diffeomorphism classes of manifolds. J. für reine und angew. math., **349**, 77-82.

Peters, S. (1987). Convergence of Riemannian manifolds. Compositio math., **62**, 3-16.

- Petersen, P. (1990). Gromov-Hausdorff convergence of metric spaces. In R. Greene & S.-T. Yau (Eds.), Differential geometry, 54, part 3 (pp. 489-504). Los Angeles: American math. society.
- Petersen, P. (1996e). Comparison geometry problem list. In M. Lovric, M. Mir-Oo, & M. Wang (Eds.), Riemannian geometry (pp. 87-115). Fields Institute monographs, American math. soc.
- Petersen, P. (1997b). Riemannian geometry. Springer.
- Petersen, P. (1997c). Convergence theorems in Riemannian geometry. In Comparison geometry MSRI publications, Cambridge University Press.
- Petersen, P., & Wei, G. (1996b). Relative volume comparison with integral curvature bounds. .
- Petersen, P., & Wei, G. (1996c). Analysis and geometry on manifolds with integral Ricci curvature bounds. .
- Petersen, P., Shteingold, S., & Wei, G.-F. (1996a). Comparison geometry with integral curvature bounds. .
- Petersen, P., Wei, G., & Ye, R. (1997a). Control geometry via smoothing. .
- Petroll, D. (1996) Existenz und Transversalität von homoklinen und heteroklinen Orbits beim geodätischen Fluss. Thesis, Universität Freiburg.
- Pinkall, U., & Sterling, I. (1989). On the classification of constant mean curvature tori. Annals of math., 130, 407-451.
- Pinsky, M. (1989). Inverse questions in stochastic differential geometry. In Probabilities, . Singapour, de Gruyter.
- Pinsky, M. (1990). Feeling the shape of a manifold with Brownian motion. In Barlow & Bingham (Eds.), Stochastic analysis , Cambridge University press.
- Pogorelov, A. (1973). Extrinsic geometry of convex surfaces. American mathematical society.
- Poincaré, H. (1905). Sur les lignes géodésiques des surfaces convexes. Trans. Amer. Math. Soc., 6, 237-274.
- Prasad, G. (1989). Volumes of S-arithmetic quotients of semi-simple Lie groups. Publications math. IHES, 69, 91-117.
- Preissmann, A. (1942-43). Quelques propriétés globales des espaces de Riemann. Comm. math. Helvetici, 15, 175-216.
- Prüfer, F., Tricerri, F., & Vanhecke, L. (1996). Curvature invariants, differential operators and local homogeneity. Transactions of the A M S., 348, 4643-4651.
- Przytycki, F. (1980). An upper estimation of the topological entropy of diffeomorphisms. Inventiones math., 59, 205-213.

- Pu, P. (1952). Some inequalities in certain nonorientable manifolds. Pacific J. Math., 2, 55-71.
- Püttmann, T. (1997). Improved pinching constants of odd-dimensional homogeneous spaces, Bochum University.
- Rademacher, H. (1994a). On a generic property of geodesic flows. Math. Annalen, 298, 101-116.
- Rademacher, H. (1994b). The Fadell-Rabinowitz index and closed geodesics. J. London math. soc., 50, 609-624.
- Ratcliffe, J. (1994). Foundations of hyperbolic manifolds. Springer.
- Rauch, H. (1951). A contribution to differential geometry in the large. Annals of math., 54, 38-55.
- Ray, D., & Singer, I. (1971). R-torsion and the Laplacian on Riemannian manifolds. Advances in math., 7, 145-210.
- Real, C. (1996). Métriques d'Einstein-Kähler sur les variétés à première classe de Chern positive. C. R. Acad. Sc. Paris, 322, 461-464.
- Reshetnyak, Y. (1993). Geometry IV. Encyclopedia of math. sciences 70, Springer.
- Ritoré, M., & Ros, A. (1992). Stable constant mean curvature tori and the isoperimetric problem in three space form. Commentarii math. helvetici, 67, 293-305.
- Robert, G. (1994) Invariants topologiques et géométriques reliés aux longueurs des géodésiques et aux sections harmoniques de fibrés. Grenoble.
- Rong, X. (1993). The limiting eta-invariant of collapsed 3-manifolds. J. of differential geometry, 37, 535-568.
- Rong, X. (1995). Rationality of geometric signatures of complete 4-manifolds. Inventiones math., 120, 513-554.
- Rong, X. (1996a). On the fundamental group of manifolds of positive sectional curvature. Annals of math., 143, 397-411.
- Rong, X. (1996b). The almost cyclicity of the fundamental group of positively curved manifolds. Inventiones math., 126, 47-64.
- Rong, X. (1997). Positive curvature, local and global symmetry, and fundamental group.
- Rosenberg, J., & Stolz, S. (1994). Manifolds of positive scalar curvature. In G. Carlsson, R. Cohen, & W.-C. Hsiang J. (Eds.), Algebraic topology and its applications (pp. 241-267). Springer.
- Rudnick, Z., & Sarnak, P. (1996). The behaviour of eigenstates of arithmetic hyperbolic manifolds. .

Ruh, B. (1985). Krümmungstreue Diffeomorphismen Riemannscher und pseudo-Riemannscher Mannigfaltigkeiten. Math. Zeitschrift, **189**, 371-391.

Ruh, E. (1971). Curvature and differentiable structures on spheres. Commentarii math. helvetici, **46**, 127-136.

Ruh, E. (1982). Almost flat manifolds. J. of differential geom., **17**, 1-14.

Sachs, R., & Wu, H. (1977). General relativity for mathematicians. Springer.

Sakai, T. (1978). Cut-loci of compact symmetric spaces. In Minimal submanifolds and geodesics, (pp. 193-208). Tokyo: Kaigai publications, Tokyo.

Sakai, T. (1996). Riemannian geometry. American math. soc.

Salamon, S. (1982). Quaternionic manifolds. Symposia math., **XXVI**, 139-151.

Salamon, S. (1989). Riemannian geometry and holonomy groups. Longman scientific and technical.

Salamon, S. (1996). On the cohomology of Kähler and Hyper-Kähler manifolds. Toology, **35**, 137-155.

Sambusetti, A. (1996). An obstruction to the existence of Einstein metrics on 4-manifolds. C. R. Acad. sciences Paris, **322**, 1213-1218.

Sampson, J. (1986). Some applications of harmonic maps to Kähler geometry. Contemporary math., **49**, 125-134

Santalo, L. (1976). Integral geometry and geometric probability. Addison-Wesley.

Sarnak, P. (1995). Arithmetic Quantum Chaos. In Isr. math. conf. (pp. 183-236).

Sarnak, P. (1997a). Extremal geometries. Contemporary math., **201**, 1-7.

Sarnak, P. (1997b). Values at integers of binary quadratic forms No. Princeton University.

Sarnak, P. (1997c). Spectra and eigenfunctions of the Laplacian No. Princeton University.

Savo, A. (1996a). Une méthode de symétrisation et quelques applications, Comptes rendus Acad. Sc. Paris, **322**, 861-864

Savo, A. (1997). A mean-value lemma and applications.

Schmidt, E. (1948-1949). Die Brunn-Minkowskische Ungleichungen und ihr Spielgebild sowie die isoperimetrische der Kugel in der euklidischen und nichteuklidischen Geometrie I , II. Math. Nachrichten, **1, 2**, 81-157, 171-244.

Schneider, R., & Wieacker, J. (1993). Integral geometry. In Handbook of convexity (pp. 1349-1390). North-Holland.

Schnirelman, A. (1974). Ergodic properties of eigenfunctions (in Russian), Usp. math. Nauk, 29, 181-182.

Schoen, R. (1991). Recent progress on non-linear problems in geometry. J. of differential geometry, surveys.

Schoen, R., & Yau, S.-T. (1979). On the structure of manifolds with positive scalar curvature. Manuscripta math., 28, 159-183.

Schoen, R., & Yau, S.-T. (1981). Proof of the positive mass theorem. Communications in math. physics, 79, 231-260.

Schouten, J. (1954). Ricci-calculus. Springer.

Schwartz, R. (1995). The quasi-isometry classification of rank one lattices. Publications math. IHES, 82, 133-168.

Seiberg, N., & Witten, E. (1994). Monopoles, duality and chiral symmetry breaking in N=2 supersymmetric QCD. Nuclear physics B, 341, 581-640.

Semmes, S. (1996). Where the buffalo roam : infinite process and infinite complexity No. M/96/77). I H E S.

Serre, J.-P. (1951). Homologie singulière des espaces fibrés. Annals of Math., 54, 425-505.

Sha, J., & Yang, D. (1991). Positive Ricci curvature on the connected sums of $S^n \times S^m$. J. of differential geometry, 33, 127-137.

Sha, J.-P., & Yang, D. (1993). Positive Ricci curvature on compact simply connected 4-manifolds. In R. Greene & S.-T. Yau (Ed.), Differential geometry, 54, part 3 (pp. 529-538). Los Angeles: American math. soc. 1993

Shen, Z. (1996). Complete manifolds with nonnegative Ricci curvature and large volume growth. Inventiones math., 125, 393-404.

Shi, W.-X. (1989). Deforming the metric in complete manifolds. J. of differential geometry, 30, 225-301.

Shikata, Y. (1967). On the differentiable pinching problem. Osaka math. J., 4, 279-287.

Shimizu, Y., & Shudo, A. (1995). Polygonal billiards : correspondence between classical trajectories and quantum eigenstates. Chaos, solitons and fractals, 5, 1337-1362.

Shiohama, K. (1983). A sphere theorem for manifolds of positive Ricci curvature. Transactions of A M S, 257, 811-819.

Shiohama, K. (1990). Recent developments in sphere theorems. In R. Greene & S.-T. Yau (Ed.), Differential geometry, 54 (pp. 551-576). Los Angeles: American math. soc. 1993

Shiohama, K., & Tanaka, M. (1992). Cut-loci and distance spheres in Alexandrov surfaces. In A. Besse (Ed.), Table ronde de géométrie

différentielle, 1 (pp. 531-560). Luminy, Séminaires et congrès 1, Société math. France.

Shioya, T. (1992). Geometry of total curvature. In A. Besse (Ed.), Table ronde de géométrie différentielle, 1 (pp. 561-600). Luminy, Séminaires et congrès 1, Société math. France.

Simon, L. (1983). Lectures on geometric measure theory. Canberra: Australian national university.

Simons, J. (1962). On the transitivity of holonomy systems. Annals of math. 76, 213-234.

Sinai, Y. (1976). Introduction to ergodic theory. Math. notes 18, Princeton Univ. Press.

Singer, I. (1960). Infinitesimally homogeneous spaces. Communications in pure and applied math., 13, 685-697.

Singer, I., & Thorpe, J. (1969). The curvature of 4-dimensional Einstein spaces. In Global analysis, papers in honour of K. Kodaira (pp. 355-365). Princeton University Press.

Siu, Y.-T. (1980). The complex analyticity of harmonic maps and the strong rigidity of compact Kähler manifolds. Annals of math., 112, 73-111.

Siu, Y.-T., & Yau, S.-T. (1980). Compact Kähler manifolds of positive bisectional curvature. Inventiones math., 59, 189-204.

Skorniakov, L. (1955). Metrization of the projective plane in connection with a given system of curves. Izvestiya Akad. Nauk SSSR, 19, 471-482.

Sommese (1975). Quaternionic manifolds, Math. Annalen, 212, 191-214

Spatzier, R. (1990). Riemannian manifolds with completely integrable geodesic flows. In R. Greene & S.-T. Yau (Ed.), Differential geometry, 54, 3 (pp. 599-608). Los Angeles: American math. soc. 1993

Spivak, M. (1970). A comprehensive introduction to differential geometry. Publish or Perish.

Steenrod, N. (1951). The topology of fiber bundles. Princeton university press.

Stolz, S. (1992). Simply connected manifolds of positive scalar curvature. Annals of math., 136, 511-540.

Stroock, D. (1996). Some thoughts about Riemannian structures on path space. Gazette Soc. math. France, 68, 31-45.

Struwe, M. (1988). Plateau's problem and the calculus of variations. Princeton University Press.

Study, E. (1905). Kürzeste Wege im komplexen Gebiete. Math. annalen, 60, 321-377.

Sullivan, D. (1977). Infinitesimal computations in topology. Publications math. IHES, 47, 269-332.

Sunada, T. (1985). Riemannian coverings and isospectral manifolds. Annals of math., 121, 169-186.

Suyama, Y. (1995). Differentiable sphere theorem by curvature pinching. Tohoku math. J., 47, 15-29.

Svarc (1955). Doklady Akad. Nauk SSSR, 105, 32-34.

Synge, J. (1936). On the connectivity of spaces of positive curvature. Quarterly J. math. Oxford, 7, 316-320.

Szabo, Z. I. (1991). A short topological proof for the symmetry of 2-point homogeneous spaces. Inventiones math., 106, 61-64.

Taimanov, I. (1993). On the existence of three non self-intersecting closed geodesics on manifolds homeomorphic to the 2-sphere. Russian Acad. Sci. Izv. Math., 40, 565-590.

Taimanov, I. (1997). On totally geodesic embeddings of 7-dimensional manifolds into 13-dimensional positively curved manifolds. , Math. sbornik.

Takeuchi, M. (1979). On the conjugate loci and cut-loci of compact symmetric spaces II. Tsukuba Journal of math., 3, 1-29.

Tanno, S. (1980). A characterization of canonical sphere by the spectrum. Math. Zeitschrift, 175, 267-274.

Taubes, C. (1992). The existence of anti-self dual conformal structures. J. of differential geometry, 36, 163-253.

Terng, C.-L. (1990). Recent progress in submanifold geometry. In R. Greene & S.-T. Yau (Ed.), Differential geometry, 54, part 1 (pp. 439-484). Los Angeles: American math. soc. 1993.

Terng, C.-L., & Thorbergsson, G. (1995). Submanifolds geometry in symmetric spaces. J. differential geometry, 42, 665-718.

Thorpe, J. (1969). Some remarks on the Gauss-Bonnet integral. J. math. mech., 18, 779-786.

Thurston, W. (1978). Geometry and topology of 3-manifolds. Princeton University.

Tian, G. (1990). On Calabi's conjecture for complex surfaces. Inventiones math., 101, 101-172.

Tian, G. (1991). On one of Calabi's problem. In Proceedings of symposia in pure math. 52, Bedford, E., d'Angelo, J., Greene, R., & Krantz, S. (Ed.) (volume 2, pp. 543-556). American math. soc. 1991

Tian, G. (1996). Kähler-Einstein metrics with positive scalar curvature No.

Tits, J. (1955). Sur certaines classes d'espaces homogènes de groupes de Lie. Mémoires de l'Académie des Sciences, XXIX, Bruxelles

Toledo, D. (1997). Rigidity theorems in Kähler geometry and fundamental groups of varieties. .

Tondeur, P. (1988). Foliations on Riemannian manifolds. Springer.

Toponogov, V. (1959). Riemann spaces with curvature bounded below. Uspehi Mat. Nauk, 14.

Toponogov, V. (1964). Riemannian spaces which contain straight lines. Americ. math. soc. translations, 37, 287-290.

Tricerri, F., & Vanhecke, L. (1983). Homogenous structures on Riemannian manifolds. Cambridge university press.

Tricerri, F., & Vanhecke, L. (1986). Variétés riemanniennes dont le tenseur de courbure est celui d'un espace symétrique riemannien irréductible. C.R. Acad. Sc. Paris, 302, 233-235.

Tsukamoto, C. (1981). Infinitesimal Blaschke conjecture on projective spaces. Ann. scient. Ec. norm. sup., 14, 339-356.

Tuschmann, W. (1996). Collapsing, solvmanifolds and infrahomogeneous spaces. Geometriae dedicata.

Vérovic, P. (1996) Entropie et métriques de Finsler. Joseph Fourier, Grenoble I.

Vigneras, M.-F. (1980a). Arithmétique des algèbres de quaternions. LNM 800, Springer.

Vignéras, M.-F. (1980b). Variétés riemanniennes isospectrales et non isométriques. Annals of math., 112, 21-32.

Ville, M. (1985a). On $(1/4)$ -pinched 4-dimensional manifolds of negative curvature. Annals of global analysis and geometry, 3, 329-336.

Ville, M. (1987). Sur le volume des variétés riemanniennes pincées. Bull. soc.math. France, 115, 127-139.

Vinberg, E. B. (1993a). Geometry II. Encyclopedia of math. sciences 22, Springer.

Vinberg, E. B. (1984). The absence of crystallographic groups of reflections in Lobachevskii spaces of large dimension. Trans. Mosc. Math. Soc., 1985, 75-112.

Voisin, C. (1996). Symétrie miroir. Société math. France.

Wall, C. T. C. (1977). Geometric properties of generic differentiable manifolds. In P. a. d. Carmo (Eds.), Geometry and topology (pp. 707-774). Springer.

Walters, P. (1982). An introduction to ergodic theory. Springer.

- Wang, H. (1952). Two-point homogeneous spaces. Annals of math., **55**, 177-191.
- Wang, H.-C. (1972). Topics on totally discontinuous groups. In W. Boothby & G. Weiss (Eds.), Symmetric spaces (pp. 459-487). Marcel Dekker.
- Wang, M. (1982). Some examples of homogeneous Einstein manifolds in dimension seven. Duke math. j., **49**, 23-28.
- Wang, M. (1992). Einstein metrics and quaternionic Kähler manifolds. Math. Zeitschrift, **210**, 305-326.
- Wang, M., & Ziller, W. (1986). Einstein metrics with positive scalar curvature. In Curvature and topology of Riemannian manifolds Springer.
- Warner, F. (1965). The conjugate locus of a Riemannian manifold. American j. of math., **87**, 575-604.
- Weil, A. (1926). Sur les surfaces à courbure négative. C. R. Acad. sc. Paris, **182**, 1069-1071.
- Weil, A. (1958). Variétés kählériennes. Hermann.
- Weinstein, A. (1967). On the homotopy type of positively pinched manifolds. Archiv der math., **18**, 523-524.
- Weinstein, A. (1970). Sur la non densité des géodésiques fermées. C. R. Acad. Sciences Paris, **271**, 504.
- Weinstein, T. (1996). An introduction ot Lorentz surfaces. Walter de Gruyter.
- Weiss, M. (1993). Pinching and cooncordance theory. J. of diffential geometry, **38**, 387-416.
- Weitzenböck, R. (1923). Invariantentheorie. Groningen: Noordhoff.
- Wells, R. (1980). Differential analysis on complex manifolds. Springer.
- Wente, H. (1986). Conterexample to a conjecture of H. Hopf. Pacific j. of math., **121**, 193-243.
- Weyl, H. (1911). Über die asymptotische Verteilung der Eigenwerte. Nach. d. Königl. Ges. d. Wiss. zu Göttingen, 110-117.
- Whitehead, H. (1935). On the covering of a complete space by geodesics. Annals of math., **36**, 679-704.
- Whitney, H. (1936). Differentiable manifolds. Annals of math., **37**, 645-680.
- Wilking, B. (1997). dissertation, Münster, Germany.
- Willmore, T. (1993). Riemannian geometry. Clarendon press Oxford.

Wirtinger, W. (1936). Eine Determinantenidentität und ihre Anwendung auf analytische Gebilde und Hermitesche Massbestimmung. Monatshefte für math. und physik, 44, 343-365.

Wolf, J. (1972). Spaces of constant curvature (second ed.). Berkeley: J. Wolf.

Wu, H. (1987). The Bochner technique in differential geometry. Math. reports, London:

Wu, J.-Y. (1990). Hausdorff convergence and sphere theorem. In R. Greene & S.-T. Yau (Ed.), Differential geometry, 54, part 3 (pp. 685-692). Los Angeles: American math. soc.

Yamabe, H. (1960). On a deformation of Riemannian structures on compact manifolds. Osaka math. J., 12, 21-37.

Yamaguchi, T. (1988). Homotopy type finiteness theorems for certain precompact families of Riemannian manifolds. Proc. American math. soc., 102, 660-666.

Yamaguchi, T. (1991). Collapsing and pinching under a lower curvature bound. Annals of math., 133, 317-357.

Yamaguchi, T. (1992). A convergence theorem in the geometry of Alexandrov spaces. In A.-L. Besse (Ed.), Table ronde de géométrie différentielle, (pp. 601-642). Luminy, Séminaires et congrès 1, Société math. France.

Yang, C. (1980). Fibre bundles and the physics of the magnetic monopoles. In W. Hsiang (Eds.), The Chern symposium Berkeley: Springer.

Yang, D. (1992). Convergence of Riemannian manifolds with integral bounds on curvature I and II. Ann. Scientifiques E N S., 25, 75-105 and 179-199.

Yau, S.-T. (1975). Harmonic functions on complete Riemannian manifolds. Comm. pure and appl. math., 28, 201-228.

Yau, S.-T. (1977). On Calabi's conjecture and some new results in algebraic geometry. Proceed. nat. acad. sciences USA, 74, 1798-1799.

Yau, S.-T. (1978). On the Ricci curvature of a compact Kähler manifold and the complex Monge-Ampère equation I. Communic. pure and applied math., XXXI, 339-411.

Yau, S.-T. (1982). Problem section. In S.-T. Yau (Eds.), Seminar on differential geometry (pp. 669-706). Annals of math. studies 102, Princeton Univ. Press.

Yau, S.-T. (1987). Nonlinear analysis in geometry. L'enseignement math., 33, 109-158.

Yau, S.-T. (1990). Open problems in geometry. In R. Greene & S.-T. Yau (Ed.), Differential geometry, 54, part 3 (pp. 1-28). Los Angeles: American math. soc. 1993

Yau, S.-T., & Zheng, F. (1991). Negatively $(1/4)$ -pinched Riemannian metric on a compact Kähler manifold. Inventiones math., 103, 527-535.

- Yomdin, Y. (1987). Volume growth and entropy. Israel J. of Math., 57, 285-300.
- Yu, G. (1997). Zero-in-the-spectrum conjecture, positive scalar curvature and asymptotic dimension. Inventiones math., 127, 96-126.
- Zamfirescu, T. (1996). Points joined by three shortest paths on convex surfaces. .
- Zelditch, S. (1987). Uniform distribution of eigenfunctions on compact hyperbolic surfaces. Duke math. J., 55, 919-941.
- Zelditch, S. (1992). Selberg trace formula and equidistribution theorems. Memoirs of the AMS, 96, vol. 465, Providence: American math. Soc.
- Ziller, W. (1982). Geometry of the Katok example. Ergodic theory and dyn. systems, 3, 135-157.
- Zoll, O. (1903). Über Flächen mit Scharen geschlossener geodätischer Linien. Math. Annalen, 57, 108-133.