Prospects for precise electroweak physics at LHC

Esben Klinkby
for the ATLAS and CMS collaborations

DIS 2009, 26-30 April 2009, Madrid

Duke
UNIVERSITY

- ▶ At tree level in the SM 3 parameters are sufficient to predict the EW sector: G_F , α , $\sin(\Theta_W)$.
- ▶ Precision EW physics look for deviations from this, introduced via:

- **Loops:**

$$\Rightarrow M_{W,tree}^2 = \pi\sqrt{2}\frac{\alpha}{G_F} \left(1 - \sqrt{1 - \frac{4\pi\alpha}{\sqrt{2}G_F M_Z^2}}\right)^{-1}$$

$$G_F = \frac{\pi\alpha}{\sqrt{2}M_W^2 \sin^2\theta_W} \cdot \frac{1}{1 - \Delta r}$$

$$\Delta r = \Delta r^t + \Delta r^H$$

$$\Delta r^t \approx \frac{3G_F m_t^2}{8\sqrt{2}\pi^2} \left(\frac{\cos^2\theta_W}{\sin^2\theta_W} \right)$$

$$\Delta r^H \approx \frac{11G_F M_W^2}{24\sqrt{2}\pi^2} \left(\ln \frac{M_H^2}{M_W^2} - \frac{5}{6} \right)$$

- **Couplings** that differ from standard model predictions. E.g.

- ▶ Check SM & search for new physics via modifications of M_W , m_t and M_H relations & couplings.

Electroweak characteristics at LHC

In general:

- Cross-sections increase with $\sqrt{s}$
- Expected instantaneous luminosity $10^{33} \text{ cm}^{-2} \text{ s}^{-1}$
(for most electroweak measurements)
- Many analyses benefit from large statistics (14TeV):

- ▶ **Top** - Cross section: $\sim 800 \text{ pb}$.
→ 0.8M events in 1 fb^{-1} .
→ 7.5K (5K) semileptonic for 1 b-tag (2 b-tag).
- ▶ **Single Top** – Cross sections 1/3 of $t\bar{t}$ bar
- ▶ **W** - NNLO Cross section: 40 nb ($e+\mu$)
→ $\sim 6 \text{ M}$ reconstructed events in 1 fb^{-1} ($e+\mu$).
- ▶ **DiBoson** Cross sections in the range 15-450pb
→ 1–1000 observed events in 1 fb^{-1} .
- ▶ **Forward-Backward asymmetry**
→ Based on Z sample $\sim 200 \text{ K}$ selected in 1 fb^{-1} .

top pair: production & decay

► Production:

All hadronic

- Huge background
- Complicated combinatorics
- 350.000 events in 1fb^{-1}

All leptonic

- Low background
- Simple combinatorics
- 50.000 events in 1fb^{-1}

But very **complex** events

all jets 44%

dileptons 9%

τ +jets 15%

μ +jets 15%

e+jets 15%

Semi leptonic

- Compromise
- 250.000 events in 1fb^{-1}

top mass: semileptonic channel

- ▶ Selecting jets for the W hypothesis:

After requiring $|M_{jj} - M_W| < 30 \text{ GeV}$, minimize *in situ* :

$$\chi^2 = \frac{(M_{jj}(\alpha_{E_{j1}}, \alpha_{E_{j2}}) - M_W^{\text{PDG}})^2}{(\Gamma_W^{\text{PDG}})^2} + \frac{(E_{j1}(1 - \alpha_{E_{j1}}))^2}{\sigma_1^2} + \frac{(E_{j2}(1 - \alpha_{E_{j2}}))^2}{\sigma_2^2}$$

In each event

- ▶ Additional cut: M_{jj} in mass window of $\pm 2\Gamma_{M_W} = 4.2 \text{ GeV}$

- ▶ Fitted with Gaussian + 3rd order pol.

- ▶ Result: $M_{\text{top}} = 175 \pm 0.2_{(\text{stat})} \text{ GeV} (1 \text{ fb}^{-1})$.

- ▶ Precision given by jet scale accuracy

I.e. 0.2 GeV if
JES is known
to 1%

Systematic uncertainty

Light jet energy scale	0.2 GeV/%
b jet energy scale	0.7 GeV/%
ISR/FSR	$\approx 0.3 \text{ GeV}$
b quark fragmentation	$\leq 0.1 \text{ GeV}$
Background	negligible
Method	0.1-0.2 GeV

Result: $M_{\text{top}} = 175 \pm 0.2_{(\text{stat})} \pm 1_{(\text{sys})} \text{ GeV with } 1 \text{ fb}^{-1}$.

Semileptonic Selection

- Exactly one isolated lepton:
 $|\eta| < 2.5, p_t^e > 25 \text{ GeV}, p_t^\mu > 20 \text{ GeV}$
- Missing $E_t > 20 \text{ GeV}$
- At least 4 jets with $p_t > 40 \text{ GeV}$

- Two of which are b-tagged

Backgrounds

- Combinatorial
- Single top
- W+jets & Z+jets
- ttbar: fully hadronic & dilepton
- QCD & DiBosons

Hadronic
top decay
(avoids ν -
reco)

top mass: leptonic and hadronic channels

Leptonic channel

- Kinematical reconstruction
- Clean signal: $S/B=12$
- Main systematics: JES, Kinematical hypothesis

- Result with 1fb^{-1} :

$$M_{\text{top}} = 178.5 \pm 1.5_{\text{(stat)}} \pm 4.2_{\text{(sys)}} \text{ GeV}$$

Hadronic channel

- Using likelihood techniques on masses and angles to define pairing.
- Main systematics: QCD background, JES, ISR/FSR

- Result with 1fb^{-1} : M_{inv} [GeV/c²]

$$M_{\text{top}} = 175.0 \pm 0.6_{\text{(stat)}} \pm 4.2_{\text{(sys)}} \text{ GeV}$$

W mass: Characteristics at LHC

Production:

Decay mode	Branching fraction
$e^+ \nu_e$	$10.75 \pm 0.13\%$
$\mu^+ \nu_\mu$	$10.57 \pm 0.15\%$
$\tau^+ \nu_\tau$	$11.25 \pm 0.20\%$
$q\bar{q}'$	$67.60 \pm 0.27\%$

For precision measurements only the $W \rightarrow e\nu$ & $W \rightarrow \mu\nu$ channels are useful.

Observables: lepton p_t & M_t . Systematics differ

Precision measurements requires extremely good measurement/knowledge of :

Systematics: Experimental sources

Theoretical sources

Environmental sources

- Lepton energy scale & linearity
- Missing Et calibration
- Backgrounds

- PDF
- FSR (effects the lepton directly)
- $y(W)$, $p_t(W)$, and ISR (effects the lepton though the W)

- Underlying event
- Pileup
- Beam crossing angle

Most calibration performed on well-known Z events

W mass: Template fitting

GENERAL IDEA:

A set of lepton p_T distributions corresponding to different input values of M_W .

STEP 1: Validate method of handling det. effects

- Fit $E_{\text{rec}}/E_{\text{truth}}$
- Smear truth according to fit parameters
- Check that smeared truth matches reco $\checkmark$

Fitted $E_{\text{rec}}/E_{\text{truth}}$ in bins $30 \text{ GeV} < p_t < 40 \text{ GeV}$

STEP 2: Calibrate on Z events

- Scale

- Technique works $\checkmark$

STEP 3: Transport to W events

$15 \text{ pb}^{-1}: M_W = 80.466 \pm 110_{\text{(stat)}} \pm 114_{\text{(exp)}} \pm 25_{\text{(PDF)}} \text{ MeV}$

W mass: Long term perspectives

Best fit corresponds to: $\alpha=0.9958\pm0.0003$ $\sigma=0.0207\pm0.0003$ for 200pb^{-1}

Since the impact on the W mass measurement goes like:

$$\frac{\partial M_W}{\partial_{\text{rel}}\alpha_l} \sim 800 \text{ MeV}/\% \quad \frac{\partial M_W}{\partial_{\text{rel}}\sigma_l} = 0.8 \text{ MeV}/\%$$

the error estimated on the W mass from scale and resolution uncertainty is (at 10fb^{-1}):

$$\delta M(\alpha) = \frac{\delta\alpha}{\alpha} \cdot \frac{\partial M}{\partial_{\text{rel}}\alpha} \sqrt{\frac{\mathcal{L}_{\text{sample}}}{\mathcal{L}_{\text{final}}}} = \frac{0.0003}{0.9952} \cdot 800 \text{ MeV}/\% \cdot \sqrt{\frac{200 \text{ pb}^{-1}}{10 \text{ fb}^{-1}}} \approx 4 \text{ MeV}$$

$$\delta M(\sigma) = \frac{\delta\sigma}{\sigma} \cdot \frac{\partial M}{\partial_{\text{rel}}\sigma} \sqrt{\frac{\mathcal{L}_{\text{sample}}}{\mathcal{L}_{\text{final}}}} = \frac{0.0003}{0.0207} \cdot 0.8 \text{ MeV}/\% \cdot \sqrt{\frac{200 \text{ pb}^{-1}}{10 \text{ fb}^{-1}}} \approx 1 \text{ MeV}.$$

▶ **LONG TERM PERSPECTIVES:** Energy dependent lepton scale and resolution

● Scale

● Resolution

Seems possible to get a relative precision of 2×10^{-5} for the scale and 2×10^{-3} for resolution

W mass: Breakdown of systematics (10fb⁻¹)

Eur.Phys.J.C57:627-651,2008.

Sys unc.	Method of evaluation	Error [MeV]
W rapidity	Using the strong correlation between Z and W rapidity + LHC improvement	1
W width	Templates study + Assume W width uncertainty reduced with LHC	0.5
W p _t	From Z (+Drell Yan), exploiting the lever arm induced by the large Z width.	3
QED	PHOTOS study + Assume existence of high precission event generators	<1
Scale & lin.	From energy dependent calibration using Z events	4
Resolution	----- -----	1
Efficiency	Tag and probe	4.5(e), 1(μ)
W→τν	Template	2
Z→l(l)	----- -----	0.3
Z→ττ	----- -----	0.1
QCD	Tevatron extrapolation	0.5
Pileup&U.E.	Assuming 2% precision in hadronic energy flow	1(e), 0(μ)
BeamCross.	Generator study	0.1
Total		~7(e), 6(μ)

Triple gauge bosons couplings (TGC) : Characteristics at LHC

- ▶ Cross sections typically 10 times higher than Tevatron
⇒ tens to hundreds events in the first fb^{-1} .
- ▶ Trilinear gauge boson couplings measured:
 - Directly from ZW, WW & ZZ cross sections
 - Boson p_t
 - Production angle
- ▶ Charged TGC: WWZ & WW γ : allowed in SM
⇒ Study WW ($\sigma=112\text{pb}$) & WZ ($\sigma=48\text{pb}$) processes
- ▶ Neutral TGC: ZZZ, ZZ γ & Z $\gamma\gamma$: forbidden in SM tree level
⇒ Study ZZ processes
- ▶ Modelled by 5 independent CP conserving couplings that don't break gauge invariance
- ▶ Complementary to direct searches for new physics
In fact background to Higgs and beyond SM physics

Triple gauge bosons: WZ selection

Z selection

- Reconstruct ee or μμ pair with: 50 GeV < M_{II} < 120 GeV
- Veto additional Z candidates

W reconstruction

- Associate 3rd lepton the W decay
- Require neutrino indirectly so that M_T(W) > 50 GeV

$$S_L = \sqrt{2 \ln Q}, \quad Q = \left(1 + \frac{N_S}{N_B}\right)^{N_S + N_B} e^{-N_S}.$$

5σ observation with less than 350pb⁻¹.
Expect observation of all di-boson processes with less than 1fb⁻¹.

Forward backward asymmetry

- In $pp \rightarrow Z/\gamma \rightarrow ee$ events, electroweak neutral current violates parity and leads to an asymmetry in the polar emission angle of the electron in the ee -restframe:

$$\frac{1}{\sigma} \frac{d\sigma}{d\cos\theta} = \frac{3}{8} N_c \left[1 + \frac{4}{3} A_{FB} \cos\theta + \cos^2\theta \right]$$

- Can be used to determine $\sin^2\Theta_{\text{eff}}$. Important for SM fit.
- Measurement assumption: the quark directions equals that of the Z boost.

Event Selection - ATLAS

- Electron trigger: $p_t^e > 20 \text{ GeV}$
- Dilepton rapidity > 1
- $|M_{(e+e^-)} - M_Z| < 6 \text{ GeV}$
- $E_{\text{tMiss}} < 20 \text{ GeV}$
- Either: $|\eta_1^e|, |\eta_2^e| < 2.5$ or $|\eta_1^e| < 2.5$ & $|\eta_2^e| < 4.9$

- Counting problem
- Prospects of measuring $\sin^2\Theta_{\text{eff}}$ with a stat precision of 10^{-4} with 100fb^{-1}
- Comparable systematic error dominated by PDF uncertainty ($\sim 2 \cdot 10^{-4}$)

Conclusions

- ▶ At LHC the challenge of precision electroweak measurements is a challenge of reducing syst.
- ▶ In the absence of unforeseen complications:
 - $\delta M_{\text{top}} = 1 \text{ GeV}$ & $\delta M_W = 7 \text{ MeV}$ seem reachable with 10fb^{-1} (after many years of hard work)
 - Measure anomalous TGC deviation from zero at the percent level
 - $\sin^2\Theta_{\text{eff}}$ measured at the 10^{-4} level
 - Direct measurement of V_{tb} at the $\sim\text{few } \%$ level.
- ▶ The electroweak section of the Standard Model will be even more constrained than presently.
- ▶ Useful not only for cross checking the model, but also to constrain extensions to the Standard Model

Backup slides: references

► W mass measurement

- ATLAS CSC note: CERN-OPEN-2008-020
- CMS Note 2006-061
- Eur.Phys.J.C57:627-651,2008.
- ATL-PHY-PUB-2006-007

► Top mass measurement

- ATLAS CSC note: CERN-OPEN-2008-020
- CMS Note 2006-066
- CMS Note 2006-077

► Forward backward asymmetry

- ATLAS CSC note: CERN-OPEN-2008-020

► Triple gauge couplings

- ATLAS CSC note: CERN-OPEN-2008-020
- CMS PAS EWK-08-003

► Single Top

- ATLAS CSC note: CERN-OPEN-2008-020

Backup slides: detectors

B field: 2 T solenoid + 4T toroid

3.8 T solenoid

Inner tracker: $|\eta|$ coverage
 $\sigma(p_T)/p_T$ at $p_T=100$ GeV

2.5
3.8%

2.5
1.5%

EM calorimeter: $|\eta|$ coverage
 $\sigma(E)/E$

3.2
 $9\%/\sqrt{E} + 0.5\%$

3.0
 $3\%/\sqrt{E} + 0.25\%$

HAD calorimeter: $|\eta|$ coverage
 $\sigma(E)/E$ (EM+HAD combined)

4.9
 $70\%/\sqrt{E} + 3.3\%$

5.2
 $70\%/\sqrt{E} + 8\%$

Muon system: $|\eta|$ coverage
 $\sigma(p_T)/p_T$ at $p_T=1$ TeV

2.7
7%

2.4
5%

top mass: semileptonic channel: systematics

Systematic uncertainty	Effect on M_{top}	Method of evaluation
Light jet energy scale	0.2GeV/%	Multiplying jet momenta by scaling factors $\Rightarrow$ linear dependence of M_{top} . Reduced due to the W mass constraint. Prospects of 1% precision with 1fb^{-1}
b jet energy scale	0.7GeV/%	Same procedure. Will ultimately be determined from Z+jets, but in the beginning from the light JES + MC correction modelling the difference in the scales.
ISR/FSR	$\approx 0.3\text{GeV}$	Estimated from shifts in M_{top} in AcerMC ttbar samples with varying ISR,FSR parameters as well as $\Lambda(\text{QCD})$.
b quark fragmentation	$\leq 0.1\text{GeV}$	Varying the Peterson parameter within its uncertainty
Background	negligible	Shape will be measured from data – variations of the size have no notable effect.

W mass: Impact of scale and resolution

Figure 5: Left: bias on m_W , $m_W^{fit} - m_W^{true}$, as a function of the relative bias on α_ℓ , $\delta\alpha_\ell = (\alpha_\ell^{fit} - \alpha_\ell^{true})/\alpha_\ell^{true}$. Right: bias on m_W as function of the resolution bias, $\delta\sigma_\ell = (\sigma_\ell^{fit} - \sigma_\ell^{true})/\sigma_\ell^{true}$. A linear dependence is observed in each case, with $\partial m_W / \partial_{rel} \alpha_\ell = 800 \text{ MeV}/\%$ and $\partial m_W / \partial_{rel} \sigma_\ell = 0.8 \text{ MeV}/\%$.

W mass: Impact of scale and resolution

† Differentiate in energy (i.e consider lepton energy bins i, j).

Fit for every pair configuration (i, j) :

† $M_{ij}^2 = E_i E_j (1 - \cos\theta) ; (1 + \beta_{ij})^2 M_{ij}^2 = (1 + \alpha_i) E_i (1 + \alpha_j) E_j (1 - \cos\theta)$

† $\Rightarrow \beta_{ij} \sim (\alpha_i + \alpha_j)/2 ; \sigma_{ij}^2/M^2 = \sigma_i^2/E_i^2 + \sigma_j^2/E_j^2 ;$ write this for all (i, j)

† and solve the linear system (least squares) to get the α_i and σ_i^2

▶ Templates from Z events : Scaled observable method

- – using lepton p_t as observable
- Remove randomly 1 e from $Z \rightarrow ee$
- Rescale the observable: $X_v = p_t / M_v$
- Weight by $R(X) = \frac{d\sigma^W / dX_W}{d\sigma^Z / dX_Z}$, $X_v = \frac{p_t^e}{M_v}$, $V = W, Z$
- Which depends on theory, selection and det effects
- Apply W selection on Z events using the scale.
e.g. Missing $E_t > 29 \text{ GeV} \times M_W / M_Z$

Results: $\pm 40_{\text{(stat)}} \pm 40_{\text{(exp)}} \pm 40_{\text{(theory)}} \text{ MeV with } 1\text{fb}^{-1}$

▶ Templates from Z events : Morphing

- Using M_t as observable
- Scaling the Z event rather than the observable

Results: $\pm 40_{\text{(stat)}} \pm 64_{\text{(exp)}} \pm 20_{\text{(theory)}} \text{ MeV with } 1\text{fb}^{-1}$

▶ Template fitting – calibrate using Z templates: next slides

Triple gauge bosons: Sensitivity at LHC

Limit on anomalous TGC (95%CL) for WWZ assuming $\Delta\kappa_Z = g^Z_1$. Systematics errors included.

single top

- ▶ With 2M ($10^{33} \text{ cm}^{-2} \text{ s}^{-1}$ at 14TeV) events produced per year, measurements of the individual cont. to the single top cross section seem feasible.
- ▶ Comparison with SM predictions provides a crucial test (sensitive to W' or charged Higgs)
- ▶ Direct measurement of V_{tb} at the $\sim \text{few } \%$ level.
- ▶ Complex analysis: Huge $t\bar{t}$ and $W+\text{jets}$ bkg.
Require good knowledge of JES
and well-understood b-tagging tools
 $\Rightarrow$ Use likelihood or BDT techniques
- ▶ 3σ evidence achievable with a few fb^{-1}
and 20% precision on cross section with 10fb^{-1} .