ATLAS Discovery Potential of the Standard Model Higgs Boson

Christian Weiser
University of Freiburg

(on behalf of the ATLAS Collaboration)

14th Lomonosov Conference on Elementary Particle Physics
Moscow, August 2009

Content

- Introduction
- Higgs search channels in ATLAS
 - $H \rightarrow \gamma\gamma$
 - VBF $H \rightarrow \tau\tau$
 - $H \rightarrow b\bar{b}$
 - $H \rightarrow ZZ$
 - $H \rightarrow WW$
- Combined sensitivity
- Summary

What do we already know?

- Direct searches at LEP

$$M_H > 114.4 \text{ GeV}/c^2 \quad (95\% \text{ CL})$$

- Direct searches at the TEVATRON

$$\text{Exclude } 160 \text{ GeV}/c^2 < M_H < 170 \text{ GeV}/c^2 \quad (95\% \text{ CL})$$

- electroweak precision measurements

$$M_H < 163 \text{ GeV}/c^2 \quad (95\% \text{ CL})$$

$$(191 \text{ GeV}/c^2 \text{ incl. LEP Limit})$$

Higgs Boson Production

(i) Gluon-Fusion

(ii) Vector Boson Fusion (VBF)

(iii) Associated Production (W/Z, tt)

Higgs Boson Decays

Search Channels

Main channels studied:

Low mass:

- $H \rightarrow \gamma\gamma$
- $qqH, H \rightarrow \tau\tau$
- $ttH, H \rightarrow bb$
- $W/Z H, H \rightarrow bb$ **NEW!**

Also medium (& high) mass:

- $H \rightarrow WW^{(*)}$
- $H \rightarrow ZZ^{(*)}$

For low mass: Combination of large production cross section (gg-fusion) and large BR ($H \rightarrow bb$) does not work because of QCD background

Higgs Studies in ATLAS

Studies presented here:

- Realistic (at least we believe ...) GEANT4 based detector description simulation
- Real reconstruction algorithms
- Trigger simulation
- Effects of additional minimum bias events (*Pile-Up*), misalignment etc. studied
- Improved generators (NLO etc.)
- Background estimation from data in many cases

For all details see **CERN-OPEN-2008-020**

Focus in this talk:

Discovery potential in the low and medium mass range ($m_H \leq 200 \text{ GeV}$) @ $\sqrt{s} = 14 \text{ TeV}$

To be (re)studied:

Parameter determination (Mass, Spin, CP, Couplings,)

Heavy Higgs bosons

$$H \rightarrow \gamma\gamma$$

No direct coupling
tiny BR ($\approx \%$)
 $\sigma \times \text{BR} \approx 100 \text{ fb}$

Backgrounds:

- **Irreducible $\gamma\gamma(+X)$**
($\sigma \approx 30 \text{ pb}$ for $p_{T,\gamma} > 25 \text{ GeV}$, $|\eta| < 2.5$
 $80 \text{ GeV} < m_{\gamma\gamma} < 150 \text{ GeV}$)
- **Reducible $\gamma+\text{jet}$, $\text{jet}+\text{jet}$**
(γj : $\sigma \approx 180 \text{ nb}$ for $p_{T,\gamma} > 25 \text{ GeV}$
 jj : $\sigma \approx 480 \text{ } \mu\text{b}$ for $p_T > 25 \text{ GeV}$)

Need excellent

- Photon identification
- Photon-jet separation
- Mass resolution

$\approx 50\%$ of events:
at least 1 photon conversion
 $\rightarrow$ reconstruct from e^+ , e^- tracks
in Inner Detector

$$H \rightarrow \gamma\gamma$$

Inclusive (ask for photons only) analysis:

Discovery Potential (counting)

$\gamma\gamma + 1,2$ jet analyses (enhance VBF):

Vector Boson Fusion $qqH, H \rightarrow \tau\tau$

Signature of Vector Boson Fusion (VBF)

Production process:

- Two hard forward/backward "Tagging Jets" with large rapidity gap and high invariant mass
- Jet activity suppressed in central part of the detector $\rightarrow$ Central Jet Veto
- Large discrepancies between generators

Rapidity Gap

Central Jet Veto

Main Backgrounds:

- $Z (\rightarrow \tau\tau) + \text{jets}$
- $W (\rightarrow \tau\nu) + \text{jets}$
- $t\bar{t}$

Vector Boson Fusion $qqH, H \rightarrow \tau\tau$

τ decay channels:

- Lepton-lepton (ll)
- Lepton-hadron (lh)
- [Hadron-hadron (hh)]

Dominating
Background

$Z \rightarrow \tau\tau$:

Estimate on data
from $Z \rightarrow \mu\mu$
events with $\mu \rightarrow \tau$
transformation

Invariant mass computation:
(> 1 neutrino in final state)
"Collinear Approximation"
(Crucial: MET resolution)

hadron-hadron final state:
No reliable estimate of QCD
background without data
 $\rightarrow$ No expected significance quoted

$H \rightarrow b\bar{b}$

$t\bar{t}H$

a promising search channel some years ago:

Access to top-Higgs Yukawa coupling!

Complex final state:

Now:

Main Backgrounds:
 $ttbb$, $ttjj$

- Need precise background normalization!
- Has to come from data!
- Pile-Up: impact on selection efficiency and mass resolution

→ $t\bar{t}H$ has disappeared from latest sensitivity plots!

W/Z H

NEW!

The most important channels at the
TEVATRON at low mass!

But: Signal to background ratio less
favourable at the LHC

$S/\sqrt{B}$	2.1
S/B	1.3%

Follow idea of J.Butterworth et al.
[PRL 100:242001,2008]:

Select events ($\approx 5\%$ of cross section),
in which H und W bosons have large
transverse momenta: $p_T > 200 \text{ GeV}$

-> b-Quarks in one "fat" Jet

+ Acceptance (more central in detector)
+ Lepton-Identification, B-Tagging

W/ZH

ATL-PHYS-PUB-2009-088

Analyse jet structure:

$$L^{int.} = 30 \text{ fb}^{-1} : \frac{S}{\sqrt{B}} = 3.0$$

$$M_H = 120 \text{ GeV}$$

$$\frac{S}{\sqrt{B}} = 1.5$$

$$\frac{S}{\sqrt{B}} = 1.6$$

Combined: $\frac{S}{\sqrt{B}} = 3.7$

(Pile-Up not yet included)

- S/B much better than for ttH
- Different backgrounds for different channels
- Still good sensitivity including systematics
(e.g. $S/\sqrt{B} = 3.0$ for 15% uncertainty on all backgrounds)

$H \rightarrow ZZ$

"Golden Channel":

- Clear, narrow mass peak on top of background
- Covers wide mass range

Backgrounds:

- Irreducible: ZZ
- Reducible: Zbb , $t\bar{t}$

→ Lepton isolation and impact parameters

Effect of *Pile-Up* ($10^{33} \text{ cm}^{-2}\text{s}^{-1}$) and *Cavern Background*:

Loss in signal efficiency of $\approx 10\%$
(studied for $m_H = 130 \text{ GeV}$)

$H \rightarrow ZZ$

$H \rightarrow WW$

- Accessible for dominant production process (gg fusion) AND decay mode ($m_H > 135 \text{ GeV}$)
- Final state considered: $H \rightarrow WW \rightarrow e\nu\mu\nu$
- Two analyses: + 0 jet (gg fusion), +2 jets (vector boson fusion)
- 2 neutrinos $\rightarrow$ no mass peak reconstructable, only transverse mass
- Use angular correlation between leptons:

Fit transverse mass and transverse momentum of The WW system in $\Delta\Phi$ signal and control regions

Combined Discovery Potential

- 5 σ discovery seems possible with 10 fb⁻¹ for $m_H > 127$ GeV
- Adding the W/Z H, $H \rightarrow b\bar{b}$ channel will improve situation for low masses

Exclusion

2 fb^{-1} to exclude $m_H > 115 \text{ GeV}$ @ 95% CL

Conclusions

- Much more realistic detector simulations, Monte Carlo generators, reconstruction tools etc. have improved the quality of the predictions for ATLAS Higgs boson searches significantly
- Strong effort has been put in studying data driven methods to become less dependent on Monte Carlo generators
- A discovery (or exclusion) should be possible over the full mass range from 115 GeV - 1 TeV
- However, the low mass region (just above the LEP limit) is very challenging and several channels may contribute