П ОУ-12
9

ACADEMY OF SCIENCES OF THE USSR
P. N. LEBEDEV PHYSICAL INSTITUTE

I.E. Tamm Department of Theoretical Physics

High energy physics
and cosmic rays

Preprint No 92

STATE OF THE FALSE VACUUM AT FINITE
TEMPERATURE: THEORY AND APPLICATIONS

A.D. Linde

CERN LIBRARIES, GENEVA

CM-P00067554

Moscow - 1980

**STATE OF THE FALSE VACUUM AT FINITE
TEMPERATURE: THEORY AND APPLICATIONS**

A.D.Linde

Lebedev Physical Institute, Moscow, USSR

The theory of the first-order phase transitions in spontaneously broken gauge theories is investigated and some applications of this theory for cosmology and for the elementary particle physics are discussed.

It is known that in most of spontaneously broken gauge theories symmetry was restored in the early universe due to the high-temperature effects [1-3]. In the course of the universe expansion temperature decreased, and at several critical values of temperature T_{c_i} a series of phase transitions has taken place after which the initial symmetry of the theory was broken down to $SU(3)_c \times U(1)_{EM}$. If the critical temperature T_{c_i} is much greater than the values of the elementary particle masses, which appear after the phase transition, then the transition typically is a second-order one [1]. If, however, some particles after the phase transition at T_{c_i} acquire masses of the same order or greater than T_{c_i} , then the phase transition becomes a strongly first-order one [2,3].

Typically (for more or less natural relations between coupling constants) symmetry breaking in the Weinberg-Salam model is a second-order phase transition. However in the theories with a larger content of elementary particles the critical temperatures become smaller than the masses, which appear after the phase transitions, and the corresponding phase transitions are of the first order. In particular, contrary to the usual belief, the phase transitions, which take place in grand unified theories at $T_c \sim 10^{13} - 10^{15}$ GeV, always are strongly first-order ones [4].

A first-order phase transition in a spontaneously broken theory is a transition from a metastable vacuum state to a stable one, or, in a terminology of Coleman [5], from the false vacuum to the true vacuum. The lifetime of the metastable

vacuum states in gauge theories usually is extremely large [6,7], which may lead to many interesting consequences for cosmology [3,8,9] and for the elementary particle physics [7]. Recently the theory of the decay of the false vacua in gauge theories became of a particular interest, since, as it was suggested in [10], the large lifetime of the metastable vacuum states in gauge theories may help us to solve the urgent problem of primordial magnetic monopoles [11], which should be copiously produced during the phase transitions in grand unified theories [10].

A theory of the decay of the false vacuum in spontaneously broken theories at zero temperature has been first suggested by Kobzarev, Okun' and Voloshin [6], and then it has been considerably developed and improved by Coleman and Callan [5,12]. Then the results of the work of Coleman [5] have been extended to the case $T \neq 0$ in our paper [7]. Here we would like to give some further comments concerning the fate of the false vacua at finite temperatures and to discuss several implications of our results for cosmology and for the elementary particle physics.

To be concrete, we shall consider phase transitions in the Higgs model, though most of our results will remain correct for more complicated models.

Effective potential $V(\varphi)$ in the Higgs model at $T=0$ is given by [13]

$$V(\varphi) = -\frac{\mu^2}{2}\varphi^2\left(1 - \frac{3\alpha^2}{\lambda}\right) + \frac{\lambda}{4}\varphi^4\left(1 - \frac{9\alpha^2}{2\lambda}\right) + \frac{3\alpha^2}{4}\varphi^4 \ln \frac{\lambda\varphi^2}{\mu^2} \quad (1)$$

Here φ is the real classical scalar Higgs field, α is the fine structure constant. Renormalized parameters μ^2 , φ_0 and λ are fixed by the following relations: $\varphi_0 = \mu/\sqrt{\lambda}$;

$$V'(\varphi_0) \equiv \frac{dV}{d\varphi}(\varphi=\varphi_0) = 0 ; \quad V''(\varphi_0) \equiv \frac{d^2V}{d\varphi^2}(\varphi=\varphi_0) = 2\mu^2 \quad (2)$$

From eq.(1) it follows that at $\lambda = 3\alpha^2$ the curvature of $V(\varphi)$ at $\varphi=0$ vanishes: $V''(0)=0$. In this case the Higgs model is exactly equivalent [13] to the "massless" scalar electrodynamics of Coleman and Weinberg [14]. With a further decrease of λ effective potential acquires an additional minimum at $\varphi=0$. At $\frac{3}{2}\alpha^2 < \lambda < 3\alpha^2$ the old minimum at $\varphi=\varphi_0$ is more deep than the local minimum at $\varphi=0$. Therefore if the system initially was in the state $\varphi=0$ (and it was actually so due to the symmetry restoration in the early universe [1-3]), then the first-order phase transition may take place from the false vacuum $\varphi=0$ to the true vacuum $\varphi=\varphi_0$.

The decay of the false vacuum proceeds by spontaneous generation and subsequent expansion of bubbles of the new phase with $\varphi \neq 0$. The generation of bubbles is a tunneling process. Indeed, the volume energy of the bubble $\sim -\zeta^3$, where ζ is its size. However the surface energy of the bubble $\sim +\zeta^2$, and therefore only the bubbles above some critical size can spontaneously appear in the vacuum.

Following [5,12], we shall describe the barrier tunneling as a classical motion in the imaginary time direction, i.e. in

the Euclidean space. To calculate the probability of the bubble creation one should first of all find the trajectory of the tunneling, i.e. one should solve equation of motion

$$\square \varphi = \frac{d^2 \varphi}{dt^2} + \Delta \varphi = V'(\varphi) \quad (3)$$

with the boundary condition $\varphi=0$ for $\bar{x}^2 + t^2 \rightarrow \infty$. The tunneling probability per unit time per unit volume is given by

$$\Gamma/V = A \cdot e^{-S_4(\varphi)}, \quad (4)$$

where $S_4(\varphi)$ is the action, corresponding to the solution φ , and A is a subexponential factor [12]

$$A = \left(\frac{S_4}{2\pi} \right)^2 \cdot \left(\frac{\det' [-\square + V''(\varphi)]}{\det [-\square + V''(0)]} \right)^{-\frac{1}{2}} \quad (5)$$

Here $\det'$ indicates that the four zero eigenvalues of the operator $-\square + V''(\varphi)$ are to be omitted when computing the determinant; the contribution $\left(\frac{S_4}{2\pi} \right)^{1/2}$ of each zero mode gives the factor $\left(\frac{S_4}{2\pi} \right)^2$ in (5) [12].

To obtain a complete answer one should summarize the contributions to Γ/V from all possible trajectories, but in fact only the simplest $O(4)$ -symmetric solution $\varphi(\bar{x}^2 + t^2)$ is relevant, since it provides a minimal value of $S_4(\varphi)$.

One should note, however, that in our universe all phase transitions have taken place not in the vacuum at $T=0$, but at very high temperatures, which greatly increased the rate of tunneling. To describe the tunneling at $T \neq 0$ one should remind, that quantum statistics of bosons at $T \neq 0$ is formally equivalent to quantum field theory in the Euclidean space-time periodic in the "time" direction β with the

period $\frac{1}{T}$ [15].

Now it is clear how the previous considerations will be modified at $T \neq 0$ [7]. At $T=0$ the solution of eq. (3), corresponding to the minimal action $S_4(\varphi)$ (the simplest instanton in the model (I)), is the $O(4)$ -symmetric bubble of size $\sim \tau(0)$, fig. 1a. At $T \ll \tau(0)^{-1}$ the solution will become a series of such bubbles, placed at a distance $\frac{1}{T}$ from each other in the "time" direction, fig. 1b. At $T \sim \tau(0)^{-1}$ the bubbles become overlapping, fig. 1c. Finally, at $T \gg \tau(0)^{-1}$ (and this is the most interesting case for us) the solution becomes a cylinder with the "time" slice being the $O(3)$ -symmetric bubble of some new size $\tau(T)$. As a result, the action $S_4(\varphi)$ in (4), (5) at high temperatures is to be replaced by $S_3(\varphi)/T$, where $S_3(\varphi)$ is the action, corresponding to the $O(3)$ -symmetric bubble [7]. To compute $S_3(\varphi)$ one should find the $O(3)$ -symmetric solution of equation

$$\Delta \varphi = \frac{dV(\varphi, T)}{d\varphi} \equiv V'(\varphi, T) \quad (6)$$

with the boundary condition $\varphi \rightarrow 0$ at $\bar{x}^2 \rightarrow \infty$. Here $V(\varphi, T)$ is the temperature-dependent effective potential [1-3]. A complete expression for the tunneling probability per unit time per unit volume at high temperatures is given by

$$\Gamma_V = T \left(\frac{S_3(\varphi, T)}{2\pi T} \right)^{\frac{3}{2}} \left(\frac{\det' [-\Delta + V''(\varphi, T)]}{\det [-\Delta + V''(0, T)]} \right)^{-\frac{1}{2}} e^{-\frac{S_3(\varphi, T)}{T}}. \quad (7)$$

Here, as before, $\det'$ indicates that the three zero eigenvalues

of the operator $-\Delta + V''(\varphi, T)$ are to be omitted when computing the determinant. The contribution of zero modes gives the factor $\left(\frac{S_3(\varphi, T)}{2\pi T}\right)^{\frac{3}{2}}$ in (9), whereas the factor T appears due to the periodicity $\frac{1}{T}$ of the Euclidean space-time at $T \neq 0$.

Usually it is impossible to find a solution of eq. (6) analytically, and one can obtain the solution and the corresponding action S_3 only by means of computer calculations [7]. ~~determinants is even more difficult,~~ Computation of the V but in most cases it is sufficient to have a rough estimate of the order of magnitude of the subexponential factor in (7). Numerical investigation of the first-order phase transitions in a number of models shows, that the tunneling probability (7) is extremely small if the curvature of the effective potential $V(\varphi)$ at the point, corresponding to the false vacuum, is not negative ($V''(\varphi) \geq 0$) [7, 16]. On the other hand, for the phase transition to the true vacuum to occur in most part of the expanding universe, the rate of the bubble generation and expansion should be large as compared with the universe expansion rate [9, 10]. Therefore if the tunneling rate is sufficiently small, then the phase transition could never take place in most part of the universe. An account taken of

the universe expansion rate leads also to some modification of the usual stationary-space Lagrange equations for scalar fields. This modification is small but nevertheless important since it leads to a slight stabilization of the false vacua. A complete investigation of the

first-order phase transitions in the expanding universe ^{contains many} technical points, and therefore here we shall merely present our main results without proofs. ^{*)} The details of the investigation together with a more precise and complete formulation and discussion of our statements will be contained in a separate publication.

1. A phase transition from the false ^{vacuum} to the true vacuum takes place in the expanding universe only when the curvature of the effective potential $V(\varphi, T)$ becomes negative at the point φ , corresponding to the false vacuum. This means, that the first-order phase transitions proceed only when the supercooled false vacuum becomes unstable rather than metastable. Consequently, the strongly first-order phase transitions proceed with a great release of energy, which may considerably heat up the universe [3] (see also [8-10]). In particular, a typical phase transition in grand unified theories, which occurs at $T_c \sim 10^{13}-10^{15}$ GeV, is a strongly first-order one [4], and therefore it looks as a great explosion in all the universe. Such a phase transition takes place at the time $t \sim 10^{-35}$ sec after the Big Bang, and for the obvious reasons we shall call it the Grand Bang [4]. Note, that the cosmological baryon production [17] starts just after the Grand Bang [4].

2. If the curvature of the effective potential at $T=0$ is not negative ($V''(\varphi) \geq 0$) in the false vacuum, then the phase transition to the true vacuum could never take place in the course of the universe expansion. In particular,

*) These results are related to symmetry breaking at the grand unification mass scale. For the Weinberg-Salam model one should write every

in the theories with the Coleman-Weinberg dynamical symmetry breaking [14] the symmetry initially was restored ($\varphi=0$), and the dynamical symmetry breaking actually could never take place since in such theories $\frac{d^2 V}{d\varphi^2} = 0$ at $\varphi=0, T=0$.

3. If the curvature of the effective potential in the false vacuum at $T=0$ is negative but very small, then with a decrease of temperature from $T=\infty$ to $T=0$ it becomes negative only at a very small temperature. In this case, the supercooling of the false vacuum becomes extremely large and the phase transition occurs only at a very small temperature (by a proper choice of parameters one can make this temperature as small as one wishes). In particular, the critical temperature of the Grand Bang may be much lower than the typical value $T_c \sim 10^{14}$ GeV, and in this case the density of monopoles produced during the Grand Bang would be extremely small, which would solve the primordial monopole problem [11, 10]. Note, that to decrease T_c there is no need in any particular hierarchy of symmetry breakings in grand unified theories, suggested in [10]. Indeed, even in the simplest Higgs model (1) T_c becomes extremely small for $0 < \lambda - 3\alpha^2 \ll \alpha^2$. This example, however, is a manifestation of a general rule: Critical temperature could be made much less than its typical value in a given theory only if some peculiar and very unnatural relations between coupling constants hold. Therefore the above mentioned "solution" of the primordial monopole problem seems unnatural and undesirable^{x)}.

^{x)} Professors Einhorn and Sato have informed me, that they

Fortunately, however, there exists a much simpler solution of the primordial monopole problem. As it is argued in [18], at the temperatures greater than the temperature of the phase transition in the Weinberg-Salam model ($T \gtrsim 10^2 \text{GeV}$) magnetic monopoles have been confined. The confined monopoles have been rapidly annihilating, and practically none of them could survive [18]. This question and some other problems touched upon in the present paper will be discussed in a series of subsequent publications on phase transitions in gauge theories and cosmology.

also have found, that the investigation of the first-order phase transitions in gauge theories does not lead to a natural solution of the primordial monopole problem.

References

1. D.A.Kirshnits, JETP Lett. 15(1972) 529;
D.A.Kirshnits and A.D.Linde, Phys. Lett. 42B (1972) 471;
S.Weinberg, Phys. Rev. D9 (1974) 3357;
L.Dolan and R.Jackiw, Phys. Rev. D9 (1974) 3320;
D.A.Kirshnits and A.D.Linde, Zh. Eksp. Teor. Fiz.
67 (1974) 1263, (JETP 40 (1975) 629).
2. D.A.Kirshnits and A.D.Linde, Ann. Phys. (N.Y.) 101
(1976) 195.
3. A.D.Linde, Rep. Progr. Phys. 42 (1979) 389.
4. A.D.Linde, Grand Bang, Lebedev Phys. Inst. preprint (1980).
5. S.Coleman, Phys. Rev. D15 (1977) 2929.
6. M.B.Voloshin, I.Yu.Kobzarev and L.B.Okun, Yad. Fiz.
20 (1974) 1229.
7. A.D.Linde, Phys. Lett. 70B (1977) 306.
8. G.Lasher, Phys. Rev. Lett. 42 (1979) 1646.
9. K.Sato, Nordita preprint (1980).
10. J.Preskill, Phys. Rev. Lett. 41 (1979) 1365;
A.H.Guth and S.-H.Tye, Phys. Rev. Lett. 44 (1980) 631;
M.B.Einhorn, D.Stein and D.Toussaint, Univ. of Michigan
preprint UM HE 80-1 (1980).
G.Lazarides, M.Magg and Q.Shafi, CERN preprint
Ref. TH.2856 - CERN (1980).
11. Ya.B.Zeldovich and M.Yu.Khlopov, Phys. Lett. 72B (1978)
239.
12. C.G.Callan and S.Coleman, Phys. Rev. D16 (1977) 1762.

13. A.D.Linde, JETP Lett. 23 (1976) 64.
14. S.Coleman and E.Weinberg, Phys. Rev. D7 (1973) 1888.
15. E.S.Fradkin, Proc. Lebedev Phys. Inst. 29 (1965) 7.
16. A.D.Linde, Lebedev Phys. Inst. preprint No 132 (1979);
Phys.Lett. 92B (1980) 119.
17. A.D.Sakharov, Pisma ZhETF 5 (1967) 32;
V.A.Kuzmin, Pisma ZhETF 13 (1970) 335;
M.Yoshimura, Phys. Rev. Lett. 41 (1978) 281;
S.Dimopoulos and L.Susskind, Phys. Rev. D18 (1978) 4500;
S.Weinberg, Phys. Rev. Lett. 42 (1979) 859.
18. A.D.Linde, Confinement of monopoles at high temperatures,
Lebedev Phys. Inst. preprint (1980).

Fig 1.

Solutions of eq. (3) at finite temperature. a) $T = 0$; b) $T \ll z_0^{-1}$; c) $T \sim z_0^{-1}$; d) $T \gg z_0^{-1}$. Dashed regions contain classical field φ .

To make figures more clear we have drawn the bubbles as if the thickness of their walls were much less than the bubble size. Actually, however, the wall thickness is very large in all cases, in which the tunneling rate is not negligibly small. This means, in particular, that the thin-wall approximation [5,6] is absolutely inapplicable to the investigation of the first-order phase transitions in realistic gauge theories.

Т - 09088

Подписано в печать 6 июня 1980 года

Заказ № 320. Тираж 100 экз. п/л 0.7

Отпечатано в Редакционно-издательском
отделе ФИАН СССР

Москва, В-312, Ленинский проспект, 53