Ref.TH.3055-CERN

CALCULATING $K \rightarrow 2\pi$

G. Nardulli

Istituto di Fisica, Università di Bari
and
Istituto Nazionale di Fisica Nucleare
Sezione di Bari

and

G. Preparata

CERN - Geneva
and
Istituto di Fisica, Università di Bari

A B S T R A C T

We calculate the long-distance contributions to $K \rightarrow 2\pi$ decays. We show that : (i) the $\Delta I = \frac{1}{2}$ rule emerges from well-known properties of hadronic interactions, and (ii) short-distance physics plays an insignificant rôle.

Ref.TH.3055-CERN

6 April 1981

Since the discovery of strange particles, the vast difference (typically a factor of 20) between the $\Delta I = \frac{1}{2}$ and the $\Delta I = \frac{3}{2}$ non-leptonic decay amplitudes has been one of the major puzzles of the theory of weak interactions. After more than a decade of unsuccessful attempts, it was realized ^{1),2)} that the origin of the so-called $\Delta I = \frac{1}{2}$ rule might lie in the peculiar property of the hadronic interaction of strongly suppressing "exotic" amplitudes (such as $\Delta I = \frac{3}{2}$). This observation implied that, even though the non-leptonic Hamiltonian explores space-time regions down to dimensions of the order of the W-boson Compton wave length $1/m_W$ ³⁾, non-leptonic physics appeared essentially determined by long-distance (~ 1 Fermi) dynamics. And, as a matter of fact, in a quark theory with free-field short-distance singularities it was shown that ²⁾ the product of the two weak currents appearing in the non-leptonic Hamiltonian is perfectly finite at short distances, due to the vanishing of the leading $(1/x^2)$ singularity.

However, it was K. Wilson ³⁾ who first pointed out that, due to possible anomalous dimensions, one might expect some important contributions at short-distances, and in particular the $\Delta I = \frac{1}{2}$ enhancement might have partly such an origin.

With the discovery of asymptotic freedom in QCD, Wilson's conjecture was remarkably confirmed by the calculations carried out by two independent groups ⁴⁾. No reliable quantitative estimate could, however, be made of the short-distance effects, due to the lack of adequate computational tools for the long-distance effects buried in the local operators appearing in the short-distance expansion.

In this letter we wish to develop such techniques and bring further insight into this problem, by performing a dynamical calculation of the long-distance contributions to the non-leptonic Hamiltonian of the $K \rightarrow 2\pi$ decays ^{*}).

Let us consider the amplitude :

$$\begin{aligned} \mathcal{A}(K^0 \rightarrow \pi^+ \pi^-) &= \langle \pi^+ \pi^- | H_{NL} | K^0 \rangle = \\ &= g_W^2 \sin \theta \cos \theta \int d^4x \Delta^{\mu\nu}(x, m_W) \langle \pi^+ \pi^- | T(J_\mu^{4i2}(x) J_\nu^{4i5}(0)) | K^0 \rangle = \\ &= g_W^2 \sin \theta \cos \theta \int \frac{d^4q}{(2\pi)^4} \frac{1}{m_W^2 - q^2} T(q). \end{aligned} \quad (1)$$

^{*}) An early attempt along these lines is in Ref. 5).

Before discussing its short-distance behaviour, we should separate the amplitude $T(q)$ into its disconnected and connected parts respectively, i.e., (see Fig. 1) :

$$T(q) = T_{\text{dis}}(q) + T_{\text{conn}}(q). \quad (2)$$

It is obvious that the Fourier transform of the disconnected part $T_{\text{dis}}(q)$ is regular when $x \rightarrow 0$, and furthermore can be calculated precisely from our knowledge of semi-leptonic physics. Such a calculation was performed a long time ago by Feynman ⁶⁾. Separating $T_{\text{dis}}(q)$ into its $\Delta I = \frac{1}{2}$ and $\Delta I = \frac{3}{2}$ parts, we find [we give in parentheses the experimental values of the amplitudes $a^{\frac{1}{2}}(K^0 \rightarrow \pi^+\pi^-)$ and $a^{\frac{3}{2}}(K^0 \rightarrow \pi^+\pi^-)$ ⁷⁾]

$$a_{\text{dis}}^{\frac{1}{2}}(K^0 \rightarrow \pi^+\pi^-) = 3.70 \cdot 10^{-8} \text{ GeV} \quad (2.71 \cdot 10^{-7} \text{ GeV}) \quad (3)$$

$$a_{\text{dis}}^{\frac{3}{2}}(K^0 \rightarrow \pi^+\pi^-) = 1.85 \cdot 10^{-8} \text{ GeV} \quad (.86 \cdot 10^{-8} \text{ GeV}). \quad (4)$$

Thus we see that : (i) the disconnected contributions are comparable (no $\Delta I = \frac{1}{2}$ enhancement), and (ii) they are a factor ~ 7 too small and a factor ~ 2 too large as compared with the experimental $\Delta I = \frac{1}{2}$ and $\Delta I = \frac{3}{2}$ amplitudes, respectively.

Turning to the Fourier transform of the connected amplitude $T_{\text{conn}}(q)$, its short-distance behaviour is governed by ³⁾ [we use the familiar SU(3) notation] :

$$\begin{aligned} & \langle \pi^+\pi^- | T(J_\mu^{1+2}(x) J_\mu^{4-5}(0)) | K^0 \rangle \xrightarrow{x \rightarrow 0} \\ & \rightarrow \langle \pi^+\pi^- | \{ C_-(x) O_-(0) + C_+(x) O_+(0) \} | K^0 \rangle, \end{aligned} \quad (5)$$

where the "normal-ordered" operators $O_\pm(0)$ are given by

$$O_\mp(0) = \frac{1}{2} : J_\mu^{1+2}(0) J_\mu^{4-5}(0) \mp (J_\mu^3(0) + \frac{1}{\sqrt{3}} J_\mu^8(0) + \sqrt{\frac{2}{3}} J_\mu^0(0)) J_\mu^{6-7}(0) :, \quad (6)$$

and the coefficient functions in QCD are (in the four-flavour approximation) ⁴⁾

$$\begin{aligned} C_-(x) &= \left[1 - \frac{25g^2(\mu)}{24\pi^2} \log \mu \sqrt{-x^2} \right]^{.48} \\ C_+(x) &= \left[1 - \frac{25g^2(\mu)}{24\pi^2} \log \mu \sqrt{-x^2} \right]^{.24}. \end{aligned} \quad (7)$$

Note that in a world where the short-distance behaviour is "canonical" (i.e., like in free-field theory)²⁾ $C_+(x) = C_-(x) = 1$. As argued in Ref. 3), Eqs. (1) and (5) imply

$$a_{\text{conn}}(K^0 \rightarrow \pi^+\pi^-) \simeq \frac{G}{\sqrt{2}} \omega_0 \theta \sin \theta \langle \pi^+\pi^- | [C_-(\frac{1}{m_W}) Q_-(0) + C_+(\frac{1}{m_W}) Q_+(0)] | K^0 \rangle \quad (8)$$

PCAC, through reduction of one π , allows us to write (8) as

$$a_{\text{conn}}(K^0 \rightarrow \pi^+\pi^-) \simeq -\frac{i}{2f_\pi} \frac{G}{\sqrt{2}} \omega_0 \theta \sin \theta \left\{ \langle \pi^+ | [Q_5^+, C_-(\frac{1}{m_W}) Q_-(0) + C_+(\frac{1}{m_W}) Q_+(0)] | K^0 \rangle + \langle \pi^- | [Q_5^-, C_-(\frac{1}{m_W}) Q_-(0) + C_+(\frac{1}{m_W}) Q_+(0)] | K^0 \rangle \right\} \quad (9)$$

Due to the well-known remarkable precocity of short-distance behaviour, Eq. (5) is expected to already represent fairly accurately the current-product for $|x| \lesssim 1/\mu$ ($\mu \simeq 1$ GeV). Thus, setting $x = x_0$ with $1/\mu = \sqrt{-x_0^2}$, we may write

$$\begin{aligned} \langle \pi | : J_\mu^A(0) J_\mu^B(0) : | K \rangle &\simeq \langle \pi | T(J_\mu^A(x_0) J_\mu^B(0)) | K \rangle \\ &= \int \frac{d^4 q}{(2\pi)^4} e^{iqx_0} T_{K\pi}^{AB}(q) \Big|_{\text{conn}} \simeq \int \frac{d^4 q}{(2\pi)^4} f_{Q_0}(q) T_{K\pi}^{AB}(q) \Big|_{\text{conn}}; \quad (10) \end{aligned}$$

where $f_{Q_0}(q)$ is unity for small q and cuts off the integral for $|q^2| > Q_0^2 = O(\mu^2)$. Equation (10) is the starting point of our calculation. Note that due to the suppression of the high q^2 region, the normal product is totally dominated (as it must be) by the long-distance, small q^2 contributions^{*)}. Following Cottingham⁸⁾ we write the integral in Eq. (10) as :

$$\begin{aligned} &\int \frac{d^4 q}{(2\pi)^4} f_{Q_0}(q) T_{K\pi}^{AB}(q) \Big|_{\text{conn}} = \\ &= \frac{1}{4(2\pi)^4} \int_0^{Q_0^2} dQ^2 \int_C dq_0 \sqrt{Q^2 + q_0^2} \int_{-1}^{+1} dz \int_0^{2\pi} d\varphi T_{K\pi}^{AB}(Q^2, q_0, z, \varphi) \Big|_{\text{conn}} \quad (11) \end{aligned}$$

*) Our experience with precocious short-distance dominance leads us to expect that the theoretical error made by going through the steps of Eq. (10) should not exceed 10%.

where we have chosen to cut off the integral over space-like $q^2 = -Q^2$, at the value $Q_0^2 \sim O(\mu^2)$, which should lie in the range $1 \div 2 \text{ GeV}^2$. The contribution of the pole terms (see Fig. 2) to our integral can be easily evaluated by making use of available information on pseudoscalar meson form factors. As for the "continuum" contribution we calculate it by shifting the contour C to C_M (see Fig. 3) ($M \gtrsim 3 \text{ GeV}$) and taking advantage of the fact that for such values of q_0 the "Compton-like" amplitude should be Regge-behaved, i.e.,

$$T_{K\pi}^{AB}(q_0, \dots) \xrightarrow{q_0 \gtrsim M} \beta^{AB}(\dots) (q_0)^{\alpha_{K^{**}}(0)}, \quad (12)$$

where $\alpha_{K^{**}}(0) \simeq 0.25$ is the intercept of the K^{**} trajectory, and the residues β^{AB} can be obtained from factorization and $SU(3)$ symmetry⁹⁾. Finally, finite energy sum rules¹⁰⁾ enable us to calculate reliably the integral over the cuts of the amplitude $T_{K\pi}^{AB}(q)$. Note that the Regge contribution is absent for the $\Delta I = \frac{3}{2}$ amplitude, due to the very low intercept of "exotic" Regge trajectories: this constitutes the most striking difference that the dynamics of hadronic interactions induces between the $\Delta I = \frac{1}{2}$ and the $\Delta I = \frac{3}{2}$ amplitudes.

Our results for the matrix elements of $O_{\pm}(0)$ are reported in Table 1, where the pole and the continuum contributions have been given separately and for different values of Q_0^2 . From this Table we can immediately write down the results for the connected amplitudes as

$$a_{\text{conn}}^{1/2}(K^0 \rightarrow \pi^+\pi^-) = c_+ \left(\frac{1}{m_w}\right) \begin{Bmatrix} 4.47 \\ 7.16 \\ 9.92 \end{Bmatrix} + c_- \left(\frac{1}{m_w}\right) \begin{Bmatrix} 5.81 \\ 8.93 \\ 12.03 \end{Bmatrix} \cdot 10^{-8} \text{ GeV} \quad (13)$$

and

$$a_{\text{conn}}^{3/2}(K^0 \rightarrow \pi^+\pi^-) = c_+ \left(\frac{1}{m_w}\right) \begin{Bmatrix} -0.67 \\ -0.88 \\ -1.06 \end{Bmatrix} \cdot 10^{-8} \text{ GeV}, \quad (14)$$

where the three different sets of values are for $Q_0^2 = 1, 1.5, 2 \text{ GeV}^2$, respectively.

We should now like to make a few comments. As the entries of Table 1 show, only the continuum contribution exhibits definite sensitivity to the cut-off parameter Q_0^2 . However, a value of $Q_0^2 \simeq 2 \text{ GeV}^2$ appears totally reasonable, while a value $Q_0^2 \simeq 1 \text{ GeV}^2$ would definitely be too small, due to the particular way in

which we have introduced the cut-off. On the other hand, the pole terms, the only significant contribution to the $\Delta I = \frac{3}{2}$ amplitude, are fairly insensitive to Q_0^2 and their uncertainty cannot exceed 15%. It should also be stressed that the large entries corresponding to the Regge dominated continuum carry some uncertainty (determination of residues, K^{**} intercept, assumed equality of axial and vector contributions at $Q^2 = 0$) which we cannot pretend to be much less than 30%. With this in mind, the main conclusions of our work can be read off from Table 2 ^{*)}.

Whereas small values (≤ 0.5 GeV) of Λ_{QCD} give a perfectly adequate description of a_1^2, a_2^2 seems to require Λ_{QCD} values less than 0.1 GeV. In particular with "canonical" short-distance singularities ($\Lambda = 0$) we obtain an extremely good description of $K \rightarrow 2\pi$ decay.

Even though it is not our intention to claim that asymptotic freedom, with the presently accepted values of Λ , is ruled out by our analysis, we would like to draw attention to the following interesting fact which emerges from reading Table 2. Our numbers show no evidence that the time-honoured $\Delta I = \frac{1}{2}$ rule has anything to do with short-distance physics ; quite the contrary. They indicate that the bizarre difference in behaviour between the K^0 and the K^+ has its origin solely in well-known long-distance properties of the hadronic interactions. And this, we believe, is the most significant result of our work.

^{*)} The details of our calculation shall be reported in a future publication.

I \ I	I = $\frac{1}{2}$			I = $\frac{3}{2}$		
	$Q_0^2 = 1 \text{ GeV}^2$	$Q_0^2 = 1.5 \text{ GeV}^2$	$Q_0^2 = 2 \text{ GeV}^2$	$Q_0^2 = 1 \text{ GeV}^2$	$Q_0^2 = 1.5 \text{ GeV}^2$	$Q_0^2 = 2 \text{ GeV}^2$
0						
O_+^{pole}	$2.1 \cdot 10^{-9}$	$2.52 \cdot 10^{-9}$	$2.90 \cdot 10^{-9}$	$-0.67 \cdot 10^{-8}$	$-0.88 \cdot 10^{-8}$	$-1.06 \cdot 10^{-8}$
O_-^{pole}	$1.55 \cdot 10^{-8}$	$2.02 \cdot 10^{-8}$	$2.40 \cdot 10^{-8}$	0	0	0
$O_+^{\text{cont}} = O_-^{\text{cont}}$	$4.26 \cdot 10^{-8}$	$6.91 \cdot 10^{-8}$	$9.63 \cdot 10^{-8}$	~ 0	~ 0	~ 0

Table 1 - Contributions to the matrix elements of the "normal ordered" operators $O_{\pm}(0)$ [see Eq. (5)] from poles and continuum for different isospin and different cut-off's Q_0^2 . Note that our results have been multiplied by $G/\sqrt{2} \cos\theta \sin\theta$. Units are GeV.

$\Lambda_{\text{QCD}} \text{ GeV}$	0.4	0.1	0	Exp.
$a \cdot 10^8 \text{ GeV}^{-1}$				
$a^{\frac{1}{2}}(K^0 \rightarrow \pi^+ \pi^-)$	34.2	29.4	25.7	27.1
$a^{\frac{3}{2}}(K^0 \rightarrow \pi^+ \pi^-)$	1.09	0.97	0.79	0.86

Table 2 - The results of our calculation for $a^{\frac{1}{2}}(K^0 \rightarrow \pi^+ \pi^-)$ and $a^{\frac{3}{2}}(K^0 \rightarrow \pi^+ \pi^-)$ for $Q_0^2 = 2 \text{ GeV}^2$, and different values of Λ_{QCD} . $\Lambda = 0$ corresponds to "canonical" short-distance behaviour.

REFERENCES

- 1) Y. Hara - Progr.Theor.Phys. (Kyoto) 37 (1967) 710.
- 2) S. Nussinov and G. Preparata - Phys.Rev. 175 (1968) 2180.
- 3) K. Wilson - Phys.Rev. 179 (1969) 1499.
- 4) G. Altarelli and L. Maiani - Phys.Letters 52B (1974) 351 ;
M.K. Gaillard and B.W. Lee - Phys.Rev.Letters 33 (1974) 108.
- 5) G. Preparata - Phys.Letters 34B (1971) 412.
- 6) R.P. Feynman in "Symmetries in Elementary Particle Physics", Academic Press (N.Y.) 1965, A. Zichichi Ed., p. 111.
- 7) T.J. Devlin and J.O. Dickey - Revs.Modern Phys. 51 (1979) 237.
- 8) W.N. Cottingham - Ann.Phys. (N.Y.) 25 (1963) 424.
- 9) W.P. Hesse et al. - Phys.Rev.Letters 25 (1970) 613 ;
J.G. Rushbrooke and B.R. Webber - Physics Reports 44 (1978) 1 ;
A.C. Irving and R.P. Worden - Physics Reports 34 (1978) 117.
- 10) For a review, see : V. De Alfaro, S. Fubini, G. Furlan and G. Rossetti -
"Currents in Hadron Physics", North Holland (Amsterdam-London) 1973.

FIGURE CAPTIONS

- Figure 1 Equation (2) in diagrammatic form.
- Figure 2 The pole diagrams of $T_{k\pi}^{AB}(q)$.
- Figure 3 The contours of integration C and C_M and the q_0 singularity structure of the "continuum" amplitude $T_{k\pi}^{AB}(q)$.

fig.1

fig.2

fig.3