Commissioning of the ATLAS Muon Trigger with Beam Collisions at the LHC

Alexander Oh

University of Manchester
on behalf of the ATLAS Collaboration

2010 Nuclear Science Symposium,
Medical Imaging Conference and
17th Room Temperature Semiconductor Detector Workshop

October 30, 2010 – November 6, 2010 – Knoxville, Tennessee

The ATLAS detector

- Multi purpose detector
 - Calorimetry
 - Tracking
 - Muon Identification

Detector: Muon Spectrometer

- Fast trigger chambers:
 - Resistive Plate Chambers (RPC) (barrel, $|\eta| < 1.05$)
 - Thin Gap Chambers (TGC) (end-caps, $1.05 < |\eta| < 2.4$)
- Precision chambers:
 - Monitored Drift Tubes (barrel and end-cap, $|\eta| < 2.4$)
 - Cathode Strip Chambers (forward region $2.0 < |\eta| < 2.7$)
- Toroidal magnetic field of 0.5 T
 - barrel and two end-cap toroids
 - 8 fold geometry
 - complex field

Detector: Inner Detector

- Silicon Tracker
 - Pixels (3 layers)
 - Strips (4 barrel layers, 9 end-cap layers)
 - Coverage $|\eta| < 2.5$
- Transition Radiation Tracker
 - Coverage $|\eta| < 2.0$
- Solenoidal magnetic field of 2.0 T

Muon Reconstruction

Objective:

Muon identification in a wide momentum range from GeV (Onia) to TeV (BSM).

Make use of all detector components and use different strategies:

- Muon Spectrometer Standalone (MS) reconstruction
- MS and Inner Detector (ID) combined fit
 - Outside-in
 - Inside-out
- Taggers
 - ID track is matched to MS segments
 - ID track is matched to a MIP in calorimeter

Trigger

- **Objective:**
Reduce from 40 MHz bunch crossing rate to ~200 Hz event rate to tape.

Three stage trigger system:

- L1 custom hardware
 - Require coincidence in eta phi view.
 - p_T estimation based on geometrical roads.
- HLT software
 - L2: Pattern recognition, LUT based p_T estimate
 - EF: Full muon reconstruction available
 - Reconstruction based on muon detectors only or in combination with inner detector tracks for both L2 and EF available.

Trigger

- Objective:
Reduce from 40 MHz bunch crossing rate to ~ 200 Hz event rate to tape.

Three stage trigger system:

- L1 custom hardware**
 - Require coincidence in eta phi view.
 - p_T estimation based on geometrical roads.
- HLT software**
 - L2: Pattern recognition, LUT based p_T estimate
 - EF: Full muon reconstruction available
 - Reconstruction based on muon detectors only or in combination with inner detector tracks for both L2 and EF available.

Performance measurement

- Performance of trigger algorithms is determined relative to offline reconstructed muons.
 - The offline and online muons are geometrically matched.
 - Only matched muons are considered.
- Efficiencies are determined relative to the previous trigger level.
 - L1 rel to offline
 - L2 rel to L1
 - EF rel to L2

L1 Trigger

- Efficiency determined by looking at
 - Inclusive muons relative to offline.
 - Tag & Probe method on J/Psi and Z events.

$$\text{efficiency} = \frac{\text{\#offline muons triggered by L1}}{\text{\#offline reconstructed muons}}$$

- Trigger Efficiencies at L1
 - barrel 75%
(limited by geometrical acceptance)
 - end-cap 95%

L2 Trigger: Muon Spectrometer only

- Resolution measured relative to off-line combined muons.
- Efficiency determined by looking at
 - Inclusive muons relative to offline and L1.

$$\text{efficiency} = \frac{\text{\#offline muons triggered by L2}}{\text{L1 \& \#offline reconstructed muons}}$$

- Trigger Efficiencies at L2 >98%

L2 Trigger: combined

- Resolution measured relative to off-line combined muons.
- Efficiency determined by looking at
 - Inclusive muons relative to offline and L2 MS.

$$\text{efficiency} = \frac{\text{\#offline muons triggered by L2}}{\text{L2MS} \&\& \text{\#offline reconstructed muons}}$$

- Trigger Efficiencies at L2 >98%

EF Trigger: Muon Spectrometer only

- Trigger Efficiencies at EF > 98%

$$\text{efficiency} = \frac{\text{\#offline muons triggered by EF}}{\text{L2 \& \#offline reconstructed muons}}$$

EF Trigger: combined

- Resolution measured relative to off-line combined muons.
- Trigger Efficiencies at EF >98%

$$\text{efficiency} = \frac{\text{\#offline muons triggered by EF}}{\text{L2 \& \#offline reconstructed muons}}$$

Trigger Rates and Evolution

- Muon rates measured in data.
 - Fit function accounts for cosmic radiation and beam induced events.
 - “Steps” due to change in filling scheme, more acceptance for cosmic radiation events.
- Extrapolation to higher luminosity.
- Determines evolution of trigger strategy.

Summary

- Performance of muon trigger was studied with first collision data.
- Good agreement between data and MC.

- Many more results to come soon with the full data set!