Signatures of collective flow in high-multiplicity pp collisions

Adam Kisiel*

Physics Department, CERN, CH-1211, Genève 23, Switzerland

(Received 8 December 2010; published 31 October 2011)

A blast-wave parametrization, including a full set of hadronic resonances, is used to model a small system, with total particle multiplicity comparable to the one measured in the high-multiplicity pp collisions at the Large Hadron Collider (LHC). Calculations are performed for three cases: with negligible, moderate, and strong radial flow on the blast-wave hypersurface. We investigate the effects of flow on inclusive p_T spectra as well as on 1D and 3D femtoscopic radii for pions. Special emphasis is put on the role of pions from resonance decays. In particular we show that they magnify the flow effects present in the blast-wave stage and significantly influence the shape of the correlation functions. A specific observable, the $R_{\text{out}}^E/R_{\text{side}}^G$ ratio is proposed as a sensitive probe of the collective effects. Model results for the high multiplicity pp collisions, for scenarios with small and large radial flow are compared. Several features of soft particles production in the medium multiplicity pp events at the LHC are identified as sensitive probes of the collective behavior of the medium.

DOI: [10.1103/PhysRevC.84.044913](https://doi.org/10.1103/PhysRevC.84.044913)

PACS number(s): 25.75.Dw, 25.75.Ld, 13.85.Ni, 25.40.Ve

I. INTRODUCTION

The pp program at the Large Hadron Collider (LHC) has successfully started, and has already produced a large set of minimum-bias measurements from the experiments [1–8]. Of particular interest is the fact that the multiplicity range measured at $\sqrt{s} = 7$ TeV overlaps with the one measured in peripheral CuCu collisions at ultrarelativistic energies of $\sqrt{s_{NN}} = 62$ and 200 GeV at relativistic heavy-ion collider (RHIC). This allows for comparison of observables between $p + p$ and heavy-ion collisions at the same measured final state multiplicity $dN_{ch}/d\eta$. In heavy-ion collisions, $dN_{ch}/d\eta$ is the scaling variable for many soft physics observables, including the inclusive particle p_T spectra, pion femtoscopic radii [9], strange particle yields, etc. It is therefore interesting whether a high-multiplicity pp collision resembles a heavy-ion one. Ideally we would like to identify features of soft particles production in the medium multiplicity pp events, which would be sensitive probes of the collective behavior of the medium.

This question has become even more relevant with the recent observation of the long-range η ridge in high multiplicity pp collisions by the CMS Collaboration experiment [10]. There are attempts to explain the phenomenon in the color glass condensate framework [11] or as an elliptic flow [12]. Both of these critically depend on the observation of radial flow (or space-momentum correlation) in such collisions. A recent study with EPOS model [13] suggests that such flow has actually been seen in the ALICE pion femtoscopy data [4]. Similar topics have been investigated in the framework of the rescattering model [14]. A comparison of pion femtoscopic radii in pp and heavy-ion collisions at lower energy $\sqrt{s} = 200$ GeV have been carried out by the STAR experiment [15]. Scaling between radii in pp and heavy-ion collisions has been observed vs. the pair momentum, suggesting that there might be a common physics mechanism, i.e., radial flow, driving them in both cases [16].

The hydrodynamic scenario is well applicable to heavy-ion collisions at RHIC. It produces specific space-momentum correlation patterns, which are commonly referred to as flow. The system created in the heavy-ion collision expands rapidly outwards, showing a strong radial flow, which is observed in the modification of the single particle inclusive p_T spectra shape. The spectrum is an observable depending only on the momenta of the particles, so its connection to space-time can only be indirectly inferred. To directly access space-time information we employ femtoscopic techniques. The fall of the “femtoscopic radii” (sometimes called “HBT radii”) with particle’s m_T can be interpreted as the decrease of “lengths of homogeneity,” a direct consequence of radial and longitudinal flow [17]. The simultaneous description of the p_T spectra and the femtoscopic radii dependence on m_T in heavy-ion collisions can be achieved with the “blast-wave” parametrization [18–20]. In this work we test what results such an approach brings when applied to small systems. We characterize the size of the system by the average charged particle multiplicity per unit of pseudorapidity that it produces. This means that simple “blast-wave” models cannot be used, as abundances of each particle species are free parameters there (this limitation applies both to $p + p$ and heavy-ion collisions). In contrast the THERMINATOR model [21] has both necessary features: it can use the traditional “blast-wave” hypersurface and provides absolute particle multiplicities.

The usage of the THERMINATOR makes it possible to test the impact of the strongly decaying resonances. As it was shown in [20], their introduction increases the apparent size of the system produced in heavy-ion collisions by approximately 1 fm and leads to the development of long-range tails in the emission function of pions. These effects were important but not dominant in the AuAu system (with the average size of 6 fm), but they may dominate in a pp collision where we expect the initial system to have a size of the order of 1 fm.

The paper is organized as follows. In Sec. II we briefly introduce the THERMINATOR model and its blast-wave variant. In Sec. III we describe the choice of parameters for the three simulations that were performed. In Sec. IV we describe the

* adam.kisiel@cern.ch

results for the inclusive transverse momentum p_T spectra and the one-dimensional femtoscopic simulations and explain the features observed there. In Sec. V we show the results of the 3D femtoscopic analysis and propose the most robust and sensitive observables that can be used to search for collectivity in the pp collisions at large multiplicities, e.g., at the LHC. In Sec. VI we discuss the origin of the effects seen in previous sections.

II. THERMINATOR MODEL AND FEMTOSCOPIC FORMALISM

In this work we use the THERMINATOR model [21], more specifically its variant that uses the blast-wave freeze-out hypersurface with the possibility to adjust the space-time correlation at the freeze-out hypersurface. For a detailed description of the model, its full theoretical basis and formalism see [20]. In short, average per-event particle abundances are calculated from the chemical model, where the temperature T and bariochemical potential μ_B are model parameters, and isospin μ_I and strange μ_S potential are calculated from conservation laws. For each event the number of particles of each type (we use 381 particle types taken from the Particle Data Group [22]) is randomly generated from a Poissonian distribution. Then each particle is put on the freeze-out hypersurface according to the generalized blast-wave prescription [20]:

$$\begin{aligned} & \frac{dN}{dyd\varphi p_\perp dp_\perp d\alpha_\parallel d\phi d\rho} \\ &= \frac{1}{(2\pi)^3} (\tau + a\rho) [m_\perp \cosh(\alpha_\parallel - y) - a p_\perp \cos(\phi - \varphi)] \\ & \times \left\{ \exp \left[\frac{\beta m_\perp}{\sqrt{1 - v_\perp^2}} \cosh(\alpha_\parallel - y) \right. \right. \\ & \left. \left. - \frac{\beta p_\perp v_\perp}{\sqrt{1 - v_\perp^2}} \cos(\phi - \varphi) - \beta\mu \right] \pm 1 \right\}^{-1}, \end{aligned} \quad (1)$$

where a , τ , $\rho_{\max}$, are parameters of the model. $\rho_{\max}$ is the transverse size of the system, τ is the proper time of emission, and a determines the transverse position (ρ) vs. time (t) correlation at the hypersurface. Of particular importance is the form of the radial flow velocity, which depends on particle emission position:

$$v_\perp = \frac{\rho/\rho_{\max}}{h_T + \rho/\rho_{\max}}, \quad (2)$$

where h_T is a parameter which controls the amount of flow in the system. Large values of h_T give small flow, small values of h_T give large flow, growing semilinearly from 0 at the center of the source to the maximum value of $1/(1 + h_T)$ at the edge. y , p_T , and φ are the components of the momentum of the particle (rapidity, transverse momentum, and azimuthal angle, respectively), while $\alpha_\parallel$, ρ , and ϕ are the corresponding coordinates of the emission point. As the last step, all unstable particles are propagated and decayed, in cascades if necessary, until only stable particles remain. The emission point of the particle is either its creation point on the hypersurface (we

call such particles “primordial”) or a decay point of its parent resonance.

Final state particle rescatterings after the emission are not included. We have estimated that for the particle densities in this work, each particle would rescatter on the average 0.8 times (the estimate was done for pions with realistic pion-pion scattering cross section). Most of the rescattering points would be very close to the original emission point, so the inclusion of rescattering would not change the emission shapes significantly.

Further in the work we show results of the analysis of the identical pion femtoscopic correlations (sometimes called “Bose-Einstein correlations” or “HBT correlations”). All the analyses were performed using the standard “two-particle” method: pions from the same event were combined into pairs, and assigned a weight equal to their wave-function squared, to form the signal. The pair wave function contained only the relevant symmetrization; final state coulomb and strong interactions were ignored as for pions they are small and are not important for the determination of the femtoscopic radii. Then, pions from different events were combined into pairs to form the background. The correlation function was constructed by dividing the signal by the background. It was then fitted with various functional forms, which will be described in detail in Sec. IV. For an extensive description of the two-particle method and the formalism see [20].

III. SIMULATION DESCRIPTION

We performed the simulations with the blast-wave model with the following parameters: transverse size of the freeze-out hypersurface $\rho_{\max} = 1.5$ fm, proper time of freeze-out $\tau_0 = 2.0$ fm/c, freeze-out temperature $T_f = 165.6$ MeV, bariochemical potential $\mu_B = 28.5$ MeV. The freeze-out hypersurface was slightly sloped in the $\rho - t$ plane ($a = -0.2$). The three simulations differed only in the value of the h_T parameter; the “no-flow” had $h_T = 10$, the “moderate flow”: $h_T = 1$ and the “strong flow”: $h_T = 0.5$. The resulting flow velocity

FIG. 1. (Color online) Average common transverse flow of primordial pions [see Eq. (3)] as a function of the distance to the central axis of the source ρ . “No flow” results are shown as blue circles, “moderate flow” as red triangles, “strong flow” as green squares. The input model flow velocity profiles [see Eq. (2)] are shown as dashed lines.

profiles can be seen in Fig. 1 as dashed lines. The common radial flow present in the model results in strong space-momentum $\rho - p_T$ correlations of the produced particles, which is reflected in the the common radial flow:

$$\langle p_T \parallel \rho \rangle = \left\langle \frac{p_T \rho \cos(\phi - \varphi)}{\rho} \right\rangle, \quad (3)$$

where averaging is done over particles. The results are shown in Fig. 1 as symbols. The plot illustrates that the “no flow” scenario has very little $\rho - p_T$ correlations, while in the “strong flow” scenario this correlation is significant. For each of the scenarios we have simulated 2×10^6 events.

IV. RESULTS

Fixing the model parameters, as described above, fixes the average absolute multiplicity of particles per event. The resulting $\langle dN_{ch}/d\eta \rangle$ for the “strong flow” sample was 8.3. This is about twice the average multiplicity reported by the ALICE Collaboration at $\sqrt{s} = 900$ GeV and is above the average multiplicity for $\sqrt{s} = 7$ TeV [2–4]. In cases with smaller flow the total energy in the system is smaller, therefore the average multiplicities slightly smaller and are 7.2 for the “moderate flow” case and 6.1 for the “no flow” case. One could make them bigger by increasing the initial size parameters. We chose not to do it, in order to easily compare systems which differ in only one of the parameters—the transverse flow magnitude. More involved studies, which would aim to describe not only the relative trends but also the absolute magnitude of the radii will need to adjust all the parameters. We leave such studies for further works.

The impact that the radial flow has on particle spectra is shown in Fig. 2. Only for the simplest case of “no flow” and primordial particles only [shown in panel (d)] the spectra for all particle types are exponential. We fit the distributions

FIG. 2. (Color online) Single particle inclusive m_T spectra for pions (green squares), kaons (red triangles), and protons (blue circles). Upper panels (a) and (b) show simulations with “strong flow,” lower panels (c) and (d) with “no flow.” Left panels (a) and (c) show all particles, right panels (b) and (d) only primordial particles.

TABLE I. Inverse slope parameters of the single particle spectra shown in Fig. 2, in MeV.

	pions	kaons	protons
	all		
“strong flow”	234 ± 1	323 ± 4	414 ± 8
“no flow”	157 ± 1	168 ± 2	157 ± 3
	primordial		
“strong flow”	285 ± 2	360 ± 6	465 ± 2
“no flow”	183 ± 1	181 ± 3	176 ± 8

with an exponential and extract their inverse slope, shown in Table I. Even in the “no flow” case the simple addition of strongly decaying resonances obscures the picture, as seen in panel (c). Resonances tend to populate the lower m_T regions more abundantly than higher m_T . There are relatively more resonances decaying into pions and protons than into kaons. As a result, the inverse slope parameter is higher for pions and protons and lower for kaons. For primordial particles in “strong flow” in panel (b) we see, as expected, that the thermal “exponential” shapes are modified by radial flow: pions get a “convex” shape, while kaons and protons develop a positive curvature. All are less steep than in the “no flow” case, resulting in the larger inverse slope parameter. Adding resonance decay products, shown in panel (a), decreases the inverse slope parameter by 50 MeV, so any attempt to extract the flow velocity and temperature from the inverse slope parameters must fully take into account the resonance contribution. In Fig. 3 the kaon to pion and proton to pion ratios vs. p_T are shown. For K/π both flow and addition of resonances decrease the ratio. This comes from the larger contribution of resonance decays to pions. In contrast, for p/π only flow decreases the ratio, as the relative contributions of resonances to both pions and protons are similar. In summary, the particle spectra retain the general characteristics associated with radial flow even when combined with resonance decays in a small system, but resonance decays do alter the quantitative estimates of temperature significantly. In addition our simple model does not include the hard processes which will contribute to all particle’s spectra at larger p_T and complicate the picture further. Therefore making strong conclusions about

FIG. 3. (Color online) Kaon to pion (a) and proton to pion (b) ratios vs. p_T . Green dashed lines show primordial particles for “no flow,” red long-dashed show all particles for “no flow,” blue dot-dashed show primordial for “strong flow,” violet full line show all for “strong flow.”

radial flow from particle spectra alone is not trivial and is model dependent.

Another observable which was extensively used in conjunction with p_T spectra as a signature of collective flow is the pion femtoscopic radius of the system, more specifically its monotonic decrease with growing pair momentum $k_T = |\mathbf{p}_{T,1} + \mathbf{p}_{T,2}|/2$. The first measurement of such a dependence at the LHC was reported in Ref. [4]. The minimum-bias measurement shows no significant dependence, however the combined multiplicity vs. k_T analysis shows that the behavior of “low” and “high” multiplicity events is different. The “low” multiplicity (M) events show an increase of pion femtoscopic radius R_{inv} at low k_T and then a fall at high k_T . The “high” M events on the other hand suggest a decrease of the R_{inv} radius with k_T , although it is also consistent with a flat dependence. New results from the LHC, especially from collisions at the higher energy of $\sqrt{s} = 7$ TeV will deliver results with better precision and will show if the trend continues at higher multiplicities.

We have obtained the pion femtoscopic correlation functions using the two-particle method described in Sec. II. We divide the sample into six k_T bins: (0.12, 0.2), (0.2, 0.3), (0.3, 0.4), (0.4, 0.5), (0.5, 0.6), and (0.6, 0.7) GeV/c. We begin by analyzing the one-dimensional correlation functions. Following the experiments [4,7], we fit them with two functional forms, the Gaussian:

$$f_G(q_{\text{inv}}) = 1 + \lambda \exp(-R_{\text{inv}}^G q_{\text{inv}}^2), \quad (4)$$

and the exponential:

$$f_E(q_{\text{inv}}) = 1 + \lambda \exp(-R_{\text{inv}}^E q_{\text{inv}}). \quad (5)$$

An example of the fits is shown in Fig. 4. We see a behavior similar to the one reported by the LHC experiments: the correlation function is better described by an exponential. The Gaussian fit is worse, but it characterizes the overall width of the correlation function.

In Fig. 5 we plot the results of the one-dimensional analysis vs. k_T for the three scenarios of flow. The Gaussian and properly scaled exponential fit results are consistent. In the “no flow” scenario the R_{inv}^G is growing with k_T , similar to low multiplicity collisions at the LHC. On the other hand, as flow

FIG. 4. (Color online) One-dimensional pion femtoscopic correlation function for the “strong flow” case, k_T bin of (0.2, 0.3) GeV/c. Triangles are the results of the simulation, dashed line is a Gaussian fit, dotted line is the exponential fit.

FIG. 5. (Color online) Overall one-dimensional femtoscopic radius R_{inv} of the system. “No flow” are shown as blue circles, “moderate flow” as red triangles, “strong flow” as green squares (the latter two shifted slightly for clarity). Gaussian radii R_{inv}^G are shown as closed symbols, exponential radii R_{inv}^E as open symbols.

develops, the R_{inv}^G becomes flat vs. k_T for “moderate flow” and has a negative slope for “strong flow.” In other words, from the comparison of the R_{inv}^G vs. k_T trends between data and the model, we observe a development of radial flow with the increase of multiplicity in the pp collisions at the LHC. It would be interesting to see if the data at higher multiplicities at $\sqrt{s} = 7$ TeV continue the trend and develop an even stronger k_T dependence. The results are intriguing enough to attempt a more in-depth explanation of their origin. We aim to propose more strict tests, which would make the conclusions from such comparisons less model dependent.

V. 3D BERTSCH-PRATT ANALYSIS

We have performed the 3D analysis of the simulated correlation functions. We have used the standard Bertsch-Pratt decomposition of the relative momentum into the “long” (along the beam axis), the “out” (along the pair transverse momentum), and the “side” (perpendicular to the other two) directions. We have used the longitudinally co-moving system (LCMS) reference frame, where the longitudinal momentum of the pair vanishes. We have analyzed two classes of pions: the “primordial” only and “all” particles. The latter includes the “primordial” plus those from strongly decaying resonances. In Fig. 6 we show the correlation function, separately in *out*, *side*, and *long* directions. We attempted to fit both the 1D projections and the full 3D function with combinations of Gaussian, exponential, and Lorentzian shapes. These three were chosen because they have an analytic relation between correlation function and pair emission function forms. We found that (a) for primordial particles a 3D Gaussian is the best fit, (b) for all particles the *out* and *long* direction is best fitted with an exponential, while *side* remains Gaussian. This is true both for 1D projections (shown in the figure) as well as for a full 3D fit. Even though the fit in Fig. 6 is best out of three forms tried, it is still not perfect. Nevertheless, the width of the function is adequately estimated, which is enough for the purpose of this paper. The investigation of the details of the shape of the correlation functions we leave for future studies.

FIG. 6. (Color online) Projections of the three-dimensional correlation function for the “strong flow” simulation for $k_T(0.3, 0.4)$ GeV/c. Dashed lines show fits to 1D projections. (a) shows the *out* projection with an exponential fit. (b) shows *side* with Gaussian, (c) shows *long* with exponential. To plot the projections, other components were integrated in the range $|q_{\text{other}}| < 0.03$ GeV/c.

Based on the study of correlation function shapes, for primordial particles a Gaussian 3D ellipsoid, with different sizes in all directions is a reasonable description of the correlation:

$$C_G(\mathbf{q}) = 1 + \lambda \exp \left(-R_{\text{out}}^G q_{\text{out}}^2 - R_{\text{side}}^G q_{\text{side}}^2 - R_{\text{long}}^G q_{\text{long}}^2 \right), \quad (6)$$

where R_{out}^G , R_{side}^G , and R_{long}^G are Gaussian radii in *out*, *side*, and *long* directions respectively, λ is the strength of the correlation. For all particles we have used the following fit functional form:

$$C_E(\mathbf{q}) = 1 + \lambda \exp \left(-R_{\text{out}}^E |q_{\text{out}}| - R_{\text{side}}^G q_{\text{side}}^2 - R_{\text{long}}^E |q_{\text{long}}| \right), \quad (7)$$

where R_{out}^E and R_{long}^E are exponential radii in *out* and *side* directions. Similarly to the one-dimensional case, the radii

FIG. 7. (Color online) k_T dependence of the three-dimensional radii for the primordial particles. Blue circles are “no flow” simulations, red triangles are “moderate flow,” green squares are “strong flow.” (a) shows R_{out}^G , (b) shows R_{side}^G , (c) shows R_{long}^G , (d) shows $R_{\text{out}}^G/R_{\text{side}}^G$ ratio.

obtained from Gaussian and exponential fits are not directly comparable, as they are parameters of different functional forms.

In Fig. 7 we show the Gaussian femtoscopic radii obtained from the analysis of “primordial” particles. We concentrate on the transverse dynamics. For the R_{out}^G radius we observe an expected behavior: the “no flow” scenario shows a flat k_T dependence, the slope grows with increasing flow and is significant for “strong flow.” In addition, the size gets *smaller* with increasing flow; the well-known mechanism of the “lengths of homogeneity” is at work here. R_{side}^G dependence is weakly changed by flow; stronger flow creates a small slope. The change in behavior is best visible in the $R_{\text{out}}^G/R_{\text{side}}^G$ ratio. For “no flow” it is large, at least 1.0–1.5, for “strong flow” it is falling, and smaller, below 1.0. The $R_{\text{out}}^G/R_{\text{side}}^G$ ratio has an advantage that it is, to a limited degree, independent of scale. The absolute values of the transverse radii are directly depending on the assumed system size (the ρ_{max} parameter), while the ratio is less dependent on it, and its qualitative behavior should not change as we change ρ_{max} in a range which would produce multiplicities observed in pp collisions.

For primordial particles we have identified the $R_{\text{out}}^G/R_{\text{side}}^G$ ratio as a sensitive, scale-independent, probe of the amount of collectivity in the system. However, the majority of pions are coming from resonance decays, which strongly influence spatial and temporal characteristics of the emission process. It

FIG. 8. (Color online) k_T dependence of the three-dimensional radii for all particles. Blue circles are “no flow” simulations, red triangles are “moderate flow,” green squares are “strong flow.” (a) shows R_{out}^E , (b) shows R_{side}^G , (c) shows R_{long}^E , (d) shows R_{out}^E/R_{side}^G ratio.

is therefore crucial to investigate how they modify the signal present for “primordial” particles. In Fig. 8 we show the fit results for all pions. We stress that inclusion of resonances changes the emission patterns visibly, which is reflected in the change of the fit functional form. R_{out}^E for “no flow” is larger than for “primordial” and develops a k_T slope. In contrast, the “strong flow” R_{out}^E grows little: R_{out}^E grows less as flow grows. That is surprising, as the “length of homogeneity” argument seems to hold for particles from resonances, even though they do not come directly from a “collective” medium. Also the negative slope of the R_{out}^E dependence on k_T alone is not a good signature of collectivity as such behavior also develops for “no flow.” R_{side}^G shows the behavior opposite to R_{out}^E . For “no flow” it is similar to “primordial” case, while for “strong flow” it grows. As a consequence, again in striking contrast to the behavior of R_{out}^E , R_{side}^G grows more as radial flow increases. Again the R_{out}^E/R_{side}^G ratio is the most sensitive to these changes, and for all particles it shows an even more dramatic difference between flow scenarios: the value for the ratio is up to a factor of 1.5 larger for “no flow” as compared to “strong flow.” In summary, we show that a small value of R_{out}^E/R_{side}^G ratio, close to unity, is a signal of emission of all particles from a collective medium. The value is monotonically dependent on the strength of flow: the stronger the flow, the lower the R_{out}^E/R_{side}^G ratio. In addition, the inclusion of resonance decay products *magnifies* this flow effect.

VI. ORIGINS OF FEMTOSCOPIC RADII BEHAVIOR

The identification of the R_{out}^E/R_{side}^G ratio as a sensitive probe of collective particle emission in systems resembling pp collisions at the LHC has consequences for experiments. It allows to test a hypothesis of a creation of a collective system in such collisions, which produces large particle multiplicities. In addition, the femtoscopic pion analysis is practically a “minimum-bias” measurement and does not require advanced triggering or acceptance corrections. It can therefore be performed quickly, with the data samples that the LHC experiments already have on tape. It is therefore critical to understand why such effect arises and how general their emergence is; in other words how model dependent is the link between R_{out}^E/R_{side}^G ratio and system collectivity.

The separation between two particles in the *out* direction is

$$r_{out} = x_1 - x_2 + \beta(t_2 - t_1), \quad (8)$$

where $x = \rho p_T \cos(\phi - \varphi)/p_T$ are the transverse emission point coordinates of the first and the second particle, projected on the pair velocity, t_1 and t_2 are the emission times of the particles, and β is the pair velocity (we neglect the differences between particles velocities since in order to be femtoscopically correlated their relative velocity must be small). We order the particles in a pair in such a way that $t_2 > t_1$, the third term in the formula reflects the fact that the first particle has already traveled some distance before the second was born. The definition of the *side* direction is that the pair velocity in this direction vanishes, so the formula for the separation is simpler:

$$r_{side} = y_1 - y_2, \quad (9)$$

where $y = [\rho p_T \sin(\phi - \varphi)]/p_T$ are the transverse emission point coordinates of the first and second particle, now perpendicular to the pair velocity. x coordinates are on the x axis of the top panels of Figs. 9 and 10, y are on the x axis of the bottom panels and t is on the y axis of all panels. The $R_{out}^{E,G}$ and R_{side}^G femtoscopic radii have the meaning of variances of *out* and *side* distributions, respectively (conventionally divided by $\sqrt{2}$). They are proportional to a combination of the single particle space-time distribution moments:

$$R_{out}^{E,G} \approx \sigma_x + \beta \sigma_t - C(x, \beta t), \quad (10)$$

$$R_{side}^G \approx \sigma_y, \quad (11)$$

where σ_x , σ_y , and σ_t are the variances of the x , y and t distributions respectively and $C(x, \beta t)$ is the term that accounts for a possible space-time correlation in the emission process. We note that C contributes to the size with a negative sign: it means that positive $x - t$ correlation *decreases* the *out* radius [23,24].

In Fig. 9 the space-time characteristics of the emission process for the “no flow” scenario is shown. In (a) and (d) are primordial particles. In this case both out and side directions show similar size, which does not depend on time. Also, the mean emission points are not shifted. Panels (b) and (e) show the behavior for resonance decay products. There is a small correlation between mean x and time. Also, out and side spreads grow with time, but only slightly. As a consequence,

FIG. 9. (Color online) Correlation between space and time in particle emission from the “no flow” model. (a), (b), and (c) show “outwards” direction [x from Eq. (8)]: projection on particles’ velocity. (d), (e), and (f) show “sideways” [y from Eq. (9)] direction: projection perpendicular to the velocity. (a) and (d) show primordial particles, (b) and (e) resonance products, (c) and (f) all particles. The graphs overlaid on the plots show the mean and spread of the distribution in a time bin.

for all particles shown in (c) and (f), the out emission point is weakly correlated with time and the side spread grows only slightly.

In Fig. 10 we present a similar study for the “strong flow” scenario. We concentrate on the differences between this scenario and the “no flow” one. For primordial particles, the out mean position is shifted by flow. The spread is slightly smaller than side; hence the smaller than one ratio for green points in panel (a) of Fig. 7. For resonance products we see more striking differences. The out mean position is strongly positively correlated with time. The spread also grows with time, but less than for the side, where the growth is more pronounced than for the “no flow” case. As a result, for all particles, the out mean position is positively correlated with time, while the side spread grows stronger than in the “no flow” case.

We can now understand the dramatic change of the $R_{\text{out}}^E/R_{\text{side}}^G$ ratio between “no flow” and “strong flow” scenarios. For “no flow”, σ_x and σ_y are similar, as can be seen in panels (c) and (f) of Fig. 9. But R_{out}^E is further increased by σ_t and the $x - t$ correlation is too small to counterbalance it. As a result $R_{\text{out}}^E/R_{\text{side}}^G$ is large. In short, resonance decays introduce significant spread in emission times, which acts to increase R_{out}^E . The fact that resonance decay points are distributed according to the exponential decay time multiplied by velocity also explains why the shape of the correlation in the *out* direction changes dramatically, while it stays Gaussian in *side*, where the velocity vanishes.

In the “strong flow” scenario we note that even for primordial particles, seen in panel (a) and (d) of Fig. 10, σ_x is smaller than σ_y —the effect of flow of primordial pions. The effect is further increased by resonances products. Because of the existence of common radial flow in the system, when resonances are born on the freeze-out hypersurface, they are already flying to the outside of the system. Since they decay after some time, the emission points of the daughter pions will, on average, be further apart compared to the “no flow” case. This is nicely illustrated as the strong growth of the σ_y with time on panel (e) of Fig. 10. The result is a significant increase of R_{side}^G , as compared to the “no flow” case. One expects the same argument to hold for σ_x , and to some extent it is seen in panel (b) of Fig. 10. However, the growth is not as strong as for σ_y —this is because the resonance emission points are also governed by flow, so their parent σ_x is smaller than σ_y , just like for primordial pions on panels (a) and (d). But more importantly, radial flow of resonances also results in much stronger (as compared to “no flow” case) positive $x - t$ correlation, seen most clearly in panel (b). This correlation also dominates when we consider both primordial and resonance daughter pions in panel (c). This will further decrease the R_{out}^E parameter, due to the $C(x, \beta t)$ term in Eq. (10). As a result, the $R_{\text{out}}^E/R_{\text{side}}^G$ ratio is small. In short, radial flow of primordial pions and resonances produces several effects which all combine together to bring the $R_{\text{out}}^E/R_{\text{side}}^G$ ratio down, even below unity if the flow is strong enough. The stronger the flow, the lower the ratio.

FIG. 10. (Color online) Correlation between space and time in particle emission from the “strong flow” model. (a), (b), and (c) show “outwards” direction: projection on particles’ velocity. (d), (e), and (f) show “sideways” direction: projection perpendicular to the velocity. (a) and (d) show primordial particles, (b) and (e) resonance products, (c) and (f) all particles. The graphs overlaid on the plots show the mean and spread of the distribution in a time bin.

We also now understand why one-dimensional R_{inv} radii dependence on k_T changes as flow develops. R_{inv} is approximately equal to the quadratic average of γR_{out} , R_{side} , and R_{long} , where the γ factor in front of R_{out} comes from the boost (with pair velocity) from the LCMS to pair rest frame, where R_{inv} is determined. For “no flow” case R_{out} is flat with k_T in LCMS, so the growth of the boost as k_T increases drives R_{inv} up. For cases with flow, the growth of the boost is countered by the fall of R_{out} resulting from flow. And since the other two radii also decrease with k_T , the slope of the R_{inv} dependence on k_T becomes negative.

As a final note we stress that the model employed here is a simple hydrodynamics-inspired parametrization. We have shown that one of its particular features—an $\rho - p_T$ correlation, also known as radial flow, produces specific signatures in single particle spectra as well as 1D and 3D pion femtoscopic radii. However it does not address the question of where such correlations come from. One possibility is that they come from a “medium” that is described by hydrodynamic equations, as suggested in Ref. [13], but there may be other mechanisms that produce such correlations.

VII. CONCLUSIONS

We have performed simulations with the THERMINATOR model, employing emission of primordial particles from a blast-wave hypersurface, and including the propagation and

decay of a full set of hadronic resonances. The parameters of the calculations have been adjusted, so that they produce multiplicities observed in high multiplicity pp collisions at LHC energies. Three sets of parameters have been used, which differ only by the strength of the radial flow (the $\rho - p_T$ correlation) in the system.

The radial flow modifies the particle spectra in the expected way, creating a convex shape for pions and positive curvature for kaons and protons. However, addition of resonance decays significantly modifies these shapes, influencing the extracted slope parameter and making a direct extraction of flow velocity from spectra measurements alone complicated and model dependent.

Results of one-dimensional femtoscopic analysis show that with no flow, the 1D R_{inv}^G pion femtoscopic radius is expected to grow with pair momentum k_T . In contrast, as flow is increased, the dependence changes to a decrease with k_T . Therefore, within the frame of this model the results of the ALICE Collaboration on multiplicity vs. k_T dependence can be interpreted as signature of the development of radial flow with increasing per-event particle multiplicity.

Three-dimensional pion femtoscopic radii show that the shape of the correlation is exponential in the *out* and *long* directions due to contributions from resonance decay products, while in *side* the shape remains Gaussian. Radial flow (or lack of it) for primordial pions and resonances has opposite effect on R_{out}^E and R_{side}^G radii. The first grows less as flow develops,

which is a result of a combination of two effects: (a) decrease of lengths of homogeneity for primordial pions and resonances and (b) strong $x - t$ correlations coming from resonances' flow-ordered velocities. R_{side}^G grows more as flow develops, the effect of the preferential propagation of the resonances to the outside of the source in the presence of flow. We propose to use the $R_{\text{out}}^E/R_{\text{side}}^G$ ratio as a probe of the amount of radial flow (e.g., velocities ordering) in the small, resonance-dominated system, produced in the high multiplicity collisions at the LHC. Any

analysis aiming to search for collective effects in such systems should combine at least the three measurements: identified particle p_T spectra, 1D and 3D femtoscopic radii.

ACKNOWLEDGMENTS

The author would like to thank Yiota Foka and Dariusz Miśkowiec for helpful comments and discussions.

-
- [1] K. Aamodt *et al.* (ALICE Collaboration), [Eur. Phys. J. C **65**, 111 \(2010\)](#).
 - [2] K. Aamodt *et al.* (ALICE Collaboration), [Eur. Phys. J. C **68**, 89 \(2010\)](#).
 - [3] K. Aamodt *et al.* (ALICE Collaboration), [Eur. Phys. J. C **68**, 345 \(2010\)](#).
 - [4] K. Aamodt *et al.* (ALICE Collaboration), [Phys. Rev. D **82**, 052001 \(2010\)](#).
 - [5] K. Aamodt *et al.* (ALICE Collaboration), [Phys. Lett. B **693**, 53 \(2010\)](#).
 - [6] V. Khachatryan *et al.* (CMS Collaboration), [JHEP **02** \(2010\) 041](#).
 - [7] V. Khachatryan *et al.* (CMS Collaboration), [Phys. Rev. Lett. **105**, 032001 \(2010\)](#).
 - [8] V. Khachatryan *et al.* (CMS Collaboration), [Phys. Rev. Lett. **105**, 022002 \(2010\)](#).
 - [9] M. A. Lisa, S. Pratt, R. Soltz, and U. Wiedemann, [Annu. Rev. Nucl. Part. Sci. **55**, 357 \(2005\)](#).
 - [10] V. Khachatryan *et al.* (CMS Collaboration), [JHEP **09** \(2010\) 091](#).
 - [11] A. Dumitru *et al.*, [Phys. Lett. B **697**, 21 \(2011\)](#).
 - [12] P. Bozek, [Eur. Phys. J. C **71**, 1530 \(2011\)](#).
 - [13] K. Werner, I. Karpenko, T. Pierog, M. Bleicher, and K. Mikhailov, [Phys. Rev. C **83**, 044915 \(2011\)](#).
 - [14] T. J. Humanic, [Phys. Rev. C **76**, 025205 \(2007\)](#).
 - [15] M. Aggarwal *et al.* (STAR Collaboration), [Phys. Rev. C **83**, 064905 \(2011\)](#).
 - [16] Z. Chajęcki and M. Lisa, [Phys. Rev. C **79**, 034908 \(2009\)](#).
 - [17] S. V. Akkelin and Y. M. Sinyukov, [Phys. Lett. B **356**, 525 \(1995\)](#).
 - [18] E. Schnedermann, J. Sollfrank, and U. W. Heinz, [Phys. Rev. C **48**, 2462 \(1993\)](#).
 - [19] F. Retiere and M. A. Lisa, [Phys. Rev. C **70**, 044907 \(2004\)](#).
 - [20] A. Kisiel, W. Florkowski, W. Broniowski, and J. Pluta, [Phys. Rev. C **73**, 064902 \(2006\)](#).
 - [21] A. Kisiel, T. Taluc, W. Broniowski, and W. Florkowski, [Comput. Phys. Commun. **174**, 669 \(2006\)](#).
 - [22] W. M. Yao *et al.* (Particle Data Group), [J. Phys. G **33**, 1 \(2006\)](#).
 - [23] C. M. Ko, [AIP Conf. Proc. **828**, 439 \(2006\)](#).
 - [24] S. Pratt, [Phys. Rev. Lett. **102**, 232301 \(2009\)](#).