EUROPEAN ORGANIZATION FOR NUCLEAR RESEARCH

CERN-EP/81-134
23 October 1981

NUCLEON LIFETIME EXPERIMENTS

D. Treille

CERN, Geneva, Switzerland

Talk given at the
Cargèse Summer Institute, July 1981

The possibility that the nucleon could be unstable has long been considered. However, it is clear that the advent of Grand Unified Theories (GUTs)¹⁾ has given a new impetus and clear motivations to this line of research. I would therefore like to start with a brief (and somewhat superficial) review of what GUTs predict.

1. THEORETICAL PREAMBLE

The mechanism leading to nucleon decay is readily understood by looking at the SU(5) multiplets (Fig. 1). In the matrix of the gauge bosons, we recognize the gluon octet, the SU(2) × U(1) sector, and new non-diagonal parts. When sandwiched between fermion multiplets, in which quarks and leptons co-exist, we see clearly that these non-diagonal parts will induce "cross-talk" between the quark and the leptonic world, making possible processes such as that shown in Fig. 1c. The X and Y are the leptoquarks mediating nucleon decay. They are supposed to be very heavy. If M_X is their mass [in SU(5) there is one single mass to be considered], the expression of the propagator leads to a proton lifetime varying as M_X^{-4} . So the first task is to evaluate M_X ²⁾.

This mass is the grand unification mass, at which the three evolving effective coupling constants -- of the strong and electroweak sectors -- are supposed to merge (Fig. 2a). The reason for the difference in the evolution of the three couplings is shown in Fig. 2b. This difference, naïvely stated, is simply due to the fact that to probe small structures -- or equivalently to feel the effect of loops with high masses -- we need to have a sufficient resolution. Each interaction has its own trajectory, and the aim is

- i) to project them as accurately as possible;
- ii) to match the low-energy couplings with experimental values and the Glashow-Weinberg-Salam (GWS) model.

By doing this, we are left with a single equation:

$$\frac{1}{\alpha_{\text{em}}} - \frac{8}{3\alpha_s(M_W^2)} = \frac{11}{2\pi} \ln \left(\frac{M_X^2}{M_W^2} \right).$$

Knowing the "running" strong coupling α_s , the electromagnetic coupling, and the W mass, we can calculate M_X . Taking $\alpha_s(M_W^2) = 0.14$, which corresponds to $\Lambda = 400$ MeV in the usual, but perhaps irrelevant, expression for α_s :

$$\alpha_s(Q^2) = \frac{12\pi}{(33-2f) \ln(Q^2/\Lambda^2)},$$

the result is

$$M_X = 2 \times 10^{16} \text{ GeV},$$

which is a total disaster, since the corresponding nucleon lifetime, τ_N , would be several orders of magnitude beyond reach.

However, the naïve calculation involves several approximations which represent large modifications of τ_N because one computes the logarithm of the mass and then raises M_X to the fourth power. When these approximations are corrected for²⁾ (see Table 1), the disaster turns into a triumph: M_X decreases by more than a factor of 50 and becomes $\sim 4 \times 10^{14}$ GeV. The major step came simply from renormalizing α from the Thomson limit to the correct value at M_W . From M_X , the computation of the nucleon lifetime (Fig. 3) still involves a few delicate steps, such as the evaluation of the wave function at the origin, $|\psi(0)|^2$, which governs the annihilation rate of the two quarks. The results obtained by various authors within SU(5) fall within the region

$$\tau_p = 10^{31 \pm 2} \text{ years}.$$

The error in the exponent includes the dispersion of the different results as well as the safety margin generally quoted by each group.

Regarding the branching ratios into two-body final states (Table 2), there are a few disagreements. One (which is difficult to understand -- at least by me) concerns the rate into μ 's: some authors³⁾ expect $\sim 20\%$, whilst others⁴⁾ expect almost nothing. This may be due to the treatment of phase space. Another disagreement concerns the nature of the meson accompanying the positron: generally a pion is favoured, but one model⁵⁾ claims that vector mesons will dominate.

Moreover, some authors⁶⁾ argue that three-body decay modes can be as important as two-body modes.

What we said about SU(5) is in fact valid for all models satisfying the "Desert" hypothesis. General arguments^{7,8)} involving dimensional counting for baryon-violating terms, lead to the retention of only those operators which give the following selection rules:

$$\begin{aligned}
 \text{i)} \quad \Delta B = \Delta L \quad & p \rightarrow e^+ \pi^0 \\
 & n \rightarrow e^+ \pi^- \\
 & n \not\rightarrow e^- \pi^+ \\
 \\
 \text{ii)} \quad \Delta S / \Delta B \leq 0 \quad & p \rightarrow K^+ \bar{\nu} \\
 & n \not\rightarrow K^- e^+ \\
 \\
 \text{iii)} \quad \Delta I = 1/2 \quad & \Gamma(p \rightarrow e^+ \pi^0) = \frac{1}{2} \Gamma(n \rightarrow e^+ \pi^-) \\
 & \text{etc.}
 \end{aligned}$$

Furthermore, the exchange of heavy vector bosons leads to the existence of a universal antilepton polarization. This is illustrated for O(10)⁹⁾ in Fig. 4.

However, the existence of this desert is simply an assumption. Left/right symmetric models¹⁰⁾, such as SU(16), having spontaneous violations of baryonic B and leptonic L numbers, would permit other types of nucleon decays which may be classified according to the degree of B-L violation:

$$\begin{aligned}
 \text{i)} \quad \Delta(B-L) = -4 \quad & p \rightarrow \underline{\underline{3\nu}} + \pi^+ \\
 \text{ii)} \quad \Delta(B-L) = -2 \quad & p \rightarrow \underline{\underline{e^-}} \pi^+ \pi^+ \\
 \text{iii)} \quad \Delta(B-L) = 0 \quad & \text{see above} \\
 \text{iv)} \quad \Delta(B-L) = 2 \quad & p \rightarrow \underline{\underline{3\bar{\nu}}} + \pi^+
 \end{aligned}$$

We see that class (iii) is the only one which was predicted by "desert" models. These "abnormal" proton decays imply the existence of intermediate mass scales. We may ask which mass scale would, for a given class, lead to an observable nucleon

lifetime in the 10^{30} - 10^{33} years range¹¹⁾: For $N \rightarrow 3\nu + X$, for instance, the answer is 10^3 - 10^4 GeV. Also relevant to the existence of such intermediate mass scales is the phenomenon of n - $\bar{n}$ oscillations¹²⁾, which will be described later.

To summarize, What would we like to know?

- 1) Independently of all subtleties, it is clear that the irrefutable demonstration of nucleon instability would be a fantastic adventure. Obtaining such a clear result implies that huge problems linked to the rate and the backgrounds should be solved, and discussing them will be my main objective.
- 2) Then, in order of increasing audacity, one would like to:
 - know the sign of the lepton in order to check the selection rules;
 - find its polarization;
 - obtain the exact conservation laws;
 - observe the nature of the accompanying hadronic final state¹³⁾;
 - check whether free quarks are liberated; people cannot help playing with big numbers, and it has already been noticed¹⁴⁾ that the fraction of nucleons which could have decayed since the beginning of the Universe, assuming for the proton lifetime the value found above, is of the same order of magnitude as the fraction of non-integer charges found in the Stanford experiment¹⁵⁾;
 - check whether Poincaré and Lorentz invariance holds¹⁶⁾ for processes occurring at such a wild scale of space time (10^{-28} cm).

2. THE EXPERIMENTAL PROBLEM

2.1 The general idea

Nothing is more simple in principle. We have just to observe enough nucleons during enough time: 10^3 tons contain 6×10^{32} nucleons. If $\tau = 6 \times 10^{30}$ years, we would get 100 decays/year in such a mass.

The problem is to avoid or reject backgrounds which could simulate the signal.

2.2 The signal

We are looking for a decay which in its ideal configuration is easy to imagine: a lepton with momentum balanced by a π or a vector meson and ~ 1 GeV energy deposit. However, things are not so clear.

A major problem is linked to the fact that the decaying nucleon is bound inside a nucleus. First, Fermi momentum ruins the back-to-back configuration and apparent momentum balance of the products, but the total energy is unchanged. Figure 5 shows, for instance, the resulting plot for $p \rightarrow e^+ \pi^0$ in the (acollinearity versus fractional momentum) plane¹⁷⁾. This spread is obviously important, because it allows background to creep in.

Then the decay products have to escape from the nucleus. Figure 6 gives the probability of non-reinteraction for pions in various nuclei as a function of energy. η and ω escape before decaying. The ρ does not, and should be treated as two pions. Summarizing various measurements and estimates, we can say that for nucleon decay in iron, one pion escapes from the nucleus for every two produced. If we further require that no energy loss occurs, then only one pion in three survive. More details can be found elsewhere¹⁸⁾.

Then, can one get the sign of the lepton? Implementing a sufficient magnetic field on such a volume seems to discourage the richest physicists since a respectable field (a few kilogauss) is needed. Using the delayed μ -decay is in principle the solution. However, this does not work for water since the μ^+ and μ^- have the same lifetimes because the nuclei are too small to cause much capture of μ^- ¹⁹⁾, whereas for iron, the mechanism is indeed viable: $\tau^-/\tau^+ \approx 0.07$. The problem here may be linked to the structure of the calorimeter: the low-energy positron (~ 30 MeV) has to cross iron and find its way to detectors (tubes) which are "alive", i.e. which have not been already fired by the decay itself and set into dead time.

Another possibility is to look, not at the direct μ (which according to some authors does not exist anyway), but at the μ from the π decay, for instance in

$n \rightarrow e^+ \pi^-$ ($\pi^- \rightarrow \mu^-$). What is relevant now is the difference in the behaviour of π^+ and π^- due to absorption of the latter indicated by the absence of μ 's.

Finally, can we measure the polarization of the lepton? This question is relevant only for μ 's, again through their $\mu^+ \rightarrow \dots$ decay. First we need a non-depolarizing medium, so that the stopping μ keeps its polarization. Water is all right but not iron. If x is the fraction of the μ momentum carried by the electron and P the muon polarization, the electron angular distribution goes as

- i) $1 - P \cos \theta$ if $x \approx 1$
- ii) $1 - 1/3 P \cos \theta$ if all x are retained.

The problems are evident:

- a) the nucleon decay to $\mu^+ + X$ may have a small branching ratio;
- b) we will be forced to keep all x 's so that the effect is small;
- c) measuring the angle between the μ and the decaying electron, even with poor resolution, is problematic. Čerenkov techniques (see later) claim that an angular resolution of 25° can be achieved.

2.3 Muon background

These are μ 's originating from atmospheric showers. The incident spectrum has a power law in energy. At a shallow depth, energy loss by ionization maintains this power law. However, at greater depths, only very energetic muons are present: for these, other types of energy losses (bremsstrahlung, etc.) which are energy-dependent have been at work, and the energy spectrum has become an exponential. Vertically it is $\sim A \exp(-h/\lambda)$. At an angle θ , there are two competing effects: in the atmosphere the π 's have a longer decay path, whereas in earth the shielding is increased.

The real flux through a given underground set-up²⁰⁾ depends on the exact thickness of the overlying rock. Figure 7 shows how different this can be for a flat surface or for a hemispherical one. In this calculation²⁰⁾ such rock layers were assumed for the Mont Blanc and Fréjus tunnels, respectively. Experimentally

the observed distributions do not reproduce the predictions well: it is thus demonstrated that the Mont Blanc region is not flat and the Alps at Fréjus are not hemispherical. Figure 8 shows the μ flux in the various set-ups we will consider. Note the two components of the μ flux, with the ν -induced muons dominating at great depths.

The μ 's themselves are not so dangerous: it is relatively easy to protect a detector against entering μ 's (see later). The problem is due to μ 's which do not penetrate the detector but send to it, for instance, an isolated neutron originating from a hadronic cascade. To give a dangerous background, the n should deposit ~ 1 GeV and no other hadron should penetrate. This has been computed by Grant for a $(10 \text{ m})^3$ detector²¹⁾. Clearly the problem occurs on the edge of the detector. The full rejection, obtained with good energy resolution and perfect detection of eventual accompanying hadrons, gives an event rate which corresponds to $\sim 2 \times 10^{-5}$ of the number of μ 's through the detector.

If you are deep enough, this background is so low that you can forget about it. If not, you have to protect the detector on its sides so as to veto such events. Here comes the important notion of fiducial volume. One can define the fiducial region F as the inner part of the detector, which satisfies the two conditions:

- i) all products of a nucleon decay occurring in F are contained and measured in the full detector;
- ii) no incoming neutral can reach F .

2.4 Neutrino background

Neutrinos have the same origin as muons, but they are not shielded. The angular distribution is therefore nearly isotropic, slightly peaked horizontally. The ν and $\bar{\nu}$ fluxes are equal. In the absence of ν oscillations (see later) there should be twice as many ν_μ as ν_e . The rates are given in Table 3²²⁾.

Considering $p \rightarrow e^+ \pi^0$, the two dangerous modes are:

$$(\nu_e, \bar{\nu}_e) N \rightarrow e \pi N, \quad \nu_e: \text{ charged current}$$

$$(\nu, \bar{\nu}) N \rightarrow (\nu, \bar{\nu}) \left(\pi \pi \right) N \quad \nu: \text{ neutral current .}$$

With $\Delta E/E = \pm 20\%$, each of these represents 10^{-2} events per year per ton. So if only an energy cut is applied, the equality of the signal from N decay and the ν background is obtained for:

$$\tau/B_{e+\pi^0} \sim 2 \times 10^{31} \text{ years .}$$

Further rejection can be obtained from topological requirements. In principle, a decay at rest and a fixed target ν interaction are different. We have seen that owing to Fermi momentum the topology of n decay is somewhat confused. In Fig. 5 we can see the domain to be considered. It has been claimed that nevertheless a gain of two orders of magnitude in background rejection can be expected (in water): this may be somewhat optimistic. Nuclear scattering and energy loss in the nucleus will cause good events to leave the domain, and should be considered as a source of loss.

Depending on the degree of optimism, we can say that the ultimate limit on $\tau/B_{e\pi}$ will be somewhere between 2×10^{32} and 2×10^{33} years (Fig. 9). To improve this, we must turn to drastic solutions, such as performing the experiment in a cave beneath the surface of the moon!

2.5 Other backgrounds

Radioactivity may be a problem at the trigger level for calorimeters but not further. Unknown backgrounds (see later, the Kolar events) could well be present, but on the other hand could introduce new physics.

3. STATUS OF N DECAY STUDIES

Figure 10 gives as a function of time the experimental limit on the N lifetime²³⁾. Most of the time the values quoted are model-dependent.

Radiochemical methods have been employed²⁴⁾. We use the fact that N decay inside the K nucleus will produce argon, which can be detected. For instance:

However, neutrinos set a limit on the sensitivity of this method to $\sim 10^{30}$ years.

Direct detection of N decay is mostly considered nowadays.

Figure 11 ²⁵⁾ shows the present detectors, in operation or planned. They are either calorimeters or Čerenkov set-ups.

The only results available come from the "small" calorimeters in the bottom left-hand corner. One sees that their fiducial mass, as previously defined, is rather modest; it is even more exact to say that this first generation does not really satisfy the criterion of the fiducial volume.

3.1 Three recent results

3.1.1 Homestake Mine (USA)²⁶⁾

The depth is 4400 mwe. The detector (Fig. 12) is made of water Čerenkov modules, piled up, with a wavelength shifter; the total mass is 180 t. Above and below, scintillation vetos are used against penetrating particles. The only signal which is looked for corresponds to the delayed sequence

$$\begin{array}{c} N \rightarrow \mu + \dots \\ \quad \downarrow \\ \quad e + \dots , \end{array}$$

where the μ can be prompt or originating from a π . The electron threshold is 10-15 MeV.

Candidates are those which satisfy the following requirements:

- i) no veto was recorded;
- ii) the event is not downward, i.e. the delayed electron signal is not geometrically recorded below the first signal;
- iii) not more than two water tanks ($2.3 \times 1.5 \times 1.5 \text{ m}^3$) are concerned: this has been shown to guarantee confinement with a good efficiency;
- iv) 50-600 MeV of energy are deposited.

The important mode, $e^+\pi^0$, does not contribute. Averaging over the contributing ones within SU(5) and putting in detection efficiencies, one predicts for water a net muon number per proton of 0.2-0.38. The signal obtained in 127 days of exposure is: ≤ 2 possible candidates. Interpreted as background, this sets a limit:

$$\tau \geq 1-2 \times 10^{30} \text{ years} .$$

3.1.2 Case-Western-Irvine (CWI)²⁷⁾

This exposure represents 67 tons per year. The detector consists of liquid scintillator and flash tubes at a depth of 9000 mwe in a South African mine.

One limit comes from μ stops^{*)}, again through the delayed pulse. The result is $\tau/B_\mu > 6 \times 10^{30}$ years with 90% CL. For SU(5) this means

$$\tau_N \geq 10^{30} \text{ years} .$$

For the Pati-Salam model, this would be $\tau_N \geq 2 \times 10^{30}$ years. Other, less restrictive limits are obtained from throughgoing particles.

This set-up is in fact used as a detector for decays which can occur in the surrounding rock; the philosophy is therefore the opposite of the one which relies on a fiducial volume.

3.1.3 The Kolar Gold Field experiment (KGF)²⁸⁾, India

Here the authors claim that they observe a positive result.

The detector is a calorimeter (Fig. 13) with 140 tons of iron plates (2 cm thick) and layers of proportional tubes ($10 \times 10 \text{ cm}^2$ of cross-section) situated at 7600 mwe. The mean density is therefore quite small, less than that of the surrounding rock.

The μ 's going through the detector demonstrate that the tubes have a good efficiency. Muons from ν interactions are observed at very large angles. A Monte Carlo simulated e of 460 MeV is shown in Fig. 14d. Note that owing to the large

^{*)} Stopping μ 's mostly come from the pions of the hadronic cascades induced by primary muons.

size of the tubes, it appears to be rectilinear with a good fraction of the tubes not being fired.

The three candidates are shown in Fig. 15. All three vertices are close to the edge. Each time, there is a track leaving (or entering), but according to the authors it does not look like a μ or a π because several tubes have not fired. Energy deposit does not contradict nucleon decay, but it does not tell us much either.

The authors claim that these three events cannot be explained by the usual interactions (Table 4) and that they demonstrate the instability of the proton with

$$\tau_p \approx 8 \times 10^{30} \text{ years} .$$

However, we are allowed to be sceptical. The events are never contained; the surrounding rock, within a couple of interaction lengths, represents much more weight than the calorimeter itself. There is in this experiment another class of unexplained events, the so-called Kolar events²⁸⁾. In a previous version of the experiment they appeared as decays in the air; but since now the detector is compact, they appear to come from dense matter. As they are defined, the deposited energy is in principle larger than m_p but only if they are correctly contained.

Without proving that the result is wrong, all these facts indicate that second-generation experiments with a large and true fiducial mass will really be welcome. Personally I will wait for their results before making up my mind about proton stability at the 10^{31} years level.

4. THE FUTURE

4.1 The very near future

The Mont Blanc detector²⁹⁾ is a 150 t fine-grain digital calorimeter: 1 cm thick iron plates, $(1 \text{ cm})^2$ streamer tubes with resistive cathodes and X-Y strip read-out (Fig. 16). A test module has been exposed to various beams at CERN³⁰⁾. The response to e^- indicates that for the $p \rightarrow e^+\pi^0$ decay, one may expect an average of $\sigma_E/E \sim 20\%$. The response to π^- gives

$\sim 85\%$ for π identification,

$\sim 10\%$ for π - μ ambiguity,

$\sim 5\%$ for π -e confusion.

The exposure to the CERN Proton Synchrotron ν beam is very instructive because the ν energy spectrum has a similar shape to that of the atmospheric neutrinos. But a test run of 3.5×10^{18} protons and 2 tons fiducial volume gives the same number of events (~ 600) as a one-year exposure of 2500 tons to atmospheric neutrinos.

A preliminary analysis of 180 events from the test reveals that several of them have a structure which resembles the Kolar candidates (Fig. 17): ν interactions inside or outside the module can simulate N decay candidates. After normalization to the Kolar ν luminosity, the claim is that one should expect, in the Indian-Japanese set-up, only 0.3 such background, whilst three candidates are observed. However, such an extrapolation from one experiment to the other looks to me to be rather tricky because of the very different structures of the detectors and their surroundings. Hopefully one will soon get results from the Mont Blanc detector (NUSEX) *in situ* and see whether candidates of the Kolar type are also observed.

A 31 t detector, Soudan 1, at 1750 mwe, made of layers of taconite concrete (taconite: a high iron-content flint-like rock, valuable as iron ore) and layers of gas proportional tubes (1" diameter) (Fig. 18), has recently been brought into operation in the Soudan Mine (Minnesota)³¹⁾. A proposal for a 1000 ton tracking calorimeter (MAO) in the Soudan mine is also available³¹⁾.

The Mont Blanc and Soudan detectors, in some sense, are both still first-generation because they have small (or null) fiducial volumes according to our previous definition. However, the granularity and compactness are much improved.

4.2 The more remote future

4.2.1 The Fréjus experiment³²⁾

This French-German experiment, recently approved and funded, will consist of a huge fine-grain calorimeter (Fig. 19). The size will be $6 \times 6 \times 17 \text{ m}^3$. The mass will come from iron plates, 3 mm thick. The trigger will be provided by 200 planes of Geiger tubes $[(15 \text{ mm})^2]$. The detection will be ensured by 800 horizontal and 800 vertical planes of flash tubes $[(5 \text{ mm})^2]$. The number of flash tubes hit is a linear estimator of energy. One should get for electrons a resolution of $15/\sqrt{E}$ and an $e-\pi$ separation better than 80%. The total mass will be 1500 tons, with 1000 tons in the fiducial volume. The hole for the calorimeter should be ready in autumn 1982, and some modules could already be installed at that time.

About the different types of tubes I referred to³³⁾, let me distinguish:

- self-triggered ones, such as proportional tubes, limited streamer, and Geiger tubes,
- triggered ones, such as flash tubes; and recall the two important parameters of such devices,
- τ_s , the memory time of the passage of the particle, typically 1 μs ,
- τ_R , the memory time of the discharge (if any), which can reach 100 ms.

For flash tubes, another important fact is that there is some probability of re-ignition during several seconds after a discharge. This, paradoxically enough, can cause a problem because of the high trigger rate due to ambient radioactivity.

4.2.2 The Čerenkov projects

First a few basic facts about Čerenkov emission. The light is emitted along a cone of opening angle $\cos \theta = 1/\beta n$. The number of photons emitted is

$$n_Y \sim 500 \sin^2 \theta \ell \text{ (cm)} .$$

An electromagnetic shower is composed of $e^\pm$ which emit Čerenkov light over a total track length ℓ , which for a given initial energy is a constant in units of radiation length. So materials with a long radiation length in centimetres will give more light. Water (36 cm) is better than glass (11 cm) and lead glass (2.5 cm).

Figure 20 ³⁴⁾ shows how the measurement of Čerenkov light (position and time of arrival) on the edge of a detector -- a tank of water, for example -- can give information on the position, the length, and the direction of the emitting track.

Suppose the detector is a cube of $\sim (20 \text{ m})^3$ and suppose a proton decays inside it to an $e + \pi^0$. The total length over which $e^\pm$ from the showers emit Čerenkov radiation is a few metres, and this gives typically 6×10^4 γ 's. Now suppose we neglect attenuation and collect 1% of these γ 's on the faces of the cube with 15% of photoelectric conversion efficiency. We get ~ 100 photoelectrons, i.e. a resolution in energy of $\sim 10\%$ just because of statistical fluctuation. Covering 1% of the 2400 m^2 of outer surface implies that we use more than 2000 5-inch diameter phototubes, which is a major investment.

The above parameters are basically the ones of the Irvine-Michigan-Brookhaven (IMB) detector, which has already been well described ³⁴⁾. Table 5 summarizes its expected properties. Water should be put in the tank quite soon.

I would now like to illustrate a project which is less known: the Japanese 1000 t Čerenkov detector ³⁵⁾. The phototubes (Fig. 21) are huge, with nevertheless a quite respectable transit time spread (7.0 ns). The expected transparency of the water is shown in Fig. 22. The claim of this team is that because of the large photocathode coverage, many features are much improved compared to those of the IMB detector, e.g. topology, lepton identification ($e-\mu$ separation), hadron identification (π, η, ρ, ω separation), energy resolution. This is illustrated by Fig. 23.

Another huge Čerenkov project ³⁶⁾ is the Harvard-Pennsylvania-Wisconsin (HPW) tank, installed at 1800 mwe, with 780 t of water, and veto systems on top of it. The photomultipliers here are regularly distributed throughout the whole volume of water. It is claimed that data will be recorded before autumn 1981. This team has also developed interesting ideas and prototypes for future large-area light intensifiers ³⁶⁾.

We see that in a couple of years from now, several of these big detectors, with fiducial volumes of 1000 tons or so, will be in operation. Furthermore, two widely different techniques will be competing.

On the other hand, we should remember that all this gigantic enterprise will bring a gain of at most two orders of magnitude in sensitivity. GUTs, as they stand now, give to believers the promise that this small step will bring about a major revolution. It would certainly be the right attitude to enthusiastically share that faith.

5. OTHER POSSIBILITIES OFFERED BY THESE SET-UPS

Here I would like to mention three of them.

5.1 The detection of GUTs monopoles

The phenomenology of such monopoles is well known³⁷⁾. Their mass is $\sim 10^{16}$ GeV or so. Estimates of the abundance of their production in the early Universe give generally unacceptably large numbers but with large uncertainties. During the cosmological evolution, they took part in galaxy formation. They are accelerated in the magnetic fields of the galaxies (typically 5 μ G on 5×10^{22} cm) and ejected from a galaxy when they reach $\beta \approx 3 \times 10^{-2} (M/10^{16} \text{ GeV})^{-1/2}$. Limits on their abundance can be obtained in various ways. The most stringent come from the energetics of ejection (they should not have carried away more energy than the energy stored in the galactic field) and considerations on the steadiness of the galactic field³⁸⁾. For monopoles of $\beta \approx 3 \times 10^{-2}$, with an anisotropic flux peaked in our galactic plane, the limit on the flux is

$$f \leq 10^{-7} \text{ cm}^{-2} \text{ y}^{-1} .$$

If one believes in such a limit, which is not a mandatory attitude, the first condition for a meaningful search is to cover an area larger than 1000 m² (10^7 cm²): none of the set-ups considered approaches that.

How could one detect them? To be ionizing they must have a velocity β larger than that of electrons in atoms, β_e . If $\beta \gg \beta_e$ the energy loss varies as $\sim \ln \beta^2$; if $\beta \sim \beta_e$ it varies as $\beta^2 \ln \beta^2$ ³⁹⁾. For the velocity considered, the

loss is substantial: 5 GeV/cm in H_2O for $\beta = 0.03$. A combination of time of flight and pulse height, at great depth since they are very penetrating (in spite of ionization, Zeeman, and ohmic losses, they can traverse the earth), should provide a fantastically clean signature. Direct detection of Zeeman effect looks desperate. What about sound waves in deep undersea experiments?

Search for such monopoles trapped in samples of matter⁴⁰⁾ may be one method, but we must try hard to convince ourselves that the monopoles did not have any good reasons for getting out of that sample at some time. For instance, a monopole trapped in a ferromagnetic meteorite is bound by an image force of $10 \text{ eV}/\text{\AA}$. If the meteorite is accelerated or decelerated above $g \sim 10$, the monopole will leave!

5.2 Relics

These are hypothetical heavy objects R of mass M_R and abundance n_R made in the early universe⁴¹⁾. We will consider long-lived ones which could still be around nowadays. They must, however, be of pretty low abundance. If we define $f = n_R/n_{\text{nucl}}$, considerations on the mass of the universe imposes $f \lesssim 10 (m_{\text{nucl}}/M_R)$. This rules out an asymmetric production, such as that of nucleons: they must have been produced in a rather symmetric way.

Proton decay set-ups are relevant because they can detect products originating from such relics decaying in the neighbourhood, for instance muons. The evaluation of the limit that can be given on their abundance for a given mass and lifetime is rather straightforward⁴²⁾. Figure 24^{42,43)} shows what the future "big experiments" will bring as an improvement. The striking fact is that a huge region is already forbidden. However, since here no theory is available to give a hint of the masses to expect, this huge region and the gain to be expected look desperately small compared to the decades of orders of magnitude where things could happen.

5.3 Neutrino oscillations

It has been claimed⁴⁴⁾ that large underground detectors provide a good way of observing ν oscillations. The source is still atmospheric ν 's. The length over which oscillations can occur goes from a few to 10 thousand kilometres (downward- and upward-going ν , respectively). Neutrino events to be measured are essentially charged-current elastic and Δ production: their threshold on the one hand and the drop of the ν spectrum with energy on the other, naturally selects a rather narrow band of ν energies with $\langle p_\nu \rangle \sim 0.5$ GeV. The mass region scanned is 10^{-1} - 10^{-3} eV. The direction of the charged leptons indicates that of the incoming ν reasonably well. Figure 25 gives an idea of the sensitivity of such a measurement.

6. n - $\bar{n}$ OSCILLATIONS

Oscillation of a neutron into an antineutron is a $\Delta B = 2$ process, quite reminiscent of well-known $\Delta S = 2$ processes⁴⁵⁾. It is relevant to the subject treated here because it is mediated by intermediate mass objects of the type discussed in Section 1.

To oscillate, n and $\bar{n}$ should be quasi-degenerate (except for B). There should be no magnetic energy term, and one has to decrease the terrestrial magnetic field more than one hundred times with an appropriate shield. There should be no nuclear field, and one has to use free neutrons.

Starting from a beam of neutrons, the probability of observing an antineutron is

$$P_{\bar{n}}(t) \approx \frac{t^2}{\tau_{n\bar{n}}^2},$$

where τ is a measure of the $\Delta B = 2$ coupling.

One has to maximize: $\Phi \equiv N \times t^2$, where N is the number of neutrons per second and t is the time of observation. For planned experiments⁴⁵⁾, using very slow neutrons from reactors, Φ can reach up to 3×10^{10} s; per day, one count

corresponds to $\tau_{nn} = 5 \times 10^7$ s. Preliminary results from the Laue-Langevin Institute²⁵⁾ give already

$$\tau_{nn} > 10^5 \text{ s} .$$

Connecting this result to nuclear instability, i.e. two-nucleon annihilation in decay experiments, seems to be strongly model-dependent, as shown in Fig. 26.

I would like to thank R. Barloutaud, B. Degrange, J. Ellis, B. Pattison and especially M. Koshiha for very useful discussions.

REFERENCES

- 1) J. Ellis, Grand unified theories, CERN preprint TH-2942 (1980), *in* Scottish Universities Summer School in Physics, eds. K.C. Bowler and D.G. Sutherland (Univ. Edinburgh, 1981), p. 201.
- 2) D.A. Ross, Phenomenology of unified gauge theories CERN preprint TH-3174 (1981) and The calculation of the decay rate of the proton, CalTech report Calt 68779/C1 (1980).
- 3) M.B. Gavela et al., Phys. Rev. D 23, 1580 (1981).
G. Kane and G. Karl, Branching ratios in baryon decay, Michigan University internal report (1980).
J.F. Donoghue, Phys. Lett. 92B, 99 (1980).
- 4) M. Machacek, Nucl. Phys. B159, 37 (1979).
A. Din et al., Phys. Lett. 91B, 77 (1980).
- 5) J.F. Donoghue, Phys. Lett. 92B, 99 (1980).
- 6) M.B. Wise et al., Three-body decays of the proton, Stanford report SLAC PUB 2614 (1980).
- 7) S. Weinberg, Phys. Rev. Lett. 43, 1566 (1979).
- 8) F. Wilczek and A. Zee, Phys. Rev. Lett. 43, 1571 (1979).
- 9) H. Georgi et al., μ^+ polarization in proton decay: a probe of flavour mixing in unified models, Harvard preprint HUTP 80/A064 (1980).
- 10) J.C. Pati and A. Salam, Quark lepton unification and proton decay, *in* Proc. First Workshop on Grand Unification, Durham, New Hampshire, 1980, eds. P.H. Frampton, S.L. Glashow and A. Yildiz (Math Sci Press, Brookline, 1980), p. 115.
- 11) S. Weinberg, Expectations for baryon and lepton non-conservation, same Proc. as Ref. 10, p. 347.
- 12) R.N. Mohapatra, Baryon and lepton non-conservation, Majorana neutrinos and neutron oscillations, same Proc. as Ref. 10, p. 69.

- 13) See for instance: W.B. Rolnick, Scalar operators in nucleon decay, Wayne State Univ. preprint (Detroit) (1981).
- 14) L.W. Jones, Proton lifetime, quark abundance and age of the universe: a numerical coincidence? Univ. Michigan preprint UMHE 81-27 (1981).
- 15) W. Fairbank, Talk given at the EPS Conf., Lisbon, 1981.
- 16) H.B. Nielsen and M. Ninomiya, Nucl. Phys. B141, 153 (1978).
See also J. Ellis et al., Nucl. Phys. B176, 61 (1980).
- 17) D.H. Perkins, Feasibility test for experiment on nucleon stability, Referee's report, CERN/EP/DP/mk (August 1979).
- 18) D.A. Sparrow, Phys. Rev. Lett. 44, 625 (1980).
- 19) B. Rossi, High-energy particles (Prentice Hall Inc., Englewood Cliffs, NJ, 1952), p. 163.
- 20) D.H. Perkins, μ induced backgrounds in the Mont Blanc and Fréjus tunnels, CERN/EP/DPMk (Sept. 1979).
- 21) A.L. Grant, Isolated neutron background in the Mont Blanc and Fréjus tunnels, CERN EF/ALG/ed (Dec. 1979).
- 22) E. Bellotti, Investigations of nucleon stability in Europe, *in* Unification of the fundamental particle interactions, eds. S. Ferrara, J. Ellis and P. Van Nieuwenhuizen (Plenum, New York, 1981), p. 673.
- 23) M. Goldhaber et al., Is the proton stable? Los Alamos report LA-UR/80-356 (1980).
- 24) L. Bergamasco and G. Cini, Nuovo Cimento 1C, 293 (1978).
- 25) R. Barloutaud, Talk given at the EPS Conf., Lisbon, 1981.
- 26) M. Deakyne et al., The Homestake mine nucleon decay search, paper 716 submitted to the Madison Conf., 1980.
- 27) J. Learned et al., Phys. Rev. Lett. 43, 907 (1979).

- 28) B. Sreekantan, Communication to the EPS Conf., Lisbon, 1981.
For the so-called KGF events, see: S. Miyake, Proc. 19th Int. Conf. on High-Energy Physics, Tokyo, 1978 (Phys. Soc. Japan, Tokyo, 1979), p. 433.
- 29) G. Battistoni et al., Frascati-Milan-Turin Collab., NUSEX Proposal.
See also Ref. 22.
- 30) E. Iarocci, Communication to the EPS Conf., Lisbon, 1981.
- 31) H. Courant et al., A dense detector for baryon decay, Paper 0350 submitted to the Madison Conf., 1980.
E.A. Peterson, Communication to the EPS Conf., Lisbon, 1981.
Minnesota-Argonne-Oxford Collab., Soudan 2, a 1000 ton tracking calorimeter for nucleon decay, Proposal OUNP/42/81, COO-1764-410, ANL-HEP-PR 81-12 (1981).
- 32) P. Bareyre et al., Orsay-Palaiseau-Paris-Saclay Collab. proposal, Proposition d'une expérience pour l'étude de l'instabilité du nucléon au moyen d'un détecteur calorimétrique.
- 33) For flash tubes, see:
M. Conversi and L. Federici, Nucl. Instrum. Methods 151, 93 (1978).
For streamer tubes:
G. Battistoni et al., Nucl. Instrum. Methods 164, 57 (1979).
- 34) L.R. Sulak, The Irvine-Michigan-Brookhaven nucleon decay search *in* same book as Ref. 22, p. 639.
J.C. Van der Velde, A review of proton decay experiments, Talk given at the SLAC Summer Institute on Particle Physics, July-Aug. 1980.
- 35) M. Koshiha, private communication.
- 36) D.R. Winn, A search for baryon decay: plans for the HPW water Čerenkov detector, in same Proc. as Ref. 10, p. 189.
- 37) G. Lazarides et al., Phys. Lett. 100B, 21 (1981).
M. Longo, Massive magnetic monopoles, Univ. Michigan preprint UMHE 81-33 (1981).
- 38) E.N. Parker, The origin of magnetic field, Astrophys. J. 160, 383 (1970).

- 39) K. Hayashi, Stopping power formula for a cosmological monopole, Kinki Univ. preprint.
S.P. Ahlen, Phys. Rev. D 17, 229 (1978).
- 40) P. Eberhard et al., Phys. Rev. D 4, 3260 (1971).
- 41) C.B. Dover et al., Phys. Rev. Lett. 42, 1117 (1979).
- 42) P.H. Frampton and S.L. Glashow, Phys. Rev. Lett. 44, 1481 (1980).
- 43) J. Ellis et al., Exotic relics from the big bang, CERN preprint TH-2893 (1981).
- 44) B. Cortez and L. Sulak, A long baseline ν oscillation experiment sensitive to mass difference of hundredths of an eV, Univ. Michigan preprint UMPD 80-1 (1980).
- 45) R. Wilson, Possibilities of experiments to measure neutron-antineutron mixing, same Proc. as Ref. 10, p. 215.

Table 1

Physical phenomenon	Effect on M_x
1) Correct treatment of the fermion mass threshold	$\times 2$
2) α_{em} at M_W^2	$\times 1/10$
3) Correct treatment of gauge boson mass thresholds (including the right choice of gauge)	$\left\{ \begin{array}{l} \times 2 \\ \times 1/1.7 \end{array} \right.$
4) Two-loop contributions	$\times 1/4$
5) Effect of the Higgs sector	$\times 1/1.7$
6) x, y are felt below their mass	$\times 1/2$
Total	$1/58$

Table 2a

Antilepton + meson branching ratios (from Ref. 32)

Mode		SU(6)					Bag model	
		a)	b)	Static model c)	Recoil model c)	Relativistic model c)	d)	e)
$e^+\pi^0$	p	0.33	0.37	0.36	0.40	0.38	0.08	0.31
	n	0.58	0.74	0.79	0.72	0.68	0.22	0.45
$e^+\rho^0$	p	0.17	0.02	0.02	0.07	0.10	0.18	0.21
	n	0.30	0.04	0.05	0.12	0.18	0.49	0.30
$e^+\eta$	p	0.12	0.07	0.07	0.01	~ 0	0.03	0.05
$\bar{\nu}_e\eta$	n	0.02	0.01	0.01	~ 0	~ 0	0.01	0.03
$e^+\omega$	p	0.21	0.18	0.21	0.25	0.26	0.47	0.19
$\nu_e\omega$	n	0.04	0.04	0.05	0.05	0.05	0.13	0.15
$\bar{\nu}_e\pi^0$	p	0.09	0.15	0.14	0.16	0.15	0.03	0.11
	n	0.04	0.07	0.08	0.07	0.07	0.02	0.04
$\bar{\nu}_e\rho^0$	p	0.04	0.01	0.09	0.03	0.04	0.07	0.08
	n	0.02	0.005	0.01	0.01	0.02	0.05	0.03
μ^+K^0	p	0.005	0.19	0.18	0.08	0.05	0.14	0.005
$\bar{\nu}_\mu K^0$	n	0.004	0.10	0.01	0.03	0.01	0.10	0.002
τ_n/τ_p		0.88	1.0	1.1	0.9	0.9	1.35	0.75

- a) M. Machacek, Ref. 4.
- b) M.B. Gavela et al., Ref. 3.
- c) G. Kane and G. Karl, Ref. 3.
- d) J.F. Donoghue, Ref. 5.
- e) A. Din et al., Ref. 4.

Table 2b

Current theoretical predictions for proton decay

$\langle \tau_B \rangle$ years	Lifetime range	Decay modes	Author and model
5×10^{30}	$10^{30 \pm 2}$	$e^+ + (\pi, \omega, \rho, \eta)$	Ellis, Gaillard, Rudaz, Nanopoulos [SU(5)]
10^{31}	$10^{31 \pm 2}$	μ decay suppressed	Wilczek, SU(5), SO(10)
10^{32}	-		Georgi, Quinn, Weinberg
$\sim 10^{31}$	$10^{31 \pm 2}$	e^+ decays	Goldmann, Ross, Marciano [SU(5)]
8×10^{30}	-	MIT bag Mode 2 + e^+ dec.	Donoghue [SU(5)]
-	-	Do not forget muonic decays	Weinberg, Glashow
10^{30}	$10^{29} - 10^{32}$	$3 \nu \pi, \nu \pi$	

Table 3

Neutrino events/year/100 t

Visible energy GeV	Charged current $\nu_\mu, \bar{\nu}_\mu$	Charged current $\nu_e, \bar{\nu}_e$	Neutral current	Total
0.4-0.6	2.06	0.94	1.02	3.96
0.6-0.8	1.34	0.64	0.51	2.49
0.8-1.0	1.13	0.47	0.31	1.91
1.0-1.2	0.89	0.38	0.23	1.51
1.2-1.4	0.74	0.31	0.10	1.22

Table 4

Basic data: Detector volume : 6 m × 4 m × 3.7 m
Total weight : 140 tons (iron)
Live-time : 205 days
Total No. of events : 388

Category of events	Observed no.	Estimated no.
1. Single muons penetrating the detector ($\theta = 0-55^\circ$)	360	350 ± 50
2. Side showers	2	-
3. Parallel muons ($E_\mu \sim 15$ TeV and multiplicity ≥ 2)	2	-
4. Stopping muons	2 *	-
5. ν -interactions in rock $\theta = 0-55^\circ$	Grouped under category (1)	~ 8
$\theta = 56-90^\circ$	13	13.2 ± 2.5
6. ν -interactions inside the detector	3	6 ± 2
7. Kolar events (Multiprong-wide angle-air vertex events)	1(?)	(?)
8. Unusual events (Nucleon decay candidates)	3	?

* A fraction of these events would be ν -induced muons or pions that stop inside the detector.

Table 5

Summary of the IMB Detector Properties

Size:

Total volume = 6.5×10^{33} nucleons

Fiducial volume = $2.5 \times 10^{33} = (16 \text{ m})^3$

Energy resolution:

500 mev shower $\sigma = 11\%$

300-450 mev/c $\pi^\pm, \mu^\pm$ $\sigma = 10-15\%$

Vertex localization

$\sigma = 0.5 \text{ m}$

Track directionality

No ambiguity

Angular resolution

Showers ($e^\pm, \pi^0$) $\sigma = 10-20^\circ$

Charged tracks $\sigma \lesssim 5^\circ$

$\mu \rightarrow e$ detection

$\gtrsim 80\%$ after trigger (event thresh. $\approx 75 \text{ mev}$)

Several $\mu \rightarrow e$: OK

Position correlation: 1.5 m

e direction: $\sim 25^\circ$

a)

$$A_\nu = \begin{pmatrix} v + \frac{\sqrt{2}}{\sqrt{15}} u & v & v & x^r & y^r \\ v & v + \frac{\sqrt{2}}{\sqrt{15}} u & v & x^w & y^w \\ v & v & v + \frac{\sqrt{2}}{\sqrt{15}} u & x^b & y^b \\ \bar{x}^r & \bar{x}^w & \bar{x}^b & \frac{w_0}{\sqrt{2}} - \frac{\sqrt{3}}{\sqrt{10}} u & w^+ \\ \bar{y}^r & \bar{y}^w & \bar{y}^b & w^- & -\frac{w_0}{\sqrt{2}} - \frac{\sqrt{3}}{\sqrt{10}} u \end{pmatrix}$$

b)

$$\psi_5 = \begin{pmatrix} d^r \\ d^w \\ d^b \\ e^+ \\ \bar{\nu} \end{pmatrix}_R \quad \psi_{10} = \begin{pmatrix} 0 & u_c^b & -u_c^w & -u^r & -d^r \\ -u_c^b & 0 & u_c^r & -u^w & -d^w \\ u_c^w & -u_c^r & 0 & -u^b & -d^b \\ u^r & u^w & u^b & 0 & e^- \\ d^r & d^w & d^b & -e^- & 0 \end{pmatrix}_L$$

c)

Fig. 1

a)

b)

Fig. 2

Fig. 3

μ^+ polarization from proton decay for final state in which pions are not distinguished from kaons in the final state. The horizontal scale is linear in $(m_A/m_B)^{1/2}$. The crosshatched (dotted) band corresponds to F(J) flavor mixing schemes in $O(10)$, where the bounds on each band correspond to various approximation schemes for hadronic matrix elements.

Fig. 4

Fig. 5

Fig. 6

Fig. 7

a)

b)

Fig. 8

Fig. 9

Fig. 10

Fig. 11

Fig. 12

Fig. 13

Shower inside detector

Simulated tracks

Fig. 14

X : NOT WORKING

Fig. 15

Fig. 16

Fig. 17

Schematic of Detector Concept

Fig. 18

Fig. 19

Illustration of how the arrival time of various parts of a Cerenkov cone can determine track direction in the IMB detector. Similar techniques can be used in the volume arrayed HPW detector.

Fig. 20

Scale: 1/3

PIN CONNECTONS

pin #	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18	19	20	socket #
	D	D	D	D	D	D	D	D	D	D	D	D	D	D	D	D	D	D	D	D	
	1	3	5	7	10	9	11	13	A	12	10	8	6	4	2	C	K	G	I	E	E678-20A

R1449 x (20"PMT)

浜松テレビ社 (株)

Sept. 10. 80 Kenji Sugita

Fig. 21

図3-2 20"φ光電管断面図 (鉄板比1:3)

Fig. 22

Fig. 23

Fig. 24

Fig. 25

Fig. 26

58