CERN LIBRARIES, GENEVA

CM-P00063908

MEASUREMENT OF THE $2S_{1/2}$ - $2P_{1/2}$ SPLITTING IN THE $(\mu^{-4}\text{He})^{+}$ MUONIC ION

G. Carboni^{*)}, G. Gorini^{*)}, E. Iacopini^{**)}, L. Palfy^{***)},
F. Palmonari^{**)}, G. Torelli^{*)} and E. Zavattini^{**)}

CERN-Pisa University Collaboration

ABSTRACT

The measurement of the $2S_{1/2} \rightarrow 2P_{1/2}$ energy transition in muonic helium is presented. The energy difference S^1 is found to be $S_{\text{exp}}^1 = 1381.3 \pm 0.5$ meV. This result agrees with the expected value $\tilde{S}^1 = 1381.2 \pm 0.3$ meV obtained assuming the previously measured value for the $2S_{1/2} \rightarrow 2P_{3/2}$ energy difference.

Geneva - 23 November 1977

^{*)} Istituto di Fisica dell'Università and INFN, Pisa, Italy.

^{**)} CERN, Geneva, Switzerland.

^{***)} CERN, Geneva, Switzerland and IISN, Belgium. Present address: Université Catholique de Louvain, Louvain la Neuve, Belgium.

Spectroscopy of the $n = 2$ level in the muonic ion $(\mu^{-4}\text{He})^{+}$ offers a powerful tool to test QED predictions on vacuum polarization, since the 2S-2P level splittings S^1 and S^2 (see Fig. 1) are almost completely due to this effect [1]. The level splitting S^2 has been previously measured by Carboni et al. [2], who found a value which is in good agreement with the latest calculations by Rinker [3] and Borie [4]. To complete the spectroscopy of this $n = 2$ level, we measured the level splitting S^1 between the $2S_{1/2}$ and the $2P_{1/2}$ levels in the $(\mu^{-4}\text{He})^{+}$ system using the same apparatus developed at CERN [5] and used to measure S^2 .

The latest experimental result for S^2 is [2]:

$$S_{\text{exp}}^2 = 1527.5 \pm 0.5 \text{ meV} \quad (1)$$

corresponding to

$$\lambda_{\text{exp}}^2 = 8116.8 \pm 1.5 \text{ \AA} \quad (1')$$

If the assignment for S^2 is correct, Eq. (1) can be used to predict accurately the value expected for S^1 since the difference between S^1 and S^2 does not suffer from the uncertainty due to the correction for the finite size of the helium nucleus. The expected value for S^1 , using Eq. (1) and the results of the theoretical calculations [3, 4] is:

$$\tilde{S}^1 = 1381.2 \pm 0.3 \text{ meV} \quad (2)$$

or

$$\tilde{\lambda}^1 = 8977 \pm 2 \text{ \AA} \quad (2')$$

The search for the $2S_{1/2} \rightarrow 2P_{1/2}$ transition was therefore performed exploring a region of about 40 \AA around value (2').

The experimental apparatus, mounted in the muon beam of the CERN Synchrocyclotron, is schematically shown in Fig. 2. Since a detailed description has been given elsewhere [2, 5], here we recall only the main lines of the experiment. Negative muons are stopped in a high-pressure (40 atm) helium target, where they

can form $(\mu^{-4}\text{He})^{+}$ ions in the 2S state. From previous measurements [6] we know that about 4% of the stopped muons actually land in the 2S state and that in our experimental conditions this metastable state has a lifetime of about 1.6 μsec . 500 nsec after the muon stopping signal, a 20 nsec long light pulse of wavelength λ , generated by a tunable pulsed dye-laser, is injected into the target to excite the $2S_{1/2} \rightarrow 2P_{1/2}$ transition. The transition is identified by detecting in coincidence with the laser pulse the 8.2 keV X-ray emitted in the $2P \rightarrow 1S$ decay (2P lifetime is 5×10^{-13} sec). By varying the laser wavelength λ we look then for the wavelength $\bar{\lambda}_1$ that gives the maximum yield of X-rays in coincidence with the laser pulse. After each pulse the laser wavelength is changed around the value ($2'$) in steps of 2.2 Å by tilting the diffraction grating in the laser cavity by means of a stepping motor (see Fig. 2). Wavelength read-out for each stepping motor position (SMP) is achieved using an on-line digitized spectrometer previously calibrated with the 8943.5 Å line from Cs and the 8952.2 Å line from Xe. Table 1 gives the main characteristics of the laser we used. Data taking and analysis are similar to those adopted to obtain result (1) and described in Ref. 2.

With respect to the previous measurement [2] of S^2 , in the measurement of S^1 there are the following drawbacks:

- i) the $2S_{1/2} \rightarrow 2P_{1/2}$ transition is two times less intense than the $2S_{1/2} \rightarrow 2P_{3/2}$ one owing to the different statistical weights for the two 2P states;
- ii) the dye-laser efficiency drops when increasing the wavelength (about 170 mJ average energy per pulse around λ_1 against over 270 mJ average energy per pulse at λ_2).

In Fig. 3 the results of our measurement are presented. The number of 8.2 keV X-rays detected in a 250 nsec time interval after the laser pulse is plotted against the SMP, showing a peak near the centre of the explored interval of λ . A best fit to the data with a Lorentzian line shape of width $\Gamma = 8$ Å FWHM plus a constant background gives

$$\lambda_{\text{exp}}^1 = 8976 \pm 3 \text{ \AA} \quad (3)$$

corresponding to

$$S_{\text{exp}}^1 = 1381.3 \pm 0.5 \text{ meV} , \quad (3')$$

with a $\chi^2 = 3$ for 7 degrees of freedom. The number of events under the peak $N_{\text{ev}} = 81 \pm 27$ is consistent with the one obtained for the line S^2 , taking into account the above-mentioned factors disfavoured the observation of the $2S_{1/2} \rightarrow 2P_{1/2}$ transition. The good agreement between values (3) and (2') shows that the transition observed by Carboni et al. [2] was indeed S^2 . Therefore the conclusions drawn on the validity of QED as far as concerns vacuum polarization are upheld.

REFERENCES

- [1] L.L. Foldy and E. Eriksen, Phys. Rev. 95 (1954) 1048.
E. Campani, Nuovo Cimento Letters 4 (1970) 982.
- [2] G. Carboni et al., Nuclear Phys. A278 (1977) 381.
- [3] G.A. Rinker, Phys. Rev. A14 (1976) 18.
- [4] E. Borie, Karlsruhe University Preprint, TKP 77-13 (1977).
- [5] A. Bertin et al., Nuovo Cimento 23B (1974) 489.
- [6] A. Bertin et al., Nuovo Cimento 26B (1975) 433.
- [7] J.P. Webb, F.G. Webster and E.B. Plourde, IEEE J. Quantum Electronics QE11
(1975) 114.

Table 1

Laser characteristics

Dye	: Kodak IR-140 ^{a)}
Solvent	: Dimethylsulfoxide
Molar concentration	: 7×10^{-4} M
Repetition rate	: 0.25 Hz
Mean infra-red energy per pulse	: ~ 170 mJ

a) Webb et al. [7].

Figure captions

- Fig. 1 : Scheme of the lowest energy levels of the $(\mu^{-4}\text{He})^{+}$ muonic ion.
- Fig. 2 : Schematic view of the apparatus. M = CH_2 moderators; 1,2,3,4 = plastic scintillators; 2 = anticoincidence Čerenkov counter (lucite); T = invar vessels; V = useful muon stopping volume; A_1 - A_8 = NaI(Tl) counters; DC = dye cell; DM = dielectric mirror ($R = 30\%$); DG = diffraction grating (1200 lines/mm); SM = stepping motor driving DG; D_1 and D_2 = photodiodes; C = internally gold-coated conical light-guide; W = antireflection window; S = light-beam splitters; TC = optical telescope.
- Fig. 3 : The $2S_{1/2}$ - $2P_{1/2}$ resonance signal. Abscissae: stepping motor position. Ordinates: the number of 8.2 keV X-rays detected in coincidence with the laser pulse. The full line drawn through the data is the result of a fit with a Lorentzian line shape ($\Gamma = 8 \text{ \AA}$) plus a constant background. The central wavelength for the Lorentzian line is $\lambda_1 = 8976 \pm 3 \text{ \AA}$.

Fig. 1

Fig. 3