Ref.TH.2752-CERN

PHOTOPRODUCTION OF CHARM AND BEAUTY

TU Tung-Sheng ^{*)}
CERN - Geneva

A B S T R A C T

The photoproduction of Charm and beauty are re-examined within QCD. In addition to the photon-gluon amalgamation mechanism, $\bar{q}q \rightarrow b\bar{b}$ or $c\bar{c}$ is also examined. It turns out that for beauty production at SPS photon beams, the $\bar{q}q \rightarrow b\bar{b}$ mechanism is dominant over γ -g amalgamation for Buras-Gaemers gluon distributions. The possibility of experimental measurement is briefly discussed.

^{*)} On leave of absence from the Institute of High Energy Physics, Peking, China.

Recently a bump at $M \sim 5.3$ GeV has been observed in the $\psi K\pi$ channel in the π -p collision experiment at CERN¹⁾.

The result might not be conclusive, but it is promising. There is good reason to consider this bump as a bottomed meson²⁾ ($b\bar{u}$, $\bar{b}u$, $b\bar{d}$, $\bar{b}d$, ...). Interest in searching for bottomed mesons in other processes is increasing. So far there is only little evidence for the photoproduction of charmed mesons³⁾ and charmed baryons⁴⁾. So, searching for charmed as well as bottomed mesons in photoproduction processes is still interesting. The photoproduction of heavy quark flavours has been discussed within QCD by several authors⁵⁻⁷⁾, but they only concentrate on the photon-gluon amalgamation mechanism and on diffractive production.

In this note we shall discuss another mechanism, namely $\bar{q}q \rightarrow b\bar{b}$ or $c\bar{c}$ [where the $\bar{q}$ comes from the hadronized incoming photon (see Fig. 2)], and compare it with the previous one.

What is the modern concept of a photon? Within QCD, a photon has two components: a point-like component and a hadronic component⁸⁾. For the hadronic component, in large momentum transfer processes (large p_T is not necessary in heavy-flavour production) the incoming photon firstly disassociates into partons and then interacts with the target. In that case perturbative QCD is applicable. In some cases, especially in low momentum transfer processes, the incoming photon couples directly to a bound state of quarks and gluons, for instance, ρ , ω , ϕ , ψ , T , ..., etc. This is a non-perturbative phenomenon and can be described approximately by the vector meson dominance. This is illustrated in Fig. 1. Obviously there is some overlap between the parton picture and the vector meson dominance picture, and the reason is the following. Even in a large momentum transfer process (i.e. where there is a large momentum flow along the quark lines inside the photon) the diagram (b) in Fig. 1 still can include bound states of quarks and gluons, although the probability of bound states might be small. So the summation of the contributions of (b) and (c) in Fig. 1 has some fraction of double counting. At large p_T , the contribution of the diagram (c) in Fig. 1 will fall off very

quickly. This means that the vector meson dominance part contributes mainly to the diffractive production process (near the forward direction).

In the following we shall calculate all the contributions separately.

1. THE PHOTOPRODUCTION OF BOTTOMED MESON

The two mechanisms considered are illustrated in Fig. 2. Although the $\bar{q}q \rightarrow b\bar{b}$ mechanism is proportional to $\alpha\alpha_s^2$ instead of $\alpha\alpha_s$ like γ -g amalgamation, m_b is quite large. The lower bound for the Feynman variable $x_{\min} \sim 4m_b^2/s$. When $E_\gamma^L \sim 150$ GeV (SPS photon beam), $x \geq 0.31$. In that case the gluon distribution is seriously suppressed, whilst the valence quark distribution is quite large. Therefore for SPS photon beams the $\bar{q}q \rightarrow b\bar{b}$ mechanism may be important. Let us compare it with γ -g amalgamation.

The contribution of γ -g amalgamation [diagram (9) in Fig. 2] is^{5,7,9)}

$$\sigma(\gamma p \rightarrow B\bar{B}x) = \int_{4m_b^2/s}^1 dx \hat{\sigma}(xs) G(x, Q^2) \quad (1)$$

where

$$\hat{\sigma}(xs) = \frac{\pi\alpha\alpha_s e_b^2}{xs} \left\{ (3-\beta^4) \ln \frac{1+\beta}{1-\beta} - 2\beta(2-\beta^2) \right\}$$

$$\beta = (1 - 4m_b^2/xs)^{\frac{1}{2}},$$

$e_b = -1/3$, $G(x, Q^2)$ is the gluon distribution inside the proton.

S is the total energy squared in the γp c.m. system.

With the constraint $\int_0^1 xG(x, Q^2) dx \sim 0.5$, many authors take

$$G(x, Q^2) \sim \frac{n+1}{2} (1-x)^n / x \quad (2)$$

According to QCD, the value of n should depend on the momentum transfer Q^2 . For $B\bar{B}$ production it is reasonable to take $Q^2 \sim 4m_b^2 \sim 100$ GeV² for the Buras and Gaemers parametrization¹⁰⁾.

$$\begin{aligned} \text{If } G(x, Q_0^2) \sim 3(1-x)^5/x & \quad \text{then } G(x, Q^2) \sim 8.5(1-x)^{16}/x \\ \text{If } G(x, Q_0^2) \sim 5.5(1-x)^{10}/x & \quad \text{then } G(x, Q^2) \sim 11.5(1-x)^{22}/x \end{aligned} \quad (3)$$

where $Q_0^2 \sim 1.8 \text{ GeV}^2$, $Q^2 \sim 100 \text{ GeV}^2$, and we take $\Lambda \sim 0.5 \text{ GeV}$ instead of 0.3 GeV in Ref. 10. The shape of $G(x, Q^2)$ is an open question. But the powers in Eq. (3) might be reasonable.

In Ref. 7, only the case $n = 5$ is considered; and in Ref. 6 both $n = 5, 9$ are calculated but with $S \rightarrow \infty$. Hence, for comparison, we give all the cases $n = 5, 7, 10, 16, 22$, both for SPS photon beams and for e-p colliding beams*).

The total cross-section of the $q\bar{q} \rightarrow b\bar{b}$ diagram (b) in Fig. 2 is

$$\sigma(\gamma p \rightarrow B\bar{B}x) \sim \int \int_{xyS \geq 4m_b^2} dx dy \hat{\sigma}(xyS) \sum_i \left(\bar{q}_i^{\gamma}(y, Q^2) q_i^p(x, Q^2) + q_i^{\gamma}(y, Q^2) \bar{q}_i^p(x, Q^2) \right) \quad (4)$$

where

$$\hat{\sigma}(xyS) = \frac{8\pi\alpha_s^2}{27(xyS)^3} [xyS(xyS - 4m_b^2)]^{1/2} (xyS + 2m_b^2) \quad (5)$$

$$\sum_i \left(\bar{q}_i^{\gamma}(y, Q^2) q_i^p(x, Q^2) + q_i^{\gamma}(y, Q^2) \bar{q}_i^p(x, Q^2) \right) = \frac{1}{xy} y q_{2/3}^{\gamma}(y, Q^2) \left(x u_v^p(x, Q^2) + \frac{1}{4} x d_v^p(x, Q^2) + \frac{1}{2} x S^p(x, Q^2) \right) \quad (6)$$

where $q_{2/3}^{\gamma}(y, Q^2)$ is the $2/3e$ charged quark distribution function of the photon, and u_v^p , d_v^p , S^p are valence up, down, and sea quark distribution functions in the proton, respectively.

For the numerical calculations, we use

$$\alpha_s(4m_b^2) \sim 0.3 \quad (7)$$

$$\alpha_s(Q^2) \sim 0.3 / \left[1 + 0.3 \frac{7}{4\pi} \ln(Q^2/4m_b^2) \right] \quad (8)$$

*) A proposed e-p colliding machine will have quite intense beams of nearly real photons around 5-10 GeV, Ref. 11.

For γ -g amalgamation calculations we simply take $\alpha_s(Q^2) \sim 0.3$ for SPS photon beams, and $\alpha_s(Q^2)$ in Eq. (8) for e-p colliding beams with $Q^2 \sim xS$.

For the $q\bar{q} \rightarrow b\bar{b}$ calculation, we use $\alpha_s(Q^2)$ of Eq. (8) with $Q^2 \sim xyS$ *).

We use $q_{2/3}^\gamma(y, Q^2)$ given by Llewellyn Smith⁸⁾ and neglect the gluon distribution of the photon. This is reasonable for SPS photon beams because of the constraint $xyS \geq 4m_b^2$, e.g. y cannot be smaller than $4m_b^2/S$. For e-p colliding beams the lower bound of x and y is very small ($\ll 0.1$). Unfortunately the quark distribution $q_{2/3}^\gamma(y, Q^2)$ of the photon is not reliable when $y < 0.1$. So we have to make a cut $y > 0.1$ and neglect the gluon contribution of the photon. Experimentally this cut is not so bad. To see this, let us look at the kinematics of Fig. 2b¹²⁾ in the γp c.m. system (see Fig. 3). There the area for $y \leq 0.1$ is illustrated by a dashed line and $y_T = 2k'_T/\sqrt{s} = 2p'_T/\sqrt{s}$, $y_L = 2k'_L/\sqrt{s}$, $y_j = 2p'_L/\sqrt{s}$, $s = 10800 \text{ GeV}^2$. The straight line $\theta^{\text{lab}} = 170^\circ$ is also calculated at $s = 10800 \text{ GeV}^2$. From Fig. 3 we can see that the cut $y > 0.1$ only cuts down the contribution of $\theta \leq 10^\circ$ in the backward direction in the lab. system. This is quite reasonable for colliding beams owing to geometrical restrictions. For the proton structure functions, we use the Buras-Gaemers parametrization¹⁰⁾.

All the results are given in Figs. 4 and 5.

From Fig. 4 we can see that at SPS photon beams the $q\bar{q} \rightarrow b\bar{b}$ contribution is much higher than γ -g amalgamation for Buras-Gaemers gluon distributions ($n=16, 22$). The largest value of $\sigma(\gamma p \rightarrow B\bar{B}X)$ is $\sim 1 \text{ nb}$ at $E_\gamma^L \sim 200 \text{ GeV}$ and $n = 5$ [$G(x, Q^2) \sim 3(1-x)5/x$]. But $n = 5$ is not the favourite gluon distribution at $Q^2 \sim 100 \text{ GeV}^2$. So, the value 1 nb is already overestimated.

At e-p colliding beams, the contribution of γ -g amalgamation is $\sim 20\text{--}40 \text{ nb}$ and is not sensitive to the shape of $G(x, Q^2)$, while the contribution of $q\bar{q} \rightarrow b\bar{b}$ is only about $1\text{--}2 \text{ nb}$. Thus the $q\bar{q} \rightarrow b\bar{b}$ mechanism is not important at very high energy. The reason is that at very high energy the small x region will be important.

*) There are some other choices; for instance, $Q^2 \sim \sqrt[3]{sf\bar{u}}$, $-\hat{t}$, etc.

Now we will say a few words about diffractive production.

Following Refs. 6 and 7,

$$\frac{\sigma(\gamma p \rightarrow B \bar{B} x)}{\sigma(\gamma p \rightarrow D \bar{D} x)} \sim \frac{M_\psi}{M_T} \cdot \frac{\Gamma_{T \rightarrow e^+ e^-}}{\Gamma_{\psi \rightarrow e^+ e^-}} \cdot \frac{\sigma(TN)}{\sigma(\psi N)} \quad (9)$$

If we assume¹³⁾

$$\sigma(TN)/\sigma(\psi N) \sim M_\psi^2/M_T^2 \quad (10)$$

then

$$\sigma(\gamma p \rightarrow B \bar{B} x)/\sigma(\gamma p \rightarrow D \bar{D} x) \sim 9 \times 10^{-3} \quad (11)$$

Using

$$\lim_{S \rightarrow \infty} \sigma(\gamma p \rightarrow D \bar{D} x) \sim 1.2 \mu b \quad (\text{Ref. 6}) \quad (12)$$

we get

$$\sigma(\gamma p \rightarrow B \bar{B} x) \sim 10 \text{ nb} \quad (13)$$

This is more than a factor of 10 larger than the QCD perturbative calculation.

But the perturbative QCD calculations are probably more reliable when heavy flavours are produced. In Eq. (10) the energy dependence is not specified. Also the ratio might be m_ψ^n/m_T^n , with $n \geq 3$ or $e^{-M_T^2}/e^{-M_\psi^2}$, etc. So the estimation (13) is not reliable. Therefore we can expect

$$\sigma_{\text{tot}}(\gamma p \rightarrow B \bar{B} x) \sim \text{a few nb}.$$

2. THE PHOTOPRODUCTION OF CHARM

The main difference between charm production and beauty is the lighter charm quark mass $m_c \sim 1.5 \text{ GeV}$. So, $Q^2 \sim 4m_c^2 \sim 9 \text{ GeV}^2$.

For Buras-Gaemers gluon distributions,

$$\begin{aligned} \text{if } G(x, Q_0^2) &\sim 3(1-x)^5/x & \text{then } G(x, Q^2) &\sim 5.5(1-x)^{10}/x \\ \text{if } G(x, Q_0^2) &\sim 5.5(1-x)^{10}/x & \text{then } G(x, Q^2) &\sim 8.5(1-x)^{16}/x \end{aligned} \quad (14)$$

where

$$Q_0^2 \sim 1.8 \text{ GeV}^2, Q^2 \sim 9 \text{ GeV}^2; \text{ also } \Lambda \sim 0.5 \text{ GeV}.$$

All the formulae are valid except for a few changes: $m_b \rightarrow m_c$, $e_b \rightarrow e_c = \frac{2}{3}$.

We take

$$\alpha_s(4m_c^2) \sim 0.4 \quad (15)$$

$$\alpha_s(Q^2) \sim 0.4 / \left[1 + 0.4 \frac{7}{4\pi} \ln(Q^2/4m_c^2) \right] \quad (16)$$

For γ -g amalgamation we use simply $\alpha_s(Q^2) \sim 0.4$ for SPS photon beams, and $\alpha_s(Q^2)$ of Eq. (16) for e-p colliding with $Q^2 \sim xS$. Also for $q\bar{q} \rightarrow c\bar{c}$ we use $\alpha_s(Q^2)$ of Eq. (8) with $Q^2 \sim xyS$.

The results are presented in Figs. 6 and 7.

From Fig. 6 we can see that at SPS photon beams the $q\bar{q} \rightarrow c\bar{c}$ mechanism is only important when $E_Y^L < 20 \text{ GeV}$. When $E_Y^2 > 100 \text{ GeV}$, $\sigma_{\text{tot}}(\gamma p \rightarrow D\bar{D}X) \sim 1\text{-}2 \text{ } \mu\text{b}$ for different gluon distributions.

For e-p colliding beams, from Fig. 7 we can see that the contribution of γ -g amalgamation is $\sim 1\text{-}4 \text{ } \mu\text{b}$, nearly the same as in the case of the SPS photon beam, while the contribution $q\bar{q} \rightarrow c\bar{c}$ reaches an asymptotic value of $\sim 10 \text{ nb}$.

3. DISCUSSION

We believe that the total cross-section $\sigma_{\text{tot}}(\gamma p \rightarrow B\bar{B}X)$ at SPS photon beams is several nanobarns, while when $E_Y^L > 150 \text{ GeV}$, the experimental upper limit when searching for $B\bar{B}$ is 2 events/pb¹⁴⁾. The cross-section $\sigma_{\text{tot}} \sim 1 \text{ nb}$ means that we can have 2 events/pb $\times 1 \text{ nb} \sim 2000 \text{ } B\bar{B}$ events, i.e. we have in all 4000 events of B and $\bar{B}$. Assuming that the branching ratio $B \rightarrow \psi K\pi$ is $\sim 3\%$ ²⁾, we can have a total of $\sim 120 \text{ } B$ or $\bar{B}$ events. The branching ratio of $\psi \rightarrow e^+e^-$ is $\sim 7\%$, so we only have about eight events of B or $\bar{B}$ in the $B \rightarrow \psi K\pi$ channel during the whole experiment, which is too small. Therefore, it seems difficult to look for bottomed mesons at the SPS photon beams.

For e-p colliding beams, $\sigma_{\text{tot}}(\gamma p \rightarrow B\bar{B}X)$ is ~ 10 nb, but the luminosity is much smaller than for the SPS photon beams. Therefore looking for B or $\bar{B}$ at e-p colliding beams is also difficult.

If $\sigma_{\text{tot}}(\gamma p \rightarrow B\bar{B}X) \sim 10$ nb, then it is possible to discover beauty with the present experimental limitation. If beauty is found near the forward direction (in diffractive production processes) this means that the vector meson dominance mechanism is dominant.

For charmed meson production, the cross-section is quite large, so there should not be too much difficulty. Recently, a group at CERN¹⁴⁾ has found some evidence for charmed mesons $\bar{D}^0$, F in photon production processes. They use an SPS photon beam ($\langle E_{\gamma}^L \rangle \sim 50$ GeV) and observe the $K\pi$, $K\pi\pi$, $\eta 3\pi$, and $\eta' 5\pi$ channels. More data are needed to confirm the results.

Acknowledgements

We would like to thank D. Amati, A. De Rújula, J. Ellis, H. Fritzsch and D. Treille for many helpful discussions.

REFERENCES

- 1) R. Barate et al., preprint Saclay, D Ph P E 79-17 (1979).
D. Treille, Photo- and hadroproduction of new flavours, Proc. EPS Int. Conf. on High-Energy Physics, Geneva, 1979 (in preparation).
- 2) H. Fritzsch, preprints CERN TH.2648, TH.2703 (1979).
- 3) D. Treille, private communication.
- 4) W.Y. Lee, paper submitted to the Int. Symp. on Lepton and Photon Interactions at High Energies, Hamburg, 1977.
- 5) L.M. Jones and H.W. Wyld, Phys. Rev. D 17 (1978) 759, and references therein.
- 6) H. Fritzsch and K.H. Streng, Phys. Lett. 72B (1978) 385.
- 7) H. Babcock et al., Phys. Rev. D 18 (1978) 162.
- 8) E. Witten, Nucl. Phys. B120 (1977) 189.
C.H. Llewellyn Smith, Phys. Lett. 79B (1978) 83.
R.J. De Witt, L.M. Jones, J.D. Sullivan, D.E. Willen and H.W. Wyld, Univ. Illinois, ILL-TH-78-54 (1978).
S.J. Brodsky, T. De Grand, J. Gunion and J. Weis, SLAC PUB-2199 (1978).
- 9) J.M. Jauch and R. Rohrlich, The theory of photons and electrons (Addison-Wesley, Reading, Mass., 1955), p. 299.
- 10) A.J. Buras and K.J.F. Gaemers, Nucl. Phys. B132 (1978) 249.
- 11) J.R. Ellis, K. Hübner and B.H. Wiik (eds.), CERN 78-02 (1978).
- 12) Chao-hshi Chang, Tung-sheng Tu and Chi-min Wu, CERN TH.2646 (unpublished).
- 13) J. Ellis, M.K. Gaillard, D.V. Nanopoulos and S. Rudaz, Nucl. Phys. B131 (1977) 285.
- 14) D. Treille, private communication.

Figure captions

Fig. 1 : The modern concept of the photon:

- a) point-like component
 - b) photon disassociates into partons, perturbative
 - c) photon couples directly to bound state, non-perturbative
- γ : ; g : ; quark or antiquark: .

Fig. 2 : The mechanisms for the photoproduction of charm and beauty:

- a) γ -g amalgamation
 - b) $q\bar{q} \rightarrow b\bar{b}$ or $c\bar{c}$
- Light line: light quark or light antiquark u, d, s
 Bold line: heavy quark or heavy antiquark c, b.

Fig. 3 : The kinematics of $q\bar{q} \rightarrow b\bar{b}$ or $c\bar{c}$ at $S = 10,800 \text{ GeV}^2$.

Fig. 4 : The beauty production $\sigma_{\text{tot}}(\gamma p \rightarrow B\bar{B}X)$ for SPS photon beams. The dashed line is $q\bar{q} \rightarrow b\bar{b}$; $n = 16, 22$ are the contributions of Buras-Gaemers gluon distributions. We take $G(x, Q^2) \sim [(n+1)/2](1-x)^n/x$.

Fig. 5 : Beauty production: $\sigma_{\text{tot}}(\gamma p \rightarrow B\bar{B}X)$ for e-p colliding beams.

Fig. 6 : Charm production: $\sigma_{\text{tot}}(\gamma p \rightarrow D\bar{D}X)$ for SPS photon beams.

Fig. 7 : Charm production for e-p colliding beams.

Fig. 1

Fig. 2

Fig. 3

Fig. 4

Fig. 5

Fig. 6

Fig. 7