CERN-EP/79-04
15 January 1979

RESONANCES IN $\bar{p}n$ SYSTEM FROM $\bar{p}D$ DATA

G. Alberi

Istituto di Fisica Teorica, Università di Trieste and INFN, Trieste, Italy

C. Alvear^{*)}

Department of Physics and Astronomy, University College, London, England

E. Castelli^{**)}

CERN, Geneva, Switzerland

P. Poropat and M. Sessa

Istituto di Fisica, Università di Trieste and INFN, Trieste, Italy.

L.P. Rosa and Z.D. Thomé

Instituto de Fisica and COPPE/UFRJ, Rio de Janeiro, Brazil

ABSTRACT

Old evidence of a large formation cross-section for the S resonance in the $\bar{p}n$ system is shown to be in contradiction with *other* data covering a larger mass interval.

The analysis of deuteron data is made including the rescattering of pions and antiproton.

(Submitted to Physics Letters B)

^{*)} On leave of absence from the Instituto de Fisica, UFRJ, Rio de Janeiro, Brazil.
^{**)} On leave of absence from Istituto di Fisica, Università di Trieste, Italy.

Old evidence [1] of a mesonic state (S) with mass $m_R = 1935 \pm 1$ MeV and narrow width $\Gamma = 9 \pm 4$ MeV was confirmed by two recent measurements [2,3] of the $\bar{p}p$ total and partial cross-sections. It is not yet clear, however, which is the isospin of the resonance. Because of the absence of the bump in the 0-prong and therefore in the charge-exchange cross-section [4], it is impossible to interpret the observed structure as a single, non-interfering resonance of definite isospin. Many different solutions have been proposed in order to solve this problem [5-8], but in any case there is need of new high-statistics high-resolution data [9a]. In particular it seems rather important to repeat the total cross-section experiment on the deuteron [9b] in order to settle this question, because from this experiment one can extract the $\bar{p}n$ total cross-section, which would provide direct information on the $I = 1$ channel. Unfortunately the extraction of the $\bar{p}n$ total cross-section from $\bar{p}D$ total cross-section data requires the unfolding of the Fermi motion: so it might be more practical to measure the inclusive proton spectator distribution in deuteron break-up, where the energy is unambiguously defined. However, as shown by Bizzarri et al. [10], this procedure implies a more sophisticated theoretical approach, because of three-body effects in the quasi-elastic channel at these energies. Therefore, so far only the copious annihilation channel has been explored [11] and some evidence of the S resonance has been found.

Other data on the same channel are available from an exposure of the CERN 2 m DBC (deuterium bubble chamber) to an antiproton beam [10], the incident antiproton momentum having three values: 459, 540, and 601 MeV/c at the entrance to the bubble chamber (all the details of this experiment can be found in ref. [10]).

In order to obtain a clean sample of antiproton-neutron annihilations, it is necessary to identify the events where a proton is emitted together with an odd number of charged pions. All the odd-pronged events belong to this category, the spectator proton being in this case too slow to give a visible track. As for the even-prong events, they can be attributed to $\bar{p}n$ annihilation if a positive track stops without decaying in the chamber.

This last category has been selected for our goal, and in fig. 1 the distribution of the laboratory momentum (p_s) of the proton spectator is shown. The momentum has been derived from the range of the proton by means of the range-momentum relations. This type of procedure ensures the maximum of sensitivity in the determination of the invariant mass of the $\bar{p}n$ system.

Because of the bubble chamber size and the chosen fiducial volume of the interaction points, there is some possibility that the spectator protons leave the chamber without stopping. The present choice of the fiducial volume excludes this possibility for events in which the momentum of the proton spectator is less than 200 MeV/c. Therefore for further discussion, only the events in which the proton spectator is between 100 and 200 MeV/c have been considered. In this way there is no possibility of missing events because of non-visibility of the proton spectator stopping point: this is true both at the lowest momentum (100 MeV/c -- corresponding to 0.3 cm in deuterium) and at the highest momentum (200 MeV/c -- corresponding to 3.3 cm in deuterium). These events are indicated by the shaded area in fig. 1.

The momentum of the antiproton at the interaction point (p_i) has been determined by means of the range-momentum relations from the mean value of the momentum at the entrance point. Also in this case this procedure has been chosen in order to reduce the spread of the invariant $\bar{p}n$ mass. In fig. 2 the distribution of the laboratory momentum at the interaction point is given for all the considered events (no particular significance has to be attributed to the visible structures of fig. 2 because events from three different exposures are plotted there).

The invariant mass of the $\bar{p}n$ system M_x has been computed for all selected events by means of the formula:

$$M_x^2 = 2m^2 + M^2 + 2M(E_i - E_s) - 2(E_i E_s - \bar{p}_i \bar{p}_s) , \quad (1)$$

where m is the proton or antiproton mass, M the deuteron mass, $E_i = \sqrt{m^2 + p_i^2}$ and $E_s = \sqrt{m^2 + p_s^2}$. The distribution of the value $Q = M_x - 2m$ is shown in fig. 3.

Here we take the attitude of fitting the raw data with a multiple-scattering model, assuming a parametrization of the annihilation cross-section, rather than extracting this quantity directly from the data with the weighting procedure used in ref. [11].

The two methods should be perfectly equivalent if the same assumptions are made for the deuteron dynamics. Here we have added to the single-scattering diagram -- the so-called impulse approximation -- used in ref. [11], the double scattering of the antiproton and the rescattering of the π^+ on the spectator proton^{*)}. We expect that the two last effects contribute substantially to the annihilation cross-section on the deuteron because of the high cross-sections involved in the secondary interactions.

While the antiproton rescattering interferes destructively with the single-scattering diagram, causing a suppression of the order of 50%, the pion rescattering adds incoherently, and partially compensates the antiproton rescattering. Pion rescattering is also responsible for the tail of the spectator distribution, as already suggested by a probabilistic calculation [13]. This is seen in fig. 1, where we compare the predictions of the impulse approximation with our multiple-scattering model.

With the same model we describe the mass distribution, which would have an expression of the type

$$\frac{d\sigma}{dM_x} = \frac{M}{(2\pi)^2} \frac{\lambda^{1/2}(M_x^2, m^2, m^2)}{\lambda(S, M^2, m^2)} M_x \sigma_A(M_x) \int_{-p}^{p^+} \frac{p dp}{\sqrt{m^2 + p^2}} \times \left\{ |D|^2 + \langle n^+ \rangle |E_\pi|^2 \right\}, \quad (2)$$

where

$\sigma_A(M_x)$ in the $\bar{p}n$ annihilation cross-section for the c.m. energy M_x ;

λ is the usual triangular symbol of relativistic kinematics [14];

S is the c.m. energy squared of the antiproton-deuteron system;

*) The effect of π^- rescattering and charge exchange is taken into account using isospin invariance and Pauli antisymmetrization for the deuteron. The result is very small and amounts to 6% of the π^+ rescattering. More details about the multiple-scattering model can be found in ref. [12].

p is the momentum of the proton in the final state, and p^+ , p^- are the kinematical limits on p for fixed M_x [14];

$$|D|^2 = (16\pi^3 M) \left\{ \psi_0^2(p) + \psi_2^2(p) + 2 \operatorname{Re} [\psi_0(p)F] + |F|^2 \right\},$$

F being the double scattering amplitude, and ψ_0 , ψ_2 the S- and D-wave functions of the deuteron. We use the Gaussian parametrization [15] of the Reid [16] wave function. The other term, $\langle n^+ \rangle |E_\pi|^2$, in expression (2) represents the effects of π^+ rescattering, and $\langle n^+ \rangle$ is the average multiplicity of π^+ in $\bar{p}n$ interactions. The detailed expressions of E_π and F has been given elsewhere [12]. Following ref. [11] the annihilation cross-section is given by the following expression:

$$\sigma_A(M_x) = \text{const} \left[\frac{4\pi}{q} + \frac{x_1}{q_1} \frac{\pi\Gamma_1^2}{(M_x - m_1)^2 + \Gamma_1^2/4} + \frac{x_2}{q_2} \frac{\pi\Gamma_2^2}{(M_x - m_2)^2 + \Gamma_2^2/4} \right], \quad (3)$$

q being the $\bar{p}n$ c.m. momentum. The first term of this formula is the well-known $1/v$ term which fits reasonably well the general behaviour of the $\bar{p}n$ annihilation cross-section (see ref. [10]). The other two terms of formula (3) are the Breit-Wigner forms for the two resonances found in ref. [11]: $m_1 = 1897$, $\Gamma_1 = 25$ MeV, and $m_2 = 1934$, $\Gamma_2 = 11$ MeV; x_1 and x_2 are the ratios between the maxima of the Breit-Wigner formulas and the background, given by the term $4\pi/q$. The values of x_1 and x_2 given in ref. [11] are $x_1 = 0.5$ and $x_2 = 1.3$, but these last two ratios will be discussed later in this paper.

The expression (2) is folded within the spectrum of the incident momentum, to give the events distribution:

$$\left\langle \frac{dN}{dM_x} \right\rangle = \frac{\int_{p_{i,\min}}^{p_{i,\max}} \frac{d\sigma(p_i)}{dM_x} \frac{I(p_i)}{\sigma(p_i)} dp_i}{\int_{p_{i,\min}}^{p_{i,\max}} \frac{I(p_i)}{\sigma(p_i)} dp_i}, \quad (4)$$

where p_i is the incident laboratory momentum of the antiproton and $I(p_i)$ is the experimental distribution of fig. 2 ($p_{i,\min} = 0.27$ and $p_{i,\max} = 0.61$ GeV/c). The $\bar{p}n$ annihilation cross-section has been evaluated by means of the formula

$\sigma(p_i) = \pi(R_n + \lambda)^2$ as given in ref. [10], with $R_n = 0.77$ fm, λ being the relative $\bar{p}$ -nucleon wavelength; $\sigma(p_i)$ is the best fit of the M_x average of the $\bar{p}n$ annihilation cross-section, measured with a different technique, in the same range of energy [10].

As a first step the numerical outcome of expression (4) is shown in fig. 3 for $x_1 = x_2 = 0$, both for the impulse approximation ($E_\pi = 0$ and $F = 0$ dot-dashed line), and for the present multiple-scattering model (dashed line). The theoretical curves have been normalized to the total number of events by means of the constant in front of formula (3).

The same comparison has been carried out with $x_1 = 0.5$ and $x_2 = 1.3$, as given in ref. [11]. The result is shown in fig. 3 for the multiple-scattering model (dotted line). The disagreement between our data and the presence of the m_2 resonance with $x_2 = 1.3$ is evident. Moreover, this result does not depend on the dynamics of the deuteron: the impulse approximation does not work as well as the multiple-scattering model. Also the value $x_1 = 0.5$ seems to be in disagreement with the data.

It is nevertheless interesting to put an upper limit for x_2 , compatible with the present experimental data. Two types of fit have been done: in the first case, both x_1 and x_2 were free but positive, and the result is that both x_1 and x_2 are compatible with zero. In the second case x_1 was free but positive, and a search for the maximum value of x_2 gives $x_2 = 0.26$ and x_1 compatible with zero.

From this value of x_2 , a value of 50 mb for the resonance annihilation cross-section for the m_2 signal, as given in ref. [11], is therefore excluded. In ref. [2], the resonance annihilation cross-section in the $\bar{p}p$ case is found to be 3.2 ± 1.8 mb: this analysis obviously does not exclude a value of this type for the $\bar{p}n$ channel.

The presence of an m_1 resonance with $x_1 \approx 0.5$ cannot be definitely excluded because the experimental errors are comparable with the height of the resonance. So the major discrepancy remains for the resonance m_2 ; this difference could be

explained by the fact that for ref. [11] the resonance is at the edge of the phase space, while in the present case it is at the centre of the mass spectrum. This means that, on a mass distribution equivalent to fig. 3 for the data of ref. [11], the resonance peak would appear on the right tail of the spectrum, where the number of events goes down, and could be deformed by statistical fluctuations.

Another possible source of discrepancy could be searched for in the relative momentum between the antiproton and the off-mass-shell neutron, which is different in the two cases. However, the incident energies being higher in our case than for ref. [11], the relative momentum is also higher, and therefore both resonances should be enhanced, because of the dependence on the relative momentum of the central barrier term.

Acknowledgements

We are deeply indebted to R. Bizzarri for having suggested this work, and to L. Bertocchi, M. Mandelkern and L. Montanet for stimulating discussions. Three of us (L.P.R., Z.D.T. and C.A.) are grateful to CNPq and FINEP for partial support, and to Professor Abdus Salam, the International Atomic Energy Agency, and UNESCO for hospitality at the International Centre for Theoretical Physics, Trieste, where most of this work was done.

REFERENCES

- [1] A.S. Carroll et al., Phys. Rev. Lett. 32 (1974) 247.
- [2] V. Chaloupka et al., Phys. Lett. 61B (1976) 487.
- [3] W. Brückner et al., Phys. Lett. 67B (1977) 222.
- [4] M. Alston-Garnjost et al., Phys. Rev. Lett. 35 (1975) 1685.
- [5] C.B. Dover and S.H. Kahana, Phys. Lett. 62B (1976) 293.
- [6] G.C. Rossi and G. Veneziano, Phys. Lett. 70B (1977) 255.
- [7] L.N. Bogdanova et al., ITEP 76-16 (1976).
- [8] R.L. Kelly and R.J. Phillips, RL 76-053T, 159 (1976).
- [9] M. Cresti et al., a) Proposal for a high-statistics high-resolution measurement of the total and partial $\bar{p}p$ cross-sections between 1900 and 1965 MeV total c.m. energy, CERN/TCC 76-6 (1 March 1976), T239. b) Complement to experiment T239 with a similar exposure in deuterium, CERN/PSC 77/13 (4 February 1977), PSC/P 5.
- [10] R. Bizzarri et al., Nuovo Cimento 22A (1974) 225.
- [11] T.E. Kalogeropoulos and G.S. Tzanakos, Phys. Rev. Lett. 34 (1975) 1047.
- [12] G. Alberi et al., preprint IC/78/119 (1978).
- [13] P.D. Zeman, Z. Ming Ma and J.M. Mountz, Phys. Rev. Lett. 38 (1977) 1443.
- [14] E. Byckling and K. Kajantie, Particle kinematics (John Wiley and Sons, Inc., NY, 1973).
- [15] G. Alberi, L.P. Rosa and Z.D. Thomé, Phys. Rev. Lett. 34 (1975) 503.
- [16] R.V. Reid, Ann. Phys. (USA) 50 (1968) 411.

Figure captions

- Fig. 1 : Proton spectator momentum distribution for the $\bar{p}D$ annihilation. The initial part of the spectrum is not visible, because the proton momentum is not measurable. The theoretical curves are normalized on the experimental area for $0 \leq p_s \leq 200$ MeV/c, which is given by the sum of the unseen spectators (5933 events) and the visible spectrum for $p_s \leq 200$ MeV/c (1702 events). The dashed line corresponds to the impulse approximation and the solid one to the present multiple-scattering model. In the calculation of pion rescattering, the momentum of the interacting pion is fixed and equal to 380 MeV/c.
- Fig. 2 : Spectrum of the laboratory momentum of the events with $100 \leq p_s \leq 200$ MeV/c. No particular significance has to be attributed to the visible structures because events from three different exposures are plotted here.
- Fig. 3 : Distribution of the total mass of the annihilation products of events with $100 \leq p_s \leq 200$ MeV/c. The curves (a) and (b) correspond to the absence of resonances: dot-dashed line, impulse approximation; dashed line, multiple-scattering model. The curve (c), dotted line, corresponds to the multiple-scattering model with the two $m_1 = 1897$ and $m_2 = 1934$ MeV resonances defined by all the parameters of ref. [11]. All theoretical curves are normalized to the total number of events.

Fig. 1

Fig. 2

Fig. 3