Ref.TH.2722-CERN

EMBEDDINGS OF CLASSICAL SOLUTIONS OF
 $O(2p+1)$ NON-LINEAR σ MODELS IN CP^{n-1} MODELS

A.M. Din and W.J. Zakrzewski *)

CERN -- Geneva

ABSTRACT

We study the embeddings of $O(3)$ instanton solutions and $O(2p+1)$ spherical harmonics solutions in CP^{n-1} models. They all turn out to be unstable under small complex perturbations. The embedded solutions can be interpreted as mixture states of CP^{n-1} instantons and anti-instantons.

*) On leave of absence from the Department of Mathematics, University of Durham, Durham, England.

Ref.TH.2722-CERN

13 August 1979

1. INTRODUCTION

The CP^{n-1} models¹⁾, which form a special class of non-linear σ models, have recently come under special scrutiny because of interesting properties like conformal invariance, asymptotic freedom and existence of instanton solutions for all n . The latter feature has led to investigations²⁾ of quantum fluctuations in the different topological sectors and the ensuing instanton gas structure might conceivably turn out to have some overlap with the situation in Yang-Mills theories.

The important point in such considerations is the existence of classical solutions of the Euclidean equations of motion. For the two dimensional CP^{n-1} models they are

$$D^2 z + \bar{D}z \cdot Dz = 0, \quad (1.1)$$

derivable from the Lagrangian

$$\mathcal{L}(z) = \bar{D}z \cdot Dz = \partial \bar{z} \cdot \partial z - (\bar{z} \cdot \partial z)^2 \quad (1.2)$$

where $z \equiv \{z_\alpha(x)\}$ is an n component complex field on the Euclidean space E^2 defined modulo an over-all phase factor; also $|z|^2=1$ and the covariant derivative $D = \partial - \bar{z}\partial z$.

The CP^{n-1} models have (stable) instanton solutions for all n given by

$$z_\alpha = \frac{p_\alpha(x_+)}{|p_\alpha(x_+)|} \quad (1.3)$$

where p_α is a polynomial in $x_+ = x_1 + ix_2$,

$$p_\alpha(x_+) = v_\alpha \prod_{i=1}^k (x_+ - u_\alpha^i) \quad (1.4)$$

The degree k of p_α is the instanton number and the action becomes

$$S = 2\pi k. \quad (1.5)$$

Anti-instanton solutions appear by complex conjugation $x_+ \rightarrow x_-$.

So far there is a close analogy with four dimensional Yang-Mills theories. In Yang-Mills theories it is commonly believed that there are no other classes of finite action classical solutions than the instanton solutions and therefore essentially no other solutions were to be taken into account when evaluating functional integrals.

In CP^{n-1} models this is not quite the case as will be discussed in the sequel. This stems from the fact that solutions of $O(n)$ models^{3),4),5)} may be embedded in a trivial way into CP^{n-1} models. Let us recall that $O(n)$ models are defined by the Lagrangian

$$\mathcal{L}(\varphi) = (\partial\varphi)^2, \quad (1.6)$$

where $q_i(x)$ is a real n component field on E^2 with $q^2=1$. The corresponding equations of motion are

$$\partial^2 \varphi + (\partial\varphi)^2 \varphi = 0 \quad (1.7)$$

For the case of $O(3)$ there exist instanton solutions³⁾ given by

$$w = c \frac{\prod_{i=1}^k (x_+ - a_i)}{\prod_{i=1}^k (x_+ - b_i)}, \quad (1.8)$$

where the complex field w is related to q by

$$w = \frac{q_1 + i q_2}{1 + q_3} \quad (1.9)$$

and c, a_i, b_j are complex constants. The instanton number corresponding to the

solution (1.8) is k and the action takes the value

$$S = 8\pi k \quad (1.10)$$

For the $O(2p+1)$ models there exists a set of solutions to the equations of motion^{4),5)} which when stereographically projected on the unit sphere S^2 by

$$\begin{aligned} r_\mu &= \frac{x_\mu}{1+x^2}, \quad \mu = 1, 2 \\ r_3 &= \frac{1-x^2}{1+x^2}, \end{aligned} \quad (1.11)$$

can be written as

$$q_i^{(0)} = \left(\frac{4\pi}{2p+1} \right)^{1/2} \sum_{m=-p}^p U_{im} Y_{pm} \quad (1.12)$$

where U is a certain unitary matrix and Y_{pm} are the spherical harmonics.

The solution $q^{(0)}$ given by (1.8) and (1.12) can as real $2p+1$ dimensional vectors be embedded trivially in the $2p+1$ dimensional space of the CP^{2p} models (and of course also in CP^{n-1} models, $n > 2p+1$; this is however trivial and will not be discussed explicitly). The embedded solutions are still solutions of CP^{2p} since the reality of $q^{(0)}$ implies $q^{(0)} \cdot \partial q^{(0)} = 0$ and thus for $q^{(0)}$ the equations of motion (1.1) and (1.7) are identical. Below we will discuss various properties of such embeddings. In Section II a general property of solutions of (1.1) following from energy momentum conservation is discussed. Section III contains the proof that the considered embeddings are all unstable under small complex perturbations. It is shown in Section IV that the embedded solution can be interpreted as mixture state of CP^{2p} instantons and anti-instantons. Finally, in Section V we comment briefly on some implications of the above.

II. ENERGY-MOMENTUM CONSERVATION

The problem of finding solutions to the equations of motion (1.1) is equivalent, from the mathematical point of view, to finding the harmonic maps⁶⁾, which by definition are maps giving a finite energy extremum of a certain

energy functional (i.e. the Euclidean Lagrangian). The theory of harmonic maps is far from being complete or even systematic and we will therefore in this section limit ourselves to discussing a general property of harmonic maps following from energy-momentum conservation. A similar property has turned out to be useful when proving the uniqueness of instanton solutions in the $O(3)$ model⁷⁾.

The energy-momentum tensor corresponding to the Lagrangian (1.2) is

$$T_{\mu\nu} = -\delta_{\mu\nu} \overline{Dz} \cdot Dz + \overline{D_\mu z} \cdot D_\nu z + D_\mu z \cdot \overline{D_\nu z} , \quad (2.1)$$

and it is of course a conserved quantity :

$$\partial_\mu T_{\mu\nu} = 0 . \quad (2.2)$$

Introducing the derivatives $\partial_\pm$ with respect to the variables $x_\pm$ we find that (2.2) can be rewritten in the simple form

$$\partial_- [D_+ z \cdot \overline{D_- z}] = 0 . \quad (2.3)$$

Thus $D_+ z \cdot \overline{D_- z}$ is a function of x_+ only. The finiteness of the action requires that $|D_\mu z| \rightarrow 0$ as $|x| \rightarrow \infty$ and this implies essentially that

$$D_+ z \cdot \overline{D_- z} = 0 . \quad (2.4)$$

This point can be made more rigorous taking account of possible singularities along the line of reasoning of Refs 4) and 7); we will however not pursue this further here. It might be tempting to conclude further that either

$$D_- z = 0 \quad (2.5)$$

or

$$D_+ z = 0 . \quad (2.6)$$

Since these equations are just the duality equations for respectively instantons and anti-instantons, this would have implied that these solutions would have been unique. This is not so for general n . However for $n=2$, since $\bar{z} \cdot D_{\pm} z = 0$, it is clear that (2.4) implies that at least one of the equations (2.5) and (2.6) must hold. This proves the (known) fact that the CP^1 model (which it is equivalent to the $O(3)$ model) has only instanton and anti-instanton solutions.

For $n \geq 3$ the field space turns out to be sufficiently large to accommodate non-trivial solutions besides the instantons (1.3) and anti-instantons. As discussed in the introduction such solutions arise when real $O(2p+1)$ solutions are embedded in CP^{2p} . If z is real then (2.4) reads simply

$$(\partial_+ z)^2 = 0 \quad (2.7)$$

which implies

$$(\partial_1 z)^2 = (\partial_2 z)^2 \quad (2.8)$$

and

$$\partial_1 z \cdot \partial_2 z = 0 \quad (2.9)$$

In the next two sections we will discuss some properties of such real solutions.

III. STABILITY OF THE REAL SOLUTIONS

The stability of the CP^{n-1} instantons is guaranteed by invoking the invariant topological charge. For the embedded solutions $q^{(0)}$ given in (1.8) and (1.12) the question of stability has to be checked explicitly.

Let us thus consider a z (with $|z|^2=1$) belonging to a neighborhood of $q^{(0)}$. Since z is defined only modulo a phase factor it can be written generally as

$$z = \xi^{(0)} (1 - |\psi|^2)^{1/2} + \psi \quad (3.1)$$

where ψ is a small complex fluctuation

$$\psi = \psi_1 + i\psi_2 \quad (3.2)$$

fulfilling

$$q^{(0)} \cdot \psi = 0 \quad (3.3)$$

The action to second orders in ψ turns out to be

$$S = \int d^2x \mathcal{L}(\psi) = \int d^2x \mathcal{L}_0 + V(\psi_1) + V(\psi_2) - 4 \int d^2x (\partial_\mu^{(0)} \psi_2)^2, \quad (3.4)$$

where

$$\mathcal{L}_0 = (\partial_\mu^{(0)})^2 \quad (3.5)$$

and

$$V(\psi_i) = \int d^2x [-\psi_i \square \psi_i - \mathcal{L}_0 \psi_i^2], \quad i=1,2 \quad (3.6)$$

Now if ψ is real, i.e. $\psi_2=0$ then Eq. (3.4) is just the fluctuation equation for the pure $O(2p+1)$ model, i.e. the stability depends solely on the properties of $V(\psi_1)$. For the case of the spherical harmonics solutions (1.12) it has however been found⁵⁾ that for $p \geq 2$, $V(\psi_1)$ has $(p-1)p(2p+1)$ negative energy fluctuation modes and is thus unstable. If for the same solutions we consider the case $\psi_1=0$ there will be still more negative modes since the previous argument goes through for $V(\psi_2)$ as well and the last term in (3.4) is always zero or negative.

If we now turn to the $O(3)$ solutions (1.8) one may first observe that since $q^{(0)} \cdot \psi_2 = 0$ and the equations (2.8) and (2.9) hold for $q^{(0)}$ then the last term

in (3.4) can be rewritten as

$$4 \int d^2x (\partial_f^{(0)} \psi_2)^2 = 2 \int d^2x \mathcal{L}_0 \psi_2^2 \quad (3.7)$$

Looking only at real fluctuations $\psi_2 = 0$ does not in this case lead to instability since $V(\psi_1)$ is non-negative (0(3) instantons are stable). However if we consider purely imaginary fluctuations $\psi_1 = 0$ then the term (3.7) will give a negative contribution (if $\mathcal{L}_0 \neq 0$) which may be numerically larger than $V(\psi_2)$. This is certainly so if for ψ_2 we take a zero mode of $V(\psi_2)$ as for example $\psi_2 \sim \partial_1 q^{(0)}$ or $\psi_2 \sim \partial_2 q^{(0)}$. From the solution (1.8) it is seen that there are precisely $2(2k+1)$ zero modes. All these modes become negative fluctuation modes of the embedded solution. For the case of the $p=1$ solution (1.12) it can be shown that the term (3.7) does not give rise to further negative fluctuation modes besides the 6 modes ($k=1$) discussed above.

IV. MIXTURE INTERPRETATION OF THE SOLUTIONS

In this section we will show how the considered solutions (1.8) and (1.12) can be interpreted as a mixture of CP^{n-1} instantons and anti-instantons. Let us first notice that for a real solution z the topological charge is zero. This follows from the fact that the topological charge density vanishes

$$\tilde{Q} = -\frac{1}{\pi} (|D_+ z|^2 - |D_- z|^2) = -\frac{1}{\pi} (|\partial_+ z|^2 - |\partial_- z|^2) = 0 \quad (4.1)$$

If the interpretation as a mixture is correct it must therefore contain an equal number of instantons and anti-instantons.

Consider now the solution (1.8) rewritten as

$$w = \frac{A}{B} \quad (4.2)$$

with

$$\begin{aligned} A(x_+) &= c \prod_{i=1}^k (x_+ - a_i) \\ B(x_+) &= \prod_{i=1}^k (x_+ - b_i) \end{aligned} \quad (4.3)$$

Then the relation (1.9) allows us to find explicitly the corresponding $q^{(0)}$ as

$$q^{(0)} = \frac{1}{|A|^2 + |B|^2} (AB^* + A^*B, -i(AB^* - A^*B), |B|^2 - |A|^2) . \quad (4.4)$$

To see what the interpretation of this element of CP^2 might be we introduce translated quantities

$$\begin{aligned} A_s(x_+) &= A(x_+ - s) \\ B_t(x_+) &= B(x_+ - t) \end{aligned} \quad (4.5)$$

and put

$$q_{s,t}^{(0)} = \frac{1}{N} (A_s B_t^* + A_t^* B_s, -i(A_s B_t^* - A_t^* B_s), B_s B_t^* - A_s A_t^*) , \quad (4.6)$$

where N is a normalization factor.

It is clear that $q_{s,t}^{(0)} = q^{(0)}$ when $s=t=0$. Also when $|s|, |t|, |s-t|$ are large compared to the $|a_i|$ and $|b_i|$ then in the neighbourhood $x_+ \sim s$, $q_{s,t}^{(0)}$ looks like a k instanton solution of CP^2 (as in (1.3)) and when $x_+ \sim t$ it looks like a k anti-instanton. The natural interpretation of (4.4) is therefore that of a mixture state of k instantons and k anti-instantons. As discussed in the previous section this mixture state is however not stable.

Let us now consider the embedding of the $O(2p+1)$ solution (1.12) into CP^{2p} . The spherical harmonics $\psi_{pm}(\theta, \phi) = \psi_{pm}(r_1, r_2, r_3)$ with r_i given in terms of the E^2 coordinates x_1, x_2 by (1.11) can be written as homogeneous polynomials of degree p in the variables $r_{\pm} = r_1 \pm ir_2$ and r_3 . Using (1.12) this implies that $q^{(0)}$ is made up of polynomials in x_+ and x_- , with the highest degrees being x_+^p and x_-^p , and an over-all normalization factor. Following the discussion of the $O(3)$ solution (4.4) it is then easy to see that (1.12) might be interpreted as a mixture of p instantons and p anti-instantons in CP^{2p} . For $p=1$ the solution (1.12) is just an $O(3)$ one-instanton so this interpretation agrees with the previous one.

For the solution (4.4) it is interesting to recall that the action was found in (1.10) to be $s = 8\pi k$. The constituents of this mixture state are k instanton

and a k anti-instanton which each according to (1.5) has an action $2\pi k$. We thus see that the interaction energy is $4\pi k$ (corresponding to repulsion).

V. DISCUSSION

Having studied various properties of the embedded solutions (1.8) and (1.12) it is natural to ask what their effect will be on calculations of quantum fluctuations around instanton solutions (1.3). Naïvely one would say that classical solutions which are not stable give rise to unstable quantum states and thus it does not make sense to study quantum fluctuations around such states. However one may also argue that while positive energy fluctuation modes could be treated with standard methods, it is not yet known how to take into account negative modes with an appropriate mathematical apparatus.

Another interesting point is that one normally does quantum fluctuation calculations separately in the instanton and anti-instanton sectors. In a dilute gas approximation one would then simply add the instanton and anti-instanton contributions arguing that if these are far apart they are approximate solutions of the equations of motion. The discussion at the end of Section IV however shows that there exists a system of instantons and anti-instantons which is a solution of the equations of motion such that the interaction energy is equal to the sum of the individual energies. This would seem to indicate that it cannot in general be correct just to add individual instanton and anti-instanton contributions.

To illuminate this point further it would be interesting if one could find new solutions to the equations of motion which were not necessarily real and thus eventually go outside the topological charge zero sector to which the considered real solutions belong.

ACKNOWLEDGEMENTS

We thank Drs. M. Lüscher and R. Stora for some useful discussions.

REFERENCES

1. H. Eichenherr, Nucl. Phys. B146 (1978) 215;
E. Cremmer and J. Scherk, Phys. Lett. 74B (1978) 341;
V. Golo and A. Perelomov, ITEP preprint (1978).
2. A. D'Adda, P. di Vecchia and M. Lüscher, Nucl. Phys. B146 (1978) 63;
V.A. Fateev, L.V. Frolov and A.S. Schwarz, ITEP preprint (1979);
B. Berg and M. Lüscher, DESY preprint (1979);
A.M. Din, P. di Vecchia and W.J. Zakrzewski, CERN preprint TH-2645 (1979)
(to be published in Nucl. Phys.).
3. A.A. Belavin and A.M. Polyakov, JETP Lett. 22 (1975) 245.
4. W.D. Garber, S. Ruijsenaars and E. Seiler, "On finite action solutions
of the non-linear σ model", Princeton Univ. preprint (1978).
5. A.M. Din and W.J. Zakrzewski, CERN preprint TH-2721 (1979).
6. J. Eells and L. Lemaire, A report on harmonic maps, Bull. London Math.
Soc. 10 (1978) 1.
7. G. Woo, J. Math. Phys. 18 (1977) 1264.