CERN-EP/80-44
9 April 1980

RESULTS ON THE MEASUREMENT OF K-SERIES X-RAYS
FROM ANTIPROTONIC HYDROGEN

M. Izycki^{*)}, G. Backenstoss and L. Tauscher

Institut für Physik, Universität Basel, Basel, Switzerland

P. Blüm, R. Guigas, N. Hassler, H. Koch and H. Poth

Kernforschungszentrum und Universität Karlsruhe,
Institut für Kernphysik, Karlsruhe, Germany

B. Emmoth, K. Fransson and A. Nilsson

Research Institute of Physics, Stockholm, Sweden

P. Pavlopoulos^{**)} and K. Zioutas^{***)}

CERN, Geneva, Switzerland

ABSTRACT

The X-ray spectrum emerging from a liquid-hydrogen target when stopping anti-protons was measured with high resolution in order to determine the strong interaction shift and width of the atomic 1s level. Owing to strong Stark quenching only weak structures could be observed. The upper limit of the yield of any one of the K-series transitions is 3.5×10^{-4} (95% C.L.) per stopped antiproton, assuming no broadening of the 1s level. Possible identifications of the weak lines with the K-series transitions are discussed.

(Submitted to Zeitschrift für Physik A)

^{*)} Now at Warsaw University, Poland.

^{**)} Now visitor at CERN from Basel.

^{***)} Now visitor at CERN from Thessaloniki University, Greece.

1. INTRODUCTION

The measurement of the strong interaction effects in hadronic atoms (shift and broadening of levels) provides a tool for studying this interaction at energies near to threshold. This possibility has in particular been exploited for pionic atoms. Measurements on antiprotonic atoms are more scarce because of the low intensity of the available beams. So far a number of light elements down to the $\bar{p}\text{He}$ system have been studied (see [1,2] and references therein). However, the study of antiprotonic hydrogen is of particular importance, since it offers the possibility of studying the bare $\bar{p}$ -nucleon interaction. Connected to this question is also the possible existence of quasi-nuclear bound states just below threshold [3,4]. Unfortunately, such experiments are more difficult compared to those with antiprotonic light elements, because the radiative transitions which occur between levels with low n ($n \leq 4$) are strongly quenched by collisional Stark effect [5,6]. Nevertheless, there have been experimental searches at CERN [7] and at Brookhaven [8] for the X-rays from the $\bar{p}p$ system (protonium). The results of our experiment to look for K-series X-rays from $\bar{p}p$ are presented and discussed in this paper.

2. INTERACTION OF SLOW ANTIPROTONS WITH LIQUID HYDROGEN

Antiprotons stopped in a hydrogen target are captured into atomic orbits, forming an exotic atom in a highly excited state. The de-excitation processes of the $\bar{p}p$ system are described in ref. [9] and cascade intensities have been calculated [10]. The radiative rate for all but the very low levels ($n \leq 4$) is much smaller than the rate of non-radiative de-excitation, e.g. dissociation of the H_2 molecules and external Auger effect. The $\bar{p}p$ system essentially behaves as a small and neutral object and it has no difficulty in penetrating atomic regions where strong electric fields occur. This causes a Stark mixing of $n\ell$ and $n\ell'$ states of protonium. Owing to s-state admixture the annihilation channel is wide open and only a small fraction (10^{-4} - 10^{-2}) of $\bar{p}$ arrives at the level $n = 4$ in liquid hydrogen. The observable intensity of the radiative transitions depends then on the density of the hydrogen, and on the $\bar{p}p$ strong interaction including the existence of deeply bound quasi-nuclear states.

The strong interaction will also shift and broaden the energy levels. The splitting of the triplet and singlet ground states originating from the strong interaction can be expected to be of the order of hundreds of eV [11,12]. The electromagnetic energies of the np-1s transitions are 9.37, 11.10, 11.71 and 12.49 keV for $n = 2, 3, 4$, and ∞ , respectively. Fine-structure splitting of the 2p, 3p, and 4p levels is smaller than 2 eV and can be neglected in our experiment.

The 2p-1s X-rays of $\bar{p}p$ cannot be identified by their energy, because the energy shift caused by the strong interaction is unknown. However, the energy differences between the K-lines are not affected by strong interaction, since the shifts of the p-levels are negligible. Therefore the line pattern provides a good signature for the K-series.

3. EXPERIMENTAL SET-UP

In this experiment we have chosen to use Si(Li) detectors in order to measure with high resolution the widths and shifts of the 1s level, and possibly see the splitting of the level into the singlet and triplet states. As the Si(Li) detectors have small surfaces, a liquid-hydrogen target had to be used to get a reasonable stop density of antiprotons even at the expense of an increased Stark effect. For sufficiently large s-level shifts, however, losses might be small.

3.1 Beam, telescope

The experiment was done at the k_{19} low-momentum electrostatically separated beam of the CERN Proton Synchrotron (PS). The beam line was 17 m long and the momentum was $800 \text{ MeV}/c \pm 2\%$ and the production targets consisted of W or Al. The antiprotons were slowed down in a carbon moderator to about $260 \text{ MeV}/c$. This results in a momentum spread equal to a width of the range curve of 2 g/cm^2 carbon. The beam had a burst length of 400 ms with a repetition rate of 2.4 s. The telescope, consisting of five scintillation counters and two Čerenkov counters, is shown in fig. 1. The antiproton trigger was defined by $S_1 \bar{C}_1 \bar{C}_2 S_2 S_3 S_4 S_5$. Typical rates per burst were $250 \bar{p}_{\text{stop}}$ and about 70000 π^- passing through the target. The rejection power of the telescope for pions was about 10^4 .

Charged particles emerging from the annihilation were detected by seven scintillation counters covering about 70% of the solid angle. For the definition of a stopped antiproton the coincidence of the telescope trigger with at least two of these counters was required.

3.2 Target

Considerable efforts were made in order to produce a target enclosure which is transparent for X-rays down to ~ 5 keV and contains a minimum of material to avoid interfering peaks in our spectra.

The liquid hydrogen was kept in a cylindrical mylar vessel of 16 cm length and 12 cm diameter, equivalent to 1.13 g/cm^2 of liquid hydrogen (see fig. 1). The vessel was made of $150 \text{ }\mu\text{m}$ mylar and it was wrapped in 100 layers of $6 \text{ }\mu\text{m}$ aluminized mylar foil (superinsulation). This insulation allows for filling with liquid helium as well. The surrounding vacuum tank was made of a 2 mm thick titanium cylinder covered at the entrance and the exit of the beam by $150 \text{ }\mu\text{m}$ mylar windows. The five Si(Li) diodes viewed the target through very thin mylar windows ($35 \text{ }\mu\text{m}$) and only two layers of superinsulation. The total window thickness was thus $2 \times 35 \text{ }\mu\text{m}$ mylar + $2 \times 6 \text{ }\mu\text{m}$ aluminized mylar. The detailed configuration is shown in fig. 2.

3.3 Si(Li) detectors

Four of the five Si(Li) diodes used in the experiment were built as twins mounted in the same cryostat and the fifth one was an ordinary single detector. The sensitive volumes were all of the same size ($80 \text{ mm}^2 \times 5 \text{ mm}$) and they were mounted in a cryostat with an end-cup of aluminium. The detectors were placed 90 mm away from the beam axis. They were tuned to give almost identical response, so it was possible to add spectra directly. The stability of the energy calibration and the response of the system were regularly checked with radioactive sources of ^{241}Am and ^{57}Co . In addition, the performance was checked in several runs on $\bar{p}\text{He}$. The energy resolution of the system under normal beam conditions was about 350 eV at 5 keV. The time resolution was about 5 ns and 20 ns at 20 keV and 5 keV, respectively.

To get the distribution of annihilation vertices in the liquid-hydrogen target, the beam profile together with the range curve were mapped with a proton beam of the same energy as that of the antiprotons stopped in the target. In addition, the geometry dependence of the X-ray absorption in our target-detectors set-up was studied by using radioactive sources placed in different positions inside the mylar vessel. This was done after the experiment was finished and the flat front face of the target vessel was cut away.

Combining these pieces of information, the total efficiency as a function of energy could be obtained by integrating over the target volume. This procedure has been described elsewhere [13]. The final result for one of the detectors is shown in fig. 3.

3.4 Data handling

The energy of an X-ray and its time of arrival relative to the $\bar{p}$ -trigger were recorded for each detector separately event by event on magnetic tape via a PDP-11 computer.

The two-dimensional energy-time spectrum was extended to 33 keV and to 400 ns (see fig. 4). The timing window was chosen such that the area of the prompt peak contains 99% of all events above background. This timing procedure was also applied to the $\bar{p}\text{He}$ spectra. It was verified that the intensities of lines in the energy spectrum were not reduced and no artificial structure was produced.

The rest of the events, outside the time window, were also used to create a spectrum. In this spectrum no peaks at all appeared and it had roughly the same shape as the continuous background of the prompt spectrum.

4. RESULTS

4.1 Background peaks

The final prompt X-ray spectrum obtained for 2.3×10^8 $\bar{p}$ stopped in the target is shown in fig. 5, where a 5-point smoothing was applied in order to reduce statistical fluctuations. The analysis was done on the original data, which are shown in figs. 6 and 7 together with a typical fit of lines in the displayed energy region. These peaks are understood to originate from X-rays of $\bar{p}$ and

secondary π^- stopping in the mylar vessel, in the superinsulation around the target, and in the Al housing of the diodes. The peak at 23.1 keV is presumably produced by secondary π^- stopped in the copper of the cooling fingers of the diodes. The identification of these strong peaks gives a good check of the energy calibration and resolution in the final spectrum. However, the energy and the resolution calibration were based on the measurements with radioactive sources of ^{241}Am and ^{57}Co , under beam conditions.

Once one has identified one of the peaks as belonging to a certain transition in a cascade, then necessarily also other members of the cascade have to be present in the spectrum. Their energies were calculated and their intensities estimated by cascade calculations. These parameters were put into the fitting procedure. The results are listed in table 1, where it is indicated which parameters are locked together.

4.2 Continuous background

The background in our spectrum cannot come from charged particles, since they will release large energies in the detector. Thus we are left with the electromagnetic radiation which is prompt in time relative to a stopped antiproton and can be of three different kinds:

- i) Low-energy radiation coming through the detector windows and depositing all its energy in the Si(Li) detector.
- ii) Low-energy radiation coming from the Al housing and other construction parts and depositing all its energy in the Si(Li) detector.
- iii) High-energy radiation coming from any direction and depositing a very small amount of its energy by Compton scattering.

In order to distinguish between (i) and (ii), data were taken where the window of one of the diodes was covered by an 8.8 mg/cm^2 foil of Pd and a 7.1 mg/cm^2 foil of Ni. By looking for the strong variation in absorption below and above the K-edges and looking for the $K_{\alpha}(\text{Pd})$ lines, we can make the following statements about the background.

Firstly, the background in our spectrum is influenced very little by the foils, which means that only a minor contribution to the spectrum originates from

(i). Secondly, when introducing the foils there is no change in the high-energy part of the spectrum, which gives a hint that case (ii) does not contribute significantly. The thicknesses of the foils were of the same size as the absorption length for 20 keV X-rays and their solid angle was about π , which is the same as for the Al housing around the Si(Li) detector.

Another result from the foil experiment is that a small peak from $K_{\alpha}(\text{Pd})$ is visible, which means that foreign atoms in front of the detector will create electronic K_{α} peaks (see Section 4.3.2).

Another experimental fact comes from the $K^{-}p$ experiment [14], which was done in the same set-up as described in this paper. The background has the same shape in the $K^{-}p$ spectrum, but the number of events is around half that in the $\bar{p}p$ spectrum for the same number of stopped charged particles. This eliminates the stopping process as the main contributor to the background (e.g. delta-electrons) and it also fits the fact that the $\bar{p}p$ annihilation creates between two and three times more charged pions and hard gamma-rays than the $K^{-}p$. Process (iii) is now left as the main contributor to our final spectrum.

4.3 Weak structures

4.3.1 Candidates for protonium

In the energy region where the $\bar{p}p$ K-series is expected only weak structures are present. From the statistics we can deduce an upper limit of the yield of 3.5×10^{-4} per stopped $\bar{p}$ (95% C.L.) for the strongest member of the K-series, assuming a line width comparable with the instrumental line width.

A more detailed analysis revealed the existence of small peaks in the energy region 5-15 keV, see fig. 6. Four of them have energy differences corresponding to a possible $\bar{p}p$ K-series pattern. The fit procedure was performed in two ways (the results are given in table 2):

- i) Free energies and intensities for the peaks.
- ii) Fixed energy differences (taken from calculations of electromagnetic energies of the 2p, 3p, and 4p states) and free intensities. The reason not to use the transitions from the ∞p states is that the peak width is uncertain owing to its complexity and therefore the energy difference is uncertain.

In both fits a fixed detector resolution was used for the peak widths and a second-order polynomial was used to fit the background. In the second case the whole energy region was scanned by introducing the expected $\bar{p}p$ K-series pattern in steps of 0.4 keV. There was only one response, the energy and intensity values of which are given in table 2, note a. The errors are the statistical ones obtained from the fitting procedure. From energy calibration a minor error of ± 30 eV has to be added. The peaks originating from transitions in πAl , $\bar{p}C$, $\bar{p}O$ and $\bar{p}Al$ are in accordance with the energy calibration. The 7-6 $\bar{p}C$ transition has a calculated energy of 6.125 keV and it will disturb the position and intensity of the 6.36 keV line. From cascade calculation and comparison with the 6-5 $\bar{p}C$ transition at 10.15 keV an upper limit of the intensity for the 7-6 $\bar{p}C$ transition is obtained. The effect of introducing this peak to the fit is small (see table 2, note b).

Everything said above is valid for fits with fixed widths equal to the detector resolution. As soon as a Lorentzian width is introduced in the line shape the fits do not converge. All results are in agreement with the assumption $\Gamma_{1s} \leq 300$ eV.

The yield per stopped antiproton has been obtained by taking into account the effective experimental detection efficiency $\eta(6.4 \text{ keV}) = 3.2 \times 10^{-3}$. The distribution of stopped antiprotons and the X-ray absorption in liquid hydrogen have been integrated over the target volume.

A special test with Monte Carlo simulated spectra was done to estimate the probability of obtaining, as a statistical fluctuation, a pattern of two lines of a given intensity separated by a given energy. The test was done with 1000 spectra having the same statistics and continuous background shape as the experimental one. The probability of obtaining the two strongest of the observed lines by chance is of the order of 5×10^{-3} .

4.3.2 Other origins than protonium

As found from the Monte Carlo calculation, there is a very small chance to build up this line pattern from fluctuations in the background. Nuclear gamma-rays are also ruled out as an explanation as no candidate has been found and the 30.64 ± 0.01 keV transition in ^{28}Al cannot be verified in our spectrum.

A possibility, however, is electronic X-rays, which show such a regularity in energy differences. Elements such as Cr, Fe, Cu, and Zn, with K_{α} energies equal to 5.41, 6.40, 8.05, and 8.64 keV, respectively, have to be checked for their possible occurrence close to our Si(Li) detectors. No construction materials were made out of these elements, but of course some of them may contain these elements as impurities. The host material must have $Z < 22$; otherwise its K_{α} or L_{α} lines should have been seen in the spectrum. Element analysis was performed for the mylar, the superinsulation and the Al housing of the detector. Two methods were used, the Rutherford back-scattering technique and the proton-induced X-ray technique, both being performed with 1.8 MeV protons from the van de Graaff accelerator of the Research Institute of Physics, Stockholm. The first-mentioned technique involves measuring the energy spectrum of the back-scattered protons from the object for a fixed geometry. In this way one gets the impurities relative to the number of host atoms. This method works fine for $Z_{\text{imp}} > Z_{\text{host}}$. The second method is based on the fact that K vacancies are created by the incoming protons and the relative X-ray intensities are giving the relative impurity abundances in the sample. The results from these studies can be seen in table 3, along with the factory specification. The conclusion drawn from the test is that the impurities in the Al housing cannot be excluded from being the origin of the interesting line pattern on the basis of relative intensities, provided one disregards the Cr line. Now the question arises as to whether there are enough atoms of the impurities to create these peaks? From the short run with the Pd and Ni foils in front of one of the diodes, one knows, for a given geometry, how many atoms are needed to give rise to the K_{α} (Pd) peak with the measured intensity. By transforming this information to the geometry of the Al housing and the number of imbedded Cu and Fe atoms, the expected contribution to our final $\bar{p}p$ spectrum can be calculated. The result is that the yield is a factor of ten too small to be able to create the peaks of interest. However, there is an additional production of secondary particles in the thick Al housing. This contribution is very difficult to estimate and might increase the number of K-vacancies to a level which is high enough to produce a line pattern of the intensity found in our $\bar{p}p$ spectrum.

On the other hand, the line corresponding to Fe-K_α does not show up in the $\text{K}^- \text{H}$ spectrum [14], although even half of its intensity in the present spectrum corresponding to the reduced background in the kaonic spectrum should be visible there. This fact makes the above interpretation of the pattern less convincing.

5. CONCLUSIONS AND DISCUSSION

From the measurements presented here the following conclusions can be drawn:

- i) Under the following, not very restrictive, assumptions an upper intensity limit for the most intensive member of the K-series can be derived: provided the natural broadening of the line is less than 300 eV and the splitting of the 1s level due to the spin dependence of the strong interaction effect is also not larger than this, an upper limit for the presence of any transition in liquid hydrogen to the 1s level can be given, i.e.

$$Y < 3.5 \times 10^{-4} \text{ per stopped } \bar{p} \text{ (95\% C.L.) .}$$

There exist several theoretical calculations of the yield of $\bar{p}\text{H}$, but the results depend strongly on the assumption regarding the $\bar{p}p$ interaction [10,15, 16]. The two extreme predictions are: $Y \sim 10^{-5}$, by Kaufmann and Pilkuhn [15] under the assumption of very strong p absorption ($\Gamma_{2p} \sim 100 \Gamma_{\text{rad}}$; $\Gamma_{\text{rad}} \approx 4 \times 10^{-4}$ eV); and $Y \sim 10^{-2}$, obtained by Markushin [16] for a certain set of strongly bound $\bar{p}p$ states. The recent cascade calculation of Borie and Leon [10] suggests a yield of the order of $Y \sim 10^{-4}$ for a repulsive shift $\epsilon_{1s} = -1$ keV and a width $\Gamma_{1s} \geq 1$ keV and $\Gamma_{2p} = 0.004$ eV. For a narrower width ($\Gamma_{1s} = 250$ eV) and a stronger shift ($\epsilon_{1s} = -2500$ eV), the same authors predict a K_α yield of $Y = 2 \times 10^{-3}$.

- ii) A line pattern typical for the K-series in $\bar{p}\text{H}$ as far as the energy spacing is concerned could be traced in our spectrum. Its probability of being produced by pure statistical fluctuations is 5×10^{-3} . If this pattern is attributed to the K-series of $\bar{p}\text{H}$ it would involve a repulsive shift of the 1s level of -3.04 keV to smaller binding energies as compared with its purely electromagnetic value. However, it should be pointed out that the assignment

of the 3 lines of the pattern to the electronic K_{α} X-rays of Fe, Cu, and Zn cannot be excluded definitely, although also this alternative assignment contains some inconsistencies as described above. Furthermore, it should be noted that there are estimates of the intensities of the pattern based on cascade calculations [10], which are different from the ones observed here in a sense that the maximum intensities should be found in the region of the 5p-1s and 6p-1s transitions and the series limit also pronounced. Direct evidence for such enhanced intensities of high Δn transitions, stronger than predicted by present standard cascade models, were also observed in $\bar{p}\text{He}$ [2]. It is obvious that a shift $\epsilon = -3.04$, combined with a width of $\Gamma_{1s} \lesssim 300$ eV, is not expected by any standard model calculation, although not excluded by a model providing strongly bound states near threshold.

- iii) If one takes the intensity estimates of some cascade calculations [10] seriously one would expect the peak intensities between the 4p-1s and 7p-1s transitions. In this context it may be worth while to note that the 6-5 transition in $\bar{p}\text{C}$ has an energy of 10.15 keV, but its intensity could be easily overestimated in comparison with the 5-4 transition of $\bar{p}\text{C}$ at 18.72 keV. As a consequence, together with an unidentified line at 9.60 keV (table 1) there remains a broad complex between 9.60 and 10.10 keV. If related to the cascade maximum of the relative K X-ray yield at np-1s and $4 \leq n \leq 7$ as calculated in ref. [10], a shift of -2.1 keV for the 1s level and a yield of $Y = 4 \times 10^{-4}$ for the total complex would result. However, the statistical significance of such an assignment being based only on one weak structure would be very low (< 90%).

We are grateful to L. Mazzone and his group for providing us with the hydrogen target. This work was supported by the Bundesministerium für Forschung und Technologie, Federal Republic of Germany, and the Swiss National Science Foundation.

REFERENCES

- [1] H. Poth, G. Backenstoss, I. Bergström, P. Blüm, J. Egger, W. Fetscher, R. Guigas, R. Hagelberg, N. Hassler, C.J. Herrlander, M. Izycki, H. Koch, A. Nilsson, P. Pavlopoulos, H.P. Povel, K. Rolli, I. Sick, L. Simons, A. Schwitter, J. Sztarkier and L. Tauscher, Nucl. Phys. A294 (1978) 435.
H. Poth, Compilation of X-ray data from hadronic atoms, Physics Data No. 14-1 Fachinformationszentrum Karlsruhe (1979).
- [2] H. Poth, R. Abela, G. Backenstoss, P. Blüm, W. Fetscher, R. Hagelberg, M. Izycki, H. Koch, A. Nilsson, P. Pavlopoulos, L. Simons and L. Tauscher, Phys. Lett. 76B (1978) 523.
- [3] P. Pavlopoulos, G. Backenstoss, P. Blüm, K. Fransson, R. Guigas, N. Hassler, M. Izycki, H. Koch, A. Nilsson, H. Poth, M. Suffert, L. Tauscher and K. Zioutas, Phys. Lett. 72B (1978) 415.
- [4] I.S. Shapiro, Phys. Rep. 35 (1978) 129.
- [5] T.B. Day, G.A. Snow and J. Sucher, Phys. Rev. Lett. 3 (1959) 61 and Phys. Rev. 118 (1960) 864.
- [6] M. Leon and H. Bethe, Phys. Rev. 127 (1962) 636.
- [7] E.G. Auld, H. Averdung, J.M. Bailey, G.A. Beer, B. Dreher, H. Drumm, K. Erdman, U. Gastaldi, E. Klempt, K. Merle, K. Neubecker, C. Sabev, H. Schwenk, V.H. Walther, R.D. Wendling, B.L. White and W.R. Wodrich, Phys. Lett. 77B (1978) 454.
- [8] R.E. Welsh, Proc. 7th Int. Conf. on High Energy Physics and Nuclear Structure, Zurich, 1977 (Birkhäuser, Basel, 1977), p. 95.
- [9] O.D. Dalkarov, B.O. Kerbikov and V.E. Markushin, Sov. J. Nucl. Phys. 25 (1977) 455.
- [10] E. Borie and M. Leon, Univ. Karlsruhe preprint TKP 79-7 (1979), to be published in Phys. Rev. A.
- [11] O.D. Dalkarov and V.M. Samoilov, JETP Lett. 16 (1972) 249.
- [12] R.A. Bryan and R.J.N. Phillips, Nucl. Phys. B5 (1968) 201.
- [13] L. Labhardt, Bestimmung der Efficiency einer Targetanordnung, Diplomarbeit in Experimentalphysik, Basel Universität, 1977, unpublished.

- [14] M. Izycki, G. Backenstoss, L. Tauscher, P. Blüm, R. Guigas, N. Hassler,
H. Koch, H. Poth, K. Fransson, A. Nilsson, P. Pavlopoulos and K. Zioutas,
Results of the search for K-series X-rays from kaonic hydrogen, preprint
CERN-EP/80-45, submitted to Z. Phys. A.
- [15] W.B. Kaufmann and H. Pilkuhn, Phys. Rev. C 17 (1978) 215.
- [16] V.E. Markushin, preprint ITEP-164, Moscow (1976).

Table 1

Energies and intensities of all lines fitted in the energy region 5 keV to 32 keV. The half-width of the lines was fixed at the value of the detector resolution.

	Transition	Energy*) (keV)	Area $\pm$ fit error	Possible explanations
$\bar{p}C$	5-4	18.722 ± 0	475 ± 53 a)	
	6-4	28.90 ± 0	164 ± 40	
	6-5	10.15 ± 0	147 ± 49	
	7-5	16.243 ± 0	36 ± 5 a)	
$\bar{p}O$	6-5	18.442 ± 0	198 ± 84 b)	
	7-5	29.551 ± 0	97 ± 44 b)	
	7-6	11.110 ± 0	< 80	
	8-6	18.24 ± 0	21 ± 9 b)	
$\bar{p}Al$	7-6	30.15 ± 0.04	565 ± 157 c)	
	8-7	19.38 ± 0.12	87 ± 43 d)	
	9-7	32.897 ± 0	81 ± 23 c)	
	9-8	13.35 ± 0	39 ± 20 d)	
	10-8	22.90 ± 0	10 ± 3 c)	
πAl	4-3	30.49 ± 0.04	313 ± 40 e)	
	5-4	14.04 ± 0.07	139 ± 46	
	6-4	21.70 ± 0	22 ± 3 e)	
Not identified		5.50 ± 0.08	144 ± 57	K_{α} (Cr)
		6.37 ± 0.07	160 ± 54	$\bar{p}p$ candidate 2-1
		8.06 ± 0.08	113 ± 51	$\bar{p}p$ candidate 3-1
		8.36 ± 0.18	62 ± 53	$\bar{p}p$ candidate 4-1
		9.60 ± 0.12	82 ± 50	$\bar{p}p$ candidate ∞ -1
		16.34 ± 0.17	62 ± 47	πSi 5-4 16.33 keV
		17.08 ± 0.10	104 ± 46	
		17.95 ± 0.13	87 ± 46	Due to asymmetric line profile
		18.39 ± 0	112 ± 65 f)	πC 3-2
		23.09 ± 0.07	134 ± 42	πCu 7-6 23.02 keV
		24.827 ± 0	38 ± 27 f)	πC 4-2
		25.29 ± 0.12	84 ± 41	πN †) 3-2 25.10 keV
		27.93 ± 0.12	88 ± 39	$\bar{p}Ti$ 10-9 27.86 keV
		31.22 ± 0.08	136 ± 40	

*) Zero in the error means locked to calculated energy.

†) Boron nitride is a constituent of the Si(Li) detector.

a-f) The relative intensities are locked when the areas have one and the same letter.

Table 2

Energies and intensities of the X-ray line pattern.
The width of the lines was fixed at the value
of the detector resolution.

E_{exp}	Total number of counts	Yield (per stopped $\bar{p}$)
6.36 ± 0.07	160 ± 54	2.5×10^{-4}
6.33 ± 0.06 a)	162 ± 55 a)	
6.49 ± 0.08 b)	138 ± 54 b)	
8.06 ± 0.08	113 ± 51	1.6×10^{-4}
8.06 ± 0.06 a)	124 ± 52 a)	
8.36 ± 0.18	62 ± 53	0.8×10^{-4}
8.68 ± 0.06 a)	80 ± 52 a)	
9.60 ± 0.12	82 ± 50	1.0×10^{-4}

a) Values from a fit with fixed energy differences.

b) Values from a fit where the 7-6 $\bar{p}C$ transition
was inserted with the calculated values:
6.125 keV and 57 counts.

Table 3

Comparison between experimental results in the low-energy region
and the calculated relative intensities expected from
electronic X-rays originating from the Al housing of the detector.

Element	K_{α} (keV)	Present experiment		Al housing		
				Abundance in Al (%)		Relative intensity ^{a)}
		E_x (keV)	No. of counts	Factory specification	Control analysis b)	
Cr	5.41	5.50 ± 0.08	149 ± 58	0.04-0.35	0.184 ± 0.006	29
Fe	6.40	6.36 ± 0.07	160 ± 54	0.7	0.51	160
Cu	8.05	8.06 ± 0.18	113 ± 51	0.15-0.40	0.301 ± 0.012	189
Zn	8.64	8.36 ± 0.18	62 ± 53	0.25	0.049 ± 0.004	43
-		9.60 ± 0.12	82 ± 50			

a) Normalized to the experimental area of the 6.40 keV peak.

b) The errors of the relative abundances are shown. The errors relative to
the number of Al atoms are about 20%.

Figure captions

- Fig. 1 : Layout of the target-detector system including the telescope arrangement.
- Fig. 2 : A magnified section of the configuration between the target and one of the Si(Li) detectors.
- Fig. 3 : The X-ray efficiency for one of the 0.80 cm^2 detectors in the experimental configuration, taking into consideration the distribution of stopped antiprotons in the liquid-hydrogen target.
- Fig. 4 : Typical energy-time spectra for two of the detectors. The solid lines show the selected time window.
- Fig. 5 : The final prompt X-ray spectrum when a 5-channel smoothing has been applied.
- Fig. 6 : Low-energy region of the final spectrum together with the fit of peaks and a second-order background. The possible $\bar{p}p$ pattern is fitted with locked energy differences.
- Fig. 7 : The high-energy region of the final spectrum together with the fit of peaks and a second-order background.

Fig. 1

Fig. 2

Fig. 3

Fig. 4

Fig. 5

Fig. 6

Fig. 7