[REDACTED]
CERN LIBRARIES, GENEVA

CM-P00059821

CERN - Data Handling Division
DD/79/9
H.N. Mülthei*)
B. Schorr
22 October 1979

ERROR ANALYSIS FOR A SPECIAL X-SPLINE

*) On leave of absence from the University of Mainz.

1. INTRODUCTION

Clenshaw and Negus [1] have recently defined a new cubic spline. It is a generalization of the conventional cubic spline in which the second derivative is allowed to have discontinuities at the inner knots. The magnitudes of these discontinuities are related to those of the third derivatives at the inner knots in such a way that one free parameter is introduced at each inner knot. The new spline is called a (cubic) X-spline. As an application, the X-spline is used in [1] for the classical interpolation problem where function values $y_j = y(x_j)$ at the knots x_j , $j = 0(1)n$, are to be interpolated, together with the end conditions that the first derivative at x_0 and x_n must be equal to given values $y'_0 = y'(x_0)$ and $y'_n = y'(x_n)$, respectively. A sufficient condition for the existence of the interpolating X-spline is given in [1]. The condition still leaves freedom in the choice of the additional parameters. Clenshaw and Negus exploit this fact to construct two special interpolating X-splines which are of practical importance:

- a) the interpolating X-spline minimizing an estimate of the truncation error;
- b) the interpolating X-spline minimizing the computational labour.

The X-spline (a) yields smaller errors than the conventional cubic spline if the knots are unequally spaced. The coefficients of the X-spline (b) can be computed by a simple two-term recurrence relation, whereas in the general case a tri-diagonal system of equations has to be solved.

There are two drawbacks to the minimal labour X-spline (b). It has larger errors than the conventional cubic spline. Secondly, there is a non-symmetry in the determination of the coefficients in the following sense. If we denote by m_j the derivatives of the X-spline at the knots, it turns out that the value y'_n does not affect the computation of any of the unknown derivatives m_j of the X-spline at the inner knots. Of course, as Clenshaw and Negus point out, if the X-spline is used for plotting, where high precision is not important, the minimum labour X-spline is preferable to the conventional cubic spline or to the X-spline (a).

In this paper we combine two special X-splines of the type (b) to obtain an interpolating spline w which itself is not always an X-spline. This procedure eliminates the above-mentioned non-symmetry in the determination of the coefficients of the spline (b). The coefficients of w can be computed by two-term recurrence relations, one running forwards, the other backwards. The computational labour for w , in the case of piecewise equally spaced knots, will still be less than that of the conventional cubic spline. The major part of the paper consists of the error analysis of w . One of the main results is that, in the case of piecewise equally spaced knots, w has essentially the same error estimate as the conventional cubic spline.

2. THE CUBIC X-SPLINE

Throughout this paper we use the following notation:

$$I_n = \{1, 2, \dots, n\}, \quad I_n^0 = I_n \cup \{0\}, \quad n \in \mathbb{N}.$$

According to Clenshaw and Negus a (cubic) X-spline is defined as follows.

Definition:

The function s is an X-spline on the interval $[a, b]$ with knots

$$x_0, x_1, \dots, x_n \quad (a = x_0 < x_1 < \dots < x_n = b)$$

and parameters $\alpha_1, \alpha_2, \dots, \alpha_{n-1}$, iff it has the following three properties:

- i) In each subinterval $[x_{j-1}, x_j]$, $j \in I_n$, s is a polynomial of third degree at most, denoted by s_j say;
- ii) $s \in C^1[a, b]$;
- iii) $s_{j+1}''(x_j) - s_j''(x_j) = \alpha_j \{s_{j+1}'''(x_j) - s_j'''(x_j)\}$, $j \in I_{n-1}$.

From property (iii) we see that, for $\alpha_j = 0$, $j \in I_{n-1}$, the X-spline s is an element of $C^2[a, b]$. Hence, s is a conventional cubic spline with the knots x_j , $j \in I_n^0$, in this case.

Using the notation $m_j = s'(x_j)$, $j \in I_n^0$, Clenshaw and Negus show that for given $y_j = y(x_j)$, $j \in I_n^0$, and for given boundary conditions

$$m_0 = y'_0 = y'(x_0), \quad m_n = y'_n = y'(x_n), \quad (2.1)$$

the interpolating X-spline can be represented, for $x \in [x_{j-1}, x_j]$, by

$$s_j(x) = \frac{m_{j-1}(x_j - x)^2(x - x_{j-1})}{h_j^2} - \frac{m_j(x - x_{j-1})^2(x_j - x)}{h_j^2} + \frac{y_{j-1}(x_j - x)^2[2(x - x_{j-1}) + h_j]}{h_j^3} + \frac{y_j(x - x_{j-1})^2[2(x_j - x) + h_j]}{h_j^3}, \quad j \in I_n, \quad (2.2)$$

where m_j , $j \in I_{n-1}$ are determined by the equations

$$\begin{aligned} & \beta_j(1 - \beta_j - k_j \beta_j)m_{j-1} + [2 + k_j - 2\beta_j(1 + k_j)]m_j + (1 - \beta_j)^2(1 + k_j)m_{j+1} \\ & = (3 + 2k_j)(1 - \beta_j)^2 y_{j+1} + \beta_j(3 - 3\beta_j - 2\beta_j k_j) y_j, \quad j \in I_{n-1}, \end{aligned} \quad (2.3)$$

with

$$\begin{aligned} h_j &= x_j - x_{j-1}, \quad y_j = \frac{y_j - y_{j-1}}{h_j}, \quad j \in I_n, \\ \beta_j &= \frac{h_{j+1}}{h_j + h_{j+1}}, \quad k_j = \frac{3\alpha_j}{h_{j+1}}, \quad j \in I_{n-1}. \end{aligned}$$

A sufficient condition for the existence of a unique solution of the system (2.3) is given by (see [1])

$$0 \leq 1 + k_j \leq \frac{1}{\beta_j}, \quad j \in I_{n-1}. \quad (2.4)$$

Taking the equality sign on the left-hand side of (2.4) we obtain

$$k_j = -1, \quad j \in I_{n-1}.$$

Setting $k_j = -1$ in (2.3) yields, after a few simple manipulations,

$$m_j = -\beta_j m_{j-1} + (1-\beta_j)^2 Y_{j+1} + \beta_j (3-\beta_j) Y_j, \quad j \in I_{n-1}. \quad (2.5)$$

The simple recurrence relation (2.5) together with the additional boundary conditions (2.1) determine Clenshaw and Negus's minimal labour X-spline, which we denote by u .

Taking the equality sign on the right-hand side of (2.4) we obtain

$$k_j = \frac{1}{\beta_j} - 1, \quad j \in I_{n-1}.$$

Inserting these k_j in (2.3) yields

$$m_j = -(1-\beta_j) m_{j+1} + (2+\beta_j)(1-\beta_j) Y_{j+1} + \beta_j^2 Y_j, \quad j \in I_{n-1}. \quad (2.6)$$

The recurrence relation (2.6) together with (2.1) determine another minimal labour X-spline, which we denote by v .

The X-splines u and v are in general different. Both are interpolating splines. The computation of the unknown coefficients m_j for the spline u runs from left to right, whereas for the spline v it runs in the opposite direction. This suggests that it might be possible to balance out the disadvantages due to favouring one direction rather than the other by using an appropriate combination of the two interpolating splines u and v . For equally spaced knots, or for piecewise equally spaced knots, the additional computational labour is almost negligible. We therefore consider, for any constant τ , the convex sum

$$w = w_\tau = \tau u + (1-\tau)v.$$

w is not necessarily an X-spline, the reason being that the property (iii) in the above definition is not always fulfilled. If the jumps of the third derivative of w at the inner knots are different from zero, w is always an X-spline. This problem is discussed at the end of Section 3 in the case of equally spaced knots. For

the error analysis of w , however, the property (iii) is not necessary. Before we discuss the choice of the constant τ , and as a preliminary to the detailed error analysis in Section 3, we need a few auxiliary results.

Let $y \in C^5[a, b]$ and let $y_j^{(i)} = y^{(i)}(x_j)$, $i = 1, 2, 3, 4$; $j \in I_{n-1}$. Clenshaw and Negus show that, for any interpolating X-spline,

$$m_j = y'_j + A_j y_j^{(4)} + O(h^4), \quad h \rightarrow 0, \quad j \in I_{n-1},$$

with $A_j = O(h^3)$, $h = \max_{j \in I_n} h_j$ and

$$\begin{aligned} & \beta_j(1-\beta_j-k_j\beta_j)A_{j-1} + [2 + k_j - 2\beta_j(1+k_j)]A_j + (1-\beta_j)^2(1+k_j)A_{j+1} \\ &= \frac{(h_j+h_{j+1})^3}{4!} \beta_j(1-\beta_j)^2[1 - 2\beta_j(1+k_j)], \quad j \in I_{n-1}, \end{aligned} \quad (2.7)$$

$$A_0 = A_n = 0.$$

Setting $k_j = -1$ in (2.7), we find that for spline u ,

$$A_j = -\frac{h_{j+1}}{h_j + h_{j+1}} A_{j-1} + \frac{h_j^2 h_{j+1}}{4!}, \quad j \in I_{n-1}, \quad (2.8)$$

$$A_0 = A_n = 0.$$

Setting $k_j = (1-\beta_j)/\beta_j$, we find that for spline v ,

$$B_j = -\frac{h_j}{h_j + h_{j+1}} B_{j+1} - \frac{h_j h_{j+1}^2}{4!}, \quad j \in I_{n-1}, \quad (2.9)$$

$$B_0 = B_n = 0,$$

where $m_j = y'_j + B_j y_j^{(4)} + O(h^4)$ for v .

From (2.8) and (2.9) and the definition of w we immediately obtain

$$m_j = y_j + C_j y_j^{(4)} + O(h^4) \text{ for } w \text{ with}$$

$$C_j = C_j(\tau) = \tau A_j + (1-\tau) B_j, \quad j \in I_n^0. \quad (2.10)$$

We now consider the case of equally spaced knots. We then obtain from (2.8) and (2.9):

$$A_j = -\frac{1}{2} A_{j-1} + \frac{h^3}{4!}, \quad j \in I_{n-1}, \quad (2.11)$$

$$B_j = -\frac{1}{2} B_{j+1} - \frac{h^3}{4!}, \quad j \in I_{n-1}. \quad (2.12)$$

From (2.11) and (2.12) and the fact that $A_0 = B_0 = A_n = B_n = 0$, we easily find that

$$B_j = -A_{n-j}, \quad j \in I_n^0. \quad (2.13)$$

For A_j and C_j we find from (2.11), (2.10) and (2.13):

$$A_j = \frac{h^3}{3 \cdot 4!} \left(2 - \frac{(-1)^j}{2^{j-1}} \right), \quad j \in I_{n-1}, \quad (2.14)$$

$$C_j = \frac{h^3}{3 \cdot 4!} \left\{ 4\tau - 2 + (-1)^{j+1} \left[\frac{\tau}{2^{j-1}} + (1-\tau) \frac{(-1)^{n+1}}{2^{n-j-1}} \right] \right\}, \quad j \in I_{n-1}, \quad (2.15)$$

$$C_0 = C_n = 0.$$

3. ERROR ANALYSIS OF THE SPLINE w

To minimize the error $e = y - w = y - \tau u - (1-\tau)v$ we may determine τ such that

$$E(\tau) = \int_a^b e^2(x) dx$$

is minimal. Setting $E'(\tau) = 0$ we obtain

$$\tau = \tau_{\text{opt}} = \frac{\int_a^b [y(x) - v(x)][u(x) - v(x)] dx}{\int_a^b [u(x) - v(x)]^2 dx},$$

where we assume that the denominator is larger than zero.

The quantity τ_{opt} requires a substantial computing effort. We therefore determine an optimal τ which is independent of y by minimizing the maximum error constant, which we now define.

Let $y \in C^5[a, b]$. It follows from (5.6) [1] that at $x = x_{j-1} + \theta h_j$, $j \in I_n$, $\theta \in [0, 1]$, the error can be written as

$$e_j(x) = h_j^4 y_j^{(4)} K_j(\theta) + O(h^5), \quad h \rightarrow 0, \quad j \in I_n,$$

where

$$K_j(\theta) = K_j(\theta; n, \tau) = \frac{1}{4!} g(\theta) r_j(\theta), \quad (3.1)$$

$$g(\theta) = \theta(1-\theta), \quad (3.2)$$

$$r_j(\theta) = g(\theta) + c_j \theta + d_j, \quad (3.3)$$

$$c_j = \frac{4!}{h_j^3} (C_j + C_{j-1}), \quad (3.4)$$

$$d_j = -\frac{4!}{h_j^3} C_{j-1}, \quad (3.5)$$

with C_j given by (2.10).

In the following we use the maximum norm exclusively. On $[x_{j-1}, x_j]$ we define the error constant as

$$\eta_j = \eta_j(w_\tau) = \|K_j\|, \quad j \in I_n. \quad (3.6)$$

The maximum error constant is then

$$\eta = \eta(w_\tau) = \max_j \eta_j. \quad (3.7)$$

An obvious choice of τ is that value which minimizes η . Using the notation

$$\eta_{\text{opt}} = \frac{1}{4!} \|g^2\| = \frac{1}{4!} g^2\left(\frac{1}{2}\right) = 1/384 \approx 0.0026, \quad (3.8)$$

it is shown in [1] that

$$\eta \geq \eta_{\text{opt}}$$

for any interpolating X-spline; $\eta = \eta_{\text{opt}}$ is valid if $C_j = 0$, $j \in I_n^0$, as can be deduced from (3.1) to (3.8) in the same way as in [1].

For $n = 2$ the parameter τ can be determined such that $\eta = \eta_{\text{opt}}$ for w . From (2.8) to (2.10) we easily find, by setting $C_1 = 0$, that

$$\tau = \frac{h_2}{b - a}.$$

It can be shown by determining $m_1(w_\tau)$ that, for this τ , w_τ is the optimal X-spline considered in [1], which coincides with the conventional cubic spline for $h_1 = h_2$.

Using the same formulae it can immediately be proved that, for $n \geq 3$, there exists, in general, no τ such that $C_j = 0$, $j \in I_n^0$, since we have $n - 1$ conditions $C_j = 0$, $j \in I_{n-1}$ for the variable τ . This implies that, in general, $\eta > \eta_{\text{opt}}$ for $n \geq 3$.

It is complicated to determine τ such that $\eta = \eta(w_\tau)$ is minimal. Since, however, we want $|C_j|$ as small as possible, we determine τ such that

$$C(\tau) = \sum_{j=1}^{n-1} C_j^2 = \sum_{j=1}^{n-1} [(A_j - B_j)\tau + B_j]^2$$

takes its minimum. Setting $C'(\tau) = 0$ we obtain

$$\tau \sum_{j=1}^{n-1} (B_j - A_j)^2 = \sum_{j=1}^{n-1} (B_j - A_j) B_j. \quad (3.9)$$

For equally spaced knots we now prove that $\tau = \frac{1}{2}$, as one would expect. From (2.13) we find that

$$\sum_{j=1}^{n-1} (B_j - A_j) B_j = \sum_{j=1}^{n-1} (A_j + A_{n-j}) A_{n-j} = \sum_{i=1}^{n-1} (A_{n-i} + A_1) A_i = - \sum_{j=1}^{n-1} (B_j - A_j) A_j,$$

and therefore

$$\sum_{j=1}^{n-1} (B_j - A_j)^2 = \sum_{j=1}^{n-1} (B_j - A_j) B_j - \sum_{j=1}^{n-1} (B_j - A_j) A_j = 2 \sum_{j=1}^{n-1} (B_j - A_j) B_j .$$

Using this result in (3.9) we have $\tau = \frac{1}{2}$.

For our main result on the error constants we will need the following.

Lemma:

Let

$$\begin{aligned} P(x) &= P(x; c, d) = g(x)r(x) , & 0 \leq x \leq 1 , \\ r(x) &= r(x; c, d) = g(x) + cx + d , \\ g(x) &= x(1 - x) . \end{aligned}$$

For

$$c + d \geq 0 , \quad d \geq 0 , \quad (3.10)$$

and for

$$c + d \leq 0 , \quad 0 \leq c , \quad -\frac{1}{2} (\sqrt{2}-1) \leq d \leq 0 , \quad (3.11)$$

we have

$$\|P\| = P(x_0; c, d)$$

with

$$x_0 = x_0(c, d) = \frac{1}{2} + \frac{1}{4} c + 2\sqrt{-p} \sin \left(\frac{1}{3} \text{Arcsin } z \right) , \quad (3.12)$$

where

$$z = \frac{q}{(-p)^{3/2}} ,$$

$$p = -\frac{1}{48} [3c^2 + 4(1+c) + 8d] , \quad (3.13)$$

$$q = -\frac{c}{64} [c^2 + 2c + 4d] . \quad (3.14)$$

Proof:

First let $c = 0$ and $d \geq -\frac{1}{2}(\sqrt{2}-1)$. It is then easy to verify that the real points

$$\bar{x}_0 = \frac{1}{2} \quad \text{and} \quad \bar{x}_{\pm} = \frac{1}{2} \pm \sqrt{1+2d}$$

are the positions of the maxima of $|P|$ on the real axis. We further have

$$P(\bar{x}_0) = \frac{1}{4} \left(\frac{1}{4} + d \right) \geq \frac{d^2}{4} = |P(\bar{x}_{\pm})|.$$

It therefore follows that, in this case, $\|P\| = P(\bar{x}_0) = P(\frac{1}{2})$, where $x_0(0,d) = \bar{x}_0$.

We now assume that $c \neq 0$. We first show that for (3.10) and (3.11) there exists exactly one $x_0 \in (0,1)$ such that $\|P\| = P(x_0)$.

For (3.10) this is seen immediately, since $P(x) > 0$ for $x \in (0,1)$, $P(0) = P(1) = 0$, and since the polynomial P tends to plus infinite as $|x|$ tends to infinity.

We already know that $|P|$ takes its maximum at $x = \frac{1}{2}$ if $c = 0$ and $d \geq -\frac{1}{2}(\sqrt{2}-1)$. The graph of $r(x; c,d)$ for $x \in (0,1)$ is similar to the graph of K_{2j} in Fig. 1. This means that, for $c > 0$, we have $r(x; c,d) > r(x; 0,d)$ for $x \in (0,1)$. We can therefore conclude that there exists a unique $x_0 \in (0,1)$ such that $\|P\| = P(x_0)$ if (3.11) holds. We now show that x_0 is defined by (3.12).

Let

$$Q(x) = \frac{1}{4} P'(x) = x^3 - \frac{3}{4} (2+c)x^2 + \frac{1}{2} (1+c-d)x + \frac{d}{4}.$$

By setting

$$x = \frac{1}{2} + \frac{c}{4} + y,$$

Q is transformed into

$$R(y) = y^3 + 3py + 2q,$$

where p and q are given by (3.13) and (3.14), respectively.

From the foregoing considerations it is trivial to show that P always has exactly three extrema, of which the middle one is to be found. The polynomial R therefore has exactly three real roots, given according to the general theory of cubic equations (see [2]) by

$$y_0 = -2\sqrt{-p} \cos \left(\frac{\pi}{2} + \frac{1}{3} \operatorname{Arccsin} z \right),$$

$$y_1 = 2\sqrt{-p} \cos \left(\frac{\pi}{6} + \frac{1}{3} \operatorname{Arccsin} z \right),$$

$$y_2 = -2\sqrt{-p} \cos \left(-\frac{\pi}{6} + \frac{1}{3} \operatorname{Arccsin} z \right),$$

with $z = q/(-p)^{3/2}$. We may conclude from what we saw above that x_0 is the middle root of the three roots of Q. We therefore have to choose that y_i which has the smallest modulus. Since

$$\frac{1}{3} |\operatorname{Arccsin} x| \leq \frac{\pi}{6}$$

for any x, it is easy to see that y_0 has the smallest modulus. Finally, since

$$x_0 = \frac{1}{2} + \frac{c}{4} + y_0 \quad \text{and} \quad \cos \left(\frac{\pi}{2} + \alpha \right) = -\sin \alpha$$

the lemma is proved.

We are now able to give an explicit expression for the error constants η_j in (3.6) if the knots are equidistant and if $\tau = \frac{1}{2}$. For this case it can be verified by (3.4), (3.5) and (2.15) that

$$\left. \begin{aligned} c_j &= \frac{(-1)^j}{3} \left(\frac{1}{2^j} + \frac{(-1)^n}{2^{n-j}} \right), & j \in I_{n-1}, \\ d_1 &= 0, \\ d_j &= \frac{(-1)^{j+1}}{3} \left(\frac{1}{2^{j-1}} + \frac{(-1)^{n+1}}{2^{n+1-j}} \right), & j = 2(1)n-1. \end{aligned} \right\} \quad (3.15)$$

In addition, it can be seen that $\eta_j = \eta_{n+1-j}$ for $1 \leq j \leq N = \left[\frac{n+1}{2} \right]$, where $[x]$ denotes the integer part of x.

Theorem:

Let $\tau = \frac{1}{2}$, $n \geq 2$, and let the knots be equally spaced. We then have, for $(n, j) \neq (3, 2)$,

$$\eta_j = K_j \left(\frac{1}{2} + \frac{1}{4} c_j + 2\sqrt{-p_j} \sin \left(\frac{1}{3} \operatorname{Arcsin} z_j \right) \right), \quad j \in I_N,$$

where K_j is given by (3.1) and

$$z_j = \frac{q_j}{(-p_j)^{3/2}},$$

$$p_j = -\frac{1}{48} [3c_j^2 + 4(1+c_j) + 8d_j],$$

$$q_j = -\frac{c_j}{64} [c_j^2 + 2c_j + 4d_j],$$

with c_j and d_j given by (3.15).

For $(n, j) = (3, 2)$ we have

$$\eta_2 = \frac{1}{4} \eta_{\text{opt}}.$$

The proof of the theorem in the general case can be given by using the lemma. In fact, for j odd, the coefficients c_j and d_j satisfy the assumptions (3.10), and, for $j \neq 2$ even, they satisfy the assumptions (3.11).

For the exceptional case $(n, j) = (3, 2)$, where $c_2 = 0$ and $d_2 = -\frac{1}{4} < -\frac{1}{2}(\sqrt{2}-1)$, direct evaluation of K_2 shows that

$$\eta_2 = \|K_2\| = \left| K_2 \left(\frac{1}{2} \pm \frac{1}{2} \sqrt{1 + 2d_2} \right) \right| = \frac{1}{4} \eta_{\text{opt}}.$$

Remark:

A more detailed evaluation of p_j and q_j shows that, for all j ,

$$|z_j| = \left| \frac{q_j}{(-p_j)^{3/2}} \right| \ll 1,$$

which means that we can use

$$\tilde{\eta}_j = K_j \left(\frac{1}{2} + \frac{1}{4} c_j - \frac{2}{3} \frac{q_j}{p_j} \right) = \frac{1}{4!} P \left(\frac{1}{2} + \frac{1}{4} c_j - \frac{2}{3} \frac{q_j}{p_j}; c_j, d_j \right) \quad (3.16)$$

as an approximation of η_j . Numerical examples show that very good approximations are obtained, as is demonstrated by the following table:

n	$10^{11} \times \max_j e_j$
3	40
5	5.9
15	3.3
20	3.3

The following corollary to our theorem shows the monotonicity of the error constants η_j , and it enables us to determine immediately the maximal error constant η .

Corollary 1:

For equally spaced knots, and $\tau = \frac{1}{2}$, we have

$$\text{for } n \geq 2: \quad \eta_j > \eta_{\text{opt}} > \eta_i \quad (3.17)$$

with $1 \leq i, j \leq N$, i even and j odd;

$$\text{for } n \geq 5: \quad \eta_j > \eta_{j+2} > \eta_{\text{opt}} > \eta_i > \eta_{i+2} \quad (3.18)$$

with $1 \leq i, j \leq N-2$, i even and j odd.

Because of the detailed proof of the lemma which was given above, we confine our demonstration of the corollary to the following hint.

Looking at the curves in Fig. 1, where the qualitative behaviour of K_j and

$$K_{\text{opt}}(\theta) = \frac{1}{4!} \theta^2(1-\theta)^2, \quad 0 \leq \theta \leq 1,$$

is sketched, it is easy to see, by inspection, that the inequalities (3.17) and (3.18) must hold.

It now follows from (3.17) for $n = 2, 3, 4$, and from (3.18) for $n \geq 5$, that

$$\eta = \max_j \eta_j = \eta_1 .$$

This means that the largest errors are to be expected near the points a and b , depending on $y_j^{(4)}$.

We now compare the error constants of the splines u and w_τ for $\tau = \frac{1}{2}$ and equally spaced knots. For simplicity we assume that n is large. We then have the following corollary 2 to our theorem.

Corollary 2:

For equally spaced knots, n large, and $\tau = \frac{1}{2}$, we have:

$$\eta_1(w) \approx 0.42 \eta_1(u) . \quad (3.19)$$

If, in addition, $1 \ll j \leq N$, we have:

$$\eta_j(w) \approx \eta_{\text{opt}} \approx 0.62 \eta_j(u) . \quad (3.20)$$

Proof:

For n large we find from (3.15) that $c_1 \approx \frac{1}{6}$ and $d_1 = 0$. From our theorem and from (3.16) we then obtain

$$\eta_1(w) \approx \frac{1}{4!} P\left(0.5353; \frac{1}{6}, 0\right) \approx 0.0035 .$$

It is shown in [1] that $\eta_1(u) \approx 0.0084$, from which (3.19) follows immediately.

According to (3.15) we may set $c_j = d_j = 0$ for n large and $1 \ll j \leq N$. From the theorem and from (3.16) we then find that

$$\eta_j(w) \approx \frac{1}{4!} P\left(\frac{1}{2}; 0, 0\right) = \eta_{\text{opt}} .$$

Since it is shown in [1] that

$$\eta_j(u) \approx 0.0042 = \eta_{\text{opt}}/0.62 ,$$

corollary 2 is proved.

Finally, we briefly discuss another property of the splines u and w in which w is superior to u . We consider the jumps of the second and the third derivatives of these splines at the inner knots.

For any X-spline s we use the notation

$$\begin{aligned} \Delta_j''(s) &= s_{j+1}''(x_j) - s_j''(x_j) , \\ \Delta_j'''(s) &= s_{j+1}'''(x_j) - s_j'''(x_j) . \end{aligned} \quad j \in I_{n-1} ,$$

Let $y \in C^5[a, b]$. Using (2.2), the relation $m_j = y_j' + A_j y_j^{(4)} + O(h^4)$, and Taylor's theorem, it can be shown that, for equidistant knots,

$$\Delta_j'''(s) = \frac{6}{h^2} \left\{ \frac{h^3}{3!} + A_{j+1} - A_{j-1} \right\} y_j^{(4)} + O(h^2) , \quad j \in I_{n-1} . \quad (3.21)$$

Using (2.14) and (2.13), respectively, in (3.21) yields:

$$\Delta_j'''(u) = \left(1 - \frac{(-1)^j}{2^{j+2}} \right) h y_j^{(4)} + O(h^2) , \quad (3.22)$$

$$j \in I_{n-1} ,$$

$$\Delta_j'''(v) = \left(1 - \frac{(-1)^{n-j}}{2^{n-j+2}} \right) h y_j^{(4)} + O(h^2) . \quad (3.23)$$

Since, by definition, we have, for any X-spline s ,

$$\Delta_j''(s) = \alpha_j \Delta_j'''(s) , \quad j \in I_{n-1} ,$$

and since $\alpha_j = -h/3$ for u and $\alpha_j = h/3$ for v , we obtain from (3.22) and (3.23):

$$\Delta_j''(u) = -\frac{1}{3} \left(1 - \frac{(-1)^j}{2^{j+2}} \right) h^2 y_j^{(4)} + O(h^3) , \quad (3.24)$$

$$j \in I_{n-1} ,$$

$$\Delta_j''(v) = \frac{1}{3} \left(1 - \frac{(-1)^{n-j}}{2^{n-j+2}} \right) h^2 y_j^{(4)} + O(h^3) . \quad (3.25)$$

Finally, for $w = w_{\frac{1}{2}} = \frac{1}{2}(u+v)$ we obtain from (3.22) and (3.25):

$$\Delta_j''(w) = \frac{(-1)^j}{24} \left(\frac{1}{2^j} - \frac{(-1)^n}{2^{n-j}} \right) h^2 y_j^{(4)} + O(h^3), \quad (3.26)$$

$$j \in I_{n-1},$$

$$\Delta_j'''(w) = \left(1 - \frac{(-1)^j}{8} \left[\frac{1}{2^j} + \frac{(-1)^n}{2^{n-j}} \right] \right) h y_j^{(4)} + O(h^2). \quad (3.27)$$

In the asymptotic formulae (3.22) to (3.27) the absolute value of the coefficient of $y_j^{(4)}$ is maximal for $j = 1$ and/or $j = n-1$. It is therefore informative to consider the following quotients for large n and $y^{(4)}(a) \neq 0$:

$$\frac{\Delta_1''(w)}{\Delta_1''(u)} \approx \frac{1}{18}, \quad \frac{\Delta_1'''(w)}{\Delta_1'''(u)} \approx \frac{17}{18}.$$

This implies that especially with respect to the jumps of the second derivative, the spline w is superior to the spline u .

Furthermore, we may conclude from (3.27) that

$$\Delta_j'''(w) \neq 0, \quad j \in I_{n-1}, \quad n \gg 1,$$

for $y^{(4)}(x) \neq 0$, $x \in [a, b]$. This gives, as indicated in Section 2, a sufficient condition for w to be an X-spline.

To summarize our error analysis of the spline w , we may say that the interpolating spline $w_{\frac{1}{2}}$ is more precise than the minimal labour X-spline u of Clenshaw and Negus. Its computation is slightly more expensive than that of u , but nevertheless less expensive than the computation of the conventional cubic spline if the step sizes h_j do not change too often. Finally, for equally spaced knots, and away from the neighbourhood of the endpoints, the spline $w_{\frac{1}{2}}$ has essentially the same precision as the conventional cubic spline which, in this case, has the smallest possible error constant η_{opt} .

4. NUMERICAL EXAMPLES

We compare, for two different functions, the conventional cubic spline s , which we denote by s_c , with the splines u , $w_{\frac{1}{2}}$, and $w_{\tau_{\text{opt}}}$, respectively.

We begin with the example which was considered by Clenshaw and Negus, and where eight values of $\sin x$ at $x = 0(\pi/9) \pi/3, \pi/3(\pi/18) 5\pi/9$ are interpolated. The results are shown graphically in Figs. 2 and 3, in which plots of the absolute difference between the splines s_c , u , $w_{\frac{1}{2}}$, $w_{\tau_{\text{opt}}}$, and y are shown. In Fig. 2 it can be seen that u and s_c have the same maximum error, while the maximum error of $w_{\frac{1}{2}}$ is smaller. In addition, the spline $w_{\frac{1}{2}}$ is much better than the spline u in the first two intervals. Figure 3 shows a comparison between $w_{\frac{1}{2}}$ and $w_{\tau_{\text{opt}}}$, where τ_{opt} was computed numerically according to the formula given in Section 3. We see that, apart from the maximum error peak, $w_{\frac{1}{2}}$ is normally better than $w_{\tau_{\text{opt}}}$. In other words, the additional work required to determine τ_{opt} is not justified in this case.

In the second example, six values of $y = x \sin 2\pi x$ at $x = j/5$, $j = 0, 1, \dots, 5$ were interpolated. Using the same notation as for Figs. 2 and 3, the results for this example are shown in Figs. 4 and 5. Here again $w_{\frac{1}{2}}$ is superior to u and almost as good as s_c . The behaviour of $w_{\frac{1}{2}}$ and $w_{\tau_{\text{opt}}}$ is similar to that of the first example.

Acknowledgement

The authors wish to thank C. Schneider, Mainz, for helpful comments.

REFERENCES

- [1] Clenshaw, C.W. and Negus, B.: The Cubic X-Spline and its Application to Interpolation, J. Inst. Maths. Applics. 22, 109-119 (1978).
- [2] Carr, G.S.: Formulas and Theorems in Pure Mathematics. New York, N.Y.: Chelsea Publ. Comp. 1970.

Prof. Dr. H.N. Mülthei
Johannes Gutenberg - Universität
Fachbereich Mathematik
Saarstrasse 21
D-6500 Mainz
West-Germany

Dr. B. Schorr
CERN
CH-1211 Geneva 23
Switzerland

Fig 1: Qualitative behaviour of the K-functions

Fig. 2 : $y = \sin x$

Fig. 3 : $y = \sin x$, $\tau_{opt} = 0.6200$

Fig. 4 : $y = x \cdot \sin 2 \pi x$

Fig. 5 : $y = x \cdot \sin 2 \pi x$, $\tau_{opt} = 0.5507$