Ref.TH.2988-CERN

GENERAL PARAMETERIZATION OF SCALE BREAKING

I.S. Barker

Preston Polytechnic

C.S. Langensiegen

University of Manchester

G. Shaw^{*)}

CERN--Geneva

ABSTRACT

A general parameterization of non-singlet structure functions is developed and used to analyse present data in the context of leading order QCD. The extension to non-leading effects is also discussed.

^{*)} On leave from University of Manchester.

1. INTRODUCTION AND METHOD

In this paper we present a general analytic parameterization of the scale breaking in deep inelastic lepton scattering predicted by asymptotically free field theories, and use it to extract values of the QCD scale breaking parameter Λ from experimental data. For the moment, we concentrate mainly on a discussion of non-singlet structure functions to leading order in QCD. Higher order terms in $\alpha_s(Q^2)$ and higher twist terms can be incorporated, and we indicate briefly how this can be done. The extension of our method to singlet structure functions, while in principle straightforward, is somewhat more complicated and will be discussed elsewhere.

For an arbitrary structure function $\Delta(x, Q^2)$, the predictions of QCD^{*)} are most easily stated in terms of the moments defined by^{**)}

$$M_i(Q^2) = \int_0^1 dx x^i \Delta(x, Q^2) \quad (1.1)$$

However, these are not the most convenient quantities from a phenomenological point of view. Firstly, experiments with complete kinematic coverage are required at each Q^2 . Secondly, the higher moments especially emphasize the $x \approx 1$ region, where non-asymptotic effects (resonances, higher twist etc) are most important^{***)}. And thirdly, different moments do not represent independent pieces of experimental information, but are highly correlated.

For these and other reasons, it is better to invert Eq. (1.1) and compare the QCD predictions directly with the measured structure functions in those kinematic regions where data are available and where the theory is believed to be valid. Two main approaches have been tried. The first is to make some simple ansatz which has the anomalous dimensions predicted by the theory. Writing

$$t = \ln \left[\ln(Q^2/\Lambda^2) / \ln(Q_0^2/\Lambda^2) \right] \quad (1.2)$$

a popular choice is

^{*)} An up-to-date review of both theory and phenomenology is given by Buras¹⁾.

^{**)} This is frequently written $M_{i+2}(Q^2)$. The notation (1.1) is much more convenient for what follows.

^{***)} For example, at $Q^2 = 9(\text{GeV}/c)^2$, the requirement $W \geq 2 \text{ GeV}$ often used in definitions of the 'scaling region' corresponds to $0 < x \leq 0.74$.

$$\Delta(x, Q^2) = c x^{\beta(t)} (1-x)^{\alpha(t)} \quad (1.3)$$

where $\alpha(t)$ and $\beta(t)$ are linear functions. Buras and Gaemers²⁾ then found that if they enforced QCD behaviour on the first two moments, the higher moments had a Q^2 dependence within 2 per cent of that required. It remains, however, an open question whether the very restrictive form chosen at the reference Q_0^2 is correct.

Alternatively, Abbott and Barnett³⁾ assume a similar form at a reference Q_0^2

$$\Delta(x, Q_0^2) = c x^{\beta} (1-x)^{\alpha} \quad c, \alpha, \beta \text{ constant} \quad (1.4)$$

and deduce the behaviour at other Q^2 by numerical integration of the Altarelli-Parisi evolution equation⁴⁾. However, the results can still only be as good as the initial ansatz at $Q^2 = Q_0^2$, and one lacks an analytic expression enabling the structure function to be easily computed elsewhere.

In this paper, our approach will be to obtain explicit solutions to the evolution equation in the form of a rapidly convergent series of orthogonal polynomials. In principle, any structure function can be described to arbitrary accuracy by such a series. The coefficients at arbitrary Q^2 are related to those at $Q^2 = Q_0^2$ by a simple matrix equation. We thus have an analytic expression at all Q^2 capable of describing arbitrary structure functions according to QCD.

We start by assuming that at $Q^2 = Q_0^2$, the structure function can be described by an expression of the form

$$\Delta(x, Q_0^2) = x^{\beta} (1-x)^{\alpha} \sum_{k=0}^N a_k(Q_0^2) \theta_k^{\alpha\beta}(x) \quad (1.5)$$

where x is the usual scaling variable and $\theta_k^{\alpha\beta}(x)$ are Jacobi polynomials of order k

$$\theta_k^{\alpha\beta}(x) = \sum_{j=0}^k C_{kj}(\alpha, \beta) x^j \quad (1.6)$$

defined to satisfy

$$\int_0^1 dx x^{\beta(1-x)^{\alpha}} \theta_k^{\alpha\beta}(x) \theta_l^{\alpha\beta}(x) = \delta_{kl} \quad (1.7)$$

As N becomes arbitrarily large, Eq. (1.5) is completely general, and in particular the freedom to increase N can compensate for injudiciously chosen values of the constants^{*)} α, β .

Our problem now is to deduce the form of $\Delta(x, Q^2)$ implied by Eq. (1.5) and the appropriate evolution equation for arbitrary Q^2 . To do this, we first expand $\Delta(x, Q^2)$ in an infinite series

$$\Delta(x, Q^2) = x^{\beta(1-x)^{\alpha}} \sum_{k=0}^{\infty} a_k(Q^2) \theta_k^{\alpha\beta}(x) \quad (1.8)$$

We will, of course, truncate this sum at some finite $M > N$; in practice, for sensibly chosen α and β , convergence is rapid as will be seen below. For the moment, we note that the Q^2 dependence is entirely contained in the 'Jacobi moments'

$$\begin{aligned} a_k(Q^2) &= \int_0^1 dx \Delta(x, Q^2) \theta_k^{\alpha\beta}(x) \\ &= \sum_{j=0}^k c_{kj}(\alpha, \beta) M_j(Q^2) \end{aligned} \quad (1.9)$$

obtained by inverting Eq. (1.8), using Eqs. (1.1), (1.6) and (1.7). In first order QCD, the Cornwall-Norton moments are given by the familiar expressions¹⁾

$$M_j(Q^2) = f_j(t) M_j(Q_0^2) \quad (1.10)$$

where
$$f_j(t) = \exp(-\xi_j t / 2\pi b) \quad (1.11)$$

$$b = (33 - 2N_f) / 12\pi \quad (1.12)$$

and for the non-singlet structure functions we consider

^{*)} Alternatively, for N finite, α and β can also be treated as variable parameters (see below).

$$\xi_j = \frac{4}{3} \left(\frac{1}{2} - \frac{1}{(j+2)(j+3)} + 2 \sum_{m=2}^{j+2} \frac{1}{m} \right) \quad (1.13)$$

N_f is the number of active quark flavours. Finally, we use Eq. (1.10) together with the inverse of Eq. (1.6)

$$x^j = \sum_{i=0}^j d_{ji}(\alpha, \beta) \theta_i^{\alpha\beta}(x) \quad (1.14)$$

to rewrite Eq. (1.9) as

$$a_k(Q^2) = \sum_{j=0}^k \sum_{i=0}^j f_j(t) c_{kj}(\alpha, \beta) d_{ji}(\alpha, \beta) a_i(Q_0^2) \quad (1.15)$$

This equation is the analogue of Eq. (1.10) for the Jacobi moments defined in Eq. (1.9). Explicit forms for the coefficients $c_{kj}(\alpha, \beta)$, $d_{ji}(\alpha, \beta)$ are given in the Appendix. Since the Jacobi moment $a_k(Q^2)$ depends only on the k^{th} and lower moments at the reference $Q_0^2 [a_i(Q_0^2), i \leq k]$, this result, together with Eq. (1.7), yields an explicit expression for $\Delta(x, Q^2)$ in terms of the $N + 4$ non-vanishing parameters α, β, Λ and $a_i(Q_0^2)$, $i \leq N$.

Of course, the utility of this expression for $\Delta(x, Q^2)$ depends upon the rapidity of convergence of Eq. (1.8) previously alluded to. Some insight into this can be gained by considering the incorporation of quark model sum rules. For the non-singlet structure functions, these are the Gross-Llewellyn Smith sum rules^{*)}

$$\frac{1}{3} \int_0^1 \frac{dx}{x} x F_3(x, Q^2) = 1 \quad (1.16)$$

and the Gottfried sum rule

$$\int_0^1 \frac{dx}{x} \Delta F_2(x, Q^2) = C \quad (1.17)$$

*) We give the forms appropriate to leading order QCD. For modifications in higher order see Section 3.2.

where xF_3 is used to denote $1/2(xF_3^{vp} + xF_3^{vn})$, and ΔF_2 to denote $F_2^{ep} - F_2^{en}$. The constant $C = 1/3$ on the assumption that the sea quark distribution of the nucleon is isospin symmetric, which should be an adequate approximation in what follows. Representing either of these structure functions by Eq. (1.8), we can transform Eqs. (1.16) and (1.17) into an infinite series

$$\int_0^1 \frac{dx}{x} \Delta(x, Q^2) = \sum_{k=0}^{\infty} w_k(\alpha, \beta) a_k(Q^2) \quad (1.18)$$

where the coefficients $w_k(\alpha, \beta)$ can be found in the Appendix. The convergence of these quark sum rules gives a useful guide to the number of terms in the series Eq. (1.8) to retain in the computation. In practice, we found that for $\alpha = 3$, $\beta = 0.5$, retaining terms up to $\theta_{10}^{\alpha\beta}(x)$ gave the integrals Eqs. (1.16) and (1.17) to within ± 0.001 over the whole Q^2 range studied.

The arrangement of the rest of this paper is as follows: in Section 2 we describe fits to experimental data on xF_3 and ΔF_2 ; in Section 3 we discuss the method further including its extension to non-leading order and draw some conclusions. Finally, the Appendix is devoted to a collection of useful properties of the Jacobi polynomials and some remarks on computation.

2. FITS TO DATA

We now consider fits to xF_3 and to ΔF_2 . Neutrino data and charged lepton data are discussed separately for two reasons.

First, values of Λ extracted from different processes are not strictly comparable in lowest order QCD. This is because the expression for Λ in terms of α_s contains a singularity at $\alpha_s = 0$ ⁵⁾, so that higher orders are needed to give meaning to the perturbation expansion. In higher order, however, one has to specify the renormalization scheme^{*)}. Hence, the values of Λ extracted from the two processes should be regarded as separate effective Λ 's.

Secondly, the nature of the two data sets differs substantially. The SLAC data⁷⁾, which dominate the charged lepton data, are limited by systematic^{**)}

*) See Ref. 6) for a discussion of this point.

**) A useful short review of systematic errors is given by Atwood⁸⁾.

rather than statistical errors. On the other hand, because of the low counting rate, statistical errors are more important for the neutrino data. This in turn means that the data are given in relatively large x bins, and for these data we felt it necessary to integrate over the x bins rather than simply evaluate $\Delta(x, Q^2)$ at the centre.

2.1. xF_3 Fits

The data points considered were the 72 xF_3 measurements of the BEBC-GGM ⁹⁾ and the CDHS ¹⁰⁾ collaborations for which $W > 2$ GeV, and $Q^2 > 2$ (GeV/c)². There is considerable overlap between the kinematic ranges of the experiments and no serious incompatibility between them. We therefore felt justified in fitting both data sets simultaneously. However, fits to the separate sets and fits with the minimum Q^2 value increased will also be considered.

In all the fits discussed, we fixed Q_0^2 at 4.5 (GeV/c)², α at 3.0, β at 0.5 and N_f at 3. We did try fits in which α, β were allowed to vary, but they resulted in no noticeable change in χ^2 , nor did α, β stray far from their original values. The fits did not significantly change on setting $N_f = 4$.

After some experiment, we found that at most $a_0(Q_0^2)$, $a_1(Q_0^2)$ and $a_2(Q_0^2)$ were required to be non-vanishing in order to obtain a good description of the data^{*)}. We also found that, if it was not explicitly imposed, the Gross-Llewellyn-Smith sum rule was automatically satisfied to within five per cent^{**)}, both for QCD fits ($\Lambda \neq 0$) and the free parton model ($\Lambda = 0$). We therefore imposed this sum rule exactly in subsequent fits by calculating $a_2(Q_0^2)$ in terms of $a_0(Q_0^2)$, $a_1(Q_0^2)$. Thus, three parameters are sufficient to describe the data. The last is of course Λ , or alternatively,

$$\alpha_s(Q_0^2) = [b \ln(Q_0^2/\Lambda^2)]^{-1} \quad (2.1)$$

The values of the parameters obtained from fits to all data with $W > 2$ GeV, $Q^2 > 2$ (GeV/c)², and to all data with $W > 2$, $Q^2 > 6$, are given in Table 1.

*) That is, the data at all Q^2 , with $xF_3 \equiv \Delta(x, Q^2)$ calculated for $Q^2 \neq Q_0^2$ using Eq. (1.8) truncated after 10 terms, as discussed in Section 1. This is further discussed in Section 3.1.

***) The normalization errors on the experiments are seven per cent (BEBC-GGM) and eight per cent (CDHS).

The corresponding fits are shown in Fig. 1, and plots of the best fit χ^2 versus Λ ^{*)} are shown in Fig. 2. Although exact scaling fits ($\Lambda = 0$) and QCD fits ($\Lambda \neq 0$) both have satisfactory χ^2 values, the latter are preferred at the 3 s.d. level ^{**) [i.e., $\chi^2(\Lambda = 0) > \chi_{\min}^2 + 9$]. As can be seen from Fig. 3, which shows the Q^2 evolution of the structure function, the crucial region for this is at the larger x values, where QCD gives a much better description of the trend of the data (see Fig. 1).}

Turning to the parameter values, it is reasonable to prefer the larger Λ values resulting from the $Q^2 > 6$ fit, since this should be much less sensitive to non-leading effects which may play a role at low Q^2 . The scope for such effects is considerably restricted by the cut $W > 2$ GeV, which excludes the large x region at low Q^2 , where target mass and higher twist terms may be important ^{***)}. Nonetheless, they may still play a role.

To investigate this further, we adopted the following procedure. Suppose the data are adequately described by lowest order QCD. Then, independent of the lower bound $Q^2 > Q_{\min}^2$ imposed, we ought to arrive at the same parameter values. Non-leading effects would show up as an apparent change in the parameter values. The results of such an investigation are shown in Fig. 4. Clearly, there is no reason to suppose that non-leading effects are important above $Q^2 = 4$ (GeV/c)², but there does seem to be some need for them to describe the data for $2 < Q^2 < 4$ (GeV/c)². However, some caution is required, since we have combined two experiments. As can be seen in Fig. 3, the region $0.2 \lesssim x \lesssim 0.6$ is important for the delineation of QCD scale breaking effects. In this region, the data for $Q^2 < 6$ (GeV/c)² stem entirely from the BEBC-GGM experiment, whereas the higher Q^2 data are dominated by CDHS. Thus, the possibility arises that the effect might be an artefact of small differences in systematic error between experiments, which will not be properly taken into account by a statistical fitting procedure. To investigate this, we refitted the two

^{*)} That is, the best fit χ^2 obtained by minimizing on the other parameters at each fixed Λ .

^{**) The errors on Λ are highly non-parabolic, as can be seen in Fig. 2. Also, χ^2 is sensitive to extremely small changes in Λ near $\Lambda = 0$, largely due to the logarithmic relation between it and the coupling $\alpha_s(Q^2)$. For these reasons, it is in some ways preferable to quote $\alpha_s(Q_0^2)$.}

^{***)} In particular, target mass effects, described by the formalism of Nachtmann¹¹⁾, are largely confined to this excluded region.

experiments separately for $Q_{\min}^2 = 2,4 \text{ (GeV/c)}^2$. There are slight differences in the coefficients $a_{0,1}(Q_0^2)$ between the two experiments, and both experiments give some indication of non-leading effects with the same sign as before. However, in view of the error bars, this is not compelling. It is interesting that, if present with the indicated sign, those effects reduce the scale breaking predicted by QCD alone, so that neglecting them would lead one to slightly underestimate Λ or $\alpha_s(Q_0^2)$. In either case, we prefer to base our estimate of Λ on the fits with $Q_{\min}^2 = 4,6 \text{ (GeV/c)}^2$ suggesting a range of

$$0.25 \leq \alpha_{\text{eff}}(Q_0^2) \leq 0.82 \quad (2.2a)$$

corresponding to

$$0.13 \leq \Lambda_{\text{eff}} \leq 0.91 \quad (2.2b)$$

To sum up, the xF_3 data favour QCD over scaling, and are fully consistent with the Gross-Llewellyn Smith sum rule. There is evidence for non-leading effects for $Q^2 < 4 \text{ (GeV/c)}^2$, but it is by no means compelling. Λ_{eff} is poorly determined, lying in the range (2.2b).

$$2.2. \quad \underline{\Delta F_2(x, Q^2) = F_2^p(x, Q^2) - F_2^n(x, Q^2)}$$

The data set^{*)} fitted was the same as that used in the moments analysis of Duke and Roberts¹²⁾, and has been fully discussed by these authors. It comprises the SLAC electron scattering⁷⁾ and the CHIO muon scattering data¹³⁾, binned into 100 x values at each of eight Q^2 values in the range $2.5 \leq Q^2 \leq 22.5 \text{ (GeV/c)}^2$, giving a total of 367 data points with $W > 2 \text{ GeV}$. In some fits, the Q^2 range was reduced to $6.5 < Q^2 \leq 22.5 \text{ (GeV/c)}^2$ (with 207 data points) to allow for possible non-leading effects, as in the neutrino case.

Samples of the data set are shown in Figs. 5a, 5b and 5c. It is dominated to a very large extent by the SLAC data^{**)}, and is characterized by very large statistical errors in the small x region $0 < x \leq 0.2$, and very small statistical errors in the large x region $0.4 \leq x \leq x_c$, where x_c is the kinematic cut-off corresponding to $W = 2 \text{ GeV}$. It is obviously this latter region which

*) We would like to thank R.G. Roberts for providing this data set.

**) This would not be true for the singlet structure function $F_2^p + F_2^n$ at small x .

primarily determines the parameters in any fit. Consequently, we found that the Gottfried sum rule Eq. (1.17), which is particularly sensitive to the small x region, was often quite badly violated in fits in which it was not explicitly imposed. Imposition of the sum rule therefore provides a useful constraint on the otherwise poorly determined small x region, and in all fits described below it was explicitly imposed.

As in the neutrino case, we fixed Q_0^2 at $4.5 (\text{GeV}/c)^2$, and α, β at 3.0 and 0.5 respectively. If α is also allowed to vary, it takes up values close to 3, with no significant improvement in χ^2 or change in the parameter values. In contrast, the small x data are not sufficiently precise to determine β . For the moment we keep it fixed at 0.5. Fits were then performed with $a_0(Q_0^2)$, $a_1(Q_0^2)$ as parameters, $a_2(Q_0^2)$ fixed by the sum rule and $a_i(Q_0^2) = 0$ for $i > 2$; and with $a_0, a_1, a_2(Q_0^2)$ as parameters, $a_3(Q_0^2)$ fixed by the sum rule and $a_i(Q_0^2) = 0$ for $i > 3$. (That is, with $N = 2, 3$ respectively.) In all case, the series Eq. (1.8) was truncated at $M = 10$ for all $Q^2 \neq Q_0^2$ as discussed in Section 1 and again in Section 3. The resulting χ^2 and parameter values are shown in Table 2, and sample fits are shown in Figs. 5a, 5b and 5c. The quality of the fits is good despite the apparently high χ^2 which takes only statistical errors into account^{*)}. This is illustrated by Fig. 5 and particularly by Fig. 5b for $Q^2 = 7.5 (\text{GeV}/c)^2$, which in terms of χ^2 per data point is the worst of all our fits. As can be seen, most of the χ^2 contribution comes from relatively few points, which lie off the trend of the other data and which are essentially ignored by the fit.

On comparing the fits for a given $Q_{\min}^2$, we see that the fits with $N = 3$ are somewhat improved over those with $N = 2$, albeit not very decisively. On comparing the fits with $Q_{\min}^2 = 2, 6$ for each N value, we see that the parameter values for $N = 2$ suggest little scope for non-leading effects. In contrast, comparison of the more flexible fits with $N = 3$ suggests that they may be more important. More detailed study, similar to that undertaken in the neutrino case, supports the conclusion that non-leading effects play a role at the lowest Q^2 values, at least at the level where fits with $N = 3$ are superior to those with

^{*)} Systematic errors on the SLAC electron data are typically of order five per cent on the proton, deuteron structure functions, whereas those for the CHIO data are of order ten per cent. For a discussion, see Atwood⁸⁾.

$N = 2$. This is confirmed by fits in which β is also allowed to vary^{*)}.

In conclusion therefore, we have fitted the existing data under various assumptions, obtaining Λ values in the range

$$0.27 \leq \Lambda_{\text{eff}} \leq 0.64 \quad (2.3a)$$

corresponding to a coupling constant in the range

$$0.34 \leq \alpha_{\text{eff}}(Q^2) \leq 0.58 \quad (2.3b)$$

at the reference $Q_0^2 = 4.5 \text{ (GeV/c)}^2$. There is evidence in the data for deviations from leading order behaviour at the lowest Q^2 values [always, of course, with $W > 2 \text{ (GeV/c)}^2$]. As in the neutrino case, the sign of the indicated effects is such that restricting the data to the higher Q^2 values [$Q^2 > 6 \text{ (GeV/c)}^2$] in order to reduce their influence favours the larger Λ values in the above range rather than the smaller.

3. DISCUSSIONS AND CONCLUSIONS

3.1. Polynomial Reconstruction and Polynomial Fitting

For any given complete polynomial basis, it is always possible to find the polynomial moments and hence evaluate the structure functions, provided the Cornwall-Norton moments are already known. This technique, which we call polynomial reconstruction, exists in two main forms. The first exploits Bernstein polynomials, which are positive definite and represent an approximate delta function $\delta(x - \bar{x})$, so that the polynomial moment itself corresponds to the structure function averaged over a bin centred around $x = \bar{x}$ ¹⁴⁾. However, the approximate end point behaviour at $x : 0,1$ is not incorporated, so that relatively high order polynomials are needed for accurate reconstruction. An alternative

*) For $Q_{\text{min}}^2 = 6 \text{ (GeV/c)}^2$, varying β in the $N = 2$ fit leads to $\beta = 1.00$, and χ^2 and Λ values similar to those of the $N = 3$ fit with $\beta = 0.5$. No improvement is achieved in the latter by allowing β to vary. In contrast, for $Q_{\text{min}}^2 = 2$ and $N = 2$, the value $\beta = 0.19$ is obtained with a low Λ value (0.27) and an improvement in χ^2 (to 666) stemming mostly from the lowest Q^2 values 2.5, 3.5 (GeV/c)^2 . The fits for $N = 3, Q_{\text{min}}^2 = 2$ are very similar, with $\chi^2 = 664$. In our view, this is evidence for derivations from leading order QCD at low Q^2 , rather than from the low β values obtained. In the absence of low x data to determine β reasonably directly, we feel that it is best to restrain β to the value 0.5 suggested by Regge arguments at low x .

is to choose a set of orthogonal polynomials, and in particular the choice of Jacobi polynomials has been advocated by Parisi and Sourlas¹⁵⁾. They point out that, given the Cornwall-Norton moments^{*}, the structure function can be obtained in principle by using Eqs. (1.8) and (1.9), and that in practice a good approximation should be obtained by retaining relatively few terms in Eq. (1.8), since the behaviour at the boundaries $x \approx 0.1$ can be approximately built into the weight function

$$\omega(x) = x^\beta(1-x)^\alpha \quad (3.1)$$

used to define the polynomials in Eq. (1.7). Indeed, one feature of their discussion which we will return to below is that their polynomial index α is a Q^2 dependent parameter

$$\alpha(t) = \alpha(t_0) + \frac{16(t-t_0)}{33-2N_f} \quad (3.2)$$

chosen so that the i dependence of the Cornwall-Norton moments $M_i(Q^2)$ as $i \rightarrow \infty$ ¹⁶⁾ is automatically incorporated.

This method, or any other form of polynomial reconstruction obtained by changing the weight function $\omega(x)$ presupposes a prior Cornwall-Norton moment analysis of the experimental data, with all the disadvantages that this implies. Our aim is to eliminate these, and our approach is based on noting that, for any set of orthogonal polynomials, one can derive an inverse expansion corresponding to Eq. (1.14). Hence, a QCD prediction for the evolution of the polynomial moments, analogous to Eq. (1.15), is easily derived. This offers two possibilities:

- (a) Polynomial moment analysis analogous to Cornwall-Norton moment analysis in which one fits moments projected out using orthogonal polynomials rather than powers. Such moments are less sensitive to the structure functions near $x = 1$, and allow a much more realistic discussion of errors, since (unlike Cornwall-Norton moments) each non-zero polynomial moment represents an independent, and, in the limit of precise data over the whole range of x , uncorrelated piece of information.
- (b) Polynomial fitting in which the evolution equation (1.15) is combined with the (truncated) series Eq. (1.8) to perform a direct fit to structure function data, in terms of the parameters $A, a_k(Q_0^2)$, $k = 0, 1, \dots, N$. This is the

^{*}) Which, for illustrative purposes, they obtain using a specific form at $Q^2 = 3.5 \text{ (GeV/c)}^2$ with known parameters.

approach adopted in this paper. Jacobi polynomials are particularly appropriate for this purpose^{*)} since the corresponding weight function Eq. (3.1) is such as to lead to rapid convergence. In this respect, the adequacy of the parameterization Eq. (1.5) at the reference $Q^2 = Q_0^2$ for given N, α, β can be investigated by studying the fit quality under various assumptions. The practicality of the method then rests on the accuracy of truncated forms of the exact series Eq. (1.8) over the kinematic range of the data considered. In this respect, one could attempt to improve the convergence of the series, at the cost of complicating the method, by allowing α, β to be Q^2 dependent as in Eq. (3.2) when the evolution equation (1.15) becomes

$$a_k(Q^2) = \sum_{j=0}^k \sum_{i=0}^j f_j(t) C_{kj}(\alpha(Q^2), \beta(Q^2)) d_{ji}(\alpha(Q_0^2), \beta(Q_0^2)) a_i(Q_0^2) \quad (3.3)$$

However, this modification is unnecessary in principle and in practice. At a given Q^2 , the series Eq. (1.8) converges for all^{**)} $0 < x < 1$ for arbitrary α, β ; and for reasonable fixed α, β values, the rate of convergence is sufficient for all practical purposes.

In practice, most of our fits have been performed for $\alpha = 3, \beta = 0.5$, with polynomials upto $\theta_{10}^{\alpha\beta}(x)$ retained. One test of the accuracy of this approximation in reproducing $\Delta(x, Q^2)$ has already been dealt with in Section 1, in our discussion of the sum rules Eqs. (1.16) and (1.17). This test is particularly sensitive to behaviour at small x . In contrast, the Cornwall-Norton moments $M_i(Q^2)$ for large i are sensitive to the large x values. To investigate this, we have numerically evaluated the $i = 15, 20$ moments of our neutrino fits. Although such behaviour is only imposed exactly for $i \leq 10$, we find that plots of $\ln M_{15,20}(Q^2)$ against t are linear, and the measured slopes over the fitted Q^2 range are within one per cent of the QCD predictions. For these and other^{***)} reasons, we therefore conclude that the convergence of the series Eq. (1.8) and the accuracy of truncating it at $M = 10$ is, as stated above, more than sufficient for all practical purposes.

*) And for polynomial moment analysis.

**) But not necessarily at $x = 0, 1$.

***) For example, we have repeated the ΔF_2 fits ($Q^2 > 2, N = 2$) for M in Eq. (1.8) increasing from 8 to 12. The changes in fit and parameters are negligible, the best fit A changing from 0.463 to 0.464 for example.

3.2. Fits and Developments

In Section 2 we have reported the results of fits to the published xF_3 and ΔF_2 data using the above formalism and assuming that the observed scale breaking is described by leading order QCD. The ranges of Λ_{eff} are given in Eqs. (2.2) and (2.3) and are consistent with previous work¹⁷⁾, but with a greater tolerance for lower values of Λ_{eff} . This probably reflects the greater flexibility allowed in the shape of the structure function at the reference Q_0^2 . Since the essential predictions of QCD concern changes in the shapes of structure functions, it is obviously desirable to have a formalism such as ours which can accommodate arbitrary initial configurations in a systematic way.

We also found indications of deviations from leading order behaviour in the range $2 < Q^2 < 6 \text{ (GeV/c)}^2$, $W > 2 \text{ (GeV)}$, although the evidence for this was surprisingly weak. Of course, this does not exclude higher twist terms being substituted for QCD over a limited range of Q^2 , and it is well known that the SLAC data are consistent with this¹⁷⁾. It is important therefore, from the point of view of the analysis of future, more extended, data sets, to consider extensions of the method beyond the leading log approximation of QCD. The discussion falls naturally into two parts.

(a) Second order in $\alpha_s(Q^2)$

As noted in Section 2, the inclusion of such effects is necessary if Λ is to be defined in a process independent manner⁵⁾. This is easily done in the present context by replacing Eq. (1.10) by a second order expression. Adapting the formulae compiled by Buras¹⁾, an appropriate substitution is

$$f_j(t) \rightarrow f_j(t) \left[1 + \eta_j^p \left(\frac{1}{\ln(Q^2/\Lambda^2)} - \frac{1}{\ln(Q_0^2/\Lambda^2)} \right) - \zeta_j \left(\frac{\ln \ln(Q^2/\Lambda^2)}{\ln(Q^2/\Lambda^2)} - \frac{\ln \ln(Q_0^2/\Lambda^2)}{\ln(Q_0^2/\Lambda^2)} \right) \right] \quad (3.4)$$

Here η_j^p depends on second order calculations of anomalous dimensions and of the coupling constant in the chosen renormalization scheme. The index p refers to the type of structure function, and ζ_j depends on the coupling constant alone¹⁾. In addition, the Gross-Llewellyn Smith sum rule Eq. (1.16) is modified to become

$$\frac{1}{3} \int_0^1 \frac{dx}{x} x F_3(x, Q^2) = 1 - [\pi b \ln(Q^2/\Lambda^2)]^{-1} \quad (3.5)$$

with b given by Eq. (1.12). The Gottfried sum rule integral Eq. (1.17) also becomes weakly Q^2 dependent in second order, but in this case the effect is very small, and in most circumstances can be neglected¹⁸⁾.

(b) Target mass and higher twist effects

Since these effects are most important at high x and low Q^2 they should be severely curtailed by the exclusion of resonance region data $W < 2$ (GeV). This is certainly true of target mass effects, which can be explicitly calculated using the formalism of Nachtmann, and which are largely confined to the resonance region. Since the Altarelli-Parisi equations, or any other forms of perturbative QCD, are inapplicable in this region, it can only be included in fits on the basis of duality arguments. In view of the fact that the scale breaking effects one hopes to characterize are typically of order 10 - 15 per cent (see Fig. 3), this does not seem to us a reliable procedure, and it is one of the advantages of the present method that the data in such regions can consistently be excluded from the analysis. However, we note that, if desired, any QCD fit can easily be extrapolated into this region, incorporating target mass effects^{*}), and compared, in some suitable average sense, with the resonance region data.

More serious from our standpoint is the suggestion that other 'higher twist' effects are important outside this region, and may indeed account for much of the observed scale breaking^{17),19)}. For example, Abbott et al.¹⁷⁾ have argued that a reasonable parameterization of such effects is

$$\Delta(x, Q^2) = \Delta^{QCD}(x, Q^2) \left[1 + \frac{\mu_1}{(1-x)Q^2} + \frac{\mu_2}{(1-x)^4 Q^4} \right] \quad (3.6)$$

and shown that scale breaking in the SLAC electron scattering data can be accounted for by QCD alone^{**)} ($\Lambda \neq 0, \mu_1 = \mu_2 = 0$), or 'higher twist' effects alone ($\Lambda = 0, \mu_1 \mu_2 \neq 0$).

^{*}) By using, for example, Eqs. (3.21) - (3.23) of Abbott, Atwood and Barnett¹⁷⁾.

^{**)} Using the Abbott and Barnett³⁾ approach of numerical integration of the Altarelli-Parisi equations.

Whether or not the large Q^{-4} term required in the latter fit is dynamically plausible, it is clear that much smaller higher twist effects could appreciably affect the determination of Λ . Within the present data set, this ambiguity cannot be resolved. However, when data over a wider Q^2 range become available, it will clearly be desirable to incorporate both QCD and higher twist effects into the analysis. This is clearly straightforward in our approach ^{*)} using Eq. (3.6).

We therefore conclude that our method not only offers a flexible and efficient approach to the leading order QCD analysis of structure function data without the disadvantages of other methods discussed in Section 1, but it can also be easily adapted to include the most important non-leading effects. This will obviously be vital in analyses of the more extensive and precise data expected from current experiments.

ACKNOWLEDGEMENTS

It is a pleasure to thank many of our colleagues for useful discussions and help during this work, and in particular Dick Roberts (who also provided us with the charged lepton data set), Mike Pennington and Terry Sloan. We would also like to thank Larry Abbott and John Ellis for reading the manuscript and suggesting several improvements incorporated into the paper. Finally, one of us, (I.S.B.) would like to thank Arthur Clegg and Sandy Donnachie for their hospitality at Lancaster and Manchester Universities while this work was in progress.

^{*)} Or in any other which fits structure function data directly.

APPENDIX : JACOBI POLYNOMIALS

The Jacobi polynomials have been extensively discussed by Szego²⁰⁾, but the normalization and orthogonality interval used there is inconvenient for our purposes. We compare our definitions with some widely used alternatives in Table A.1.

The $\Theta_k^{\alpha\beta}(x)$ are defined by Eq. (1.7) supplemented by the condition that $\Theta_k^{\alpha\beta}(0)$ is positive. It is convenient to define an auxiliary polynomial $H_k^{\alpha\beta}(x)$ by

$$\Theta_k^{\alpha\beta}(x) = N_k^{\alpha\beta} H_k^{\alpha\beta}(x) \quad (A.1)$$

where^{*)}

$$N_k^{\alpha\beta} = \Theta_k^{\alpha\beta}(0) = (\beta+1)_k \left\{ \frac{(2k+\alpha+\beta+1) \Gamma(k+\alpha+\beta+1)}{\Gamma(k+\alpha+1) \Gamma(k+\beta+1) k!} \right\}^{1/2} \quad (A.2)$$

The $H_k^{\alpha\beta}(x)$ are essentially identical to the polynomials used by Courant and Hilbert²¹⁾ (see Table A.1), and satisfy the simple Rodrigues formula

$$(\beta+1)_k H_k^{\alpha\beta}(x) = \frac{1}{x^\beta (1-x)^\alpha} \left(\frac{d}{dx} \right)^k \left\{ x^{\beta+k} (1-x)^{\alpha+k} \right\} \quad (A.3)$$

Equation (A.3) is particularly useful in obtaining the expansion coefficients

$$C_k(\alpha, \beta) = (-1)^j \binom{k}{j} N_k^{\alpha\beta} \frac{(k+\alpha+\beta+1)_j}{(\beta+1)_j} \quad (A.4)$$

of Eq. (1.6), as well as the less well-known expansion coefficients

$$d_{ji}(\alpha, \beta) = (-1)^i (j-i+1)_i N_i^{\alpha\beta} \frac{B(j+\beta+1, i+\alpha+1)}{(\beta+1)_i} \quad (A.5)$$

of Eq. (1.14). To compute the evolution of the Jacobi moments according to Eq. (1.15), we need the matrix

$$P_{ji}^k(\alpha, \beta) = \begin{cases} C_{kj}(\alpha, \beta) d_{ji}(\alpha, \beta) & k \geq j \geq i \\ 0 & \text{otherwise} \end{cases} \quad (A.6)$$

Its non-vanishing elements are

*) The notation $(a)_n \equiv a(a+1)(a+2)\dots(a+n-1)$ as usual.

$$P_{ji}^k(\alpha, \beta) = \frac{N_i^{\alpha\beta}}{N_k^{\alpha\beta}} (-1)^{i+j} \frac{(2k+\alpha+\beta+1)(i+\beta+1)_{k-i}(i+j+\alpha+\beta+1)_{k-i}}{(j-1)!(k-j)!(i+j+\alpha+\beta+1)(i+\alpha+1)_{k-i}} \quad (A.7)$$

Apart from the normalization constants $N_i^{\alpha\beta}$, $N_k^{\alpha\beta}$, this equation contains only simple multiplicative factors. All the square root and Γ factors are absorbed into the $N_i^{\alpha\beta}$. From a computational point of view, it is therefore convenient to rewrite Eq. (1.8) as

$$\Delta(x, Q^2) = x^\beta (1-x)^\alpha \sum_{i=0}^{\infty} \hat{a}_i(Q^2) H_i^{\alpha\beta}(x) \quad (A.8)$$

where

$$\hat{a}_i(Q^2) = N_i^{\alpha\beta} a_i(Q^2) \quad (A.9)$$

Turning to the computation of the polynomials themselves, we found that the power series expansion Eq. (1.6) is numerically unsatisfactory beyond the first few orders because of rounding errors. We therefore generated them via the recurrence relations

$$\begin{aligned} & (k+\alpha+\beta+1)(k+\beta+1)(2k+\alpha+\beta) H_{k+1}^{\alpha\beta}(x) = \\ & (2k+\alpha+\beta+1) [2k(k+\alpha+\beta+1) + (\alpha+\beta)(\beta+1) - x(2k+\alpha+\beta+2)(2k+\alpha+\beta)] H_k^{\alpha\beta}(x) \\ & - k(k+\alpha)(2k+\alpha+\beta+2) H_{k-1}^{\alpha\beta}(x) \end{aligned} \quad (A.10)$$

together with

$$H_0^{\alpha\beta}(x) = 1 \quad H_1^{\alpha\beta}(x) = 1 - \left(\frac{\alpha+\beta+2}{\beta+1} \right) x \quad (A.11)$$

Finally, we turn to the formulae needed for the Gottfried and Gross-Llewellyn Smith sum rules Eq. (1.16) and (1.17). We need to evaluate integrals of the general type

$$I_k(\alpha, \beta, \gamma) = \int_0^1 dx x^\gamma (1-x)^\alpha \theta_k^{\alpha\beta}(x) \quad (A.12)$$

We obtain

$$I_k(\alpha, \beta, \gamma) = (-1)^k N_k^{\alpha\beta} B(k+\alpha+1, \gamma+1) \frac{(\gamma-\beta-k+1)_k}{(\beta+1)_k} \quad (A.13)$$

provided $\gamma > -1$. The $W_k(\alpha, \beta)$ defined in Eq. (1.18) correspond to the special case

$$W_k(\alpha, \beta) \equiv I_k(\alpha, \beta, \beta-1) = N_k^{\alpha\beta} k! \frac{B(k+\alpha+1, \beta)}{(\beta+1)_k} \quad (A.14)$$

TABLE I

	Scaling	QCD $Q^2 \geq 2$ (GeV/c) ²	QCD $Q^2 \geq 6$ (GeV/c) ²
Λ (GeV/c ²)		0.167 ^{+0.175} _{-0.124}	0.55 ^{+0.35} _{-0.32}
$a_0(Q_0^2)$	0.990 ± 0.013	1.138 ^{+0.050} _{-0.049}	1.25 ^{+0.12} _{-0.09}
$a_1(Q_0^2)$	0.105 ± 0.015	-0.048 ^{+0.055} _{-0.061}	-0.18 ^{+0.11} _{-0.15}
$a_2(Q_0^2)$	-0.047	-0.054	-0.047
		Variance Matrix ^{*)}	
$\sigma_{\Lambda\Lambda}^2$		2.32×10^{-2}	1.15×10^{-1}
$\sigma_{\Lambda 0}^2$		7.07×10^{-3}	3.31×10^{-2}
$\sigma_{\Lambda 1}^2$		-8.12×10^{-3}	-4.12×10^{-2}
σ_{00}^2	1.57×10^{-4}	2.39×10^{-3}	9.90×10^{-3}
σ_{01}^2	-6.13×10^{-5}	-2.58×10^{-3}	-1.20×10^{-2}
σ_{11}^2	2.25×10^{-4}	3.23×10^{-3}	1.53×10^{-2}

*) For entries not involving Λ , the errors are exactly parabolic. However, there are noticeable deviations from parabolic behaviour when Λ is not fixed. The above represents a parabolic approximation.

Table 1 : Parameter values and variance matrices for the neutrino fits described in the text. The parameter $a_2(Q_0^2)$ is adjusted so that the Gross-Llewellyn Smith sum rule is exactly satisfied, the tabulated value corresponding to the central values of $a_0, a_1(Q_0^2)$.

TABLE 2

N	2	2	3	3
$Q_{\min}^2 (\text{GeV}/c)^2$	2	6	2	6
Deg. of Freedom	364	204	363	203
χ^2	732	417	714	393
Λ	0.464	0.391	0.325	0.580
$a_0(Q_0^2)$	0.161	0.167	0.157	0.182
$a_1(Q_0^2)$	-0.082	-0.081	-0.079	-0.084
$a_2(Q_0^2)$	0.035	0.028	0.030	0.025
$a_3(Q_0^2)$			0.006	-0.012

Table 2 : Chi-squared and parameter values corresponding to the $\Delta F_2(x, Q^2)$ fits described in the text. In each case, $a_N(Q_0^2)$ is determined by the requirement that the Gottfried sum rule be exactly satisfied. The chi-squared values take into account statistical errors only (see text). Statistical errors on the parameters are available on request, but are small compared to the differences between fits, which gives a more reasonable estimate of the uncertainties. For Λ , they vary from 0.04 to 0.07 depending on the fit.

TABLE A.1

Author	Notation	Interval	Relation to $H_i^{\alpha\beta}(x)$
Szego	$P_i^{(\alpha,\beta)}(x)$	$[-1,1]$	$P_i^{(\alpha,\beta)}(x) = (-1)^i \frac{(\beta+1)_i}{i!} H_i^{\alpha\beta}\left(\frac{x+1}{2}\right)$
Courant and Hilbert	$G_i(p,q,x)$	$[0,1]$	$G_i(p,q,x) = H^{p-q,q-1}(x)$
Abramowicz and Stegun	$G_i(p,q,x)$	$[0,1]$	$G_i(p,q,x) = (-1)^i \frac{(\beta+1)_i}{(\alpha+1)_i} H^{p-q,q-1}(x)$

Table A.1 : Relationship between the Jacobi polynomials defined by Eqs.(1.7) and (A.1) and those of Szegő²⁰⁾, Courant and Hilbert²¹⁾ and Abramowicz and Stegun²²⁾. The latter authors give two definitions, the first coinciding with that of Szegő.

REFERENCES

1. A.J. Buras, Rev. Mod. Phys. 52 (1980) 199.
2. A.J. Buras and K.F.J. Gaemers, Nucl. Phys. B132 (1978) 249.
3. L.F. Abbott and R.M. Barnett, Ann. Phys. (NY) 125 (1980) 276.
4. G. Altarelli and G. Parisi, Nucl. Phys. B126 (1977) 298.
5. M. Bacé, Phys. Lett. 78B (1978) 132.
6. J. Ellis, Proc. 1979 International Symposium on Lepton and Photon Interactions at High Energy, Fermi National Accelerator Laboratory, p.142.
7. A. Bodek et al., Phys. Rev. D20 (1979) 1471;
M. Mestayer et al., SLAC-rep-214 (1979);
J.S. Poucher et al., Phys. Rev. Lett. 32 (1974) 118; SLAC-PUB-1309 (1973);
W.B. Atwood et al., Phys. Lett. 64B (1976) 479; SLAC-PUB-185 (1975);
E.D. Bloom et al., SLAC-PUB-653;
S. Stein et al., Phys. Rev. D12 (1975) 1884;
G. Miller et al., Phys. Rev. D5 (1972) 528.
8. W.B. Atwood, Lectures at the 1979 SLAC Summer Institute, SLAC-PUB 2428 (1979).
9. P.C. Bosetti et al., Nucl. Phys. B142 (1978) 1.
10. J.H. de Groot et al., Z. Phys. C1 (1979) 142.
11. O. Nachtmann, Nucl. Phys. B63 (1973) 237; B78 (1974) 455.
12. D.W. Duke and R.G. Roberts, Nucl. Phys. B166 (1980) 248.
13. B.A. Gordon et al., Phys. Rev. D20 (1979) 2645.
14. F.J. Yndurain, Phys. Lett. 74B (1978) 68;
A. Gonzalez-Arroyo, C. Lopez and F.J. Yndurain, Nucl. Phys. B153 (1979) 161.
15. G. Parisi and N. Surlas, Nucl. Phys. B151 (1979) 421.
16. D.J. Gross, Phys. Rev. Lett. 32 (1974) 1071.
17. L.F. Abbott, W.B. Atwood and R. M. Barnett, Phys. Rev. D22 (1980) 582.
18. W.A. Bardeen, A.J. Buras, D.W. Duke and T. Muta, Phys. Rev. D18 (1978) 3998;
A.J. Buras, FNAL report 80/79-THY (1980).
19. A. Donnachie and P.V. Landshoff, Phys. Lett. 95B (1980) 437.
20. G. Szegő, 'Orthogonal Polynomials', American Mathematical Society Colloquium Publications, Vol. 23 New York 1959.
21. R. Courant and D. Hilbert, 'Methods of Mathematical Physics', Interscience, New York 1953.
22. M. Abramowicz and I.A. Stegun, 'Handbook of Mathematical Functions', Dover, New York 1965.

FIGURE CAPTIONS

- Figures 1a and 1b : QCD (solid line) and scaling (dash-dot line) fits to the BEBC-Gargamelle (solid circles) and CDHS (open circles) data. Also shown is a fit to the data with $Q^2 \geq 6 \text{ (GeV/c)}^2$ (broken line) where it is distinguishable from the $Q^2 \geq 2 \text{ (GeV/c)}^2$ fit.
- Figure 2 : Comparison of χ^2 as a function of Λ_{eff} for fits to the neutrino data with $Q^2 \geq 2 \text{ (GeV/c)}^2$ (solid line) and $Q^2 \geq 6 \text{ (GeV/c)}^2$ (dotted line), with 69 and 54 degrees of freedom respectively.
- Figure 3 : Evolution of the QCD fit to xF_3 with $Q^2(\Lambda = 0.167)$. Curves are shown for $Q^2 = 3, 30$, and 300 (GeV/c)^2 (solid lines) and for $Q^2 = 10$ and 100 (GeV/c)^2 (broken lines).
- Figure 4 : Variation of the fitted parameter values with the lower bound on Q^2 for the combined data set. Also shown are the results of fitting the CDHS data alone (ϕ) and the BEBC-Gargamelle data alone (ϕ).
- Figures 5a, b and c : Sample QCD fits to the $\Delta F_2(x, Q^2)$ data with $Q_{\text{min}}^2 = 2 \text{ (GeV/c)}^2$, $N = 2$ (see text) for $Q^2 = 3.5, 7.5, 12.5 \text{ (GeV/c)}^2$. The limit $0 < x < x_c$ corresponds to the cut $W > 2 \text{ GeV}$.

Fig. 1a

Fig. 1b

Fig. 2

Fig. 3

Fig. 4

Fig. 5a

Fig. 5b

Fig. 5c