Ref.TH.3168-CERN

THE ELECTRIC QUADRUPOLE MOMENT OF THE Ω^-

Jean-Marc Richard

Division de Physique Théorique
IPN, Orsay, France

and

CERN, Geneva, Switzerland

ABSTRACT

We estimate the electric quadrupole moment of the Ω^- within current non-relativistic quark models. With only one-gluon exchange as a source of tensor forces we get typically a value $Q_\Omega = 0.004 \text{ fm}^2$ which is probably very hard to measure experimentally.

1. INTRODUCTION

One-gluon exchange (OGE) is commonly accepted as providing a satisfactory description of the mass splittings within hadron multiplets^{1),2)}. This property is considered as one of the successes of QCD. In fact, our understanding of the spin-dependent forces between quarks is far from being completely satisfactory. In the ground-state multiplets, like the $J/\psi - \eta_c$ system, one tests only the spin-spin components of OGE. Moreover, fitting the N to Δ mass difference in current models requires either a troublesome large α_s ³⁾ or the help of the pionic cloud^{4),5)}. The spin-orbit splitting of P wave mesons and baryons is not described by OGE only and one has also to include the Thomas precession term arising from confinement^{2),6)}. To the order m_Q^{-2} in recoil corrections, there is another spin-dependent force in OGE, namely the tensor component. Its effect in light baryons has been reviewed in Ref. 7). The use of a non-relativistic quark model without pionic corrections in that sector is, however, controversial. In heavy hadron spectroscopy, the strength and the range of the tensor potential are not yet measured in great detail. The knowledge of all spin-triplet P states of charmonium allows one, in principle, to extract the tensor contribution to the mass shifts⁸⁾

$$\langle P | V_T | P \rangle = \frac{5}{72} \{ -2 M(^3P_0) + 3 M(^3P_1) - M(^3P_2) \} \simeq 10 \text{ MeV} \quad (1)$$

However, the S to D matrix elements are only vaguely related to the measured properties of the $J/\psi, \psi', \psi'', \dots$ vector mesons. For the leptonic widths, in particular, the Pietschmann-Thirring-Van Royen-Weisskopf type of formulae $\Gamma \propto |\phi(0)|^2$ does not work too well, so relativistic and/or QCD corrections⁹⁾ shadow the effect of tensor forces. Moreover, the mixing with continuum states, which presumably occurs¹⁰⁾, competes with the tensor force due to OGE to induce some S wave components into the $\psi''(3.772)$ and to explain its leptonic width. In fact, the leptonic decay depends on the wave function at only one point, the origin, i.e., precisely where the non-relativistic potential picture is the most questionable. It is preferable to consider quantities integrated over a large range of the wave function. The analogy with Nuclear Physics is here of great help, as often in constituent quark physics. In the deuteron, the long-range tensor potential due to one-pion exchange is tested with a very good accuracy from the asymptotic D/S ratio and the quadrupole moment¹¹⁾. Other nuclei, like ${}^7\text{Li}$ also provide rich information on the nuclear tensor force. Similarly, the knowledge of the quadrupole deformation of the hadrons in their ground state would be very useful for the phenomenological determination of the tensor interaction between quarks.

Indeed, some years ago, Sternheimer and Goldhaber¹²⁾ suggested measuring the electric quadrupole moment Q_Ω of the Ω^- by studying the hyperfine splitting of the levels of the exotic Ω^- nucleus atoms. In Ref. 12) the authors concluded that $|Q_\Omega|$ should be as large as 1 fm^2 to be detectable. Such a value seems by far too high, with respect to the predictions of standard quark models. For instance, the typical radius of a bag is 1 fm , so a one percent squashing effect would give only $Q_\Omega \approx 0.01 \text{ fm}^2$. On the other hand, conservative values were adopted in Ref. 12) for the experimental resolution of X-ray lines. It has been claimed recently that with better detectors one could lower the threshold close to 0.01 fm^2 ¹³⁾. Since a recent calculation¹⁴⁾ using a crude harmonic oscillator model gives a value at the edge of this experimental limit, it seems worthwhile to study the stability of this prediction with respect to the choice of the parameters for the inter-quark potential. Such a discussion is missing in Ref. 14), as well as any discussion on the validity of several approximations used in the details of the calculation of Q_Ω .

2. QUARK MODELS

The authors of Ref. 14) used the model of Isgur and Karl¹⁵⁾ (hereafter called IK model), which is essentially a harmonic oscillator whose parameters are adjusted by fitting the masses of the light baryons, for which a non-relativistic picture is, of course, rather speculative. Instead of considering the strange quark as the last of the light quarks, another possibility consists of attaching it to the family of the heavy quarks, where potential models have been shown to be rather successful and where non-relativistic approximations are presumably justified.

An illustration of such a strategy has been provided recently in the mesonic sector by A. Martin¹⁶⁾, who fitted the "bcs pairs", i.e., the T , J/ψ , ϕ and F families with a universal potential made of a simple power law $r^{0.1}$ which has very nice scaling properties¹⁷⁾. The success of the fit of the masses and relative leptonic widths of heavy mesons is reinforced by the fact that the quark masses used are compatible with the known values of quantities like $M_D - M_K$. Recently, this potential was applied in the baryonic sector¹⁸⁾ by adopting, with some precautions, the rule

$$V_{q\bar{q}} = \frac{1}{2} V_{q\bar{q}} \quad (2)$$

as an educated guess (hereafter this quark potential is referred to as the MR model). Solving the three-body Schrödinger equation by the method of the hyperspherical expansion led to a mass $M = 1.662 \text{ GeV}$ for the Ω^- (after spin corrections) in rather good agreement with the experimental value $M = 1.672 \text{ GeV}$.

Clearly, the models IK and MR are rather different:

- i) the former is a sharp oscillator, the latter has a quasi-logarithmic, very smooth confinement;
- ii) to fit the Δ -N splitting, the IK model uses the rather large value $\alpha_s \approx 1.7$. The model MR also has a hyperfine term of the contact type, whose strength is adjusted to the splitting $M(J/\psi) - M(\eta_c) = 0.112$ GeV. It corresponds to $\alpha_s \approx 0.4$ if identified with the spin-spin part of OGE. The effect in the mass shift of the Ω^- is the following: the model IK gives $\Delta M = 70$ MeV whereas the MR model gives only $\Delta M = 45$ MeV (in the hyperscalar approximation).

To illustrate the sensitivity of the inter-quark force used we consider a family of power law potentials with various exponents, supplemented by an OGE type of hyperfine corrections, i.e.,

$$V_{ij} = A + B r_{ij}^\beta \quad (3)$$

and

$$V_{ij}^{spin} = \frac{2}{3} \alpha_s \left(\frac{1}{6m_s^2} \right) \left\{ 4\pi \delta^3(r_{ij}) \vec{\nabla}_i \cdot \vec{\nabla}_j + \frac{3}{2r_{ij}^3} (3\vec{\nabla}_i \cdot \vec{r}_{ij} \vec{\nabla}_j \cdot \vec{r}_{ij} - \vec{\nabla}_i \cdot \vec{\nabla}_j) \right\} \quad (4)$$

The strength B is adjusted to give always the same excitation energy for positive parity states with total orbital momentum 2^+

$$E = M(2^+) - M(0^+) \quad (5)$$

when E is calculated in the hyperscalar approximation. Similarly, α_s is tuned in such a way that the spin-spin shift ΔM of the Ω^- is always the same.

For simplicity, the strange quark mass and the values of E and ΔM are taken from the MR model^{15),18)}, i.e.,

$$m_s = 518 \text{ MeV}, \quad E = 653 \text{ MeV}, \quad \Delta M = 45 \text{ MeV} \quad (6)$$

Adopting other values would result in simple changes for Q_Ω , due to the scaling properties of our potentials (3). For any exponent β , if V^{spin} is treated in first order perturbation theory, the following rule holds

$$\frac{Q'_\Omega}{Q_\Omega} = \frac{\Delta M'}{\Delta M} \frac{m_s}{m'_s} \frac{E'^2}{E^2} \quad (7)$$

Note that the deformation of the wave function follows similar rules. Consider, for instance, the IK model with non-strange and with strange quarks. Apart from rounding errors and minor non-linear corrections, the truncated expansion of Ref. 7)

$$\Delta\psi(\Delta^{++}) = 0.20 |^4S_{3/2}^* > - 0.10 |^4D_{3/2} > + 0.07 |^2D_{3/2} > \quad (8)$$

is nothing but the rescaling of the similarly truncated expansion of Ref. 14)

$$\Delta\psi(\Omega^-) = 0.15 |^4S_{3/2}^* > - 0.06 |^4D_{3/2} > + 0.05 |^2D_{3/2} > \quad (9)$$

3. CALCULATION

The quadrupole moment is directly related to the presence of D state impurities in the Ω^- wave function. Let us introduce the standard Jacobian and hyperspherical coordinates

$$\begin{aligned} \vec{\xi}_1 &= \vec{r}_2 - \vec{r}_1 & \vec{\xi}_2 &= (2\vec{r}_3 - \vec{r}_1 - \vec{r}_2) / \sqrt{3} \\ r &= (\vec{\xi}_1^2 + \vec{\xi}_2^2)^{1/2} & \Omega &= (\hat{\xi}_1, \hat{\xi}_2, \text{tg}^{-1} \frac{\xi_1}{\xi_2}) \end{aligned} \quad (10)$$

and expand the wave function.

$$\psi = \sum_n \frac{u^{(n)}(r)}{r^{5/2}} |^4S_{3/2}^{(n)} > + \sum_p \frac{w^{(p)}(r)}{r^{5/2}} |^4D_{3/2}^{(p)} > + \sum_q \frac{z^{(q)}(r)}{r^{5/2}} |^2D_{3/2}^{(q)} > \quad (11)$$

where the $|^{2S+1}\ell_J^{(n)}(\Omega) >$, $n = 0, 1, \dots$ are the successive completely symmetric hyperspherical harmonics (HH) associated with total spin S , total angular momentum ℓ , the coupling of ℓ and S generating the $J^P = 3/2^+$ of the Ω^- . The "grand" angular momentum of these HH is $L = \ell + 2n$. We shall use explicitly only

$$|^4S_{3/2}^{(0)} > = \pi^{-3/2} \chi(3/2)$$

and

$$|{}^4D_{3/2}^{(0)}\rangle = \frac{\sqrt{12}}{\pi^{3/2}} \left\{ \sqrt{\frac{2}{5}} (p_+^2 + \lambda_+^2) \chi(-1/2) - \frac{2}{\sqrt{5}} (p_+ p_3 + \lambda_+ \lambda_3) \chi(1/2) \right. \\ \left. + \sqrt{\frac{2}{15}} (p_3^2 + p_+ p_- + \lambda_3^2 + \lambda_+ \lambda_-) \chi(3/2) \right\} \quad (12)$$

where $p_{\pm} = \mp(p_x \pm ip_y)/\sqrt{2}$ and $\chi(3/2) = \uparrow\uparrow\uparrow$, $\chi(1/2) = 1/\sqrt{3}(\uparrow\uparrow\uparrow + \uparrow\uparrow\downarrow + \uparrow\downarrow\uparrow)$, $\chi(-1/2) = 1/\sqrt{3}(\uparrow\uparrow\downarrow + \uparrow\downarrow\downarrow + \downarrow\uparrow\uparrow)$ are the spin $3/2$ wave functions.

Before spin corrections one has only S state and the hyperscalar approximation ($u^{(n)} = 0$ for $n \geq 2$, $n = 1$ being absent in any case) is exact for the harmonic oscillator and very good elsewhere, as discussed in Ref. 18) for the case $\beta = 0.1$ and in Ref. 19) for the case of a potential derived from the bag model. So, with good accuracy, we keep only

$$\psi_0 = \frac{u_0}{r^{5/2}} |{}^4S_{3/2}^{(0)}\rangle \quad (13)$$

Now, the quadrupole operator

$$Q_{22} = \sum_i q_i (3z_i^2 - \vec{r}_i^2) = \frac{e_s}{2} (3p_3^2 - \vec{p}^2 + 3\lambda_3^2 - \vec{\lambda}^2) \quad (14)$$

connects $|{}^4S_{3/2}^{(0)}\rangle$ only with $|{}^4D_{3/2}^{(0)}\rangle$, i.e., to the first order one can neglect all $L \geq 4$ D states and, of course, any spin $1/2$ (${}^2D_{3/2}$) piece. Changes in the S wave component also act only to second order. This means that we consider the correction

$$\Delta\psi = \frac{w(r)}{r^{5/2}} |{}^4D_{3/2}^{(0)}\rangle \quad (15)$$

which induces the quadrupole moment

$$Q_{22} = \langle Q_{22} \rangle = \frac{e_s}{\sqrt{10}} \int_0^\infty u_0 r^2 w dr \quad (16)$$

$e_s = -1/3$ being the charge of the strange quark. In ordinary perturbation theory, one projects $\Delta\psi$ on the eigenstates w_k of the unperturbed Hamiltonian, i.e.,

$$w(r) = \sum_k C_k w_k$$

$$C_k = \frac{\alpha_s}{3m^2} \frac{32\sqrt{2}}{\pi\sqrt{5}} \frac{\int_0^\infty w_k r^{-3} \mu_0 dr}{E(\psi_0) - E(w_k)} \quad (17)$$

Truncating such a series to the first D state by arguing that the others are suppressed by a large energy denominator or by cancellations due to the nodes in the numerator may be sometimes a very dangerous game, as has been extensively discussed in the case of muonium²⁰⁾. Summing over all hyperradial excitations in Eq. (17) is, in fact, very easy and the same as the procedure which has been used for the muonium²⁰⁾ and whose principle has been known for several years²¹⁾. This results in solving the inhomogeneous equation:

$$w'' - \frac{63}{4r^2} w + m_s [E(\psi_0) - V_0(r)] w = \frac{\alpha_s}{3m_s} \frac{32\sqrt{2}}{\pi\sqrt{5}} \frac{\mu_0}{r^3} \quad (18)$$

where

$$V_0(r) = 3A + \frac{12}{\sqrt{\pi}} B \frac{\Gamma(\frac{3}{2} + \frac{\beta}{2})}{\Gamma(3 + \frac{\beta}{2})} r^\beta \quad (19)$$

is the hyperscalar approximation to the potential (3). Due to the singular r^{-3} nature of the perturbation, it is clear that w cannot be saturated by one component of the unperturbed Hamiltonian, i.e., the expansion (17) is infinite. Indeed, at the origin, each w_k behaves like $r^{9/2}$ whereas the normalizable solution of Eq. (18) vanishes only as $r^{1/2}$. Note, however, that in computing the integral of r^2 in Eq. (16) we have the nice feature that is dominated by the first term ($k = 0$) of the expansion (17). For the harmonic oscillator one has the proportionality relation

$$\mu_0 r^2 \propto w_0 \quad (20)$$

so Q_Ω is exactly saturated by the nodeless D state component. For other exponents β , we have compared numerically the exact value of Q_Ω from Eq. (16) and the $k = 0$ approximation

$$Q^{app} = \frac{e_s}{\sqrt{10}} C_0 \int_0^\infty u_0 r^2 w_0 dr \quad (21)$$

The results in Fig. 1 show that the higher radial components play little role for Q_Ω .

Some comments on this point are in order:

- i) in the Coulombic case, Q is not at all saturated by the first term. In fact, $Q^{app}/Q \approx 0.22$ for a symmetric Coulomb three-body problem (in the hyperscalar approximation) and $Q^{app}/Q = (3^5/5 \times 2^{10}) \approx 0.05$ in the two-body case (orthomuonium). Near the origin the relation (20) always holds. The point is that in a Coulomb or nuclear potential, each state has a different exponential tail $\exp -\sqrt{-mE} r$, whereas in a confining potential, all wave functions have similar asymptotic decrease so the relation (20) holds approximatively up to rather large distances;
- ii) this discussion is somehow similar to that of Jackson²²⁾, who tried to relate $\langle S|V_T|D \rangle$ off-diagonal matrix elements to the diagonal mass-shift $\langle P|V_T|P \rangle$ of P states. Again, there is a simple relation which is exact for the harmonic oscillator and rather good for other types of confinements;
- iii) if one now compares the ${}^4D_{3/2}^{(0)}$ probability $P = \int_0^\infty w^2 dr$ with the value $P^{app} = C_0^2$ obtained with the first term only, one gets for the harmonic oscillator a ratio $P^{app}/P = 0.58$ which shows the importance of the higher components and the possible difficulties with the truncations (8) and (9) made by other authors^{7),14)}.

To close this technical discussion, we note that it would be interesting to include the spin effects better than in first order perturbation theory and, in particular, to incorporate some interplay between spin-spin and tensor forces. The former induces, indeed, some mass shifts which might influence the mixing properties of the latter. In a variational approach, this is tentatively accounted for by diagonalizing a matrix whose elements are obtained by sandwiching the Hamiltonian with very simple S and D state wave functions as in Ref. 14). To keep the matrix reasonably sized one has to truncate the series of excitations. As discussed above, this seems reasonable for the quadrupole moment of the Ω^- , but this could be very dangerous in other cases. Alternatively, one might try to regularize the spin-dependent Hamiltonian²³⁾. A cut-off parameter is therefore needed, which could be fitted elsewhere. Then v^{spin} could be treated non-perturbatively in solving the three-body problem.

4. DISCUSSION

We first notice that there is some sensitivity to the kind of potential used in the calculation, almost a factor of two between the harmonic oscillator ($\beta = 2$) and the logarithmic confinement ($\beta \rightarrow 0$), for given excitation energy E and hyperfine splitting ΔM . This is illustrated in Fig. 2, which plots the reduced quantity $q = (Q_\Omega E^2 m_s / \Delta M)$ which depends only on the exponent β of the confining potential.

Actual models have different predictions for E and ΔM . When one puts all effects together one obtains, for instance, $Q_\Omega = 0.013 \text{ fm}^2$ with model IK, if one uses their values $E = 560 \text{ MeV}$ and $\Delta M = 70 \text{ MeV}$, and only $Q_\Omega = 0.0038 \text{ fm}^2$ with model MR and the corresponding values of E and ΔM given in Eq. (6). Our simple perturbative estimate for model IK has to be compared with the value $Q_\Omega = 0.018 \text{ fm}^2$ obtained in Ref. (14) after some partial diagonalization.

In any case, any model with OGE mediated tensor force only gives a very small positive value for Q_Ω . With such a magnitude, it would be very hard to measure Q_Ω experimentally. Now, a larger quadrupole deformation can occur if there is some departure from the one-gluon exchange¹⁾. At short distances, one expects perturbative QCD to be reliable. However, the quadrupole moment depends also on the long-range interactions, where new kinds of tensor forces can arise. For instance, in a generalization of the S.U.N.Y.^{4),24)} or cloudy²⁵⁾ bag models of the nucleon, one might think that the Ω^- is not purely a sss system but also sometimes a hyperon surrounded by kaons. This is, in fact, similar to the mixing of the charmonia with the continuum states ($D\bar{D}$, $D\bar{D}^*$...) analyzed in detail by the Cornell group¹⁰⁾.

The magnitude of the mesonic effects is presently hard to estimate. Close and Horgan²⁶⁾, for instance, concluded that the charge radius of the neutron $\langle \Sigma e_i r_i^2 \rangle_n \approx -0.10 \text{ fm}^2$ cannot be explained by OGE effects, in contradiction with previous claims²⁷⁾. In an extreme model, $\langle \Sigma e_i r_i^2 \rangle_n$ would be entirely due to pionic effects, i.e., to virtual transitions $n \rightarrow (p \text{ or } \Delta^+) + \pi^- \rightarrow n$. One can now try to extrapolate to the transition $\Omega^- \rightarrow \Xi^0 + K^-$. The K^- being in a relative P wave, there is a factor $-2/5$ between $\langle \Sigma e_i r_i^2 \rangle$ and $\langle \Sigma e_i r_i^2 (3\cos^2\theta - 1) \rangle$ i.e., the induced quadrupole moment is positive. The size of the contribution is rather model dependent. Taking (too) seriously the chiral bag model would result in a suppression factor at least as effective as $e^{-2m_K R} / e^{-2m_\pi R}$ where $R \approx 1 \text{ fm}$ would be some typical bag radius. In this case, the contribution of the K meson cloud to Q_Ω would be negligible. Now the suppression factor could be less dramatic. Remember, for instance, that the large energy shift of the $\Lambda(1405)$ resonance has tentatively been explained by a mixing of the sqq quark

configuration with the $K - N$ channel²⁸⁾, suggesting that the loop corrections involving K mesons are not too much suppressed by the large mass of the K . In any case, measuring the quadrupole moment of the Ω^- or setting a limit to its value would provide very useful constraints on the composite models of baryons.

ACKNOWLEDGEMENTS

I would like to thank T.E.O. Ericson, M. Goldhaber, R.R. Horgan, J.D. Jackson, A. Martin, R.M. Sternheimer and P. Taxil for stimulating discussions.

REFERENCES

- 1) A. De Rújula, H. Georgi and S.L. Glashow, Phys. Rev. D12 (1975) 147.
- 2) For a recent review, see, e.g., H.J. Schnitzer in Proc. XVI Rencontre de Moriond, Les Arcs, France (1981), to be edited by J. Tran Thanh Van.
- 3) For instance, an $\alpha_s = 2.2$ is required in the M.I.T. bag model, T. De Grand, R.L. Jaffe, K. Johnson and J. Kiskis, Phys. Rev. D12 (1975) 2060.
- 4) F. Myhrer, G.E. Brown and Z. Xu, Nucl. Phys. A362 (1981) 317.
- 5) W.N. Cottingham, J.M. Richard and K. Tsu, Nucl. Phys. B179 (1981) 541.
- 6) L.J. Reinders, in Proc. IV. Int. Conf. on Baryon Resonances, Toronto (1980), ed. N. Isgur;
F.E. Close and R.H. Dalitz, in Proc. of the Workshop on Low and Intermediate Energy Kaon-Nucleon Physics, Rome (24 - 28 March, 1980), eds E. Ferrari and G. Violini, D. Reidel publ.)
- 7) N. Isgur, G. Karl and R. Koniuk, "D waves in the nucleon: a test of colour magnetism" (1981).
- 8) J.D. Jackson, in Proc. 1976 Summer Institute on Particle Physics, ed. M.C. Zipf, SLAC report 198 (1976) pp. 147 - 202;
M. Krammer and H. Krasemann, Acta Phys. Austriaca Supp. XXI (1979) 259.
- 9) See, e.g., A. Bradley and A. Khare, Zeit. f. Physik C8 (1981) 13 and references therein. See also, A. Donnachie, R.R. Horgan and P.V. Landshoff, Zeit. F. Physik C10 (1981) 71.
- 10) E. Eichten, K. Gottfried, T. Kinoshita, K.D. Lane and T.M. Yan, Phys. Rev. D21 (1980) 203.
- 11) For an updated study, see T.E.O. Ericson and M. Rosa-Clot, CERN preprint, in preparation.
- 12) R.M. Sternheimer and M. Goldhaber, Phys. Rev. A8 (1973) 2207.
- 13) S.P. Alliluyev and S.S. Gershtein, Moscow Technical Institute preprint (1978).
- 14) S.S. Gershtein and Yu.M. Zinoviev, IHEP preprint, Serpukhov (1980).
- 15) N. Isgur and G. Karl, Phys. Rev. D20 (1979) 1191;
K.T. Chao, N. Isgur and G. Karl, Phys. Rev. D23 (1981) 155.
- 16) A. Martin, Phys. Lett. 100B (1981) 511.
- 17) C. Quigg and J.L. Rosner, Phys. Rep. 56 (1979) 167.
- 18) J.M. Richard, Phys. Lett. 100B (1981) 515.
- 19) P. Hasenfratz, R.R. Horgan, J. Kuti and J.M. Richard, Phys. Lett. 95B (1980) 299;
Ad.T. Aerts and L. Heller, Los Alamos preprint LA-UR-81-1775 (1981).
- 20) V.G. Baryshevsky and S.A. Kuten, Phys. Lett. A67 (1978) 355;
J.M. Richard, Phys. Lett. B82 (1979) 415;
D.S. Beder, Phys. Rev. B21 (1980) 3861.

- 21) R.M. Sternheimer, Phys. Rev. 84 (1951) 244;
C. Schwartz, Ann. Phys. 6 (1959) 156.
- 22) J.D. Jackson, Phys. Lett. 87B (1979) 106.
- 23) J.S. Kang and J. Sucher, Phys. Rev. D18 (1978) 2698;
P. Ditsas, N.A. McDougall and R.G. Moorhouse, Nucl. Phys. B146 (1978) 191;
R.K. Badhuri, L.E. Cohler and Y. Nogami, Phys. Rev. Lett. 44 (1980) 1369.
- 24) M. Rho and G.E. Brown, Phys. Lett. 82B (1979) 177.
- 25) S. Theberge, A.W. Thomas and G.A. Miller, Phys. Rev. D22 (1981) 2838;
C.E. De Tar, Phys. Rev. D24 (1981) 762.
- 26) F.E. Close and R.R. Horgan, Nucl. Phys. B185 (1981) 333.
- 27) R.D. Carlitz, S.D. Ellis and R. Sant, Phys. Lett. 68B (1977) 443;
N. Isgur, G. Karl and R. Koniuk, Phys. Rev. Lett. 41 (1978) 1269.
- 28) See, e.g., R.H. Dalitz and J.G. McGinley, Oxford preprint (1980).

FIGURE CAPTIONS

Fig. 1 : Influence of higher radial excitations on the quadrupole moment. The value Q^{app} , computed with the nodeless ${}^4D_{3/2}$ state only, is compared with Q^{ex} , the result of the complete calculation, for several confining potentials $V = r^\beta$.

Fig. 2 : The dimensionless reduced quadrupole moment $q = Q_\Omega E^2 m_S / \Delta M$ as a function of the confining power β .

FIG. 1

FIG. 2