Ref.TH.3117-CERN

SUPERGRAVITY IN SUPERSPACE

Paul Howe

CERN -- Geneva

ABSTRACT

A comprehensive discussion of the formulation of supergravity theories in superspace is presented. After reviewing the general formalism and the linearized theory, we exhibit a complete solution to the Bianchi Identities for all N . This solution still contains a reducible representation of supersymmetry and further constraints are imposed to obtain off-shell conformal supergravity ($N \leq 4$). $N = 1$ and 2 off-shell Poincaré supergravities are rederived from the corresponding conformal theories. On-shell Poincaré supergravity is discussed for all N in superspaces with additional bosonic co-ordinates.

Ref.TH.3117-CERN

7 July 1981

1. INTRODUCTION

Superspace is perhaps the most natural setting for supersymmetric theories and in particular supergravity, although historically most results have been obtained in x space first and only subsequently recast into superspace language. Nevertheless, there are some features of superspace which we believe make its study worth while.

Firstly, since supersymmetry algebras are extensions of space-time algebras it is reasonable that the underlying space on which the fields of the theory are defined should also be extended. In this sense, superspace arises rather naturally and perhaps has some conceptual advantages compared with the component approach. Supergravity theories may then be given a completely geometrical interpretation in terms of the curvature and torsion of curved superspace in a way which is analogous to Einstein's treatment of general relativity. As a consequence of working in this larger space, one also maintains manifest covariance and this has some calculational applications because the supersymmetry algebra is guaranteed to close. Furthermore, the basic equations governing supergravity theories may be stated in a compact and simple form as conditions that certain parts of the torsion tensor should vanish. The consequences of these basic equations are, not surprisingly, rather complicated, but are nevertheless obtainable in a straightforward way by use of the Bianchi Identities which play a somewhat special rôle in superspace geometry as we shall discuss later.

The second and possibly more important feature is that quantum calculations become simpler in terms of supergraphs. This has been demonstrated by the explicit verification that the β function for $N = 4$ supersymmetric Yang-Mills vanishes up to three loops^{1)*}. Calculational rules have also been set up for $N = 1$ quantum supergravity³⁾, but as yet no explicit calculations have been performed with extended superfields. One reason for this is that in order to use supergraphs, it is necessary to formulate the theory first in terms of unconstrained superfields (prepotentials), and this becomes progressively more complicated as N increases. The prepotentials are now known for $N = 2$ Maxwell theory⁴⁾, linearized $N = 2$ Poincaré supergravity⁵⁾ and linearized conformal supergravity ($N \leq 4$)^{6),7)}, and a recent investigation into $N = 2$ Maxwell theory has indicated that there are some technical difficulties involved in the construction of the quantum theory⁵⁾. However, extended superfields have been used to obtain some insight into the structure of extended supergravity; this has been done by constructing higher-order invariants which may serve as counterterm Lagrangians⁸⁾⁻¹⁰⁾.

*¹⁾ This three-loop calculation was actually done first in x space, but with the aid of a computer²⁾.

In view of the fact that the three-loop $N = 4$ Yang-Mills β function is zero where there is certainly a counterterm available, it is clear that explicit calculations need to be done, or other arguments found, if $N = 8$ supergravity is also to exhibit striking quantum features as has been widely conjectured.

The programme for supergravity in superspace is therefore clear: one should construct the full theory with auxiliary fields (i.e., off-shell), solve the constraints in terms of prepotentials and then construct the Feynman rules for the supergraphs. There are, however, many difficulties still to be overcome in this programme and as yet only partial results are known. In this article we shall discuss in some detail the results on the first part of this programme that have been obtained and attempt to state clearly the problems that remain. We shall focus exclusively on "conventional" superspace first introduced by Salam and Strathdee¹¹⁾, and attempt to generalize the approach of Wess and Zumino¹²⁾ to $N = 1$ supergravity to higher N . The basic formalism is reviewed in the next section where we also show how to obtain the component fields and their supersymmetry transformations. Following that we review linearized on-shell Poincaré supergravity and linearized off-shell conformal supergravity. These results and dimensional analysis are then used to determine the constraints on the superspace torsions that we shall employ and it turns out that they are rather similar for both the conformal and Poincaré cases. This fact enables us to solve the Bianchi Identities in a general way for all N and then to specialize to either case. The conformal case is discussed first (Section 5). We are not primarily interested in conformal supergravity for its own sake, but rather because it is a convenient starting point for the construction of off-shell Poincaré theories. This point of view¹³⁾ has been emphasized previously and in Section 6 we discuss how the scheme may be implemented in superspace and rederive $N = 1$ Poincaré supergravity. In Section 7 we apply this technique to $N = 2$ supergravity, recovering firstly the "almost simple" formulation¹⁴⁾ and then the two inequivalent $(40+40)$ dimensional representations^{15),16)} which may be used in Lagrangians. The method we present here is essentially a variant on other approaches previously employed, i.e., compensating multiplets in x space^{13),16)}, precurvatures¹⁴⁾ and "type three" superspace constraints¹⁷⁾⁻¹⁹⁾, but perhaps has the merit that it is clear in principle how to generalize it to $N = 3$ and 4 . In practice, however, it is rather difficult to carry through, chiefly because the natural "compensating" supermultiplets have high spins and are not suitable for constructing representations which may be used to write down a Lagrangian. In Section 8 we switch our attention to on-shell Poincaré theories and show how to construct them for all N by employing additional bosonic co-ordinates. The additional co-ordinates (the same in number as the number of vector fields) are essentially trivial in

the sense that no field depends on them, but they have some formal advantages in that the non-compact global symmetry groups found by Cremmer and Julia²⁰⁾ emerge rather naturally. In Section 9 we give our conclusions and briefly discuss the problems remaining. There are Appendices on notations and conventions, differential forms in superspace and the non-compact global symmetry groups.

2. GENERAL FORMALISM

The superspaces we shall consider are generalizations of ordinary space-time with four bosonic co-ordinates x^m and $2N$ anticommuting co-ordinates $\theta^{\underline{M}}$ which have complex conjugates $\bar{\theta}^{\underline{M}}$. The entire set of co-ordinates is denoted by z^M . Following Ref. 21), we shall assume that there exists a preferred set of frames E^A defined throughout superspace which are related to the co-ordinate one-forms dz^M by the vielbein matrix E_M^A ,

$$E^A = dz^M E_M^A \quad (2.1)$$

The vielbein is invertible and its inverse is E_A^M where

$$E_A^M E_M^B = \delta_A^B \quad (2.2)$$

Under general co-ordinate transformations in superspace the vielbein transforms as a co-vector on the index M ,

$$\delta E_M^A = \xi^N \partial_N E_M^A + (\partial_M \xi^N) E_N^A \quad (2.3)$$

while under tangent space rotations one has

$$\delta E^A = E^B L_B^A \quad (2.4)$$

The matrix L_A^B is taken to have the following form:

$$\begin{aligned} L_{ab} &= -L_{ba} \\ L_{\alpha\beta} &= \delta_{\beta}^i L_{\alpha}^{\beta} + \delta_{\alpha}^{\beta} L^i_j \\ L_{\alpha\beta} &= \frac{1}{2} (G^{ab})_{\alpha\beta} L_{ab} \\ \bar{L}_{\dot{i}\dot{j}} &= -L^j_i \\ L_{\dot{i}\dot{\beta}} &= -\overline{(L_{\alpha\beta})} \end{aligned} \quad (2.5)$$

All other parts of L , e.g., L_{α}^b , are zero so that the tangent space group is $SL(2, \mathbb{C}) \times G$, where G is $U(N)$ or a subgroup thereof [if G is a subgroup of $U(N)$ then there will be further restrictions on L_j^i]. The reason for this choice is that the Poincaré supersymmetry algebra has the following form

$$\begin{aligned} \{Q_{\alpha}^i, \bar{Q}_{\dot{\beta}j}\} &= \delta_j^i P_{\alpha\dot{\beta}} \\ \{Q, Q\} &= [P, Q] = [P, P] = 0 \end{aligned} \quad (2.6)$$

together with the automorphisms of this subalgebra, i.e., $SL(2, \mathbb{C}) \times G$. Here Q_{α}^i are the N supersymmetry generators which are in the fundamental representation of G and transform as spinors under $SL(2, \mathbb{C})$, and $P_{\alpha\dot{\beta}}$ is the generator of translations. Rigid superspace may be formed as the coset space of the entire group divided by $SL(2, \mathbb{C}) \times G$ (just as Minkowski space is the Poincaré group divided by the Lorentz group), and the superspace we have introduced is the natural curved generalization of this. It will turn out that this superspace is also adequate for conformal supergravity even though the superconformal algebra differs from (2.6). We mention in passing that, in practice, the choice of tangent space group is largely a matter of taste. Even in conformal supergravity where there is a genuine $U(N)$ local symmetry, the theory may be written in a superspace with only $SL(2, \mathbb{C})$ in the tangent space group; the local $U(N)$ will then emerge as an additional symmetry of the (constrained) geometry¹⁹⁾. In general, however, the formalism is simplified by the inclusion of the $U(N)$ in the tangent space group since otherwise one has to contend with low dimensional fields which are not present in the multiplets one wishes to discuss (i.e., they correspond to Wess-Zumino gauges for the new symmetries). Another reason for including $U(N)$ explicitly is that it allows one to handle the non-linearities in the dimension zero scalar fields in a relatively simple way.

Consider now a p form which, for illustrative purposes, we take to have one co-vector and one contravariant vector tangent space indices,

$$\phi_B^C = E^A \wedge \dots \wedge E^A \phi_{A \dots A_p, B}^C \quad (2.7)$$

Under (2.5) one has

$$\delta \phi_A^B = -L_A^C \phi_C^B + \phi_A^C L_C^B \quad (2.8)$$

Since L_A^B is a local transformation it is necessary to introduce a connection one-form Ω_A^B in order to exterior differentiate ϕ covariantly. The exterior covariant derivative is

$$D\phi_A^B = d\phi_A^B - (-1)^p \Omega_A^C \wedge \phi_C^B + \phi_A^C \wedge \Omega_C^B \quad (2.9)$$

where Ω transforms inhomogeneously,

$$\begin{aligned} \delta \Omega_A^B &= -dL_A^B - L_A^C \Omega_C^B + \Omega_A^C L_C^B \\ &= -DL_A^B \end{aligned} \quad (2.10)$$

In a co-ordinate basis, $\Omega_A^B = dz^M \Omega_{MA}^B$ and (2.10) becomes

$$\delta \Omega_{MA}^B = -D_M L_A^B \quad (2.11)$$

In view of (2.10), it follows that Ω_A^B has the same algebraic properties as L_A^B , i.e., it is a Lie algebra valued one-form for the algebra $SL(2, \mathbb{C}) \times G$. The exterior derivative of E^A itself is the torsion two-form

$$T^A = DE^A = \frac{1}{2} E^C \wedge E^B T_{BC}^A \quad (2.12)$$

while differentiating ϕ twice one has

$$D^2 \phi_A^B = -R_A^C \wedge \phi_C^B + \phi_A^C \wedge R_C^B \quad (2.13)$$

where R_A^B is the curvature two-form,

$$R_A^B = d\Omega_A^B + \Omega_A^C \wedge \Omega_C^B = \frac{1}{2} E^D \wedge E^C R_{CD A}^B \quad (2.14)$$

Equation (2.13) is the Ricci Identity and is readily extended to any p form ϕ transforming under an arbitrary representation of $SL(2, \mathbb{C}) \times G$. Differentiating (2.12), (2.14) and utilising the Ricci Identity one finds the Bianchi Identities

$$DT^A = E^B \wedge R_B^A, \quad DR_A^B = 0 \quad (2.15)$$

Let us write

$$I^A = E^B{}_A R_B{}^A - D T^A \quad (2.16)$$

$$I_A{}^B = D R_A{}^B \quad (2.17)$$

Then, expanding out in a tangent space basis one has

$$I_{ABC}{}^D = \sum_{(ABC)} \left(R_{ABC}{}^D - D_A T_{BC}{}^D - T_{AB}{}^E T_{EC}{}^D \right) \quad (2.18)$$

$$I_{ABCD}{}^E = \sum_{(ABC)} \left(D_A R_{BCD}{}^E + T_{AB}{}^F R_{FCD}{}^E \right) \quad (2.19)$$

where $\sum_{(ABC)}$ indicates the graded cyclic sum over ABC and $D_A = E_A{}^M D_M$ [the bilinear terms arise from differentiating the basis one-forms in (2.12) and (2.14)].

Before explaining the significance of the Bianchi Identities, we shall first briefly discuss the relation between superspace and components. A straightforward way is, of course, to expand $E_M{}^A$ and $\Omega_{MA}{}^B$ in a power series in θ and $\bar{\theta}$, but although this is practicable in $N = 1$ (and has been investigated in a series of papers²²⁾), it is rather unwieldy for higher N since a real scalar superfield has 2^{4N} components in N extended supersymmetry. Furthermore, the gauge parameter superfields $L_A{}^B$ and $\xi^A (= \xi^M E_M{}^A)$ consist, except for their first components, entirely of Wess-Zumino gauges which have to be fixed if one wishes to make comparisons with components. We shall therefore focus on covariant objects, i.e., the torsion and the curvature, thus avoiding the problem of Wess-Zumino gauges altogether^{*)}. Further we shall follow Ref. 23) and define the components of a superfield S as the spinorial covariant derivatives of S evaluated at $\theta(\bar{\theta}) = 0$. Thus S has components

$$S|_{\theta=0}, D_\alpha S|_{\theta=0}, \bar{D}_{\dot{\alpha}} S|_{\theta=0}, D_\alpha D_{\beta} S|_{\theta=0} \text{ etc.} \quad (2.20)$$

This is clearly equivalent to a straightforward θ expansion but has the advantages that the component fields defined this way have well-defined transformation properties under $SL(2, \mathbb{C})$ and internal symmetry and also that our superfields will

^{*)} This is not quite true because there may be additional invariances (see Section 5).

in general be constrained so that only the first few spinorial derivatives need be evaluated. Higher components will then be vectorial derivatives of lower components or known components of other superfields. There is clearly some freedom in the definition of components in that we may choose to take certain combinations of spinorial derivatives or add in bilinears. For example, if S is a scalar superfield, the combination $i[D_\alpha^i, \bar{D}_{\dot{\alpha}i}]S|_{\theta=0}$ is a real vector field. We shall also follow Ref. 23) in defining the supersymmetry transformations by means of covariant derivatives. Thus for a superfield S with an arbitrary number of tangent space indices one has under general co-ordinate transformations (2.3)

$$\delta S^{\dots} = \xi^A D_A S^{\dots} \quad (2.21)$$

where $\xi^A = \xi^M E_M^A$ is taken to be field-independent and where we have omitted a field-dependent tangent space transformation. Then a supersymmetry transformation is defined by

$$\delta S^{\dots}|_{\theta=0} = \xi^M D_M S^{\dots}|_{\theta=0} - \bar{\xi}^{\dot{M}} \bar{D}_{\dot{M}} S^{\dots}|_{\theta=0} \quad (2.22)$$

Therefore, having constructed the components in the above fashion we automatically have their supersymmetry transformations. For the vielbein itself, (2.3) becomes

$$\delta E_M^A = D_M \xi^A + \xi^B T_{BM}^A \quad (2.23)$$

where we have again omitted a field-dependent Lorentz transformation.

In view of the above discussion, the programme for finding representations of supersymmetry appropriate for supergravity starting from the highly reducible representation furnished by E_M^A and Ω_{MA}^B is now clear: one imposes constraints on the torsions and curvatures and investigates the component contents of the covariant superfields by means of spinorial derivatives. This is where the Bianchi Identities are important; with their aid one can find out the covariant consequences of a given set of constraints which follow automatically from the definitions of the torsion and curvature without having to write out any explicit θ expansions. It is in this context that we shall speak of a "solution" to the Bianchi Identities, and it is of course important to know whether the constraints one has chosen are too strong, either in the sense that they imply unwanted equations of motion or because they lead to inconsistencies (an example of this latter problem would be eleven-dimensional supergravity with a six-form potential instead of the usual three-form one²⁴⁾).

There are two special features of the Bianchi Identities which will be important in our subsequent analysis. The first is that the first identity (2.16) allows one to solve for the curvature in terms of the torsion. This is Dragon's first theorem²⁵⁾ and may be made plausible as follows: from its definition (2.14), R_A^B has the same algebraic properties as Ω_A^B and hence as L_A^B (2.5). This implies that in many of the identities written in index form (2.18), there will be only one contribution from the curvature part, e.g.,

$$I_{\alpha\beta,c}{}^d \Rightarrow R_{\alpha\beta,c}{}^d = D_{\alpha} T_{\beta c}{}^d + D_{\beta} T_{\alpha c}{}^d + D_c T_{\alpha\beta}{}^d + \dots \quad (2.24)$$

Continuing in this vein one arrives at the quoted result. The second feature is that the identities themselves are not all independent. From (2.16) and (2.17) one derives with the aid of (2.12) and the Ricci Identity (2.13),

$$D I^A = E^B{}_A I_B^A \quad (2.25)$$

$$D I_A^B = 0 \quad (2.26)$$

Then if $I^A = 0$, (2.25) implies that $I_A^B = 0$ again because of the tangent space structure. This is Dragon's second theorem²⁵⁾ and states simply that if the first identities are satisfied then there is no further information to be derived from the second identities that cannot be derived from covariant differentiation and use of the Ricci Identity. It has been pointed out²⁶⁾ that this result may be strengthened somewhat and that not all of the first identities are independent either. However, in order to derive an identity for the original Identities, one cannot assume that the original identities are themselves satisfied, so that the restrictions on the number of independent original identities are weaker than was suggested in Ref. 26). Indeed, the analysis of the "identities for the identities" is almost as complicated as the analysis of the identities themselves (since the former are four-form as opposed to three-form equations) and we have therefore not carried out the systematic analysis of precisely which $SL(2, \mathbb{C}) \times G$ irreducible parts of the Bianchi Identities are in fact independent. Nevertheless, the following result is true: in addition to Dragon's second theorem (dependence of the second identities), the dimension two identity, $I_{ab\gamma}{}^{\delta}$, is dependent (except for $N = 1$) and there are no independent identities of dimension greater than two. We should also add that for off-shell representations for higher N not all the information one requires may be obtained from the Bianchi Identities and one is obliged to employ either the Ricci Identity or dependent Bianchi Identities.

By imposing constraints and analyzing the Bianchi Identities, one expects to find the covariant component fields in a given theory, i.e., scalars, spinors, field strengths and auxiliary fields. The gauge fields themselves are however located elsewhere. The vierbein and the gravitino fields may be found at leading order in θ in the vielbein¹²⁾,

$$E_m^a |_{\theta=0} = e_m^a(x) \quad (2.27)$$

$$E_m^\alpha |_{\theta=0} = \psi_m^\alpha(x)$$

and for conformal supergravity $[G = U(N), (SU(N) \text{ for } N = 4)]^*)$ the $(S)U(N)$ gauge fields are the leading components of Ω_m^i . However, in extended Poincaré supergravity there are $\binom{N}{2} + \binom{N}{6}$ $[\binom{N}{2} \text{ for } N = 8]$ Abelian vector fields which cannot be contained in the vielbein or the connection. These gauge fields do not have to be explicitly included in the geometry since supersymmetry will force them to appear via their field strengths in the torsions and curvatures; however, the theory is perhaps rendered more transparent by their inclusion and this may be done in a geometrical way by introducing a number of additional bosonic co-ordinates equal to the number of vector fields⁹⁾. This larger superspace may be thought of as corresponding to the modification of the algebra (2.6) by the introduction of a set of "Abelian charges".

$$\{Q_\alpha^i, Q_\beta^j\} = \epsilon_{\alpha\beta} Z^{ij}, \quad [Z, Z] = 0 \quad (2.28)$$

These Z 's are not genuine central charges (they transform under G) and no field carries such a charge. The description of this extended superspace is very similar to ordinary superspace and will be discussed in more detail in Section 8. An alternative way to accommodate the vector fields is to modify the covariant derivative by the replacement^{19), 27)}

$$D_A \rightarrow D_A + \phi_A \cdot Z \quad (2.29)$$

where the ϕ_A are gauge fields for the Abelian charges. This approach is equivalent to the introduction of extra co-ordinates but the geometrical interpretation is perhaps not quite as clear.

*) We shall actually choose $G = U(N)$ even for $N = 4$, but the $U(1)$ factor in this case is "fake".

3. LINEAR THEORY

Linearized on-shell Poincaré supergravity is described for each N by a single field strength superfield. This result is easily derivable from the full non-linear $N = 8$ superspace theory²⁸⁾ and the details were given in Refs. 8) and 9). We reproduce the results here because we wish to use them to obtain some information on the torsion constraints we shall impose.

The field content of supergravity is well known and is as follows: (the dimensions of the fields are the same as their spins)

$$\begin{aligned}
 \text{Spin } 0 & : A_{ijkl} \\
 \text{Spin } \frac{1}{2} & : \chi_{\alpha ij k}, \chi_{\alpha}^{ijklm} \\
 \text{Spin } 1 & : M_{\kappa\beta ij}, M_{\alpha\beta}^{ijklmn} \\
 \text{Spin } \frac{3}{2} & : \psi_{\alpha\beta\gamma\delta} \\
 \text{Spin } 2 & : C_{\kappa\beta\gamma\delta}
 \end{aligned} \tag{3.1}$$

All the fields are in totally antisymmetric representations of $U(N)$ [$SU(N)$ for $N = 8$] and are totally symmetric on their spinor indices. We therefore have one spin 2, N spin, $3/2$ $\binom{N}{2} + \binom{N}{6}$ spin 1, $\binom{N}{3} + \binom{N}{5}$ spin $1/2$ and $2 \times \binom{N}{4}$ spin 0. In $N = 8$ we have only $SU(8)$ and A is self-dual:

$$\bar{A}^{ijkl} = \frac{1}{4!} \epsilon^{ijklmnrs} A_{mnrs} \tag{3.2}$$

In addition, χ (five indices) and M (six indices) are related to χ (three indices) and M (two indices) by the $SU(8)$ ϵ tensor. All fields obey their equations of motion, ($s = \text{spin}$)

$$\begin{aligned}
 \partial_{\alpha}^{\beta} F_{\alpha_1 \dots \alpha_s} &= 0 \\
 \square A_{ijkl} &= 0
 \end{aligned} \tag{3.3}$$

where F is any of χ, M, ψ, C . For spins ≥ 1 (3.3) is also the (x space) Bianchi Identity for these fields which tells us that they are expressible in terms of potentials. We observe that because we are using unitary groups, there is an apparent doubling of the number of vector fields, e.g., in $N = 8$ we have 28 vectors, but $\underline{28}$ is a complex representation in $SU(8)$. However, there is no field $M_{\alpha\beta ij}$, so that the field strengths are self dual. Hence there are only 28 real spin 1 states. It is now straightforward to place these component fields in a single superfield W . For $N \geq 5$, this superfield is a scalar, W_{ijkl} , which is totally antisymmetric on its internal indices (its leading component is A_{ijkl}). It satisfies

$$D_{\alpha}^i W_{jklm} = 2 \delta_{[j}^i \chi_{\alpha klm]} \quad (3.4)$$

$$\bar{D}_{\alpha i} W_{jklm} = \bar{\chi}_{\alpha i jklm} \quad (3.5)$$

and in $N = 8$ is also self-dual

$$\bar{W}^{ijkl} = \frac{1}{4!} \epsilon^{ijklmnpq} W_{mnpq} \quad (3.6)$$

in which case (3.4) and (3.5) are no longer independent. For $N \leq 4$, W has N internal indices on which it is totally antisymmetric and $4 - N$ spinor indices on which it is totally symmetric. It satisfies

$$D_{\alpha}^i W_{\beta_1 \dots \beta_m j_1 \dots j_N} = -N (-1)^N \delta_{[j_1}^i D_{\alpha}^{\beta_1} W_{\beta_1 \dots \beta_m j_2 \dots j_N] k_1 \dots k_m} \quad (3.7)$$

$m = 4 - N$

$$\bar{D}_{\alpha i} W = 0 \quad (3.8)$$

For $N = 4$, W is again a scalar and we need a slightly stronger equation than (3.7), namely

$$D_{\alpha}^i D_{\beta}^j W_{klmn} = +6 \delta_{[k}^i \delta_{l]}^j D_{\alpha}^r D_{\beta}^s W_{mn] rs} \quad (3.9)$$

It is straightforward to verify that the components of these superfields are those given in (3.1) and that the equations of motion (3.3) hold. This is done by repeatedly differentiating with D 's and $\bar{D}$'s and using their commutation relations,

$$\{D_\alpha^i, D_\rho^j\} = 0, \quad \{D_\alpha^i, \bar{D}_{\dot{\rho}j}\} = i\delta_j^i \partial_{\alpha\dot{\rho}} \quad (3.10)$$

The component field content of off-shell conformal supergravity is as follows²⁹⁾ (dimensions in brackets):

$$\begin{aligned} \text{Spin } 0: & A_{ijh} (0); B_{ijh}{}^e (1); D^{ij}{}_{\dot{h}e} (2) \\ \text{Spin } \frac{1}{2}: & \chi_{\alpha ijh} (\frac{1}{2}); \lambda_{\alpha ij}{}^h (\frac{3}{2}) \\ \text{Spin } 1: & M_{\alpha\rho ij} (1); X_{\alpha\rho}{}^i (2) \\ \text{Spin } \frac{3}{2}: & \psi_{\alpha\rho\dot{e}i} (\frac{3}{2}); \phi_{\alpha\rho\dot{e}} (\frac{5}{2}) \\ \text{Spin } 2: & C_{\alpha\rho\dot{\gamma}\delta} (2) \end{aligned} \quad (3.11)$$

A, χ, M, ψ and C are again totally antisymmetric on their $U(N)$ indices and totally symmetric on their spinor indices, while B, λ and X are skew on their lower indices and traceless in $N = 4$. D is skew on both pairs of indices, totally traceless for $N = 4$ and satisfies $D^{ij}{}_{ij} = 0$ for $N = 3$. M is no longer a spin 1 field strength, but rather a non-gauge antisymmetric tensor and X is the field strength tensor for the gauged $U(N)$ [$SU(N)$ for $N = 4$]^{*}). As was pointed out in Ref. 27), these fields may be put into the same superfields as in the Poincaré case for $N \leq 4$, with the difference that (3.7) and (3.9) are dropped. It is necessary to impose a further constraint²⁹⁾, however, to ensure that X, ψ, ϕ and C satisfy the relevant (conformal) Bianchi Identities which are

$$\begin{aligned} \partial_\alpha{}^{\dot{\rho}} \bar{X}_{\dot{\rho}j}{}^i &= \partial_\alpha{}^{\dot{\rho}} X_{\alpha\rho}{}^i{}_j \\ \partial_\alpha{}^{\dot{\rho}} \partial_{\dot{\rho}}{}^{\dot{\gamma}} \bar{\psi}_{\dot{\gamma}i}{}^j &= \partial_\alpha{}^{\dot{\rho}} \partial_{\dot{\rho}}{}^{\dot{\gamma}} \phi_{\alpha\rho\dot{\gamma}}{}^i{}_j \\ \partial_\alpha{}^{\dot{\rho}} \partial_{\dot{\rho}}{}^{\dot{\gamma}} \bar{C}_{\dot{\gamma}i}{}^j &= \partial_\alpha{}^{\dot{\rho}} \partial_{\dot{\rho}}{}^{\dot{\gamma}} C_{\alpha\rho\dot{\gamma}\delta} \end{aligned} \quad (3.12)$$

^{*}) $N = 4$ linearized conformal theory is globally $U(4)$ covariant however.

For $N \geq 2$, this constraint is simply that D^{ij}_{kl} is real while for $N = 1$ it is necessary to impose the Bianchi Identity for the field strength, $X_{\alpha\beta}$. Hence conformal supergravity is described off-shell by a chiral superfield $W_{(\alpha_1, \dots, \alpha_m)}$ $[i_1, \dots, i_N]$ ($m = 4 - N$) satisfying the following additional constraints*)

$$\left. \begin{aligned} N=1: \quad \partial_\alpha{}^\lambda \bar{D}^{\dot{\alpha}} \bar{W}_{\lambda\dot{\beta}\dot{\gamma}} &= \partial_\alpha{}^\lambda D^{\dot{\beta}} W_{\lambda\dot{\beta}\dot{\gamma}} \\ N=2 \quad D_\alpha{}^i D_\beta{}^j W_{kl}^{\alpha\beta} \\ N=3 \quad D_{(\alpha}{}^i D_{\beta)}{}^j D^{\alpha m} W_{klm}^\beta \\ N=4 \quad D_{(\alpha}{}^i D_{\beta)}{}^j D^{\alpha m} D^{\beta n} W_{klmn} \end{aligned} \right\} \text{real} \quad (3.13)$$

We remark parenthetically that the conformal superfields may be derived very simply from supercurrents (indeed, this is the method that was first used to find the $N = 4$ theory²⁹⁾). For example, $N = 4$ Maxwell theory is described by a self-dual scalar superfield, W_{ij} ³¹⁾,

$$\bar{W}^{ij} = \frac{1}{2} \epsilon^{ijkl} W_{kl} \quad (3.14)$$

satisfying

$$D_\alpha{}^i W_{jk} = -\frac{2}{3} \delta_{[j}^i D_\alpha{}^l W_{k]l} \quad (3.15)$$

from which the equations of motion follow. The supercurrent is⁷⁾

$$J^{ij}{}_{kl} = \bar{W}^{ij} W_{kl} - \frac{1}{6} \delta_{[kl}^{ij} \bar{W}^{mn} W_{mn} \quad (3.16)$$

and is real and totally traceless by construction. It couples to the conformal prepotential V^{ij}_{kl} via a Noether coupling,

$$I_{\text{Noether}} = \int d^4x d^8\theta d^8\bar{\theta} J^{ij}{}_{kl} V^{kl}{}_{ij} \quad (3.17)$$

*) The $N = 1$ theory has been known for some time³⁰⁾.

From (3.14) and (3.16), J is in the real $\underline{20}$ of $U(4)$ and by (3.15) only transforms into a (complex) $\underline{20}$. Equation (3.17) is therefore invariant under the gauge transformation

$$\delta V^{ij}_{kl} = D^m_{\alpha} \Lambda^{\alpha ij}_{klm} - \bar{D}_{\dot{\alpha} m} \bar{\Lambda}^{\dot{\alpha}}_{kl}{}^{ijm} \quad (3.18)$$

where Λ is in the $\underline{60}$ representation of $U(4)$. In terms of V the field strength W_{ijkl} is given by

$$W_{ijkl} = \bar{D}_{\dot{\alpha} i} \bar{D}_{\dot{\beta} j} \bar{D}_{\dot{\gamma} k} \bar{D}_{\dot{\delta} l} \bar{D}^{(\dot{\gamma}}_{\dot{m}} \bar{D}^{\dot{\beta}}_{\dot{n}} \bar{D}^{\dot{\delta}}_{\dot{r}} \bar{D}^{i)}_{\dot{s}} D^m_{(\alpha} D^n_{\beta)} D^{\alpha r} D^{\beta s} V^{rs}_{pq} \quad (3.19)$$

The action for linearized conformal supergravity takes a very simple form in superspace²⁷⁾:

$$I = \int d^4x D^{i_1}_{\alpha_1} \dots D^{i_N}_{\alpha_N} D^{\alpha_1 j_1}_{\dots} \dots D^{\alpha_N j_N}_{\dots} W_{\beta_1 \dots \beta_m i_1 \dots i_N} W^{\beta_1 \dots \beta_m}_{j_1 \dots j_N} + c.c. \quad (3.20)$$

Using the prepotentials, one can straightforwardly derive the equations of motion which are for $N \geq 2$ that the D field is zero and for $N = 1$ that the spin 1 equation of motion is satisfied.

Let us now turn to the problem of finding the geometrical superspace constraints. The linearized torsions are given by

$$\begin{aligned} T_{AB}{}^C &= D_A H_B{}^C - (-1)^{AB} D_B H_A{}^C + t_{AB}{}^D H_D{}^C \\ &\quad - H_A{}^D t_{DB}{}^C + (-1)^{AB} H_B{}^D t_{DA}{}^C \\ &\quad + \Omega_{AB}{}^C - (-1)^{AB} \Omega_{BA}{}^C \end{aligned} \quad (3.21)$$

where we have expanded the full vielbein into a flat part E^0 plus the linearized field, H

$$E_M{}^A = E^0_M{}^A + E^0_M{}^B H_B{}^A \quad (3.22)$$

and similarly for the linearized connection $\Omega_{A,B}{}^C$. $D_A = (\partial_{\alpha\dot{\alpha}}, D^i_{\alpha}, \bar{D}_{\dot{\alpha} i})$ is the flat space covariant derivative satisfying (3.10) and $t_{AB}{}^C$ is the flat space torsion which is zero except for $t_{\alpha\dot{\beta}}{}^C$,

$$t_{\alpha\dot{\beta}}^c = -i \delta_j^i (\sigma^c)_{\alpha\dot{\beta}} \quad (3.23)$$

The linearized gauge transformations are

$$\begin{aligned} \delta H_A^B &= D_A \xi^B + \xi^C t_{CA}^B + L_A^B \\ \delta \Omega_{AB}^C &= -D_A L_B^C \end{aligned} \quad (3.24)$$

where ξ^A is the parameter of general co-ordinate transformations and L_A^B has the form given in Eq. (2.5), with $G = U(N)$ except for $N = 8$ where it is $SU(8)$.

We treat the conformal case first. At dimension zero, there are no fields except in $N = 4$ where we have a complex scalar. However, although one can have one real scalar in $T_{\alpha\dot{\beta}}^c$ it corresponds to a conventional constraint, i.e., we can always redefine a part of the vielbein by adding on a tensor such that this scalar is no longer present in the torsion. At dimension zero we therefore choose

$$T_{\alpha\dot{\beta}}^c = T_{\alpha\dot{\beta}}^c = 0 \quad (3.25)$$

For $N \leq 3$, we are now nearly finished. Using all possible dimension $1/2$ conventional constraints together with the Bianchi Identities $I_{\alpha\beta\gamma}^d, I_{\alpha\beta\dot{\gamma}}^d$ one finds (at dimension $1/2$)

$$T_{\alpha\dot{\beta}}^{\gamma} = T_{\alpha\dot{\beta}}^c = T_{\alpha\dot{\beta}}^{\dot{\gamma}} = 0 \quad (3.26)$$

$$T_{\alpha\dot{\beta}}^{\dot{\gamma}} = \epsilon_{\alpha\dot{\beta}} \bar{\chi}^{ijk\dot{\gamma}} \quad (3.27)$$

When we have imposed the dimension 1 conventional constraints (corresponding to $\Omega_{ab}^c, \Omega_{a\dot{b}}^{\dot{c}}$), this set of equations will turn out to describe off-shell conformal supergravity for $N \leq 3$ completely. For $N = 4$, however, things are slightly more complicated. This is partly because we have chosen a manifest $U(4)$ covariant formalism [since the linearized theory is globally $U(4)$ covariant], but we only have gauge fields corresponding to $SU(4)$. The extra local $U(1)$ in the tangent space group is therefore "fake" and it is not legitimate to choose conventional constraints corresponding to the $U(1)$ part of the connection.

It may seem odd that one is allowed to choose the $SL(2, \mathbb{C})$ connection which is also in a sense "fake" and not the $U(1)$, but the reason is as follows: for the vector part of $SL(2, \mathbb{C})$ we may for example choose $T_{ab}^c = 0$. From (3.21) one then has

$$\Omega_{a,bc} = D_b H_{ac} - D_c H_{ab} \quad (3.28)$$

i.e., $\Omega_{a,bc}$ is determined in terms of H_{ab} . For the $U(N)$ connection on the other hand, we can choose $T_{a\beta}^{j\beta} = 0$ from which one finds

$$2\Omega_a^{j\beta} = D_{\beta}^j H_a^{\beta} - D_a H_{\beta}^{j\beta} \quad (3.29)$$

But from (3.24) we can always choose a gauge in which $H_{\beta}^{j\beta}$ is zero at $\theta = 0$. Hence (3.29) serves the purpose of placing the $U(N)$ gauge field within the θ expansion of H_a^{β} and does not determine it in terms of known fields. Since the $U(1)$ in $N = 4$ is "fake", the corresponding connection will have to be determined in terms of other fields so it is clear that using it to determine conventional constraints is tantamount to choosing it twice. Indeed, in contradistinction to the $SL(2, \mathbb{C})$ case, the linearized $U(1)$ part of the curvature must vanish and hence in $N = 4$ we must impose

$$R^i_i = 0 \quad (3.30)$$

(This equation can equally well be stated in terms of torsion constants.) Having said this, there is still the possibility of having an additional dimension 1/2 spinor in a four dimensional representation in $T_{\alpha\beta}^{\dot{Y}}$ (and also in $T_{\alpha\beta}^{\dot{Y}}$). However, it is an inequivalent $U(4)$ representation to $\chi_{\alpha ijk}$ and so we set $\tilde{t}$ equal to zero. Therefore, $N = 4$ conformal supergravity will be described by Eqs. (3.25) - (3.27) and (3.30) (not all parts of the latter are independent) together with the additional dimension 1 constraints corresponding to Ω_{ab}^c and Ω_a^j (traceless). One may well ask where the dimension zero scalars have gone. From the identity $I_{\alpha\beta\gamma}^{\delta}$ one finds

$$D_{\alpha}^i \bar{\chi}_{\beta}^{jkl} = D_{\alpha}^i \bar{\chi}_{\beta}^{jkl} = 1i \bar{P}_{\alpha\beta}^{jkl} \quad (3.31)$$

A consequence of (3.30) is that P is actually the divergence of a scalar

$$\bar{P}_{\alpha\beta}^{ijkl} = \partial_{\alpha\beta} \bar{W}^{ijkl} \quad (3.32)$$

Furthermore, W_{ijkl} is chiral, so that we recover the original superfield (3.19). We observe that since W only occurs through its derivatives in the torsions (and in the θ expansion of the vielbein), the linearized geometry will be unchanged by constant shifts of the field W and this is a signal of a non-compact global symmetry group in the non-linear theory, which is known to be $SU(1,1)$ ²⁹⁾.

We now switch our attention to on-shell Poincaré supergravity. In this case we will have the same problems as in $N = 4$ conformal supergravity, only now the argument applies to the entire $(S)U(N)$ group. This means that the number of conventional constraints we can choose is limited and we will have to rely more on dimensional analysis. Examining the list of fields present in on-shell Poincaré theory given in (3.1) we again see that the only possible choices for the dimension 0 and $1/2$ torsions are (3.25) - (3.27). Furthermore, the entire linearized $(S)U(N)$ curvature is zero

$$R^i_j = 0 \quad (3.33)$$

For $N \leq 3$ this completes the discussion (apart from choosing Ω_{ab}^c) and for $N > 4$ we again find as a consequence of our equations and the Bianchi identities the equations (3.31) and (3.32). The field W_{ijkl} now satisfies (3.4) (3.5) [(3.8) and (3.9) for $N = 4$] so that we indeed have the right constraints. We notice that although the fields X_{α}^{ijklm} , $M_{\alpha\beta}^{ijklmn}$ and A_{ijkl} of (3.1) do not appear directly in the torsions, nevertheless supersymmetry forces them to appear elsewhere. However, in the non-linear case it will be more convenient to enlarge the superspace as discussed in the last section and this we shall do in Section 8. As in $N = 4$ conformal supergravity, the fact that W_{ijkl} appears only via derivatives is again a signal of a non-compact global symmetry group and this is to be expected in the light of the work of Cremmer and Julia²⁰⁾.

To conclude this section, we note that the method we have employed here to find the constraints does not rely on the details of the component results, one only needs to know the component fields which may be determined by representation theory. Having said this, we should remark that an equally valid point of view is to build up superspace from explicit component results. This technique has been used to find the constraints for off-shell $N = 1$ ^{32),33)} and $N = 2$ ³⁴⁾ Poincaré supergravity, $N = 1$ conformal supergravity³⁵⁾, on-shell $N = 2,3$ de Sitter supergravity³⁶⁾ and on-shell eleven dimensional supergravity³⁷⁾ (which was also found by the method we have used here³⁸⁾).

4. THE SOLUTION TO THE BIANCHI IDENTITIES

In the last section we have seen that the constraints on the torsions for on-shell Poincaré supergravity and off-shell conformal supergravity are very similar and we shall exploit this fact in order to solve the Bianchi Identities in some generality. In terms of covariant superfields, the description of the general superspace is relatively simple; there is a set of dimension 1 superfields, together with a superfield W whose form depends on N ,

$$G_{\alpha\dot{\beta}j}^i, H_{\alpha\dot{\beta}j}^i, N_{\alpha\beta}^{ij}, S^{ij}$$

$$W = \begin{cases} \chi_{\alpha ijk} & (\dim \frac{1}{2}), \quad N \geq 3 \\ M_{\alpha\beta ij} & (\dim 1), \quad N = 2 \\ \psi_{\alpha\beta\gamma} & (\dim \frac{3}{2}), \quad N = 1 \end{cases} \quad (4.1)$$

where W is symmetric on its spinor indices and skew symmetric on its internal indices. The fields G and H are Hermitean and for the conformal case we shall be able to use the $\Omega_a^j{}_k$ constraint to set them equal to each other, but in on-shell Poincaré supergravity they will turn out to be bilinears in the spin 1/2 fields χ . $N_{\alpha\beta}^{ij}$ is symmetric on α, β and skew symmetric on ij and S^{ij} is symmetric. Both of these fields will again be bilinears in on-shell Poincaré, but in the off-shell conformal theory they will turn out to correspond to components of the scale parameter superfield and are therefore not part of the multiplet (3.11). They will, however, be important for off-shell Poincaré theory.

The full non-linear dimension 0 and 1/2 constraint we shall adopt are exactly the same in form as the linearized ones given in the previous section. This is a consequence of the fact that we are explicitly gauging $U(N)$. They are

dim 0

$$T_{\alpha\beta}{}^c = 0 \quad (4.2)$$

$$T_{\alpha\dot{\beta}}{}^c = -i \delta_j^i (\sigma^c)_{\alpha\dot{\beta}} \quad (4.3)$$

dim 1/2

$$T_{\alpha\beta}^{\gamma} = T_{\alpha\dot{\beta}}^{\dot{\gamma}} = T_{ab}^c = 0 \quad (4.4)$$

$$T_{\alpha\beta}^{\dot{\gamma}} = \epsilon_{\alpha\beta} \bar{\chi}^{ijk\dot{\gamma}} \quad (4.5)$$

As explained in Section 3, these are not all independent and some of them correspond to conventional constraints. Furthermore, the dimension 1/2 identities $I_{\alpha\beta\gamma}^d$ and $I_{\alpha\dot{\beta}\dot{\gamma}}^d$ are satisfied by (4.2) - (4.5). We shall not impose Eqs. (3.30) and (3.33) at this stage, however, and we shall also assume that G is $U(N)$ without restriction. The only additional constraint we shall impose is the conventional one for Ω_{ab}^c ; we take

$$T_{ab}^c = 0 \quad (4.6)$$

We shall now systematically exploit the consequences of Eqs. (4.2) - (4.6) that follow from the Bianchi Identities. At dimension 1 one has the following identities

$$I_{\alpha\beta\gamma}^{\delta}, I_{\alpha\dot{\beta}\dot{\gamma}}^{\dot{\delta}}, I_{\alpha\beta c}^d \quad (4.7)$$

$$I_{\alpha\dot{\beta}\gamma}^{\dot{\delta}}, I_{\alpha\dot{\beta}c}^d \quad (4.8)$$

$$I_{\alpha\beta\gamma}^{\dot{\delta}} \quad (4.9)$$

From (4.9) one finds

$$D_{\alpha}^i \bar{\chi}_{\dot{\beta}}^{jke} = D_{\alpha}^i [\bar{\chi}_{\dot{\beta}}^{jke}] \quad (4.10)$$

and from (4.7) one obtains

$$T_{\alpha\beta}^{\dot{\gamma}} \rightarrow T_{\alpha\dot{\beta}}^{\dot{\gamma}} = i \epsilon_{\alpha\beta} \bar{M}_{\dot{\alpha}\dot{\gamma}}^{jk} + i \epsilon_{\dot{\alpha}\dot{\gamma}} (\epsilon_{\alpha\beta} S^{jk} + N_{\alpha\beta}^{jk}) \quad (4.11)$$

$$R_{\alpha\beta, \gamma\delta} = -\epsilon_{\alpha\beta} \bar{M}_{\gamma\delta}^{ij} \quad (4.12)$$

$$R_{\alpha\beta, \gamma\delta} = \epsilon_{\alpha\beta} N_{\gamma\delta}^{ij} + (\epsilon_{\alpha\gamma}\epsilon_{\beta\delta} + \epsilon_{\alpha\delta}\epsilon_{\beta\gamma}) S^{ij} \quad (4.13)$$

$$R_{\alpha\beta, \gamma}^{\quad h} = \epsilon_{\alpha\beta} \left(\delta_{\gamma}^i \delta^{jh} - \delta_{\gamma}^j \delta^{ih} - \bar{B}^{ijk} \right) + \delta_{\gamma}^i N_{\alpha\beta}^{jh} + \delta_{\gamma}^j N_{\alpha\beta}^{ih} \quad (4.14)$$

where

$$B_{ijk}^{\quad l} = \frac{1}{2} D_{\alpha}^l \chi_{ijk}^{\alpha} \quad (4.15)$$

and

$$D_{(\alpha} \chi_{\beta)jkl}^i = 3 \delta_{[\alpha}^i \delta_{\beta]}^j M_{\alpha\beta kl} \quad (4.16)$$

For $N \leq 2$, χ is zero, but for $N = 2$ M is not, it is a new independent field. The identities (4.8) imply the following results

$$T_{\alpha\beta}^{\quad \gamma} \rightarrow T_{\alpha\beta, \gamma}^{\quad j} = -i \left\{ \epsilon_{\beta\gamma} G_{\alpha}^j + \epsilon_{\alpha\gamma} G_{\beta}^j + \epsilon_{\alpha\beta} H_{\gamma}^j \right\} \quad (4.17)$$

$$R_{\alpha\beta, \gamma\delta} = \frac{1}{2} \epsilon_{\alpha\gamma} (G_{\beta\delta}^i + H_{\beta\delta}^i) + \frac{1}{2} \epsilon_{\alpha\delta} (G_{\beta\gamma}^i + H_{\beta\gamma}^i) \quad (4.18)$$

$$R_{\alpha\beta, \gamma}^{\quad h} = -J_{\alpha\beta, \gamma}^{ih} + \delta_{\gamma}^i (H_{\alpha\beta}^h - G_{\alpha\beta}^h) - \delta_{\gamma}^h (H_{\alpha\beta}^i + G_{\alpha\beta}^i) - \delta_{\gamma}^i (H_{\alpha\beta}^j + G_{\alpha\beta}^j) + \frac{1}{2} \delta_{\gamma}^h (H_{\alpha\beta}^i + G_{\alpha\beta}^i) \quad (4.19)$$

where

$$J_{\alpha\beta,he}^{ij} = \bar{\chi}_{\dot{\beta}}^{ijm} \chi_{\alpha h e m} \quad (4.20)$$

Hence, except for the case $N = 1$ where $\psi_{\alpha\beta\gamma}$ has dimension $3/2$, we have already obtained all the independent superfields. At dimension $3/2$ we have the following identities

$$I_{\alpha\beta\dot{\gamma}}^{\dot{\delta}}, I_{\alpha\beta\dot{\gamma}}^{\dot{\delta}}, I_{\alpha\beta c}^d \quad (4.21)$$

$$I_{\alpha\beta c}^{\dot{\delta}} \quad (4.22)$$

From (4.21) we find

$$T_{ab}^{\dot{\delta}} \rightarrow T_{\alpha\dot{\beta},\rho\dot{\sigma},\dot{\gamma}}^R = \epsilon_{\alpha\rho} (\bar{\psi}_{\dot{\beta}\dot{\sigma}}^R + \epsilon_{\alpha\dot{\gamma}} \bar{\psi}_{\dot{\rho}}^R + \epsilon_{\dot{\rho}\dot{\gamma}} \bar{\psi}_{\alpha}^R) + \epsilon_{\alpha\dot{\beta}} \psi_{\rho\dot{\sigma}}^R \quad (4.23)$$

The ψ functions are the irreducible parts of the gravitino field strength (so in linearized Poincaré supergravity the spin $3/2$ equations of motion are $\psi_{\alpha}^{\dot{\delta}} = \psi_{\alpha\beta\dot{\gamma}}^{\dot{\delta}} = 0$). The dimension $3/2$ curvatures are found to be

$$R_{\alpha,\rho\dot{\sigma},\dot{\gamma}\dot{\delta}}^i = -\frac{i}{4} (2\epsilon_{\alpha\rho} \psi_{\dot{\gamma}\dot{\delta}}^i + \epsilon_{\alpha\dot{\gamma}} \psi_{\rho\dot{\delta}}^i + \epsilon_{\alpha\dot{\delta}} \psi_{\rho\dot{\gamma}}^i) + \frac{3i}{4} (\epsilon_{\alpha\dot{\gamma}} \epsilon_{\rho\dot{\delta}} + \epsilon_{\alpha\dot{\delta}} \epsilon_{\rho\dot{\gamma}}) \bar{\psi}_{\dot{\rho}}^i \quad (4.24)$$

$$R_{\alpha,\rho\dot{\sigma},\dot{\gamma}\dot{\delta}}^i = i\epsilon_{\alpha\rho} \bar{\psi}_{\dot{\gamma}\dot{\delta}}^i - \frac{i}{4} \epsilon_{\dot{\rho}\dot{\gamma}} (\psi_{\alpha\dot{\sigma}}^i + \epsilon_{\alpha\dot{\sigma}} \bar{\psi}_{\dot{\rho}}^i) - \frac{i}{4} \epsilon_{\dot{\rho}\dot{\delta}} (\psi_{\alpha\dot{\sigma}}^i + \epsilon_{\alpha\dot{\sigma}} \bar{\psi}_{\dot{\rho}}^i) \quad (4.25)$$

$$\begin{aligned}
 2i R_{\alpha, \beta}^i, j_k &= 3 D_{\alpha}^i G_{\beta}^j \dot{\gamma}_k - D_{\alpha}^i H_{\beta}^j \dot{\gamma}_k - \bar{D}_{\beta k} N_{\alpha}^{ij} + \epsilon_{\alpha\beta} \bar{D}_{\beta k} S^{ij} \\
 &+ \epsilon_{\alpha\beta} \epsilon^{ij} \bar{D}_{\alpha k} \bar{M}_{\beta}^{ij} - \delta_k^i \psi_{\alpha\beta}^j - 3 \delta_k^i \epsilon_{\alpha\beta} \bar{\psi}_{\beta}^j \\
 &+ \epsilon_{\alpha\beta} \bar{\chi}_{\beta}^{ij} \bar{S}_{kl} + \epsilon_{\alpha\beta} \bar{\chi}_{\alpha}^{ij} \bar{N}_{\beta kl} - \bar{\chi}_{\beta}^{ij} M_{\alpha\beta kl}
 \end{aligned}
 \tag{4.26}$$

In addition, one finds the following relations among the spinorial derivatives of the dimension 1 superfields,

$$\bar{D}_{(\alpha k} \bar{M}_{\beta}^{ij)} - \bar{\chi}_{(\alpha}^{ij} \bar{N}_{\beta}^{kl)} = 2 \delta_{\alpha}^{[i} \bar{\psi}_{\beta}^{j]} \tag{4.27}$$

$$\begin{aligned}
 D_{(\alpha}^i G_{\beta)}^j \dot{\gamma}_k + D_{\alpha}^i H_{\beta}^j \dot{\gamma}_k - \bar{D}_{\beta k} N_{\alpha}^{ij} \\
 = -\delta_k^i \psi_{\alpha\beta}^j + \frac{1}{2} \delta_k^j \psi_{\alpha\beta}^i - \bar{\chi}_{\beta}^{ij} M_{\alpha\beta kl}
 \end{aligned}
 \tag{4.28}$$

$$\begin{aligned}
 \frac{1}{2} \epsilon^{\alpha\beta} (D_{\alpha}^i G_{\beta}^j \dot{\gamma}_k + D_{\alpha}^i H_{\beta}^j \dot{\gamma}_k) + \bar{D}_{\beta k} S^{ij} + \frac{1}{3} \bar{D}_{\beta k} \bar{M}_{\beta}^{ij} \\
 = \delta_k^i \bar{\psi}_{\beta}^j + \frac{1}{2} \delta_k^j \bar{\psi}_{\beta}^i + \bar{\chi}_{\beta}^{ij} \bar{S}_{kl} - \frac{1}{3} \bar{\chi}_{\beta}^{ij} \bar{N}_{\beta}^{kl}
 \end{aligned}
 \tag{4.29}$$

For $N = 1$, $\bar{\psi}_{\alpha\beta}^i$ is a new field but otherwise it is a derivative of $\bar{M}_{\alpha\beta}^{ij}$ via (4.27).

The other dimension $3/2$ identity (4.22) does not determine any curvatures or torsions, but merely relates the derivatives of the superfields M , N and S . One finds

$$D_{(\alpha}^i N_{\beta}^{jk)} = D_{\alpha}^{[i} N_{\beta}^{jk]} \tag{4.30}$$

$$D_{\alpha}^{(i} S^{jk)} = 0 \tag{4.31}$$

$$n_{\alpha}^{ijk} = -\frac{1}{6} D_{\alpha}^{\beta} \bar{\chi}_{\beta}^{ijk} - 2i g_{\alpha}^{ijk} \tag{4.32}$$

$$\begin{aligned}
 \tilde{n}_\alpha^{ijk} + \frac{1}{2} (D_\alpha^j S^{ki} - D_\alpha^k S^{ji}) &= i \tilde{g}_\alpha^{ijk} + i \tilde{h}_\alpha^{ijk} \\
 m_{\alpha\delta\delta}^{ijk} &= D_\alpha (\delta \bar{\chi}_\delta)^{ijk} - 4i g_{\alpha\delta\delta}^{ijk} + 2i h_{\alpha\delta\delta}^{ijk} \\
 \tilde{m}_{\alpha\delta\delta}^{ijk} &= -i \tilde{g}_{\alpha\delta\delta}^{ijk} - i \tilde{h}_{\alpha\delta\delta}^{ijk}
 \end{aligned} \tag{4.33}$$

where we have introduced the abbreviations

$$\epsilon_{\gamma\delta\epsilon} \bar{\chi}_\delta^{jkl} = (g_{\gamma\delta\delta}^{ijk} + \tilde{g}_{\gamma\delta\delta}^{ijk}) + \epsilon_{\delta\delta} (g_{\gamma}^{ijk} + \tilde{g}_{\gamma}^{ijk}) \tag{4.34}$$

and similarly for $H\bar{\chi}$,

$$\begin{aligned}
 D_\alpha^i \bar{M}_{\delta\delta}^{jk} &= m_{\alpha\delta\delta}^{ijk} + \tilde{m}_{\alpha\delta\delta}^{ijk} \\
 \frac{1}{3} \epsilon^{\alpha\beta} D_\alpha^i N_{\beta\gamma}^{jk} &= n_\gamma^{ijk} + \tilde{n}_\gamma^{ijk}
 \end{aligned} \tag{4.35}$$

The meaning of the splitting of a given field, e.g., n into n^{ijk} and $\tilde{n}^{i,jk}$ is best expressed via Young Tableaux:

$$n_\alpha^{ijk} + \tilde{n}_\alpha^{i,jk} = \begin{array}{|c|c|c|} \hline \square & \square & \square \\ \hline \end{array} + \begin{array}{|c|c|} \hline \square & \square \\ \hline \end{array} \tag{4.36}$$

Finally we have the dimension 2 identities

$$I_{abc}^d, \quad I_{ab\gamma}^{\underline{\delta}} \tag{4.37}$$

$$I_{ab\underline{\gamma}}^{\underline{\delta}} \tag{4.38}$$

However, as we pointed out in Section 2, (4.38) is only independent for $N = 1$ and will not be discussed here. In view of the fact that a complete solution to the $N = 1$ identities has already been given³⁹⁾, this is perhaps a pardonable omission. The constraint (4.6) implies that I_{abc}^d is just the normal x space cyclic identity, so we have

$$R_{ab,cd} \rightarrow R_{\alpha\dot{\alpha},\beta\dot{\beta},\gamma\dot{\gamma},\delta\dot{\delta}} = 2\epsilon_{\dot{\gamma}\dot{\delta}} R_{\alpha\dot{\alpha},\beta\dot{\beta},\gamma\dot{\gamma}} + 2\epsilon_{\gamma\delta} \bar{R}_{\alpha\dot{\alpha},\beta\dot{\beta},\dot{\gamma}\dot{\delta}} \quad (4.39)$$

$$R_{\alpha\dot{\alpha},\beta\dot{\beta},\gamma\dot{\gamma}} = \epsilon_{\alpha\beta} B_{\dot{\alpha}\dot{\beta},\gamma\dot{\gamma}} + \epsilon_{\dot{\alpha}\dot{\beta}} C_{\alpha\beta,\gamma\dot{\gamma}} + \epsilon_{\dot{\alpha}\dot{\beta}} (\epsilon_{\alpha\gamma}\epsilon_{\beta\dot{\gamma}} + \epsilon_{\alpha\dot{\gamma}}\epsilon_{\beta\gamma}) A$$

A, B and C are the irreducible parts of the Riemann tensor and correspond to the curvature scalar, the tracefree Ricci tensor and the Weyl tensor. One has

$$A = \bar{A}, \quad \bar{B}_{\alpha\beta,\dot{\gamma}\dot{\delta}} = B_{\dot{\gamma}\dot{\delta},\alpha\beta} \quad (4.40)$$

where B is symmetric on both pairs of indices, C is totally symmetric. From the remaining identity one finds.

$$\begin{aligned} D_{(\alpha}^k \psi_{\beta\delta\delta)} &= \delta_{\alpha}^k C_{\beta\delta\delta} - \gamma_{\alpha\beta\delta\delta}^k \\ D_{\delta}^k \psi_{\alpha\beta}^{\gamma} &= \frac{1}{12} X_{\alpha\beta}^k{}_{\delta} - \frac{1}{12} \gamma_{\alpha\beta}^k{}_{\delta} + \frac{1}{6} \gamma'_{\alpha\beta}{}^k{}_{\delta} \\ D_{(\alpha}^k \psi_{\beta)} &= \frac{1}{3} X_{\alpha\beta}^k - \frac{4}{3} \gamma_{\alpha\beta}^k - \frac{1}{3} \gamma'_{\alpha\beta}{}^k \\ D_{\alpha}^k \psi^{\alpha} &= 2 \gamma^k{}_{\alpha} - 2 \delta_{\alpha}^k A \\ D_{(\delta}^k \bar{\psi}_{\gamma)\dot{\alpha}\dot{\beta}} &= \delta_{\delta}^k B_{\dot{\alpha}\dot{\beta},\gamma\dot{\gamma}} - \gamma_{\dot{\alpha}\dot{\beta},\gamma\dot{\gamma}}^k{}_{\delta} \\ D_{\delta}^k \bar{\psi}^{\gamma}{}_{\dot{\alpha}\dot{\beta}} &= -2 \bar{X}_{\dot{\alpha}\dot{\beta}}^k{}_{\delta} - 2 \gamma_{\dot{\alpha}\dot{\beta}}^k{}_{\delta} \end{aligned} \quad (4.41)$$

where X is the U(N) field strength,

$$R_{ab}{}^k{}_l \rightarrow R_{\alpha\dot{\alpha},\beta\dot{\beta}}{}^k{}_l = \epsilon_{\alpha\beta} X_{\alpha\dot{\beta}}{}^k{}_l - \epsilon_{\alpha\dot{\beta}} \bar{X}_{\alpha\beta}{}^k{}_l \quad (4.42)$$

and the Y functions are given in terms of known fields by

$$\begin{aligned} Y_{\alpha\beta\gamma\delta l}^k &= -2 N_{(\alpha\beta}{}^{km} M_{\gamma\delta)ml} \\ Y_{\alpha\beta l}^k &= -2i/3 D_{\dot{\alpha}}(\alpha G_{\beta})^{\dot{k}}{}_l - 4/3 G_{\dot{\alpha}}^k(\alpha m G^{\dot{\alpha}}{}_{\beta})^m{}_l \\ &\quad - 1/3 N_{\gamma(\alpha}{}^{km} M^{\gamma}{}_{\beta)ml} - 2/3 N_{\alpha\beta}{}^{km} \bar{S}_{ml} \\ Y_{\alpha\beta l}^{k'} &= -i/2 D_{\dot{\alpha}}(\alpha H_{\beta})^{\dot{k}}{}_l + i/6 D_{\dot{\alpha}}(\alpha G_{\beta})^{\dot{k}}{}_l \\ &\quad - 1/6 G_{\dot{\alpha}}^k(\alpha m G^{\dot{\alpha}}{}_{\beta})^m{}_l - 1/2 G_{\dot{\alpha}}^k(\alpha m H^{\dot{\alpha}}{}_{\beta})^m{}_l - 1/2 H_{\dot{\alpha}}^k(\alpha m G^{\dot{\alpha}}{}_{\beta})^m{}_l \\ &\quad - 1/3 N_{\gamma(\alpha}{}^{km} M^{\gamma}{}_{\beta)ml} - 1/3 N_{\alpha\beta}{}^{km} \bar{S}_{ml} - S^{km} M_{\alpha\beta ml} \\ &= -\frac{i}{4} (D^{\alpha\dot{\alpha}} G_{\alpha\dot{\alpha}}{}^k{}_l + D^{\alpha\dot{\alpha}} H_{\alpha\dot{\alpha}}{}^k{}_l) \\ &\quad + \frac{1}{4} (G_{\alpha\dot{\alpha}}^k G^{\alpha\dot{\alpha}m}{}_l - G_{\alpha\dot{\alpha}}^k H^{\alpha\dot{\alpha}m}{}_l + 3 H_{\alpha\dot{\alpha}}^k G^{\alpha\dot{\alpha}m}{}_l + H_{\alpha\dot{\alpha}}^k H^{\alpha\dot{\alpha}m}{}_l) \\ &\quad - S^{km} \bar{S}_{ml} - 1/2 \bar{M}_{\dot{\alpha}\dot{\beta}}{}^{km} \bar{N}_{ml}^{\dot{\alpha}\dot{\beta}} - 1/6 N_{\alpha\beta}{}^{km} M_{ml}^{\alpha\beta} \\ Y_{\alpha\beta,\gamma\delta l}^{k'} &= i D_{\dot{\alpha}}(\delta G_{\beta})^{\dot{k}}{}_l + i D_{\dot{\alpha}}(\delta H_{\beta})^{\dot{k}}{}_l \\ &\quad + G_{\dot{\alpha}}^k(\delta m G_{\beta})^m{}_l + 3 G_{\dot{\alpha}}^k(\delta m H_{\beta})^m{}_l - H_{\dot{\alpha}}^k(\delta m G_{\beta})^m{}_l \\ &\quad + H_{\dot{\alpha}}^k(\delta m H_{\beta})^m{}_l - \bar{M}_{\dot{\alpha}\dot{\beta}}{}^{km} M_{\gamma\delta ml} - N_{\gamma\delta}{}^{km} \bar{N}_{\dot{\alpha}\dot{\beta} ml} \end{aligned} \quad (4.43)$$

$$\begin{aligned}
 Y_{\alpha\beta\ell}^k &= -\frac{3i}{2} D_{\alpha}(\dot{\alpha} \dot{\gamma}^{\alpha k} \dot{\beta})_{\ell} + \frac{i}{2} D_{\alpha}(\dot{\alpha} H^{\alpha k} \dot{\beta})_{\ell} \\
 &\quad - \frac{3}{2} \dot{\gamma}^{\alpha k}(\dot{\alpha} m \dot{\gamma}^{\alpha m} \dot{\beta})_{\ell} + \frac{3}{2} \dot{\gamma}^{\alpha k}(\dot{\alpha} m H^{\alpha m} \dot{\beta})_{\ell} + \frac{3}{2} H^{\alpha k}(\dot{\alpha} m \dot{\gamma}^{\alpha m} \dot{\beta})_{\ell} \\
 &\quad + \frac{1}{2} H^{\alpha k}(\dot{\alpha} m H^{\alpha m} \dot{\beta})_{\ell} + \overline{M}_{\alpha\dot{\beta}}^{km} \dot{\gamma} \dot{\beta})_{\ell} \dot{\gamma} m \ell \\
 &\quad - \overline{M}_{\alpha\dot{\beta}}^{km} \dot{\gamma} m \ell + S^{km} \overline{N}_{\alpha\dot{\beta}} m \ell
 \end{aligned} \tag{4.43}$$

cont.

This completes our presentation of the solution to the Bianchi Identities based on the constraints (4.2) - (4.6). As it stands, it is somewhat messy although the basic constraints are rather simple and there are only a few independent covariant superfields. The representation of supersymmetry defined here is still reducible and there are further constraints to be imposed before we can disentangle the component field content, and, indeed, if one wished to know exactly what representations we have in our solution it would be necessary to do a lot more calculations - this time using the Ricci Identity. However, there are already some features that are perhaps worth mentioning. Firstly, there are no equations of motion for the supergravity fields (χ, M, ψ, C) . From (4.32), we see that $D_{\alpha} \dot{\gamma}^{\beta-ijk}$ is located as a component of $N_{\alpha\beta}^{ij}$ and from (4.29) and (4.30) a similar statement is true for the spin 3/2 equations of motion, i.e., $\psi_{\alpha\beta\gamma}^i, \overline{\psi}_{\alpha}^i$. From the dimension 2 equations, $B_{\alpha\beta,\gamma\delta}^{\cdot}$ and A are also higher components of the basic superfields (4.1). However, the scalar fields themselves are not present (since $D_{\alpha} \dot{\gamma}^{\beta-ijkl}$ does not have to be a derivative of a scalar) and furthermore the field M is not as yet a field strength for a spin 1 field. For these conditions to be fulfilled, we shall need further constraints, and the presentation we have given has the advantage that we can particularize either to off-shell conformal supergravity or to on-shell Poincaré supergravity by different choices of constraint.

5. CONFORMAL SUPERGRAVITY

Full non-linear conformal supergravity has been studied in a superspace setting in $N = 1$ (40), (41) but without explicit gauging of $U(1)$. The $U(1)$ emerged as an additional invariance group of the torsion constraints and furthermore its parameter was constrained; e.g., in Ref. 40) the combined scale and $U(1)$ parameter

is a chiral superfield. More recently, a discussion of conformal supergravity for N up to 4 has been given based on the tangent space group (2.5) with $G = U(N)$ ⁴²⁾. (There are fields with spins > 2 for $N > 4$, so $N = 4$ is the maximal value of N for the conformal theories.) In this section we shall expand the somewhat terse presentation given there.

Before doing this, we remark that although the superconformal algebra is $SU(2,2|N)$, the superspace we have introduced in Section 2 is perfectly adequate for the description of conformal supergravity. This is because some of the additional parameters of $SU(2,2|N)$, i.e., the conformal and superconformal boosts, get absorbed into general co-ordinate transformations, while Weyl (scale) and S supersymmetry transformations will appear as extra invariances of the constraints as in previous approaches. These results are the natural superspace generalizations of results previously found in x space^{43),29)}. Other authors⁴⁴⁾ have used a similar superspace to us (although with explicit gauging of Weyl transformations), but no attempt was made by them to make contact with the irreducible set of conformal supergravity fields (3.11).

The constraints we shall use are the same as in the previous section, (4.2) - (4.6), but in addition we shall impose a dimension one constraint corresponding to the vector part of the $U(N)$ connection $\Omega_a^j{}_k$. We choose

$$H_{\alpha\dot{\beta}}^i = \zeta_{\alpha}^i \dot{\rho}_{\dot{\beta}} + \frac{(N-3)}{2} \delta_j^i T_{\alpha\dot{\beta}}^j \quad (5.1)$$

$$T_{\alpha\dot{\beta}}^i = \bar{\chi}_{\dot{\beta}}^{jk} \chi_{\alpha}{}_{ijk} \quad (5.2)$$

The second term in (5.1) may look a little artificial at first sight, but it is chosen so that it vanishes for $N \leq 3$. In $N = 4$ we recall from the discussion in Section 2 that we had to choose the $U(1)$ part of the connection somewhat carefully in order to guarantee (3.30), and the precise form of this term has been selected to ensure that the appropriate non-linear generalization of (3.30) holds. With the choice (5.1) the torsions of Section 4 simplify a little and we shall list them again:

dim 0

$$T_{\alpha\dot{\beta}}^c = 0 \quad ; \quad T_{\alpha\dot{\beta}}^i = -i \delta_j^i (\zeta^c)_{\alpha\dot{\beta}}$$

dim 1/2

$$T_{\alpha\beta}^{\gamma} = T_{\alpha\dot{\beta}}^{\dot{\gamma}} = T_{\alpha b}^c = 0$$

$$T_{\alpha\beta}^{\dot{\gamma}} = \epsilon_{\alpha\beta} \bar{\chi}^{ijk\dot{\gamma}}$$

dim 1

$$T_{\alpha\beta}^{\dot{\gamma}} \rightarrow T_{\alpha\dot{\beta},\rho,\delta}^{j,h} = i\epsilon_{\alpha\rho} \bar{M}_{\dot{\beta}\delta}^{jh} + i\epsilon_{\dot{\beta}\delta} N_{\alpha\rho}^{jh} + i\epsilon_{\alpha\rho} \epsilon_{\dot{\beta}\delta} S^{jh} \quad (5.3)$$

$$T_{\alpha\beta}^{\dot{\gamma}} \rightarrow T_{\alpha\dot{\beta},\rho,\delta k}^j = -2i\epsilon_{\alpha\delta} G_{\rho\dot{\beta}k}^j - i\frac{(N-3)}{12} \delta_k^j \epsilon_{\alpha\beta} J_{\rho\dot{\delta}}$$

the dimension 3/2 torsions having the same form as (4.23). The dimension one curvatures $R_{\alpha\beta,C}^D$ also remain unchanged (4.12)-(4.14), but we recall from (3.30) that in the linearized case we had to choose $R^i_i = 0$. In the non-linear case we have

$$R_{\alpha\beta}^{\gamma k} = \frac{1}{2} \epsilon_{\alpha\beta} \bar{D}_{\dot{\delta}k} \bar{\chi}^{ijk\dot{\delta}} \quad (5.4)$$

However, there is no bilinear available to set this equal to, so that our final $N = 4$ constraint is

$$D_{\alpha}^k \chi_{ijk}^{\alpha} = 0 \quad (5.5)$$

Equations (5.1) and (5.3) describe the $N \leq 3$ theories fully and with the addition of (5.5) we are now in a position to complete the discussion of $N = 4$. To avoid having to write a large number of unnecessary indices out, let us introduce the following notation ($N = 4$ only):

$$\bar{\chi}_{\dot{\alpha}i} = -\frac{1}{3!} \epsilon_{ijkl} \bar{\chi}_{\dot{\alpha}}^{jkl} \quad (5.6)$$

$$L = L^i_i, \quad \Omega = \Omega^i_i, \quad R = R^i_i$$

Under $U(1)$, $\bar{\chi}_{\alpha i}$ transforms as

$$\delta \bar{\chi}_{\alpha i} = -\frac{3}{4} \epsilon \bar{\chi}_{\alpha i} \quad (5.7)$$

Consider now the one-form P defined by

$$P = E^{\alpha} P_{\alpha} - E^{\dot{\alpha}} P_{\dot{\alpha}} \quad (5.8)$$

where $P_{\alpha\dot{\alpha}} = \frac{i}{4} D_{\alpha\dot{\alpha}} \bar{\chi}_{\alpha i}$, $P_{\dot{\alpha}i} = \bar{\chi}_{\alpha i}$

Then we claim the following equations are true

$$DP = 0 \quad (5.9)$$

$$R = 2 \bar{P} \wedge P$$

To see this let us write

$$\begin{aligned} \stackrel{(1)}{I} &= DP \\ \stackrel{(2)}{I} &= R - 2 \bar{P} \wedge P \end{aligned} \quad (5.10)$$

Then by the Ricci Identity (2.13)

$$\begin{aligned} D \stackrel{(1)}{I} &= -P \wedge \stackrel{(2)}{I} \\ D \stackrel{(2)}{I} &= -2 \bar{P} \wedge \stackrel{(1)}{I} + 2 \stackrel{(1)}{I} \wedge P \end{aligned} \quad (5.11)$$

Or, in indices

$$\sum_{(ABC)} (D_A \stackrel{(1)}{I}_{BC} + T_{AB}{}^D \stackrel{(1)}{I}_{DC} + P_A \stackrel{(2)}{I}_{BC}) = 0 \quad (5.12)$$

$$\sum_{(ABC)} \left(D_A \overset{(2)}{I}_{BC} + T_{AB} D \overset{(2)}{I}_{DC} + 2 \bar{P}_A \overset{(1)}{I}_{AC} - 2 P_A \overset{(1)}{I}_{BC} \right) = 0 \quad (5.13)$$

Now let us suppose that the dimension one parts of Eqs. (5.9) are true, i.e.,

$$\overset{(1)}{I}_{\underline{\alpha}\underline{\beta}} = \overset{(1)}{I}_{\underline{\alpha}\dot{\beta}} = \overset{(1)}{I}_{\dot{\alpha}\underline{\beta}} = \overset{(2)}{I}_{\underline{\alpha}\underline{\beta}} = \overset{(2)}{I}_{\underline{\alpha}\dot{\beta}} = 0 \quad (5.14)$$

$\overset{(2)}{I}$ is imaginary since $\bar{R} = -R$.) Then (5.12) and (5.13) show that the others are by the Ricci Identity. For example, taking the $(\alpha\beta\dot{\gamma})$ component of (5.12), we find

$$T_{\dot{\gamma}\underline{\alpha}} D \overset{(1)}{I}_{\underline{\alpha}\underline{\beta}} + T_{\dot{\gamma}\underline{\beta}} D \overset{(1)}{I}_{\underline{\alpha}\underline{\alpha}} = 0 \quad (5.15)$$

$$\Rightarrow \overset{(1)}{I}_{\underline{\alpha}\underline{\beta}} = 0 \quad (5.16)$$

In a similar way, one proves that $\overset{(1)}{I}_{\dot{\alpha}\dot{\beta}} = 0$ from the $(\alpha\beta\dot{\gamma})$ cpt. of (5.12). We can now use the fact that $\overset{(1)}{I}_{\underline{\alpha}\underline{\beta}} = \overset{(1)}{I}_{\dot{\alpha}\dot{\beta}} = 0$ to prove $\overset{(1)}{I}_{\underline{\alpha}\dot{\beta}} = 0$ and similarly for $\overset{(2)}{I}$. It therefore remains to show that (5.14) holds. We have

$$\overset{(1)}{I}_{AB} = D_A P_B - (-1)^{AB} D_B P_A + T_{AB}{}^C P_C \quad (5.17)$$

Since $P_{\underline{\alpha}} = 0$, $\overset{(1)}{I}_{\underline{\alpha}\underline{\beta}}$ becomes just

$$\epsilon^{ijk} \bar{\chi}_{\dot{\gamma}k} \bar{\chi}_{\dot{\gamma}i} = 0 \quad (5.18)$$

which is clearly true. The $\overset{(1)}{I}_{\underline{\alpha}\dot{\beta}}$ equation is

$$D_{\underline{\alpha}} \bar{\chi}_{\dot{\beta}j} = \frac{1}{4} \delta_j^i D_{\underline{\alpha}} \bar{\chi}_{\dot{\beta}k} \quad (5.19)$$

which is just Eq. (4.10), whilst the $\binom{1}{I}_{\alpha\beta}$ part is

$$\bar{D}_{\alpha i} \bar{\chi}_{\dot{\rho} j} + \bar{D}_{\dot{\rho} j} \bar{\chi}_{\alpha i} = 0 \quad (5.20)$$

which is a rewritten version of Eqs. (4.16) and (5.5). Since $P_{\alpha} = 0$, $\binom{2}{I}_{\alpha\beta}$ tells us that $R_{\alpha\beta}^k = 0$, i.e., Eq. (5.5) again. Finally, substituting (5.1) into (4.19), one finds

$$R_{\alpha\dot{\rho}}^k = 2 \bar{\chi}_{\dot{\rho} j} \chi_{\alpha}^i \quad (5.21)$$

which is just the equation $\binom{2}{I}_{\alpha\beta} = 0$. Therefore, Eqs. (5.9) are satisfied and they can be written in the form

$$d\hat{\Omega} - \hat{\Omega} \wedge \hat{\Omega} = 0 \quad (5.22)$$

where

$$\hat{\Omega} = \begin{pmatrix} \frac{1}{2} \Omega & \bar{P} \\ P & \frac{1}{2} \bar{\Omega} \end{pmatrix}$$

$\hat{\Omega}$ is therefore a Lie algebra valued one-form, the relevant group being $SU(1,1)$ ²⁹⁾. From (5.22) the $SU(1,1)$ curvature is zero and this equation therefore has the solution

$$\hat{\Omega} = v^{-1} dv \quad (5.23)$$

where v is an element of $SU(1,1)$. This is of course a three parameter group, but we may use the $U(1)$ invariance to fix one of them so that v describes two real scalar superfields as the coset $SU(1,1)/U(1)$. The leading ($\theta=0$) component of v therefore corresponds to the non-linear version of the dimension zero complex scalar field of the $N=4$ conformal theory. We observe that the chirality condition satisfied by the linearized superfield W_{ijkl} has become more complicated in the non-linear case.

Before going on to discuss the component content described by our superspace, we shall first show that the basic constraints are invariant under a larger group of transformations. We recall from Section 3 that for $N \leq 3$ only the dimension zero constraints are really independent for this purpose, since the conventional constraints merely determine the transformations of the connection and the vielbein under the new invariance. The extra constraints in $N = 4$ do not change the result which is as follows:

$$\begin{aligned} H_a{}^b &= 2\delta_a{}^b U \\ H_\alpha{}^\beta &= \delta_\alpha{}^\beta \delta_j{}^i U \quad ; \quad H_{\dot{\alpha}}{}^{\dot{\beta}} = -\delta_{\dot{\alpha}}{}^{\dot{\beta}} \delta_i{}^j U \\ H_{\underline{\alpha}}{}^{\dot{\beta}} &= H_{\dot{\alpha}}{}^\beta = H_{\underline{\alpha}}{}^b = H_{\dot{\alpha}}{}^{\dot{b}} = 0 \\ H_a{}^{\dot{\beta}} \rightarrow H_{\alpha\dot{\alpha}}{}^{\dot{\beta}j} &= 4i\delta_{\alpha\dot{\alpha}}{}^{\dot{\beta}} D_\alpha^j U \\ H_a{}^\beta \rightarrow H_{\alpha\bar{\alpha}}{}^{\beta}{}_{,j} &= -4i\delta_{\alpha\bar{\alpha}}{}^{\beta} \bar{D}_{\bar{\alpha}j} U \end{aligned}$$

$$\begin{aligned} \phi_{\alpha,\dot{\beta}\gamma}^i &= 0 \\ \phi_{\alpha,\beta\gamma}^i &= 2(\epsilon_{\alpha\beta} D_\gamma^i + \epsilon_{\alpha\gamma} D_\beta^i) U \\ \phi_{\alpha,jk}^i &= (\delta_{jk} D_\alpha^i - 4\delta_{\alpha k} D_j^i) U \\ \phi_{a,bc} &= 2\eta_{ac} D_b U - 2\eta_{ab} D_c U \\ \phi_a{}^j &= 0 \end{aligned} \tag{5.24}$$

where

$$\begin{aligned} H_A{}^B &= E_A{}^M \delta E_M{}^B \\ \phi_{AB}{}^C &= E_A{}^M \delta \Omega_{MB}{}^C \end{aligned}$$

This additional invariance may be expressed more succinctly in the form

$$\delta E_{\underline{\alpha}}^M = -U E_{\underline{\alpha}}^M \quad (5.25)$$

from which (5.24) follows via the constraints.

Here, U is a real unconstrained superfield. We may identify its leading component with ordinary scale transformations and its second component $D_{\underline{\alpha}}^i U|_{\theta=0}$ with the S supersymmetry parameter. To see this, let us choose the gauge where to leading order in θ ,

$$E_m^a|_{\theta=0} = e_m^a(x) \quad (5.26)$$

$$E_m^{\underline{\alpha}}|_{\theta=0} = \psi_m^{\underline{\alpha}}(x)$$

Then, from (5.24), we find

$$\begin{aligned} \delta e_m^a &= 2U|_{\theta=0} e_m^a \\ \delta \psi_m^{\underline{\alpha}} &= U|_{\theta=0} \psi_m^{\underline{\alpha}} - 2ie_m^b(\sigma_b)^{\underline{\alpha}\dot{\alpha}} \bar{D}_{\dot{\alpha}}^i U|_{\theta=0} \end{aligned} \quad (5.27)$$

i.e., the standard scale and S supersymmetry transformations for these fields.

The transformations of the covariant superfields are given by

$$\delta S^{ij} = 2\epsilon^{\alpha\beta} D_{\alpha}^i D_{\beta}^j U - 2U S^{ij} \quad (5.28)$$

$$\delta N_{\alpha\beta}^{ij} = -4D_{(\alpha}^i D_{\beta)}^j U - 2UN_{\alpha\beta}^{ij} \quad (5.29)$$

$$\delta G_{\alpha\beta}^i = -[D_{\alpha}^i, \bar{D}_{\beta}^j] U - 2U G_{\alpha\beta}^i \quad (5.30)$$

and

$$\delta \bar{\chi}_{\dot{\alpha}}^{ijk} = -U \bar{\chi}_{\dot{\alpha}}^{ijk} \quad (5.31)$$

$$\delta \bar{M}_{\dot{\alpha}\dot{\beta}}^{\dot{i}\dot{j}} = -2u \bar{M}_{\dot{\alpha}\dot{\beta}}^{\dot{i}\dot{j}} \quad (5.32)$$

$$\delta \bar{\Psi}_{\dot{\alpha}\dot{\beta}\dot{\gamma}}^{\dot{i}} = -3u \bar{\Psi}_{\dot{\alpha}\dot{\beta}\dot{\gamma}}^{\dot{i}} \quad (5.33)$$

We have given $\delta\chi$, δM and $\delta\psi$ separately since in $N=2$ $\chi=0$ and in $N=1$ $\chi=M=0$. Hence the superfields divide naturally into two types, one which transforms by a mere rescaling, the others transforming inhomogeneously. We may regard the superfields $\chi_{\alpha i j k}$ ($N=3,4$), $M_{\alpha\beta i j}$ ($N=2$) and $\psi_{\alpha\beta\gamma}$ ($N=1$) as the superspace generalizations of the Weyl tensor in ordinary geometry.

We shall now indicate how the components of (3.11) fit into the structure we have described. We clearly have more component fields than those listed in (3.11), but this is because we have not explicitly gauged the dilations and hence see fields corresponding to the higher components of the superfield U explicitly in the torsions. In order to compare with (3.11) therefore, we should go to a Wess-Zumino gauge for the parameter U . Alternatively, we may say that we have the components (3.11) (8+8, 24+24, 64+64, 128+128 for $N=1,2,3,4$ respectively) together with an entire scalar superfield. (In this way of counting we should remember to count the number of degrees of freedom of gauge fields by subtracting the conformal gauges, i.e., e_m^a counts 5 and ψ_m^α counts 8N). Since at dimension 2 the number of components one expects to see becomes rather large, we shall content ourselves with a discussion up to dimension 3/2; in order to do this we shall first have to simplify the dimension 3/2 results given in Section 4.

Using (5.1) one finds Eqs. (4.27) - (4.33) become

$$\bar{D}_{(\dot{\alpha}} \bar{M}_{\dot{\beta}\dot{\gamma}}^{\dot{i}\dot{j}}) - \bar{\chi}_{(\dot{\alpha}}^{\dot{i}\dot{j}} \bar{N}_{\dot{\beta}\dot{\gamma})} = 2 \delta_{\dot{\alpha}\dot{\beta}}^{\dot{i}\dot{j}} \bar{\Psi}_{\dot{\gamma}} \quad (5.34)$$

$$\begin{aligned} 2 D_{(\alpha} G_{\beta)}^j \dot{\gamma} \dot{k} - \bar{D}_{\dot{\gamma}\dot{k}} N_{\alpha\beta}^{\dot{i}\dot{j}} \\ = - \delta_{\dot{\alpha}\dot{\beta}}^{\dot{i}\dot{j}} \Psi_{\alpha\beta\dot{\gamma}} + \frac{1}{2} \delta_{\dot{\alpha}\dot{\beta}}^{\dot{i}\dot{j}} \Psi_{\alpha\beta\dot{\gamma}} - \bar{\chi}_{\dot{\gamma}}^{\dot{i}\dot{j}} M_{\alpha\beta\dot{\gamma}} \\ - \frac{(N-3)}{12} \delta_{\dot{\alpha}\dot{\beta}}^{\dot{i}\dot{j}} D_{(\alpha} J_{\beta)} \dot{\gamma} \end{aligned} \quad (5.35)$$

$$\begin{aligned}
 \epsilon^{\alpha\beta} D_\alpha^i C_{\beta\gamma}^j + \bar{D}_{\gamma h} S^j - \frac{1}{3} \epsilon^{\alpha\beta} \bar{D}_{\alpha k} \bar{M}_{\beta\gamma}^{ij} \\
 = \delta_k^i \bar{\Psi}_\gamma^j + \frac{1}{2} \delta_k^j \bar{\Psi}_\gamma^i + \bar{\chi}_\gamma^{ij} \bar{S}_{kh} \\
 + \frac{1}{3} \epsilon^{\alpha\beta} \bar{\chi}_\alpha^{ij} \bar{M}_{\beta\gamma k\ell} - \frac{(N-3)}{6} \delta_k^j \epsilon^{\alpha\beta} D_\alpha^i J_{\beta\gamma}
 \end{aligned} \tag{5.36}$$

$$D_\alpha^{(i} S^{jk)} = 0 \tag{5.37}$$

$$D_\alpha^i N_{\beta\gamma}^{jk} = D_\alpha^{[i} N_{\beta\gamma}^{jk]} \tag{5.38}$$

$$n_\alpha^{ijk} = -\frac{1}{6} D_\alpha^j \bar{\chi}_\beta^{ik} - 2i g_\alpha^{ijk} \tag{5.39}$$

$$\tilde{n}_\alpha^{ijk} + \frac{1}{3} (D_\alpha^j S^{ki} - D_\alpha^k S^{ji}) = 2i \tilde{g}_\alpha^{ijk} \tag{5.40}$$

$$m_{\alpha\beta\gamma}^{ijk} = D_\alpha(\beta \bar{\chi}_{\gamma\delta}^{jk}) - 2i g_{\alpha\beta\gamma}^{ijk} + i \frac{(N-3)}{6} J_{\alpha(\beta} \bar{\chi}_{\gamma\delta}^{jk)} \tag{5.41}$$

$$\tilde{m}_{\alpha\beta\gamma}^{ijk} = -2i \tilde{g}_{\alpha\beta\gamma}^{ijk} \tag{5.42}$$

whilst the additional constraint in $N = 4$ (5.5) implies

$$\begin{aligned}
 \epsilon^{\alpha\beta} \bar{D}_{\alpha k} \bar{M}_{\beta\gamma}^{ik} &= \bar{\chi}_\alpha^{ijk} \bar{N}_{\beta\gamma k} - 3 \bar{\chi}_{\beta\gamma}^i D_\alpha^j \chi^{\alpha j} \\
 &+ 3 \frac{i}{2} (D_\alpha^j \bar{\chi}_{\beta\gamma}^i) \chi^{\alpha j}
 \end{aligned} \tag{5.43}$$

For the purpose of counting components, the bilinear terms in these equations are irrelevant and we must remember that the functions $\psi_{\alpha\beta\gamma}^i$, $\bar{\psi}_\alpha^i$ are known as vectorial derivatives of the spin 3/2 fields (plus non-linear contributions), although they are not conformally invariant. It is now straightforward to see that we have the fields of dimension ≤ 1 of (3.11), i.e., $\chi_{\alpha ijk}$, $M_{\alpha\beta ij}$,

B_{ijk}^{ℓ} and A_{ijkl} (which appears in the non-linear case as $\mathcal{U}$). The spin 3/2 field $\psi_{\alpha\beta\gamma i}$ is also present. Therefore to complete the conformal set up to dimension 3/2 we only need $\lambda_{\alpha ij}^k$. We may identify this as

$$\lambda_{\alpha ij}^k = \frac{1}{3} \epsilon^{\alpha\gamma} D_{\beta}^R M_{\gamma ij} \quad (5.44)$$

The non-conformally invariant components are U (trace of e_m^a), $D_{\alpha}^i U$ (γ trace of ψ_m^{α}) and at dimension one $N_{\alpha\beta}^{ij}$, S^{ij} . At dimension 3/2 a scalar field has the following independent components

$$\epsilon^{\alpha\beta} D_{\alpha}^i D_{\beta}^j D_{\gamma}^R U = \begin{array}{|c|} \hline \square \\ \hline \end{array}, \quad (\frac{1}{2}, 0) \quad (5.45)$$

$$D_{\alpha}^i D_{\beta}^j D_{\gamma}^R U = \begin{array}{|c|} \hline \square \\ \hline \end{array}, \quad (\frac{3}{2}, 0) \quad (5.46)$$

$$\epsilon^{\alpha\beta} D_{\alpha}^i D_{\beta}^j \bar{D}_{\gamma k} U = \left\{ \begin{array}{|c|} \hline \square \\ \hline \end{array} \right\}_{N-1} + \square, \quad (0, \frac{1}{2}) \quad (5.47)$$

$$D_{\alpha}^i D_{\beta}^j \bar{D}_{\gamma k} U = \left\{ \begin{array}{|c|} \hline \square \\ \hline \end{array} \right\}_{N-1} + \square, \quad (1, \frac{1}{2}) \quad (5.48)$$

where we have indicated the irreducible representations of $SU(N)$ and $SL(2, \mathbb{C})$ that these components fall into. We may identify the component (5.46) with $D_{\alpha\beta\gamma}^{ijk}$ and the component (5.45) with $\tilde{N}_{\alpha}^{i,jk}$ which is related to $D_{\alpha}^i S^{jk}$ by (5.40). The other components are contained in $D_{\alpha}^i G_{\beta\gamma k}^j$ and $\bar{D}_{\gamma j} S^{ij}$ and it is not difficult to check that they occur in the representations (5.47) and (5.48). Thus we have verified our claim that the constraints we have imposed yield the component fields of (3.11) together with a real scalar superfield, at least up to dimension 3/2. At dimension 2 we do have $C_{\alpha\beta\gamma\delta}$ and $X_{\alpha\beta j}^i$ so that the only field of the set (3.11) that we have not located is D_{kl}^{ij} . It is not difficult to believe that this is present however. For example, for $N = 4$, we showed in Section 3 that the linearized superspace contained the chiral superfield, W_{ijkl} . Such a field automatically contains a D_{kl}^{ij} at dimension 2, so the only problem is whether it is real or not. However, the reality condition was imposed in order that the components X, ψ, ϕ and C should be field strengths, but in the geometrical approach they are defined this way. Hence we conclude that we do

have a full description of conformal supergravity. (It might be objected that D_{kl}^{ij} may actually be zero, however we shall show that this is not the case explicitly for $N = 1$ and 2 later.)

To summarize, then, conformal supergravity may be described in a superspace with $SL(2, \mathbb{C}) \times U(N)$ as the tangent space group in terms of a conformally covariant superfield which is $\mathcal{U}$, $\chi_{\alpha ijk}$, $M_{\alpha\beta ij}$ and $\psi_{\alpha\beta\gamma}$ for $N = 4, 3, 2, 1$ respectively together with some additional non-Weyl covariant superfields $N_{\alpha\beta}^{ij}$, S^{ij} and $G_{\alpha\beta j}^i$. The conformally covariant fields are chiral even in the non-linear case. For $N = 4$ the idea of chirality has to be extended a little as we have seen; for $N = 3$, the chirality condition is Eq. (4.10); for $N = 2$ it is Eq. (5.42) since $\tilde{g}_{\alpha\gamma\delta}^{i,jk}$ is zero in this case. For $N = 1$, the chirality of $\psi_{\alpha\beta\gamma}$ arises as a consequence of the Bianchi Identity $I_{ab\gamma}^{\delta}$ which we did not solve in Section 4. For completeness, we include its solution here,

$$\bar{D}_{\dot{\alpha}} \psi_{\alpha\beta\gamma} = 0$$

$$\bar{D}_{\dot{\alpha}} \psi_{\beta} = \frac{i}{2} D_{\beta\dot{\alpha}} \bar{S} + G_{\beta\dot{\alpha}} \bar{S} \quad (5.49)$$

$$\begin{aligned} \bar{D}_{\dot{\alpha}} \bar{\psi}_{\dot{\alpha}\dot{\beta}\dot{\gamma}} &= \left(\frac{i}{2} \epsilon_{\dot{\alpha}\dot{\beta}} D_{\dot{\gamma}\dot{\alpha}} \bar{S} + \epsilon_{\dot{\beta}\dot{\gamma}} G_{\dot{\alpha}\dot{\beta}} \bar{S} \right) \\ &+ \quad \dot{\alpha} \leftrightarrow \dot{\beta} \end{aligned}$$

The Weyl non-covariant fields undergo the transformations (5.28) - (5.30) (which also show that their leading components can be gauged away) and those higher components of these superfields which are not covariant are Wess-Zumino gauge fields for the Weyl parameter U .

6. FROM CONFORMAL TO POINCARÉ

The motivation for studying conformal supergravity is, as we mentioned in the Introduction, not primarily for the sake of the conformal theory itself, but rather because it is a convenient starting point for the construction of off-shell Poincaré supergravity theories. This approach has been emphasized by x space workers¹³⁾ where use is made of the notion of compensating multiplets. For example, in $N = 1$, one couples the basic (8+8) Weyl multiplet to a chiral multiplet (4+4) in a conformally covariant way. Then one fixes the conformal gauges to obtain the minimal (12+12) off-shell Poincaré representation⁴⁵⁾.

In $N = 2$, the Weyl multiplet has $(24+24)$ components and to fix the Weyl and $U(1)$ gauges one may add a vector multiplet with $(8+8)$ components. The resulting $(32+32)$ component multiplet^{46),16)} is the so-called minimal $N = 2$ Poincaré representation, but the associated Lagrangian leads to nonsensical equations of motion. In order to circumvent this latter problem one has to add an additional multiplet which is a linear multiplet^{16),46)} (sometimes called the tensor multiplet or the non-linear multiplet). This multiplet again has $(8+8)$ components, including a quasi-conserved vector, and may alternatively be regarded as a compensating multiplet for the local $SU(2)$ that the minimal representation possesses.

In superspace, the use of the conformal group to obtain information on Poincaré supergravity has taken a slightly different form via constraint analysis^{17),18)}. Let us recall that there are three types of constraint:

- i) Conventional constraints: these allow one to redefine the basic gauge fields, the vielbein and the connection by tensorial shifts, e.g., if Ω_{MA}^B is a connection then an equally good one is

$$\Omega'_{MA}{}^B = \Omega_{MA}{}^B + K_{MA}{}^B \quad (6.1)$$

where K should have the same symmetry properties as Ω on the A, B indices but transforms homogeneously. This allows one to choose many components of the torsion to vanish, and a well-known example is in ordinary Riemannian geometry where it is always possible to choose the torsion to be zero this way.

- ii) Representation-preserving constraints: one may enforce the requirement that certain representations of rigid supersymmetry should have natural extensions in the local case. The simplest example is the requirement that there exists a chiral scalar field, $S (\bar{D}S=0)$ in ordinary $N = 1$ superspace [i.e., without an explicitly gauged $U(1)$]. If this is to be consistent one must have

$$T_{\alpha\beta}{}^c = T_{\alpha\beta}{}^{\dot{c}} = 0 \quad (6.2)$$

iii) Conformal constraints: these arise as the requirement that the constraints should admit an additional invariance (super Weyl group) which has a parameter which is in a well-defined representation of supersymmetry. The minimal $N = 1$ superspace constraints¹²⁾, for example, are invariant under a chiral super Weyl group⁴⁰⁾ and this third type of constraint is therefore related to the component notion of compensating multiplets.

Constraint analysis has been carried out for both $N = 1$ and $N = 2$ where the basic superspaces were taken to have only $SL(2, \mathbb{C})$ as the tangent space group. In addition, Gates^{17), 19)} has proposed that the non-conventional superspace constraints should be imposed to ensure "super conformal breakdown". This author again works in a superspace without explicit local symmetry and uses an extended notion of the superconformal group (essentially defined as the invariance group of the constraints). Then, demanding that this group be restricted to scale invariance and $U(N)$, he derives some constraints for the torsions. With the aid of some type ii) constraints he is then able to discuss $N = 1$ and $N = 2$ Poincaré constraints completely and derive some partial results for higher N . In our approach to conformal supergravity, we employed type i) constraints but neither of types ii) and iii). We did not use type ii) simply because it is not clear which representations one should preserve for $N > 2$. Type iii) constraints were not employed because in our formalism [i.e., with explicit gauging of $U(N)$] they are not needed although we have seen that there is still an additional (but unconstrained) Weyl invariance. In this section we shall discuss how one obtains Poincaré theory from the conformal theory presented in Section 5 by a process that we shall call "de-gauging". This turns out to be rather similar to type iii) constraint analysis and "precurvatures"¹⁴⁾, and we shall also be able to relate it in concrete examples to the idea of compensating multiplets.

In order to "de-gauge" we write the connection in the form

$$\hat{\Omega}_A^B = \Omega_A^B + \Sigma_A^B \quad (6.3)$$

where $\hat{\Omega}_A^B$ is an $SL(2, \mathbb{C}) \times G$ connection, Ω_A^B an $SL(2, \mathbb{C}) \times H$ connection, H being a subgroup of G and Σ_A^B is the connection for the remainder of G , $(G-H)$, which will also be a group in the cases we shall discuss. The torsions and curvatures defined with respect to $\hat{\Omega}$ and Ω are related by

$$\hat{T}^A = T^A + E^B \wedge \Sigma_B^A \quad (6.4)$$

$$\hat{R}_A{}^B = R_A{}^B + D Z_A{}^B + \Sigma_A{}^C \bar{\Sigma}_C{}^B \quad (6.5)$$

where D is the covariant derivative with respect to Ω . If the $\hat{\Omega}$ Bianchi Identities are satisfied, then so are the Ω ones. In index notation (6.4) and (6.5) read,

$$\hat{T}_{AB}{}^C = T_{AB}{}^C + \Sigma_{AB}{}^C - (-1)^{AB} \Sigma_{BA}{}^C \quad (6.6)$$

$$\begin{aligned} \hat{R}_{ABC}{}^D &= R_{ABC}{}^D + D_A \Sigma_{BC}{}^D - (-1)^{AB} D_B \Sigma_{AC}{}^D \\ &\quad + T_{AB}{}^E \Sigma_{EC}{}^D + \Sigma_{AE}{}^D \Sigma_{BC}{}^E \\ &\quad - (-1)^{AB} \Sigma_{BE}{}^D \Sigma_{AC}{}^E \end{aligned} \quad (6.7)$$

The step we have made so far is of course completely trivial since we still have full local G invariance. However, the Ω geometry is now only manifestly H covariant and if we impose constraints on Σ itself the $(G-H)$ invariance will be reduced from an unconstrained superfield parameter to a constrained one, the latter being irreducible if the constraints are chosen appropriately. In this sense choosing constraints on Σ is very similar to type iii) constraint analysis. We can also look at it in terms of multiplets. As an example, consider $N = 1$. The component field content of the conformal $N = 1$ superspace was shown to consist of the Weyl multiplet $(8+8)$ plus the parameter U . If we now de-gauge the $U(1)$ the fields we will see will be $(8+8)$ plus $(U+iV)$ where iV is the $U(1)$ parameter (V real) i.e., there will be altogether $(8+8)$ plus $(16+16)$ component fields. Hence if we demand that the constraint on Σ be chosen such that the remaining reduced Weyl + $U(1)$ symmetry is chiral, we will have a $(12+12)$ component field representation of Poincaré supergravity^{*)}. To illustrate the procedure let us briefly go through the $N = 1$ example. From Sections 4 and 5, the conformal torsions are rather simple:

dim 0

$$\hat{T}_{\alpha\beta}{}^c = 0 ; \quad \hat{T}_{\alpha\dot{\beta}}{}^c = -i(\sigma^c)_{\alpha\dot{\beta}} \quad (6.8)$$

*) The $12 + 12$ has two interpretations, either as a Poincaré multiplet or as the Weyl multiplet together with the Wess-Zumino gauge fields of the superconformal parameter.

dim 1/2

$$\hat{T}_{\alpha\beta}{}^\gamma = \hat{T}_{\alpha\dot{\beta}}{}^{\dot{\gamma}} = \hat{T}_{\alpha\beta}{}^{\dot{\gamma}} = \hat{T}_{\alpha\dot{\beta}}{}^{\gamma} = 0 \quad (6.9)$$

dim 1

$$\hat{T}_{\alpha\dot{\alpha},\beta,\dot{\gamma}} = i\epsilon_{\alpha\beta}\epsilon_{\dot{\alpha}\dot{\gamma}}S \quad (6.10)$$

$$\hat{T}_{\alpha\dot{\alpha},\beta,\dot{\gamma}} = -2i\epsilon_{\alpha\beta}\epsilon_{\dot{\alpha}\dot{\gamma}}G \quad (6.11)$$

dim 3/2

$$\begin{aligned} \hat{T}_{\alpha\dot{\alpha},\beta\dot{\beta},\dot{\gamma}} &= \epsilon_{\alpha\beta}(\bar{\psi}_{\dot{\alpha}\dot{\beta}\dot{\gamma}} + \epsilon_{\dot{\alpha}\dot{\gamma}}\bar{\psi}_{\dot{\beta}} \\ &\quad + \epsilon_{\dot{\beta}\dot{\gamma}}\bar{\psi}_{\dot{\alpha}}) \\ &\quad + \epsilon_{\dot{\alpha}\dot{\beta}}\psi_{\alpha\beta\dot{\gamma}} \end{aligned} \quad (6.12)$$

The U(1) parts of the curvature are $(\hat{R}^i_j \equiv F)$,

dim 1

$$F_{\alpha\beta} = 0 \quad (6.13)$$

$$F_{\alpha\dot{\beta}} = -3G\epsilon_{\alpha\dot{\beta}} \quad (6.14)$$

dim 3/2

$$F_{\alpha,\beta\dot{\beta}} = \frac{3i}{4}(\psi_{\alpha\beta\dot{\beta}} + \epsilon_{\alpha\beta}\bar{\psi}_{\dot{\beta}}) \quad (6.15)$$

dim 2

$$F_{\alpha\dot{\alpha},\beta\dot{\beta}} = \epsilon_{\dot{\alpha}\dot{\beta}}X_{\alpha\beta} - \epsilon_{\alpha\beta}\bar{X}_{\dot{\alpha}\dot{\beta}} \quad (6.16)$$

In addition, (5.37) tells us that S is chiral

$$\hat{D}_\alpha S = 0 \quad (6.17)$$

while (5.35) and (5.36) become

$$\hat{D}_\alpha G_{\beta\gamma} = -\frac{1}{4} \psi_{\alpha\beta\gamma} \quad (6.18)$$

$$\hat{D}_\alpha G^{\alpha\beta} = \frac{3}{4} \bar{\psi}_\beta - \frac{1}{4} \bar{D}_\beta S \quad (6.19)$$

These equations are not quite the same as for ordinary Wess-Zumino superspace¹²⁾; from (6.19) we see that there is an additional dimension $3/2$ spin $1/2$ field in the theory. To implement the de-gauging we split the $SL(2, \mathbb{C}) \times U(1)$ connection, $\hat{\Omega}$, into an $SL(2, \mathbb{C})$ part Ω and a $U(1)$ part Σ so

$$\Sigma_\alpha{}^\beta = \delta_\alpha{}^\beta \Sigma, \quad \Sigma_\alpha{}^\beta = \delta_\alpha{}^\beta \Sigma, \quad \bar{\Sigma} = -\Sigma \quad (6.20)$$

Then we have

$$F_{AB} = D_A \Sigma_B - (-1)^{AB} D_B \Sigma_A + T_{AB}{}^C \Sigma_C \quad (6.21)$$

If we now choose the constraint

$$\Sigma_\alpha = 0 \quad (6.22)$$

then all the new torsions are the same as in (6.8) - (6.12) except for $T_{\alpha\dot{\alpha},\beta,\gamma}$,

$$T_{\alpha\dot{\alpha},\beta,\gamma} = -2i \epsilon_{\alpha\gamma} G_{\beta\dot{\alpha}} + 3i \epsilon_{\beta\gamma} G_{\alpha\dot{\alpha}} \quad (6.23)$$

while the $F_{\alpha\dot{\alpha}}$ equation gives ($D_\alpha = \hat{D}_\alpha$)

$$D_\alpha G^{\alpha\beta} = -\frac{1}{2} \bar{\psi}_\beta = -\frac{1}{10} \bar{D}_\beta S \quad (6.24)$$

i.e., the standard equations modulo conventions. We know that this set of

constraints has a chiral super Weyl invariance⁴⁰⁾, but it is instructive to see how it arises this way. The original equations were invariant under the Weyl transformations (5.24) and of course $SL(2, \mathbb{C}) \times U(1)$. In particular,

$$\hat{\Phi}_\alpha^{ij}{}_k = \delta \Sigma_\alpha = -3D_\alpha U - iD_\alpha V \quad (6.25)$$

where iV is the $U(1)$ parameter. Then (6.22) implies that $3U + iV = 3\bar{\Lambda}$ is the reduced chiral combined scale and $U(1)$ parameter. The non-trivial vielbein transformations are then

$$\begin{aligned} H_a{}^b &= \delta_a{}^b (\Lambda + \bar{\Lambda}) \\ H_\alpha{}^\beta &= \delta_\alpha{}^\beta (2\bar{\Lambda} - \Lambda) \\ H_{\alpha\dot{\alpha}}{}^{\beta} &= -2i \delta_\alpha{}^\beta \bar{D}_{\dot{\alpha}} \bar{\Lambda} \end{aligned} \quad (6.26)$$

as expected. It is well known that there is an alternative non-minimal version of $N = 1$ ^{47),32)}, and this may be obtained by keeping $\Sigma_\alpha = 0$, but imposing a dimension one constraint such as $S = 0$ ¹⁷⁾. In this case the remaining additional symmetry is a linear superfield having $(12+12)$ components. Whichever constraint we choose, the irreducible parameter invariance of the final geometry is just the compensating multiplet.

7. $N = 2$ POINCARÉ SUPERGRAVITY

The first off-shell version of $N = 2$ Poincaré supergravity in superspace was given by Breitenlohner and Sohnius⁴⁶⁾ who were subsequently able to find a simpler formulation¹⁴⁾. $N = 2$ superspace constraints were also derived from component results³⁴⁾ and by constraint analysis^{18),19)}. A complete solution to the Bianchi Identities in a superspace with local $SU(2)$ has been given in Ref. 48). $N = 2$ supergravity has also been discussed within the framework of chiral superspace and using this approach, Sokatchev was able to construct the action⁴⁹⁾. (The action given in Refs. 14) and 46) is really a tensor calculus one as opposed to a superspace one.) Here we shall analyze $N = 2$ by the method introduced in the last section and we shall "de-gauge" in two steps, $U(2) \rightarrow SU(2) \rightarrow 1$. Upon de-gauging the $U(1)$ factor we recover the "almost simple" version of Ref. 14) and we are able to de-gauge the $SU(2)$ in two different ways to find the standard $(40+40)$ representation¹⁵⁾ and the inequivalent $40 + 40$ which has an off-shell central charge¹⁶⁾

We begin by writing the $SL(2, \mathbb{C}) \times U(2)$ connection of Section 5 in the form

$$\hat{\Omega}_A{}^B = \Omega_A{}^B + \Sigma_A{}^B \quad (7.1)$$

where

$$\Sigma_\alpha{}^A = \delta_\alpha{}^\rho \delta_j{}^i \Sigma, \quad \Sigma_{\dot{\alpha}}{}^{\dot{A}} = \delta_{\dot{\alpha}}{}^{\dot{\rho}} \delta_{\dot{j}}{}^{\dot{i}} \Sigma, \quad \bar{\Sigma} = -\Sigma \quad (7.2)$$

We then impose the constraint

$$\Sigma_\alpha{}^i = 0 \quad (7.3)$$

The reduced symmetry group consisting of the super Weyl transformations of (5.24) and $U(1)$ transformations is then given by a chiral superfield $\Lambda = U + iV$, $D_\alpha^i \Lambda = 0$ where iV is the $U(1)$ parameter. However, a chiral superfield in $N = 2$ is still reducible and we impose a further constraint

$$S^{ij} = -\epsilon^{ik} \epsilon^{jl} \bar{S}_{kl} \quad (7.4)$$

which reduces the parameter still further

$$\epsilon^{\alpha\beta} D_\alpha^i D_\beta^j \bar{\Lambda} - \Lambda S^{ij} = \epsilon^{ik} \epsilon^{jl} (\epsilon^{\dot{\alpha}\dot{\beta}} \bar{D}_{\dot{\alpha}k} \bar{D}_{\dot{\beta}l} \Lambda + \bar{\Lambda} \bar{S}_{kl}) \quad (7.5)$$

The parameter Λ now has $(8+8)$ components and corresponds to an $N = 2$ vector multiplet, so that according to our discussion in the previous section we should expect to find that the $SU(2)$ superspace has a $(32+32)$ component representation, i.e., the minimal representation of $N = 2$ Poincaré. Using (6.4) and (6.5) it is straightforward to calculate the new torsions from the conformal torsions (5.3). They are

dim 0

$$T_{\alpha\beta}{}^c = 0, \quad T_{\alpha\beta}{}^c = -i\delta_j{}^i (\sigma^c)_{\alpha\beta}$$

dim 1/2

$$T_{\alpha\beta}{}^{\dot{\gamma}} = T_{\alpha\dot{\beta}}{}^{\dot{\gamma}} = T_{\alpha\beta}{}^{\dot{\gamma}} = T_{\alpha\dot{\beta}}{}^c = 0 \quad (7.6)$$

dim 1

$$T_{\alpha\dot{\alpha}, \beta\dot{\beta}, \gamma\dot{\gamma}}{}^{j,k} = i\epsilon^{jk} (\epsilon_{\alpha\beta} \bar{M}_{\dot{\alpha}\dot{\gamma}} + \epsilon_{\dot{\alpha}\dot{\gamma}} N_{\alpha\beta}) + i\epsilon_{\alpha\beta} \epsilon_{\dot{\alpha}\dot{\gamma}} S^{jk}$$

$$T_{\alpha\dot{\alpha},\dot{\rho},\delta k}^j = -i\delta_k^j (2\epsilon_{\alpha\dot{\rho}}G_{\dot{\rho}\alpha} + \epsilon_{\dot{\rho}\delta}G_{\alpha\dot{\alpha}}) \quad (7.6) \text{ cont.}$$

the dimension 3/2 ones being unchanged in form. In (7.6)

$$\bar{M}_{\dot{\alpha}\dot{\rho}}^{ij} = \epsilon^{ij}\bar{M}_{\dot{\alpha}\dot{\rho}}, \quad N_{\alpha\dot{\rho}}^{ij} = \epsilon^{ij}N_{\alpha\dot{\rho}} \quad (7.7)$$

$$iG_{\alpha\dot{\alpha}} = \Sigma_{\alpha\dot{\alpha}}$$

$G_{\alpha\dot{\alpha}}$ is related to $G_{\alpha\dot{\beta}j}^i$ by

$$G_{\alpha\dot{\rho}j}^i = -\delta_j^i G_{\alpha\dot{\rho}} \quad (7.8)$$

and S^{ij} satisfies (7.4).

The curvatures also simplify,

dim 1

$$\begin{aligned} R_{\alpha\dot{\rho},\delta\delta}^{ij} &= \epsilon^{ij}\epsilon_{\alpha\dot{\rho}}N_{\delta\delta} + (\epsilon_{\alpha\delta}\epsilon_{\dot{\rho}\delta} + \epsilon_{\alpha\dot{\rho}}\epsilon_{\delta\delta})S^{ij} \\ R_{\alpha\dot{\rho},\delta\delta}^{ij} &= -\epsilon_{\alpha\dot{\rho}}\epsilon^{ij}\bar{M}_{\delta\delta} \\ R_{\alpha\dot{\rho}}^{ijk}{}_{\ell} &= (\delta_{\ell}^i\epsilon^{jk} + \delta_{\ell}^j\epsilon^{ik})N_{\alpha\dot{\rho}} + \epsilon_{\alpha\dot{\rho}}(\delta_{\ell}^i S^{jk} - \delta_{\ell}^j S^{ik}) \\ R_{\alpha\dot{\rho}j,\delta\delta}^i &= -\delta_j^i (\epsilon_{\alpha\dot{\rho}}G_{\delta\delta} + \epsilon_{\alpha\delta}G_{\dot{\rho}\delta}) \\ R_{\alpha\dot{\rho}j,\delta\delta}^i &= (4\delta_{\ell}^i\delta_j^{\ell} - 2\delta_j^i\delta_{\ell}^{\ell})G_{\alpha\dot{\rho}} \end{aligned}$$

dim 3/2

$$\begin{aligned} R_{\alpha,\dot{\rho}\dot{\rho},j}^i{}_{\delta k} &= i\delta_k^i\psi_{\alpha\dot{\rho}\dot{\rho}}^j - \frac{i}{2}\delta_k^j\psi_{\alpha\dot{\rho}\dot{\rho}}^i \\ &\quad + i\epsilon_{\alpha\dot{\rho}}\left\{\delta_k^i\bar{\psi}_{\dot{\rho}}^j - \frac{1}{2}\delta_k^j\bar{\psi}_{\dot{\rho}}^i - 2\epsilon^{ij}\bar{\lambda}_{\dot{\rho}k} \right. \\ &\quad \left. + \delta_k^j\epsilon^{i\ell}\bar{\lambda}_{\dot{\rho}\ell}\right\} \quad (7.9) \end{aligned}$$

$R_{\alpha,\dot{\beta}\dot{\beta},\gamma\delta}^i$ and $R_{\alpha,\dot{\beta}\dot{\beta},\dot{\gamma}\dot{\delta}}^i$ are given in (4.24) and (4.25). The dimension 2 curvatures are given by (4.39) and (4.42) where $X_{\alpha\dot{\beta}}^k$ is now the SU(2) field strength. The dimension 3/2 relations (4.27) - (4.33) become

$$\begin{aligned}
 \bar{D}_{\alpha i} \bar{M}_{\beta \gamma} &= \epsilon_{ij} \bar{\psi}_{\alpha \beta \gamma}^j + \epsilon_{i\beta} \bar{\lambda}_{\gamma i} + \epsilon_{i\gamma} \bar{\lambda}_{\beta i} \\
 D_{\alpha}^i \bar{M}_{\beta \gamma} &= 0 \\
 D_{\alpha}^i N_{\beta \gamma} &= -\epsilon_{\alpha\beta} (\epsilon^{ij} \psi_{\gamma j} + \lambda_{\gamma}^i) + \beta \leftrightarrow \gamma \\
 \bar{D}_{\gamma k} N_{\alpha \beta} &= \epsilon_{k\alpha} \psi_{\beta \gamma}^{\dagger} \quad (7.10) \\
 D_{\alpha}^i S^{jk} &= (\epsilon^{ij} \epsilon^{kl} + \epsilon^{ik} \epsilon^{jl}) \psi_{\alpha l} + \epsilon^{ij} \lambda_{\alpha}^k \\
 &\quad + \epsilon^{ik} \lambda_{\alpha}^j \\
 D_{\alpha}^i G_{\beta \gamma} &= \frac{1}{4} \psi_{\alpha \beta \gamma}^i + \frac{1}{4} \epsilon_{\alpha\beta} \bar{\psi}_{\gamma}^i - \epsilon_{\alpha\beta} \epsilon^{ij} \bar{\lambda}_{\gamma j}
 \end{aligned}$$

There is also a field at dimension 2 (the D field of the conformal multiplet)

$$D_{\alpha}^i \lambda^{\alpha j} = \epsilon^{ij} I + \frac{1}{12} (\bar{M}_{\alpha \beta} \bar{N}^{\alpha \beta} - M_{\alpha \beta} N^{\alpha \beta}) \quad (7.11)$$

I being real^{*)}. In addition, it is not surprising to find that these constraints imply the existence of a new superspace two form F corresponding to a vector field. Its non-trivial components are

$$\begin{aligned}
 F_{\alpha \beta}^{ij} &= -i \epsilon_{\alpha\beta} \epsilon^{ij} \\
 F_{\alpha i, \beta \dot{j}} &= \epsilon_{i\beta} F_{\alpha \dot{j}} + \epsilon_{\alpha \dot{j}} \bar{F}_{\beta i} \quad (7.12) \\
 F_{\alpha \beta} &= i (N_{\alpha \beta} - M_{\alpha \beta})
 \end{aligned}$$

and it satisfies the normal spin one Bianchi Identity

$$dF = 0 \Rightarrow F = dA \quad (7.13)$$

It is straightforward to see that the geometry now describes the minimal

^{*)} We have changed notation for this field here to avoid confusion with the covariant derivative.

(32+32) component multiplet $[e_m^a, A_m, S^{ij}, G_{\alpha\dot{\alpha}}, (N_{\alpha\beta} + M_{\alpha\dot{\beta}}), \Omega_a^i, I; \psi_m^\alpha, \lambda_\alpha^i]$ where A_m is the gauge field corresponding to F . The invariance of this geometry under complex Weyl transformations takes the following form:

$$\begin{aligned} H_a^b &= \delta_a^b (\Lambda + \bar{\Lambda}) \\ H_\alpha^\beta &= \delta_\alpha^\beta \delta_j^i \Lambda \\ H_{\alpha\dot{\alpha}}, j &= -4i \delta_\alpha^\beta \bar{D}_{\dot{\alpha}j} \Lambda \\ \phi_{\alpha, j}^i &= \delta_R^j D_\alpha^i \bar{\Lambda} - 2 \delta_R^i D_\alpha^j \bar{\Lambda} \end{aligned} \quad (7.14)$$

while the fields transform as follows:

$$\begin{aligned} \delta S^{ij} &= 2\epsilon^{\alpha\beta} D_\alpha^i D_\beta^j \bar{\Lambda} - 2\Lambda S^{ij} \\ \delta M_{\alpha\beta} &= -2\bar{\Lambda} M_{\alpha\beta} \\ \delta N_{\alpha\beta} &= -2\epsilon_{ij} D_\alpha^i D_\beta^j \bar{\Lambda} - 2\Lambda N_{\alpha\beta} \\ \delta G_{\alpha\dot{\alpha}} &= i D_{\alpha\dot{\alpha}} (\Lambda - \bar{\Lambda}) - \frac{3}{2} (\Lambda + \bar{\Lambda}) G_{\alpha\dot{\alpha}} \end{aligned} \quad (7.15)$$

Λ is chiral, $D_\alpha \Lambda = 0$, and satisfies (7.5). The variation of the vector potential, A_A , satisfies

$$\begin{aligned} D_\alpha^i (\delta A_\beta^j) + D_\beta^j (\delta A_\alpha^i) &= -2i \epsilon^{ij} \epsilon_{\alpha\beta} \Lambda \\ D_\alpha^i (\delta \bar{A}_{\dot{\beta}j}) + \bar{D}_{\dot{\beta}j} (\delta A_\alpha^i) &= i \delta_j^i \delta A_{\alpha\dot{\beta}} \end{aligned} \quad (7.16)$$

We note that (7.16) is formally the same as the constraint equations for $N = 2$ supersymmetric Yang-Mills⁵⁰⁾, so this is direct evidence that Λ describes a vector multiplet.

Having obtained the "almost simple" version of $N = 2$ supergravity in two component spinor notation, we shall proceed to de-gauge the $SU(2)$. Denoting now the $SU(2)$ superspace quantities by a caret and the superspace with no local internal symmetry by Ω , etc., we have

$$\hat{\Omega}_A{}^B = \Omega_A{}^B + \Sigma_A{}^B \quad (7.17)$$

where

$$\begin{aligned} \Sigma_\alpha{}^\beta &= \delta_\alpha{}^\beta \Sigma^i{}_j ; \quad \Sigma_{\dot{\alpha}}{}^{\dot{\beta}} = \delta_{\dot{\alpha}}{}^{\dot{\beta}} \Sigma^j{}_i \\ \Sigma^i{}_j &= -\Sigma^j{}_i ; \quad \Sigma^i{}_i = 0 \end{aligned} \quad (7.18)$$

We choose

$$\Sigma_{\alpha, k}{}^{ij} = 2 \delta_k^i \rho_\alpha^j - \delta_k^j \rho_\alpha^i \quad (7.19)$$

The reduced SU(2) parameter preserving (7.19) satisfies

$$D_\alpha^{(i} L^{jk)} - \rho_\alpha^{(i} L^{jk)} = 0, \quad L^{ij} = \epsilon^{jk} L^i{}_k \quad (7.20)$$

But (7.20) is nothing but the defining equation of an $N = 2$ linear multiplet (modified by the bilinear term) and hence we expect to see $(32+32) + (8+8)$ components in this superspace. The torsions become modified via (6.6) and the dimension $1/2$ ones are now no longer zero. One has

$$\begin{aligned} T_{\alpha\beta}{}^{\gamma k}{}_{ij} &= \epsilon_{\alpha\gamma} (\delta_k^i \rho_\beta^j - 2 \delta_k^j \rho_\beta^i) + \epsilon \leftrightarrow \beta \\ T_{\alpha\beta}{}^{\gamma j}{}_{ik} &= \delta_\beta^j (\delta_k^i \rho_\alpha^k - 2 \delta_k^k \rho_\alpha^i) \end{aligned} \quad (7.21)$$

By using (6.7) one finds the following dimension one equations

$$\begin{aligned} S^{ij} &= -D_\alpha^{(i} \rho^{\alpha j)} - 3 \epsilon^{\alpha\beta} \rho_\alpha^i \rho_\beta^j \\ N_{\alpha\beta} &= 2 \epsilon_{ij} (D_\alpha^i \rho_\beta^j - \rho_\alpha^i \rho_\beta^j) \\ \epsilon_{ij} D_\alpha^i \rho^{\alpha j} &= P \\ D_{(\alpha}^{(i} \rho_{\beta}^{j)} &= 0 \end{aligned} \quad (7.22)$$

$$\begin{aligned}
 G_{\alpha\dot{\alpha}} &= -\frac{1}{4} (D_{\alpha}^i \bar{\rho}_{\dot{\alpha}i} - \bar{D}_{\dot{\alpha}i} \rho_{\alpha}^i) - \frac{1}{2} \rho_{\alpha}^i \bar{\rho}_{\dot{\alpha}i} \\
 K_{\alpha\dot{\beta}j}^i &= 2 D_{\alpha}^i \bar{\rho}_{\dot{\beta}j} + 6 \rho_{\alpha}^i \bar{\rho}_{\dot{\beta}j} \\
 &\quad - \text{trace over } i, j \\
 D_{\alpha}^i \bar{\rho}_{\dot{\beta}i} &= -2 G_{\alpha\dot{\beta}} - \rho_{\alpha}^i \bar{\rho}_{\dot{\beta}i} + i H_{\alpha\dot{\beta}}
 \end{aligned} \tag{7.22}$$

cont.

Here P (complex), $H_{\alpha\dot{\alpha}}$ (Hermitean) and $K_{\alpha\dot{\alpha}j}^i$ are new component fields, the latter being the old $SU(2)$ connection

$$\Sigma_{\alpha\dot{\alpha}}^i{}_j = i K_{\alpha\dot{\alpha}j}^i \tag{7.23}$$

In the final superspace therefore there are only two covariant superfields, ρ_{α}^i and $M_{\alpha\dot{\beta}}^{19}$, the equation of motion being $\rho_{\alpha}^i = 0$. The new singlet vector field $H_{\alpha\dot{\alpha}}$ is the vector of the additional linear multiplet and its divergence is essentially the dimension 2 field I . We have therefore recovered the standard $40 + 40$ component representation^{15), 46)} ($e_m^a, A_m, S^{ij}, P, N_{\alpha\dot{\beta}}, G_{\alpha\dot{\alpha}}, H_{\alpha\dot{\alpha}}, K_{\alpha\dot{\alpha}j}^i; \psi_m^{\alpha}, \rho_{\alpha}^i, \lambda_{\alpha}^i$).

To conclude this section we shall briefly indicate how one obtains the alternative $(40+40)$ component representation¹⁶⁾. This set of fields is built up from the minimal $(32+32)$ component set together with a scalar multiplet (hypermultiplet). Since this latter multiplet has a central charge it is necessary to introduce a new bosonic co-ordinate into the superspace. A tangent space vector X^A then has an extra component

$$X^A = (X^a, X^5) \tag{7.24}$$

where X^5 is a scalar under the tangent space group $SL(2, \mathbb{C}) \times SU(2)$. "Almost simple" supergravity is described in this space by the identification

$$T_{AB}^5 = F_{AB} \tag{7.25}$$

and all other extra torsions and curvatures are zero (i.e., $T_{A5}^5, T_{5A}^B, R_{A5,C}^D$). The fields depend trivially on the extra co-ordinate, but this situation changes when we de-gauge from $SU(2) \rightarrow 1$. We again do this by (7.18). The first of Eqs. (7.22) is modified and becomes

$$- D_{\alpha}^i \rho^{\alpha j} - 3 \epsilon^{\alpha\beta} \rho_{\alpha}^i \rho_{\beta}^j = S^j + B^j \quad (7.26)$$

where

$$\Sigma_5^i{}_j = i B^i{}_j \quad j \quad \overline{B}_j{}^i = B^i{}_j$$

We also have

$$D_{\alpha}^i \bar{\rho}_{\beta}^j + \bar{D}_{\beta}^j \rho_{\alpha}^i = i \delta_j^i H_{\alpha\beta} \quad (7.27)$$

$$D_{\alpha}^i \rho_{\beta}^j + D_{\beta}^j \rho_{\alpha}^i = \frac{1}{2} \epsilon_{\alpha\beta} \epsilon^j{}_i P$$

Then, choosing P imaginary, we find

$$dQ = 0 \quad (7.28)$$

where the one-form Q has the form

$$Q = \frac{1}{2} E^{\alpha\dot{\alpha}} H_{\alpha\dot{\alpha}} - \frac{i}{2} E^5 P + E^{\alpha} \rho_{\alpha} - E^{\dot{\alpha}} \bar{\rho}_{\dot{\alpha}} \quad (7.29)$$

Hence

$$Q = dA \quad (7.30)$$

and

$$\begin{aligned} \rho_{\alpha}^i &= D_{\alpha}^i A \\ H_{\alpha\dot{\alpha}} &= D_{\alpha\dot{\alpha}} A \\ P &= 2i D_5 A \end{aligned} \quad (7.31)$$

Therefore we have replaced the six components, $H_{\alpha\dot{\alpha}}$, P of the conventional set by the six scalars A , $B^i{}_j$, $D_5 A$ and I (no longer dependent). Notice that in terms of covariant superfields, this set is not as simple as the previous one since not all the dimension one fields are derivable from A . The superspace is therefore described by three superfields A , S^{ij} and $M_{\alpha\beta}$ and it is necessary to set both A and S^{ij} to zero to obtain the equations of motion.

To conclude this section we note that we have been able to construct both inequivalent (40+40) component representations by using two different linear multiplets obeying (7.20). The first has no central charge and is an ordinary linear multiplet, while the second has a central charge and a further constraint is necessary to make it irreducible; it turns out to be just a different way of

writing the scalar multiplet. This result is somewhat reminiscent of the situation in $N = 2$ Yang-Mills where if one works in a superspace with a central charge there are again two possible irreducible representations⁵¹⁾, the standard Grimm-Wess-Sohnius⁵⁰⁾ one and the centrally charged version of Sohnius-Stelle-West⁵¹⁾. We also remark that $N = 2$ conformal supergravity may also be described in superspaces with local $SU(2)$ or no local internal symmetry group, as well as in the $U(2)$ space of Section 5. To do this, one re-interprets the $32 + 32$ ($40+40$) representations as the ($24+24$) components of the Weyl multiplet together with the Wess-Zumino gauges of the super Weyl parameter Λ and the local $SU(2)$ parameter L^i_j (7.20). Indeed, the most economical version is the $SU(2)$ superspace since the number of Wess-Zumino gauges of Λ is rather small.

8. ON-SHELL POINCARÉ SUPERGRAVITY

The first extended on-shell supergravity theory to be obtained in superspace was the $N = 3$ theory with a gauged $O(3)$ ³⁶⁾. The method employed was that of gauge completion, i.e., building up the superspace from the component results. The full non-linear $N = 8$ theory, which has local $SU(8)$ and global E_7 symmetries, was given in Ref. 28) and although the starting point for this formulation was a superspace of the type we have been using, it was shown that supersymmetry implied several other Bianchi-type Identities including those for the vector fields. All other theories ($N < 8$) may be obtained from the $N = 8$ theory directly by truncation, but although this formulation is adequate, the theory is perhaps a little more transparent if one uses additional bosonic co-ordinates. The $N = 8$ theory was recast in this form in Ref. 9) and has also been written in terms of the covariant derivatives of Eq. (2.29)²⁷⁾; these two approaches are equivalent, but the former has a more direct geometrical interpretation and it is this version we shall present here.

The basic idea is to introduce a number of extra bosonic co-ordinates equal to the number of vector fields, i.e., $\binom{N}{2} + \binom{N}{6}$ [$\binom{N}{2}$ for $N = 8$]. Since we are working on-shell and wish to have unitary groups in the tangent space, we shall actually take these co-ordinates to be complex; this is permissible because the $(S) U(N)$ covariant spin one field strengths will be essentially self-dual, as discussed for the linearized case in Section 3. We note that the $\binom{N}{6}$ vectors are only present in $N = 6$ and to simplify the formalism a little we shall only introduce $\binom{N}{2}$ extra co-ordinates, so that the formalism will only be directly applicable to $N = 8, N \leq 5$. We shall indicate how to obtain the $N = 6$ theory by truncation.

We consider a superspace with co-ordinates $z^M = (z^M, y^{\underline{m}}, y^{\underline{m}'})$ where z^M are the co-ordinates of ordinary superspace and $y^{\underline{m}}$ are $1/2N(N-1)$ complex co-ordinates with complex conjugates $y^{\underline{m}'}$. A tangent space vector has the form

$$X^A = (X^A, X^{\underline{a}}, X^{\underline{a}'}) \quad (8.1)$$

where

$$\begin{aligned} X^A &= (X^a, X^{\underline{x}}, X^{\underline{z}}) \\ X^{\underline{a}} &= X_{ij} ; \quad X^{\underline{a}'} = \overline{X}^{\underline{j}} \end{aligned} \quad (8.2)$$

The tangent space group is again $SL(2, \mathbb{C}) \times (S) U(N)$,

$$\delta X^A = X^B L_B^A \quad (8.3)$$

where L_A^B is given by (2.5), and

$$\begin{aligned} L_{\underline{a}}^{\underline{b}} &= L^{\underline{ij}}_{\underline{kl}} = 2 \delta^{\underline{ij}}_{\underline{kl}} L^{\underline{j}}_{\underline{i}} \\ L_{\underline{a}'}^{\underline{b}'} &= \overline{(L_{\underline{a}}^{\underline{b}})} \end{aligned} \quad (8.4)$$

all others vanishing. In the large space we have a "sehrvielbein" $\delta_m^{\mathcal{A}}$ and a connection $\Omega_{m, \mathcal{A}}^{\mathcal{B}}$ from which we construct the torsions and curvatures in exactly the same way as in Section 2. We shall impose the condition that all fields depend trivially on the extra co-ordinates,

$$D_{\underline{a}} = \partial_{\underline{m}} = 0 \quad (8.5)$$

from which it follows that D_A is the covariant derivative of ordinary superspace. Furthermore, (8.5) implies via the Ricci Identity (2.13) that most of the new torsions and curvatures are zero; the only non-zero ones being T_{AB}^C , $T_{AB}^{\underline{c}}$, $T_{AB}^{\underline{c}'}$, R_{ABC}^D and $R_{AB, \underline{c}}^{\underline{d}}$ together with their complex conjugates. The non-trivial Bianchi Identities in the extended superspace are

$$I_{ABC}^D, I_{ABC}^{\underline{d}}, I_{AB \underline{c}}^{\underline{d}}, I_{AB \underline{c}}^{\underline{d}'} \quad (8.6)$$

the others being identically satisfied. I_{ABC}^D is the same as in Section 2 [Eq. (2.18)] while the others may be written in the following form:

$$I_{ABC} \xrightarrow{d} DF_{ij} = \bar{F}^{kl} \wedge P_{kl ij} \quad (8.7)$$

$$I_{ABC} \xrightarrow{d} R^{ij}_{kl} = 2 \delta^{[i}_{[k} R^{j]}_{l]} = \bar{P}^{ijmn} \wedge P_{mnkl} \quad (8.8)$$

$$I_{ABC} \xrightarrow{d'} DP_{ijkl} = 0 \quad (8.9)$$

where

$$F_{ij} = \frac{1}{2} E^D \wedge E^C T_{CD, ij} \quad (8.10)$$

$$P_{ijkl} = E^A T_{A, ij, kl} \quad (8.11)$$

Equations (8.7) - (8.9) hold on ordinary superspace since by (8.5), $\varepsilon^A = E^A$. These identities were found in Ref. 28) and the problem is therefore to find the components of the tensors F , P , T , R . Since we have already solved I_{ABC}^D in Section 4, this amounts to solving (8.7)-(8.9). A priori, the one-form P is skew on both pairs of indices, but in order for (8.8) to hold we take it to be totally antisymmetric. In addition, in $N = 8$, it is self-dual

$$\bar{P}^{ijkl} = \frac{1}{4!} \epsilon^{ijklmnpq} P_{mnpq} \quad (8.12)$$

This requirement is reasonable since in the linearized limit (8.9) will then have the solution

$$P_{ijkl} = dW_{ijkl} \quad (8.13)$$

where W_{ijkl} is the linearized superfield of Section 3 (for $N \geq 4$). For $N = 6$, we have to keep P_{ij78} when truncating from $N = 8$ ($i, j=1, \dots, 6$), so that one finds

$$DF_{ij} = P_{ijkl} \wedge \bar{F}^{kl} + 2 P_{ij78} \wedge \bar{F}^{78} \quad (N=6) \quad (8.14)$$

$$D F_{78} = P_{78 k\ell} \wedge \bar{F}^{k\ell}, \quad N=6 \quad (8.15)$$

instead of (8.7), where we note the appearance of an extra vector field strength F_{78} . P_{ij78} is not independent, for by (8.12) one has

$$P_{ij78} = \frac{1}{4!} \epsilon_{ijklmn} \bar{P}^{klmn}, \quad N=6 \quad (8.16)$$

Equation (8.8) can be written in the following form

$$\begin{aligned} R^i_j &= -\frac{1}{3} \bar{P}^{iklm} \wedge P_{jk\ell m}, \quad N=8 \\ &= -\frac{2}{3} \bar{P}^{iklm} \wedge P_{jk\ell m} + \frac{1}{12} \delta^i_j \bar{P}^{mnrs} \wedge P_{mnrs}, \quad N \neq 8 \end{aligned} \quad (8.17)$$

The solution to the dimension $1/2$ F identities [when use is made of the dimension 0 and $1/2$ torsions (4.2) - (4.5)]

$$P^i_{\alpha jk\ell m} = 2 \delta^i_{[\alpha} \chi_{\beta]jk\ell m} \quad (8.18)$$

$$P_{\alpha ij k\ell m} = \bar{\chi}_{\alpha ij k\ell m} \quad (8.19)$$

$$F_{\alpha\beta, \beta j, k\ell} = \epsilon_{\alpha\beta} \chi_{\alpha j k\ell}, \quad F_{\alpha\beta, k\ell} = 0 \quad (8.20)$$

$$\left. \begin{aligned} F'_{\alpha\beta, \beta j} &= 2 \epsilon_{\alpha\beta} \chi_{\alpha j} \\ F'_{\alpha\beta, \beta} &= 0 \end{aligned} \right\} N=6 \quad (8.21)$$

where $F' = F_{78}$ and

$$\chi_{\alpha}^{ijklm} = \epsilon^{ijklmn} \chi_{\alpha n}, \quad N=6 \quad (8.22)$$

To derive these results we assumed that the only non-vanishing dimension zero component of the F 's is given by

$$F_{\alpha\beta, k\ell}^{ij} = -2i \epsilon_{\alpha\beta} \delta_{[k\ell]}^{ij} \quad (8.23)$$

At dimension one, one then finds from (8.9)

$$D_{\alpha}^i \bar{\chi}_{\dot{\beta} j k l m n} = 5i \delta_{\dot{\beta} j}^i P_{\alpha \dot{\beta} k l m n} \quad (8.24)$$

$$\bar{D}_{\dot{\alpha} i} \chi_{\beta j k l} = 2i P_{\beta \dot{\alpha} i j k l} \quad (8.25)$$

$$D_{(\alpha}^i \chi_{\beta)}^{j k l m n} = M_{\alpha \beta}^{i j k l m n} \quad (8.26)$$

$$D_{\alpha}^i \chi^{\alpha j k l m n} = -5 \bar{\chi}_{\dot{\alpha}}^i [j k \bar{\chi}^{\dot{\alpha} l m n}]^{\dot{\alpha}} \quad (8.27)$$

$$B_{ijk}^{\dot{\ell}} = \frac{1}{2} D_{\alpha}^{\dot{\ell}} \chi^{\alpha}_{ijk} = \frac{2}{(N-4)} \bar{\chi}_{\dot{\alpha} i j k m n} \bar{\chi}^{\dot{\alpha} l m n} - \frac{3}{(N-3)(N-4)} \delta_{[i}^{\dot{\ell}} \bar{\chi}_{\dot{\alpha} j k] m n r} \bar{\chi}^{\dot{\alpha} m n r} \quad (8.28)$$

i.e., the first variations of the spinor fields. The dimension one component of F is then determined to be (from the F identities)

$$F_{\alpha \dot{\alpha}, \beta \dot{\beta}, ij} = -i \epsilon_{\dot{\alpha} \dot{\beta}} M_{\alpha \beta ij} - i \epsilon_{\alpha \beta} \bar{N}_{\dot{\alpha} \dot{\beta} ij} \quad (8.29)$$

$$F'_{\alpha \dot{\alpha}, \beta \dot{\beta}} = 2i \epsilon_{\dot{\alpha} \dot{\beta}} M_{\alpha \beta} - \frac{i}{3 \cdot 4!} \epsilon_{\alpha \beta} \epsilon_{ijklmn} \bar{\chi}_{(\dot{\alpha}}^{ijk} \bar{\chi}_{\dot{\beta})}^{lmn} \quad (N=6) \quad (8.30)$$

with

$$M_{\alpha \beta} = \frac{1}{6!} \epsilon_{ijklmn} M_{\alpha \beta}^{ijklmn} \quad (N=6) \quad (8.31)$$

and the functions S, N, H and G are determined from the dimension one part of (8.17):

$$S^{ij} = 0 \quad (8.32)$$

$$N_{\alpha \beta}^{ij} = \frac{1}{3} \chi_{(\alpha}^{ijklm} \chi_{\beta)}^{klm}, \quad N \neq 8 \\ = -\frac{1}{72} \epsilon^{ijklmnpq} \chi_{\alpha klm} \chi_{\beta npq} \quad N=8 \quad (8.33)$$

$$\left. \begin{aligned} H_{\alpha \dot{\beta} j}^i &= \frac{1}{2} J_{\alpha \dot{\beta} j}^i - \frac{1}{16} \delta_j^i J_{\alpha \dot{\beta}}^i \\ G_{\alpha \dot{\beta} j}^i &= -\frac{1}{48} \delta_j^i J_{\alpha \dot{\beta}}^i \end{aligned} \right\} N=8 \quad (8.34)$$

$$\left. \begin{aligned} H_{\alpha\beta j}^i &= \frac{1}{2} J_{\alpha\beta j}^i + \frac{1}{6} K_{\alpha\beta j}^i - \frac{1}{16} \delta_j^i J_{\alpha\beta} - \frac{1}{80} \delta_j^i K_{\alpha\beta} \end{aligned} \right\} N \neq 8$$

$$G_{\alpha\beta j}^i = \frac{7}{240} \delta_j^i K_{\alpha\beta} - \frac{1}{48} \delta_j^i J_{\alpha\beta} \quad (8.35)$$

where

$$J_{\alpha\beta j}^i = \bar{\chi}_{\beta}^{ihl} \chi_{\alpha jhl} \quad ; \quad J = J^i_i$$

$$K_{\alpha\beta j}^i = \chi_{\alpha}^{iklmn} \bar{\chi}_{\beta jklmn} \quad ; \quad K = K^i_i \quad (8.36)$$

This completes the solution to all the Bianchi Identities in the enlarged superspace; we have solved I_{ABC}^D already in Section 4, and (8.8) and (8.9) are two-form equations which by a similar argument to that given for $N = 4$ conformal supergravity in Section 5 are satisfied if their dimension one components are. The spin one identities also have rather few independent components as one may verify by employing the type of analysis we used in Section 2 in discussing identities for the Bianchi Identities [see also Ref. 31)].

It now remains to show how one may find the global symmetries. Equations (8.8) and (8.9) may be rewritten in the form

$$d\hat{\Omega} + \hat{\Omega} \wedge \hat{\Omega} = 0 \quad (8.37)$$

where the block-matrix one-form $\hat{\Omega}$ has the form

$$\hat{\Omega} = \begin{pmatrix} \Omega_{ij}^{kl} & -\bar{P}_{ij}^{kl} \\ -P_{ij}^{kl} & \bar{\Omega}_{ij}^{kl} \end{pmatrix} \quad (8.38)$$

$\hat{\Omega}$ is therefore a Lie algebra valued one-form corresponding to the group G' (see Appendix C). Equation (8.37) shows that the associated curvature is zero and hence has the solution

$$\hat{\Omega} = (d v^{-1}) \cdot v \quad (8.39)$$

where $\mathcal{U}$ is an element of the Lie group G' . For $N = 8$, this group is E_7 , for $N = 5$, $U(5,1)$ and for $N = 4$ $SU(4) \times SU(1,1)$ ²⁰⁾. For $N = 6$, (8.8) and hence (8.38) is slightly altered, the global group in this case being $SO^*(12)$ ²⁰⁾. The matrix $\mathcal{U}$ has the form ($N \neq 6$)

$$\mathcal{V} = \begin{pmatrix} u^{IJ}{}_{ij} & \bar{v}^{IJ}{}_{ij} \\ v_{IJ}{}^{ij} & \bar{u}_{IJ}{}^{ij} \end{pmatrix} \quad (8.40)$$

where each entry is antisymmetric on both pairs of indices. $(S) U(N)$ acts on the small indices in the natural way, while the global group G' acts on the capital indices by

$$\delta \mathcal{V} = - \Lambda \mathcal{V} \quad (8.41)$$

$$\Lambda = \begin{pmatrix} \Lambda^{IJ}{}_{KL} & \bar{\Sigma}^{IJKL} \\ \Sigma_{IJKL} & \bar{\Lambda}_{IJ}{}^{KL} \end{pmatrix} \quad (8.42)$$

where

$$\Lambda^{IJ}{}_{KL} = 2 \delta \begin{bmatrix} I & J \\ K & L \end{bmatrix} \Lambda^{IJ}{}_{KL} ; \quad \Lambda^I{}_J = - \bar{\Lambda}_J{}^I \quad (8.43)$$

and Σ is totally antisymmetric. In addition, in $N = 8$, Λ is traceless and Σ self-dual. We may identify $\mathcal{U}$ with part of the sechvielbein. To do this we first observe that (8.37) implies

$$\begin{pmatrix} 0 & \bar{P}^{ijhk} \\ P_{ijhk} & 0 \end{pmatrix} = - (D \mathcal{V}^{-1}) \mathcal{V} \quad (8.44)$$

But P_{ijkl} was originally one of the extra torsions

$$P_{A, ij, kl} = T_{A, ij, kl} (T_{A, b', c'}) \quad (8.45)$$

and [using (8.5)]

$$T_{A \underline{b}'}^{\underline{c}} = - (D_A \varepsilon_{\underline{b}'}^{\underline{m}}) \varepsilon_{\underline{m}}^{\underline{c}} - (D_A \varepsilon_{\underline{b}'}^{\underline{m}'}) \varepsilon_{\underline{m}'}^{\underline{c}} \quad (8.46)$$

Hence we have the result

$$U^{IJ}{}_{ij} \leftrightarrow \varepsilon_{\underline{m}}^{\underline{a}} \quad (8.47)$$

$$V_{IJ}{}_{ij} \leftrightarrow \varepsilon_{\underline{m}'}^{\underline{a}}$$

With the aid of $\mathcal{U}$ one can define the following vector field strengths

$$(F_{IJ}, \bar{F}^{IJ}) = (F_{ij}, \bar{F}^{ij}) \mathcal{U}^{-1} \quad (8.48)$$

which from (8.7) - (8.9) satisfy ordinary spin one identities

$$dF_{IJ} = 0 \quad (8.49)$$

They transform under a representation of G' (e.g., the $\underline{56}$ of E_7 for $N=8$) by

$$\delta(F, \bar{F}) = (F, \bar{F}) \Lambda \quad (8.50)$$

The sehrvielbein therefore contains not only the ordinary vielbein, but also the scalars and the spin one potentials. It has the following form

$$\varepsilon_m^{\mathcal{A}} = \begin{pmatrix} E_m^{\underline{a}} & E_{mij} & \bar{E}_m^{ij} \\ 0 & U^{IJ}{}_{ij} & \bar{V}^{IJ}{}_{ij} \\ & V_{IJ}{}_{ij} & \bar{U}_{IJ}{}^{ij} \end{pmatrix} \quad (8.51)$$

where $\mathcal{A} = (\underline{a}, ij, \bar{ij})$; $m = (\underline{m}, {}^{IJ}, {}_{IJ})$

and

$$(\bar{E}_{mij}, E_m^{ij}) = (A_{mIJ}, \bar{A}_m^{IJ}) \mathcal{U} \quad (8.52)$$

with A_{mIJ} being the $\binom{N}{2}$ vector fields.

Hence we see that on-shell Poincaré supergravities with global G' and local $(S) U(N)$ symmetries may be described rather naturally in a superspace with extra bosonic co-ordinates. For $N \leq 3$, the local $U(N)$ becomes trivial and the global symmetry is also not needed. This is because there are no scalars for $N \leq 3$, while for $N \geq 4$ they appear in the coset space form $G'/(S) U(N)$ [the number of scalars is therefore equal to $\dim G'$ minus the dimension of $(S) U(N)$, i.e., the same as in (3.1)].

To conclude this section we note that all the fields of (3.1) have now been located in superspace and by supersymmetry they must satisfy their equations of motion. However, these equations can now be read off by combining the results of Section 4 with those we have just obtained. The spin $3/2$ equations are obtainable from (4.28) and (4.29) since G and H are now known bilinears; similarly, the spin 2 equations is gotten from the explicit expressions for $B_{\alpha\beta, \gamma\gamma}$ and A given in (4.41) and (4.43). The spin $1/2$ equation is contained in (4.32) since N is also a known function. The spin 0 and spin 1 equations are obtainable from (4.32) by differentiating it with respect to D_{α}^i and $\bar{D}_{\alpha i}^{\cdot}$ respectively.

9. CONCLUDING REMARKS

We have given a rather thorough presentation of supergravity in superspace from the point of view of conventional superspace differential geometry. We have been able to obtain all off-shell conformal supergravities and all on-shell Poincaré supergravities as specializations of the general solution to the Bianchi Identities of Section 4. We have also been able to rederive off-shell $N = 1$ and $N = 2$ Poincaré supergravity theories, including the $N = 2$ centrally charged one, from the corresponding conformal theories.

As we remarked in the Introduction, there are many problems outstanding. The principal one is, of course, that of the auxiliary fields for Poincaré supergravity with $N \geq 3$. For $N \leq 4$, it might be thought that one could simply repeat the steps used to go from conformal supergravity to Poincaré supergravity for $N = 1$ and 2 . However, the success of this construction in these cases is due to the availability of (low-spin) multiplets which can be used for compensation purposes. For higher N , the off-shell conformal multiplets are known^{(6), (7)}, but they are higher spin multiplets. For example, in $N = 4$ where one does not have to compensate for the $U(1)$ since it is composite, one might try to restrict the parameters of scale and $SU(4)$ symmetries by conditions such as (7.19) which defines the reduced $SU(2)$ parameter by (7.20). The natural generalization of (7.20) to $N = 4$ is (at the linearized level)

$$D_\alpha^i L^j_R = \rho_\alpha^{ij} R + \delta_R^i \rho_\alpha^j - \frac{1}{4} \delta_R^j \rho_\alpha^i \quad (9.1)$$

where

$$L^i_j = -\bar{L}_j^i, \quad L^i_i = 0$$

and ρ^{ij}_k is in the $\underline{20}$ of $SU(4)$, ρ^i in the $\underline{4}$. Such a multiplet has some nice properties in the sense that it naturally compensates the $SU(4)$ and ρ^{ij}_k looks like a Lagrange multiplier for λ_{ij}^k (5.44). It is, however, an irreducible multiplet with a conserved spin 3 field of dimension 3⁷⁾. Similarly, one might reduce the scale parameter, U , by imposing a constraint such that

$$D_\alpha^i D^{\alpha j} U = 0 \quad (9.2)$$

This again defines an irreducible multiplet, this time with a conserved spin 4 field of dimension 4⁷⁾. Therefore, if we were to use these two multiplets in $N = 4$, we would "compensate" the conformal gauges [e.g., the $SU(4)$ vector fields would cease to be gauge fields], but the construction of a Lagrangian from a representation involving such high dimension high spin (conserved) fields would be impossible. Thus we are forced to conclude that there is no simple generalization of the $N = 1$ and 2 results.

In $N \geq 5$, the situation is more difficult because there is no conformal theory to start with, and the general situation seems to be that although one can construct combinations of representations which are irreducible if one does not impose conformal invariance, nevertheless these representations will be unsuitable for the construction of Lagrangians. With the benefit of hindsight this ought not to be too surprising in that the imposition of constraints in superspace is a way of doing group theory; it does not guarantee that the representations one finds can be employed in actions. Negative results have been obtained before, for example in $N = 4$ Yang-Mills⁵²⁾, and by explicitly trying to construct an $N = 3$ Lagrangian for Poincaré supergravity in x space⁵³⁾; if there is no way around this seeming impasse, it would be difficult to see how to proceed in the superspace programme.

One possibility is that off-shell central charges might help and indeed it has recently been suggested that an off-shell formulation of $N = 4$ Poincaré supergravity in five dimensions might be obtainable this way⁵⁴⁾. This theory would then appear to be the generalization of the alternative $N = 2$ theory¹⁶⁾ with a

"centrally charged linear multiplet". In $N = 2$ in four dimensions this multiplet turned out to be a scalar multiplet but in five-dimensional $N = 4$ it is five Sohnius-Stelle-West "vector" multiplets⁵¹⁾. (In fact, the vectors are antisymmetric tensors off-shell.) One should also bear in mind the fact that off-shell central charges do not have to be central charges at all, they could be "Abelian charges" as we discussed in Section 2, and for this reason the extended superspace used in Section 8 might be more than a formal device. However, there are drawbacks to central charges, in particular, there is no way of constructing unconstrained prepotentials which are necessary for the quantization techniques that are currently available.

In connection with the five-dimensional $N = 4$ theory just mentioned, it is interesting to observe that after dimensional reduction it does not separate into $N = 4$ supergravity and $N = 4$ Maxwell theory because of the presence of five antisymmetric tensors. It is well known that supergravity in 11 dimensions⁵⁵⁾ does not reduce straightforwardly to the $N = 8$ theory given here; one has to perform duality transformations on the antisymmetric tensors²⁰⁾. There is therefore the possibility that 11 dimensional supergravity has an off-shell extension, but it does not reduce to a four-dimensional theory which would separate into sub-theories under truncation. If this were the case then the various no-go theorems that have been discussed would be circumvented. We should also point out that even in $N = 4$ there is the possibility that one of the dimension zero scalars is an antisymmetric tensor⁵⁶⁾; if this version has an off-shell extension then it is not clear how it would be related to the $N = 4$ conformal theory where there are no gauge antisymmetric tensors.

If there is a solution to the auxiliary field problem, there would still be considerable technical problems involved in passing to the quantum theory. This is because the solutions to the constraints are rather complicated for $N > 1$; for example, in linearized $N = 2$ Poincaré theory the prepotential has dimension $-5/2$ and the associated gauge transformations are highly non-trivial⁵⁾. This is because an unconstrained superfield in extended supersymmetry contains a large number of representations most of which have to be projected out by gauge invariance^{6),57)}. These representations will, in general, contain higher spins than the supergravity multiplet and in background gauges this leads to the problem of infinite sequences of ghosts⁶⁾. Another technical problem is that of the action itself. In N extended theories, the full superspace measure has dimension $2N - 4$ which means that the Lagrangian has negative dimensions for $N \geq 2$ (there is a factor of $1/\kappa^2$ to be taken into account). For rigid supersymmetry a solution has been found to this problem⁵⁸⁾, but as yet the full theory of subintegration

in curved superspace has not been developed. We should also mention that in superspace one cannot construct the analogues of the "on-shell" Lagrangians that have been found in x space, even in principle. This is because the superspace method has strict supersymmetry whereas in x space one can demand the weaker condition that the algebra only closes when the field equations are satisfied. Imposition of strict closure for supergravity constrains the actions to vanish.

Finally, even if the auxiliary fields turn out to be non-existent, this does not imply that quantum supergravity does not make sense, and it has recently been shown that no additional complications arise in the quantization of 11-dimensional supergravity⁵⁹⁾. It may then be that the best way to proceed is to use $N = 1$ superfields, although in this case the manifest covariance of extended superspace would no longer be apparent.

APPENDIX A - NOTATIONS AND CONVENTIONS

The indices we use are as follows:

	Tangent space		Curved space	
Vector	$a, b, \dots$		$m, n, \dots$	
Spinor	$\alpha, \beta, \dots$	$\left\{ \begin{array}{l} \alpha, \beta, \dots \\ \underline{\alpha}, \underline{\beta}, \dots \end{array} \right\}$	$A, B, \dots$	$\left\{ \begin{array}{l} M, N \\ \underline{M}, \underline{N} \end{array} \right\}$
Internal	$i, j, \dots$		$\underline{\mu}, \underline{\nu}$	

There is some risk of overlap between the tangent space internal symmetry indices $i, j, \dots$ and the curved vector indices $m, n, \dots$ but it should be clear from the context which set we are referring to. A tangent space vector, X^A , therefore has components

$$X^A = (X^a, X^\alpha, X^{\dot{i}}) = (X^a, X^\alpha_i, X^{\dot{i}i}) \quad (A.1)$$

and a covector, Y_A , has components

$$Y_A = (Y_a, Y_\alpha, Y_{\dot{i}}) = (Y_a, Y_\alpha^i, Y_{\dot{i}i}) \quad (A.2)$$

The summation convention is

$$X^A Y_A = X^a Y_a + X^\alpha Y_\alpha - X^{\dot{i}} Y_{\dot{i}} \quad (A.3)$$

Under complex conjugation one has

$$X^\alpha_i \rightarrow (\overline{X^\alpha_i}) = \overline{X}^{\dot{i}i} \quad (A.4)$$

i.e., dotted (undotted) indices become undotted (dotted) and internal indices are raised (lowered). In addition, the order of fermionic quantities in a product is reversed under complex conjugation. The vectorial metric is

$$\eta_{ab} = (+1, -1, -1, -1) \quad (A.5)$$

and the two component spinor metric is

$$\epsilon^{\alpha\beta} = \epsilon_{\alpha\beta} = \begin{pmatrix} 0 & 1 \\ -1 & 0 \end{pmatrix} \quad (\text{A.6})$$

and similarly for dotted indices. Spinor indices are raised and lowered according to the rules

$$X_{\alpha i} = X_i^\beta \epsilon_{\beta\alpha} ; \quad X_i^\alpha = \epsilon^{\alpha\beta} X_{\beta i} \quad (\text{A.7})$$

Any vector index may be converted to spinor indices by the σ matrix

$$X_{\alpha i} = (\sigma^a)_{\alpha i} X_a ; \quad X_a = \frac{1}{2} (\sigma_a)^{\alpha i} X_{\alpha i} \quad (\text{A.8})$$

where

$$(\sigma^a)_{\alpha i} = (1, \vec{\sigma}) \quad (\text{A.9})$$

The matrix $(\sigma^{ab})_{\alpha\beta}$ is defined by

$$(\sigma^{ab})_{\alpha\beta} = \frac{1}{4} \left((\sigma^a)_{\alpha i} (\sigma^b)^{i\beta} - a \leftrightarrow b \right) \quad (\text{A.10})$$

and is symmetric on α, β .

For a product of two quantities one has

$$X^A Y^B = (-1)^{AB} Y^B X^A \quad (\text{A.11})$$

where the factor $(-1)^{AB}$ is +1 unless both indices are spinorial. A tensor X_{AB} is graded antisymmetric if

$$X_{AB} = -(-1)^{AB} X_{BA} \quad (\text{A.12})$$

and graded symmetric if

$$X_{AB} = (-1)^{AB} X_{BA} \quad (\text{A.13})$$

The cyclic sum over ABC of $X_{A,BC}$ which is graded antisymmetric on BC is

$$\sum_{(ABC)} X_{ABC} = X_{ABC} + (-1)^{C(A+B)} X_{CAB} + (-1)^{A(B+C)} X_{BCA} \quad (\text{A.14})$$

(Anti) symmetrizations are always defined with combinatorial factors, e.g.,

$$X_{[ij]} = \frac{1}{2} (X_{ij} - X_{ji}) \quad (A.15)$$

For the most part we work with two component indices only and express vector indices in terms of them via (A.9). For example, a (complex) vector field strength has the two component form

$$F_{\alpha\dot{\alpha},\beta\dot{\beta}} = (\sigma^a)_{\alpha\dot{\alpha}} (\sigma^b)_{\beta\dot{\beta}} F_{ab} = \epsilon_{\dot{\alpha}\dot{\beta}} M_{\alpha\beta} + \epsilon_{\alpha\beta} \bar{N}_{\dot{\alpha}\dot{\beta}} \quad (A.16)$$

where M and N are symmetric.

Therefore, if either M or N is zero, F is (anti) self-dual

$$F_{ab} = \pm \frac{i}{2} \epsilon_{abcd} F^{cd} \quad (A.17)$$

Similarly, for a spin 3/2 field $\psi_{a\alpha}$, the field strength is

$$\begin{aligned} \psi_{\alpha\dot{\alpha},\beta\dot{\beta},\gamma} &= (\sigma^a)_{\alpha\dot{\alpha}} (\sigma^b)_{\beta\dot{\beta}} (\partial_a \psi_{b\dot{\gamma}} - \partial_b \psi_{a\dot{\gamma}}) \\ &= \epsilon_{\alpha\beta} (\bar{\psi}_{\dot{\alpha}\dot{\beta}\dot{\gamma}} + \epsilon_{\dot{\alpha}\dot{\gamma}} \bar{\psi}_{\dot{\beta}} + \epsilon_{\dot{\beta}\dot{\gamma}} \bar{\psi}_{\dot{\alpha}}) \\ &\quad + \epsilon_{\dot{\alpha}\dot{\beta}} \psi_{\alpha\beta\gamma} \end{aligned} \quad (A.18)$$

where the functions on the right-hand side are irreducible under $SL(2, \mathbb{C})$.

Throughout the paper we use "geometrical" dimensions. That is, $[x] = -1$, $[\theta] = -1/2$ (mass) and the dimensions of the various fields are determined via their appearances in geometrical quantities. For conformal supergravity, the fields therefore have the correct dimensions, but in Poincaré supergravity the fields have dimension one less than canonical dimensions. This is because there should really be a factor κ multiplying H in the expansion (3.22).

APPENDIX B - DIFFERENTIAL FORMS IN SUPERSPACE

There is no problem in generalizing differential forms to superspace and we have followed the approach developed in Ref. 21). There is an alternative version⁶⁰⁾ in which one employs single rather than double grading, but the two approaches are equivalent⁶¹⁾.

The basic rule for the co-ordinate one-forms dz^M is

$$dz^M \wedge dz^N = -(-1)^{MN} dz^N \wedge dz^M \quad (B.1)$$

i.e., they anticommute except for $d\theta \wedge d\theta$. A general p form ϕ^p is given by

$$\phi^p = dz^{M_1} \wedge \dots \wedge dz^{M_p} \phi_{M_1 \dots M_p} \quad (B.2)$$

where the functions $\phi_{M_1, \dots, M_p}$ are totally (graded) antisymmetric. A p form ϕ^p and a q form ϕ^q obey the following rule

$$\phi^p \wedge \phi^q = (-1)^{pq} \phi^q \wedge \phi^p \quad (B.3)$$

as for ordinary differential forms. The exterior derivative acts on ϕ^p to give a $(p+1)$ form;

$$d\phi^p = dz^{M_1} \wedge \dots \wedge dz^{M_p} \wedge dz^N \partial_N \phi_{M_1 \dots M_p} \quad (B.4)$$

and enjoys the property that $d^2 = 0$. The exterior derivative starts from the right so that

$$d(\phi^p \wedge \phi^q) = \phi^p \wedge d\phi^q + (-1)^q d\phi^p \wedge \phi^q \quad (B.5)$$

In two component notation, many components of field strengths are obtained from stated ones by complex conjugation. For example, since $E_{\alpha\beta} = \bar{E}_{\dot{\alpha}\dot{\beta}}$ one has

$$T_{\alpha\beta}^{\dot{\gamma}} = \overline{T_{\dot{\alpha}\dot{\beta}}^{\gamma}} \quad (B.6)$$

since D is a real operator. To calculate $T_{\alpha\beta}^{\dot{\gamma}}$ from $T_{\dot{\alpha}\dot{\beta}}^{\gamma}$, for example, one may use (B.6). However, there is a slight subtlety in that

$$\overline{(E^{\dot{p}} \wedge E^{\dot{q}})} = - E^{\dot{p}} \wedge E^{\dot{q}} \quad (\text{B.7})$$

This follows from (B.1). So for example

$$T_{\dot{p}}^{\dot{q}} = - \overline{T_{\dot{q}}^{\dot{p}}} \quad (\text{B.8})$$

APPENDIX C - GLOBAL SYMMETRY GROUPS

The properties of global symmetry groups present in extended supergravity theories have been given in a form useful for physicists by Cremmer and Julia²⁰⁾, and we briefly recall their results here for convenience. The group E_7 is most easily described by giving the transformation rules of the fundamental (56) representation in a complex basis. Let x_{ij} be antisymmetric on i, j ($=1, \dots, 8$), then the basic E_7 transformation is

$$\delta x_{ij} = x_{kl} L^{kl}{}_{ij} + \bar{x}^{kl} \Sigma_{kl}{}^{ij} \quad (C.1)$$

where

$$L^{kl}{}_{ij} = 2 \delta \begin{bmatrix} k & l \\ i & j \end{bmatrix} \quad (C.2)$$

$$\Sigma_{ijkl} = \Sigma [ijkl] \quad (C.3)$$

$$\bar{\Sigma}^{ijkl} = \frac{1}{4!} \epsilon^{ijklmnpq} \Sigma_{mnpq} \quad (C.4)$$

$$\bar{L}_i{}^j = -L^j{}_i \quad (C.5)$$

$$L^i{}_i = 0 \quad (C.6)$$

In this way of writing E_7 , the splitting into the maximal compact sub-group $SU(8)$ (i.e., $L^i{}_j$) and the non-compact part represented by the 70 parameters Σ_{ijkl} emerges clearly. Equation (C.1) may be written in block matrix form:

$$\delta \begin{pmatrix} x_{ij} \\ \bar{x}^{ij} \end{pmatrix} = \begin{pmatrix} x_{kl} & \bar{x}^{kl} \end{pmatrix} \begin{pmatrix} L^{kl}{}_{ij} & \bar{\Sigma}^{kl}{}^{ij} \\ \Sigma_{kl}{}^{ij} & \bar{L}^{kl}{}_{ij} \end{pmatrix} \quad (C.7)$$

from which the structure of the Lie algebra is seen to be the same as of the matrix $\hat{\Omega}$ of (8.38). For $N = 4$ and 5 the groups $SU(4) \times SU(1,1)$ and $U(5,1)$ may be represented in the same way although the self-duality condition (C.4) and the tracelessness of L (C.6) no longer hold. For $N = 6$, one can obtain $SO^*(12)$ by truncating (C.1) but x_{78} must be retained:

$$\begin{aligned}\delta x_{ij} &= x_{he} L^{he}_{ij} + \bar{x}^{he} \Sigma_{heij} + 2\bar{x}^{78} \Sigma_{78ij} \\ \delta x_{78} &= -x_{78} L^i_i + \bar{x}^{he} \Sigma_{he78}\end{aligned}\tag{C.8}$$

where i, j now run from 1 to 6 and

$$\Sigma_{ij78} = \frac{1}{4!} \epsilon_{ijkemn} \bar{\Sigma}^{klemn}\tag{C.9}$$

Equation (C.7) now takes on a slightly more complicated form and may be obtained straightforwardly using (C.8).

REFERENCES

- 1) M. Grisaru, M. Rocek and W. Siegel, Phys. Rev. Lett. 45 (1980) 1063;
W. Caswell and D. Zanon, Phys. Lett. 100B (1980) 152.
- 2) O.V. Tarasov, and A.A. Vladimirov, Dubna preprint (1980).
- 3) M. Grisaru and W. Siegel, Caltech preprint CALT-68-816 (1981).
- 4) L. Mezincescu, JINR preprint P2-12572 (1979).
- 5) W. Siegel and S.J. Gates, Linearized $N = 2$ Supergravity in Superspace,
Caltech preprint (1981).
- 6) W. Siegel and S.J. Gates, Superprojectors, Caltech preprint (1981).
- 7) P. Howe, K. Stelle and P. Townsend, CERN preprint TH-3098 (1981).
- 8) P. Howe and U. Lindström, in "Superspace and Supergravity", eds. S.W. Hawking
and M. Rocek, C.U.P. (1981).
- 9) P. Howe and U. Lindström, Nucl. Phys. B181 (1981).
- 10) R.E. Kallosh, Phys. Lett. 99B (1981) 122; J.E.T.P. Lett. 33 (1981) 292.
- 11) A. Salam and J. Strathdee, Nucl. Phys. B76 (1974) 477.
- 12) J. Wess and B. Zumino, Phys. Lett. 66B (1977)
R. Grimm, J. Wess and B. Zumino, Phys. Lett. 73B (1978) 415.
- 13) M. Kaku and P.K. Townsend, Phys. Lett. 76B (1978) 54;
A. Das, M. Kaku and P.K. Townsend, Phys. Rev. Lett. 40 (1978) 1215;
B. de Wit, J.W. van Holten and A. van Proeyen, Phys. Lett. 95B (1980) 51.
- 14) P. Breitenlohner and M. Sohnius, Nucl. Phys. B178 (1981) 151.
- 15) E.S. Fradkin and M.A. Vasiliev, Lett. Nuovo Cimento 25 (1979) 79; Phys. Lett.
85B (1979) 47;
B. de Wit and J.W. van Holten, Nucl. Phys. B158 (1979) 530;
B. de Wit, J.W. van Holten, Nucl. Phys. B167 (1980) 186.
- 16) E. Bergshoeff, M. de Roo, J.W. van Holten, B. de Wit and A. van Proeyen in
"Superspace and Supergravity", eds. S.W. Hawking and M. Rocek, C.U.P. (1980);
B. de Wit, J.W. van Holten and A. van Proeyen, Nucl. Phys. B167 (1980) 186.
- 17) Articles by K.S. Stelle and P.C. West; S.J. Gates; W. Siegel in "Supergravity",
eds. P. van Nieuwenhuizen and D.Z. Freedman, North Holland (1979);
W. Siegel and S.J. Gates, Nucl. Phys. B163 (1980) 519;
S.J. Gates, K.S. Stelle and P.C. West, Nucl. Phys. B169 (1980) 347.
- 18) K.S. Stelle and P.C. West, Phys. Lett. 90B (1980) 393.
- 19) S.J. Gates, Harvard preprint HUTP-80/A011 (1980).
- 20) E. Cremmer and B. Julia, Nucl. Phys. B159 (1979) 141.
- 21) B. Zumino in Proc. Conference at Northern University, eds. R. Arnowitt and
P. North (MIT press) 1975.

- 22) M. Rocek and U. Lindström, Phys. Lett. 79B (1978) 217; Phys. Lett. 83B (1979) 179;
U. Lindström, A. Karlhede and M. Rocek, Stockholm preprint (1980).
- 23) J. Wess and B. Zumino, Phys. Lett. 79B (1978) 394.
- 24) H. Nicolai, P. Townsend and P. van Nieuwenhuizen, Lett. Nuovo Cimento 30 (1981) 315.
- 25) N. Dragon, Z. Phys. C2 (1979) 29.
- 26) M. Sohnius in "Superspace and Supergravity", eds. S.W. Hawking and M. Rocek, C.U.P. (1981).
- 27) W. Siegel, Nucl. Phys. B177 (1981) 325.
- 28) L. Brink and P. Howe, Phys. Lett. 88B (1979) 268.
- 29) E. Bergshoeff, M. de Roo and B. de Wit, Nucl. Phys. B182 (1981) 173.
- 30) S. Ferrara and B. Zumino, Nucl. Phys. B134 (1978) 301.
- 31) M. Sohnius, Nucl. Phys. B136 (1978) 275.
- 32) L. Brink, M. Gell-Mann, J. Schwarz and P. Ramond, Phys. Lett. 74B (1978) 336.
- 33) S. Ferrara and P. van Nieuwenhuizen, Ann. of Phys. 127 (1980) 74.
- 34) L. Castellani, S.J. Gates and P. van Nieuwenhuizen, Stony Brook preprint ITP-SB-80-9 (1980).
- 35) P. van Nieuwenhuizen and P.C. West, Nucl. Phys. B172 (1980)
- 36) L. Brink, M. Gell-Mann, J. Schwarz and P. Ramond, Phys. Lett. 76B (1978) 417.
- 37) E. Cremmer and S. Ferrara, Phys. Lett. 91B (1980) 61.
- 38) L. Brink and P. Howe, Phys. Lett. 91B (1980) 384.
- 39) R. Grimm, J. Wess and B. Zumino, Nucl. Phys. B152 (1979) 255, ibid. 80B (1978) 138.
- 40) P. Howe and R.W. Tucker, Phys. Lett. 80B (1978) 138.
- 41) W. Siegel, Phys. Lett. 80B (1979) 220.
- 42) P. Howe, Phys. Lett. 100B (1981) 389.
- 43) M. Kaku, P.K. Townsend and P. van Nieuwenhuizen, Phys. Lett. 69B (1977) 304; Phys. Rev. D17 (1978) 3179.
- 44) J. Ramao, A. Ferber and P.G. Freund, Nucl. Phys. B126 (1977) 429;
F. Mansouri and C. Schaer, Phys. Lett. 83 (1979) 327; Phys. Lett. 101B (1981) 51.
- 45) S. Ferrara and P. van Nieuwenhuizen, Phys. Lett. 74B (1978) 333;
K.S. Stelle and P.C. West, Phys. Lett. 74B (1978) 330.
- 46) P. Breitenlohner and M. Sohnius, Nucl. Phys. B165 (1980) 483.

- 47) P. Breitenlohner, Phys. Lett. 67B (1977) 49;
W. Siegel and S.J. Gates, Nucl. Phys. B147 (1979) 77.
- 48) R. Grimm in "Unification of Fundamental Particle Interactions", eds. S. Ferrara,
J. Ellis and P. van Nieuwenhuizen, Plenum Press (1981).
- 49) E. Sokatchev in "Superspace and Supergravity", eds. S.W. Hawking and M. Rocek,
C.U.P. (1981); Phys. Lett. 100b (1981) 466.
- 50) R. Grimm, M. Sohnius and J. Wess, Nucl. Phys. B133 (1978) 275.
- 51) M. Sohnius, K.S. Stelle and P.C. West in "Superspace and Supergravity", eds.
S.W. Hawking and M. Rocek, C.U.P. (1981).
- 52) W. Siegel and M. Rocek, Caltech preprint (1981).
- 53) J.G. Taylor and V.O. Rivelles, Kings College preprint (1981).
- 54) P. Howe and U. Lindström, CERN preprint TH-3075 (1981).
- 55) E. Cremmer, B. Julia and J. Scherk, Phys. Lett. 76B (1978) 409.
- 56) H. Nicolai and P.K. Townsend, Phys. Lett. 98B (1981) 257.
- 57) E. Sokatchev and V. Rittenberg, Bonn preprint HE-81-5 (1981).
- 58) P. Howe, K.S. Stelle and P.K. Townsend, CERN preprint TH-3065 (1981).
- 59) B. de Wit, P. van Nieuwenhuizen and A. van Proeyen, NIKHEF preprint (1981).
- 60) F.A. Berezin, Moscow preprint ITEP-71 (1979).
- 61) B. Zumino in "Unification of the Fundamental Particle Interactions", eds.
S. Ferrara, J. Ellis and P. van Nieuwenhuizen, Plenum Press (1980).