Search for Dark Matter Candidate with the ATLAS detector

Marie-Hélène Genest (LMU Munich)
On behalf of the ATLAS Collaboration

RICAP 2011

GEFÖRDERT VOM

Bundesministerium
für Bildung
und Forschung

A brief reminder of SUSY

Spin 1/2

Spin 0

Spin 1

Spin 0

Spin 1/2

If R-parity $(-1)^{L+3B+2S}$ is conserved:

- SM particles have $R = +1$
- SUSY particles have $R = -1$

The lightest sparticle (LSP) is stable: Dark matter candidate!

Typical signature

- Pair of gluinos/squarks produced by strong interactions
- Their decays give high- p_T jets and charginos/neutralinos
- Charginos/neutralinos decays can give leptons and the decay chain stops when the LSP is produced (R-parity conserving scenarios)
- The pair of stable LSP produced escapes the detector undetected leading to high transverse missing energy

multi-Jets + n leptons + E_T^{miss}

Standard Model backgrounds ($t\bar{t}$, W +jets, Z +jets, QCD jets and dibosons)

Muon Spectrometer ($|\eta| < 2.7$) : air-core toroids with gas-based muon chambers
Muon trigger and measurement with momentum resolution **< 10% up to $E_\mu \sim 1$ TeV**

3-level trigger
reducing the
rate to ~ 200 Hz

Hadronic calorimetry ($|\eta| < 3$): segmentation **0.1×0.1**
Fe/scintillator Tiles (central), Cu/W-LAr (fwd)
E-resolution: $\sigma/E \sim 50\%/\sqrt{E} \oplus 0.03$
Forward calorimetry: W/LAr $\sigma/E \sim 90\%/\sqrt{E} \oplus 0.07$

Length : ~ 46 m
Radius : ~ 12 m
Weight : ~ 7000 tons
 $\sim 10^8$ electronic channels
3000 km of cables

Inner Detector ($|\eta| < 2.5$, $B=2$ T):
Precise tracking and vertexing,
 $\sigma/p_T \sim 3.8 \times 10^{-4} p_T (\text{GeV}) \oplus 0.015$

Electromagnetic calorimeter: Pb-Liquid Argon
Accordion
e/ γ trigger, identification and measurement
E-resolution: $\sigma/E \sim 10\%/\sqrt{E} \oplus 0.007$
granularity : $0.025 \times 0.025 \oplus$ strips

Data accumulated in 2010

- Excellent LHC performance

- Very good detector efficiency:
 - Inner tracking detectors: 99.1% to 100%
 - Calorimeters: 90.7% to 100%
 - Muon detectors: 96.2% to 99.8%

ATLAS public results of R-parity conserving SUSY searches

Covered today

one lepton, jets, and missing transverse momentum	ArXiv:1102.2357 Phys. Rev. Lett. 106, 131802 (2011)
jets and missing transverse momentum	ArXiv:1102.5290, submitted to PLB
Multilepton final state with jets and missing transverse momentum	ATLAS-CONF-2011-039
b-tagged jets and missing transverse momentum	ArXiv:1103.4344, submitted to PLB
Dilepton final states with missing transverse momentum	ArXiv:1103.6214, submitted EPJLC Same-flavour: ArXiv:1103.6208, accepted EPJLC

Some useful variables

- $\Delta\phi(\text{jets}, E_T^{\text{miss}})$
 - Cutting on $\Delta\phi$ eliminates events in which E_T^{miss} is closely related to one of the leading jets (QCD)
- Effective mass m_{eff} (scalar sum of sel. jets & leptons p_T and E_T^{miss})
 - peaks at a value which is correlated with the mass of the pair of SUSY particles produced in the pp interaction
- The transverse mass m_T $m_T^2 \equiv 2|\mathbf{p}_T^\ell||E_T^{\text{miss}}| - 2\mathbf{p}_T^\ell \cdot \vec{E}_T^{\text{miss}}$
 - useful to remove BG in which a W decays leptonically

The 0-lepton channel

Select events with jets, missing transverse momentum and no lepton (veto e/μ)

*arXiv:1102.5290
submitted to PLB*

Defining the signal regions

	A	B	C	D
Pre-selection	Number of required jets	≥ 2	≥ 2	≥ 3
	Leading jet p_T [GeV]	> 120	> 120	> 120
	Other jet(s) p_T [GeV]	> 40	> 40	> 40
	E_T^{miss} [GeV]	> 100	> 100	> 100
Final selection	$\Delta\phi(\text{jet}, \vec{P}_T^{\text{miss}})_{\text{min}}$	> 0.4	> 0.4	> 0.4
	$E_T^{\text{miss}}/m_{\text{eff}}$	> 0.3	—	> 0.25
	m_{eff} [GeV]	> 500	> 500	> 1000
	m_{T2} [GeV]	—	> 300	—

gluino/gluino and
squark/gluino production

Low- and
high-mass
squark pair
production

Defining the signal regions

		A	B	C	D
Pre-selection	Number of required jets	≥ 2	≥ 2	≥ 3	≥ 3
	Leading jet p_T [GeV]	> 120	> 120	> 120	> 120
	Other jet(s) p_T [GeV]	> 40	> 40	> 40	> 40
	E_T^{miss} [GeV]	> 100	> 100	> 100	> 100
Final selection	$\Delta\phi(\text{jet}, \vec{P}_T^{\text{miss}})_{\min}$	> 0.4	> 0.4	> 0.4	> 0.4
	$E_T^{\text{miss}}/m_{\text{eff}}$	> 0.3	–	> 0.25	> 0.25
	m_{eff} [GeV]	> 500	–	> 500	> 1000
	m_{T2} [GeV]	–	> 300	–	–

Trigger requirements

Reject the QCD BG

Optimize for SUSY

Stranverse mass m_{T2} :

- The sum of the transverse missing momentum of the two neutralinos is know: $\vec{q}_T^{(1)} + \vec{q}_T^{(2)} = \vec{E}_T^{\text{miss}}$
- Using this constraint, calculate the stransverse mass as:

$$m_{T2}(\vec{p}_T^{(1)}, \vec{p}_T^{(2)}, \vec{p}_T) \equiv \min_{\vec{q}_T^{(1)} + \vec{q}_T^{(2)} = \vec{E}_T^{\text{miss}}} \{ \max(m_T(\vec{p}_T^{(1)}, \vec{q}_T^{(1)}), m_T(\vec{p}_T^{(2)}, \vec{q}_T^{(2)})) \}$$

Main backgrounds

W+jets: $\tau\nu$ decay or missed e/μ

Z+jets: $\nu\nu$ decay

Top pair production: τ decay of a W

Alpgen with NNLO normalization

MC@NLO

Cross-check on data:

Z(l l)+jets control sample with leptons removed to recalculate the missing transverse momentum

W $\rightarrow \mu\nu$ control sample with the muon replaced by a MC τ decay

Results

95% CL limits on cross section $\cdot$ acceptance $\cdot$ efficiency:
 1.3 pb (A), 0.35 pb (B), 1.1 pb (C) and 0.11 pb (D)

Exclusion plot

The limit does not depend strongly on $\tan\beta$, A_0 or $\text{sign}(\mu)$

The 1-lepton channel

Select events with jets, missing transverse momentum and exactly one lepton (e/μ)

ArXiv:1102.2357

Phys. Rev. Lett. 106, 131802 (2011)

Defining the signal region

The isolated one-lepton requirement suppresses QCD multijet and allows a lepton-based trigger

- Exactly one lepton (e/μ) with $p_T > 20$ GeV
 - At least 3 jets with $p_T > 60, 30, 30$ GeV
 - $\Delta\phi(\text{jets}, E_T^{\text{miss}}) > 0.2$
 - $E_T^{\text{miss}}/m_{\text{eff}} > 0.25$
 - $m_T > 100$ GeV
 - $m_{\text{eff}} > 500$ GeV
- } gluino/squark cascade decay with intermediate steps
- } Reduce the QCD BG further
- } Suppresses W+jets and tt
- } Optimize for SUSY

Main backgrounds: W+jets and tt

W control region: no b-jet

Top control region: ≥ 1 b-jet

$$\underbrace{N(\bar{t}t \text{ pred.}, SR)}_{\text{predicted events in signal region}} = \underbrace{(N(\text{fitted } \bar{t}t(\text{data}), CR))}_{\text{measured events in control region - other BG}} \times \underbrace{\frac{N(\bar{t}t(MC), SR)}{N(\bar{t}t(MC), CR)}}_{\text{extrapolation factor CR to SR}}$$

③

①

②

From data

From MC

Method cross-checked with additional control regions

Results

95% CL limits on cross section $\cdot$ acceptance $\cdot$ efficiency:
 0.065 pb (electron), 0.073 pb (muon)

Exclusion plot

parameters at GUT scale

1. Unified gaugino(scalar) mass $m_{1/2}(m_0)$
3. Ratio of H_u, H_d vevs $\tan\beta$
4. Trilinear coupling A_0
5. Higgs mass term $\text{sgn}(\mu)$

0- and 1-lepton combination

Exclude a mass of 815 GeV for $m_{\text{gluino}} = m_{\text{squark}}$

How to use the data

- For each signal region and analysis channel, the efficiency x acceptance is provided
- The LHA SUSY files are provided
 - validate your setup
 - interpret the data in your model

Summary and outlook

- The search is on!
- ATLAS has already produced many results (in some scenarios the most stringent limits to date) in various channels and more are in the pipeline
- Already 250 pb^{-1} accumulated in 2011 with a new record luminosity of $8.3 \times 10^{32} \text{ cm}^{-2} \text{ s}^{-1}$!
- In 2011: more than 1 fb^{-1} of data with luminosities on the order of $10^{33} \text{ cm}^{-2} \text{ s}^{-1}$
- Sensitivity beyond 1 TeV already for 2011

This year is the SUSY year!

Many more results...

Have a look here:

<https://twiki.cern.ch/twiki/bin/view/AtlasPublic/SupersymmetryPublicResults>

Backup slides

Details of the results

	Signal region A	Signal region B	Signal region C	Signal region D
Total SM	$118 \pm 25[\text{u}]^{+32}_{-23}[\text{j}] \pm 12[\mathcal{L}]$	$10.0 \pm 4.3[\text{u}]^{+4.0}_{-1.9}[\text{j}] \pm 1.0[\mathcal{L}]$	$88 \pm 18[\text{u}]^{+26}_{-18}[\text{j}] \pm 9[\mathcal{L}]$	$2.5 \pm 1.0[\text{u}]^{+1.0}_{-0.4}[\text{j}] \pm 0.2[\mathcal{L}]$
Data	87	11	66	2

Electron channel	Signal region
Observed events	1
Fitted top events	1.34 ± 0.52 (1.29)
Fitted W/Z events	0.47 ± 0.40 (0.46)
Fitted QCD events	$0.0^{+0.3}_{-0.0}$
Fitted sum of background events	1.81 ± 0.75
Muon channel	Signal region
Observed events	1
Fitted top events	1.76 ± 0.67 (1.39)
Fitted W/Z events	0.49 ± 0.36 (0.71)
Fitted QCD events	$0.0^{+0.5}_{-0.0}$
Fitted sum of background events	2.25 ± 0.94

SUSY production at LHC

SUSY production at LHC

If $m(\text{gluino}) \sim 500$ GeV, expect ~ 2000 gluino pairs produced for 1 fb^{-1} of data

**The cross sections for SUSY processes are way higher at LHC
High-impact results possible with first data!**

Object identification

- **Jets** (anti-Kt, R=0.4): $p_T > 20$ GeV, $|\eta| < 2.5$
 - Reject events compatible with noise or cosmics
 - Remove if $\Delta R(\text{jet}, \text{electron}) < 0.2$
- **Electrons**: $p_T > 20$ GeV, $|\eta| < 2.47$
 - Outside problematic regions of the calorimeter
 - Remove if $\Delta R(\text{jet}, \text{electron}) < 0.4$
- **Muons**: $p_T > 20$ GeV, $|\eta| < 2.4$, Sum p_T of tracks ($\Delta R < 0.2$) < 1.8 GeV
 - Remove if $\Delta R(\text{jet}, \text{muon}) < 0.4$
- **Missing transverse momentum (E_T^{miss}):**
 - sum over the transverse momentum of all jets (up to $|\eta| < 4.9$), electrons, muons and all calorimeter clusters not associated to such objects

$$\Delta R = \sqrt{(\Delta\eta)^2 + (\Delta\phi)^2}$$

ϕ : azimuthal angle around the beam pipe

$\eta = -\ln \tan(\theta/2)$ where θ is the polar angle

E_T^{miss} performance

MET resolution

QCD background: 1-lepton channel

Evaluated using the 'matrix method' which plays on the difference in isolation between the leptons in QCD events with respect to signal leptons

- **Loose** control sample with isolation criteria relaxed with respect to the **tight** SUSY selections
- Define two categories: QCD leptons (**Q**) and non-QCD leptons (**ϕ**)
- ϵ is the probability that a loose lepton is also tight

$$N_{tight}^{obs} = N_{tight}^{\phi} + N_{tight}^Q$$

$$N_{loose\ not\ tight}^{obs} = (1/\epsilon_{\phi} - 1) N_{tight}^{\phi} + (1/\epsilon_Q - 1) N_{tight}^Q$$

The quantities in **red** are measured: solve the equations and extract the number of QCD events

electron channel

$$N_{tight}^Q < 0.3$$

muon channel

$$N_{tight}^Q < 0.5$$

QCD background: 0-lepton channel

Partially data-driven estimate:
 rescale the MC in the control region
 defined by reversing the $\Delta\phi$ cut
 $< 10\%$ of the total BG
 (cross checked with two other methods)

Systematic uncertainties in the 1-lepton channel

- Background:
 - MC modeling of the E_t^{miss} and m_T distributions (variation of MC generator and internal generator parameters)
 - Finite statistics in the control regions
 - Experimental uncertainties dominated by the jet energy scale, b-tagging and luminosity uncertainties
- Signal:
 - Variation of the factorization and renormalization scales in Prospino
 - PDF uncertainties (eigenvector sets provided by CTEQ6)
 - Calculated separately for each production process

Limit setting

$$\begin{aligned} L(n, \theta^0 | \mu, b, \theta) &= P_{\text{SR}} \times P_{\text{CR}} \times P_{\text{Syst}} \\ &= P(n_S | \lambda_S(\mu, b, \theta)) \times \prod_{i \in \text{CS}} P(n_i | \lambda_i(\mu, b, \theta)) \times P_{\text{Syst}}(\theta^0, \theta) \end{aligned}$$

The statistical treatment is based on the profile LLR, defined in the usual way as

$$\Lambda(\mu) \equiv \Lambda(\mu, n, \theta^0) \equiv -2 \left(\ln L(n, \theta^0 | \mu, \hat{b}, \hat{\theta}) - \ln L(n, \theta^0 | \hat{\mu}, \hat{b}, \hat{\theta}) \right)$$

where $\hat{\mu}, \hat{b}, \hat{\theta}$ maximize the likelihood function, and $\hat{b}, \hat{\theta}$ maximize the likelihood for the specific, fixed value of the signal strength μ , and the data n, θ^0 .

- The 1-sigma exclusion band has the same meaning as in a typical Higgs analysis, which show the upper limit on the x-section (but here the x-section is mapped on the m_0, m_{12} plane): any observation of number of events in the signal region that agrees with the bkg only prediction within 68% CL, gives an observed exclusion limit that lies within the 1-sigma uncertainty band.
- It can also be seen to correspond roughly to the 68% and 99% CL exclusion lines. The limit and bands have been obtained directly with toy MC. The coverage is therefore guaranteed to be exact.