TRIGGERING ON HADRONIC TAU DECAYS: ATLAS MEETS THE CHALLENGE

WHY DO WE NEED A TAU TRIGGER?

One of the main aims of ATLAS is the search for the Higgs boson (H^0).

- Low mass Standard Model (SM) H^0 decays to two taus with high branching ratio

ATLAS was also built to search for beyond SM (BSM) physics.

- BSM models with high mass particles will generally prefer to decay into SM taus.

Taus produced in interesting physics + decay rate to hadrons > 64%
= Dedicated hadronic tau trigger required

TAUS AND QCD JETS

QCD jet events are a large background for physics with taus $\rightarrow$ QCD jets fake taus

Hadronic tau decays:

- One or three charged pions
- A tau neutrino
- Neutral pions

QCD jets also leave a similar signature in the detector, but taus can be distinguished from jets by looking for:

- Low track multiplicity
- Narrow jet cone
- A region of isolation around the centre of the decay

These differences are exploited in the ATLAS trigger to select the hadronic taus.

TRIGGERING ON TAUS IN ATLAS

- Hardware based trigger
- identifies Regions of Interest (Rois)

- Sum of energy in 2x1 electromagnetic calorimeter (EM) towers and 2x2 hadronic calorimeter (HAD) towers

- EM+HAD cluster energy must pass a required threshold, ie: L1_TAU6 $\Rightarrow$ 7GeV of energy

- A hardware prescale (PS) is applied to reduce the output rate of L1

- A L2 PS is applied depending on the trigger chain
- Tracks and calorimeter showers are reconstructed using full detector granularity within RoI
- Selections are applied to identify taus and discriminate against QCD jets

- EM Radius, a measure of jet size
- Number of tracks

- An EF PS applied to reduce output rate
- Full event is built, using the whole detector, with taus from the passing L2 RoIs firing the trigger
- Algorithms similar to the offline reconstruction are run.
- EF tau trigger selects taus and discriminates against QCD jets with enhanced precision compared to L2.

The figure to the right shows QCD jet rejection with respect to L1 for trigger chains with loose, medium or tight tau identification at L2 and EF

TAU TRIGGER EFFICIENCIES AND QCD JET REJECTION

The tau trigger efficiency (ϵ) - the conditional probability for a reconstructed offline tau to pass the tau trigger - has been measured in Monte Carlo (MC) simulations, and in data using various methods.

$Z \rightarrow \tau^+ \tau^-$ in MC

- ϵ of trigger requiring $E_T > 16\text{GeV}$ and loose tau identification (ID) - tau16_loose
- L1 - no discrimination power (see TAU TRIGGER IN HIGH LUMINOSITY) - $\epsilon \sim 1$
- L2/EF - tau ID applied to reject QCD jets - $\epsilon \sim 0.95$

$Z \rightarrow \tau_{lep}^+ \tau_{had}^-$ in Data

Tau trigger efficiency using real Z bosons decaying to two taus using a tag-and-probe method - one tau decaying hadronically, and the other leptonically.

- Tag triggered lepton
- Check for probe hadronic tau from Z
- Check tau candidate passed trigger
- $E_T > 12\text{GeV}$ - ~94%
- $E_T > 16\text{GeV}$ - ~90%

$W \rightarrow \tau_{had} \nu$ in Data

Tau trigger efficiency using real W bosons decaying to a hadronic tau and missing energy (E_{miss}) using a tag-and-probe method.

- Tag triggered E_{miss}
- Check for probe tau back-to-back with E_{miss}
- Check tau and E_{miss} came from W decay by building transverse mass (m_T)
- Check if tau candidate passed trigger

Efficiency using QCD Jets in Data

Similarities between hadronic tau decays and QCD jets were exploited to measure ϵ - If a jet looks like a tau offline, it should also look like a tau at the trigger level.

- High statistics
- Study high E_T triggers
- Tag-and-probe used to select di-jets in data
- Back-to-back and transverse momentum (p_T) balanced jets
- Tag one jet passing a jet trigger

- Look for tight tau from probe jet
- Check tau fired trigger
- After ~60GeV, for a chain with energy threshold 16GeV and loose ID - $\epsilon \sim 95\%$

QCD Jet Rejection in Data

Efficiency of 12GeV tau trigger chain comparing signal taus and QCD jets

- Agreement between data and QCD MC
- QCD jets fake taus ~30% of the time
- High fake rate used to measure ϵ of high E_T tau triggers (see Efficiency using QCD Jets)

PHYSICS WITH TAU TRIGGERS

Due to the high rates of single tau triggers, they are often combined with triggers from other signatures for use in many physics analyses, forming trigger chains like

- Di-tau triggers
- Tau + lepton triggers
- Tau + E_{miss} triggers

$W \rightarrow \tau_{had} \nu$
W decays to hadronic taus were observed in 2010 using the tau trigger [1], measuring

$55.1 \pm 10.5(stat.) \pm 5.2(syst.)$ events
The W to hadronic tau production cross-section is also being measured, using events from a combined tau + E_{miss} trigger

$H \rightarrow \tau_{had}^+ \tau_{had}^-$

A search for a SM higgs boson decaying into two hadronic taus is also underway, using a di-tau trigger requiring two taus at EF

- Tau 1: $p_T > 29\text{GeV}$ with medium ID
- Tau 2: $p_T > 20\text{GeV}$ with medium ID

TAU TRIGGER IN HIGH $\mathcal{L}$

As the instantaneous luminosity ($\mathcal{L}$) of the LHC increases, so too do the trigger rates. In order to stay within the assigned bandwidth and keep prescaling to a minimum, the triggers must be changed by either:

- Increasing energy thresholds
- Increasing background rejection and trigger efficiency

The plots below show the trigger rate for various L1 tau triggers, and the EF chains that they seed.

To maintain constant L1 rate, L1_TAU6 at $\mathcal{L} = 1 \times 10^{32} \text{cm}^{-2} \text{s}^{-1} \rightarrow$ L1_TAU11 at $\mathcal{L} = 7 \times 10^{32} \text{cm}^{-2} \text{s}^{-1}$

Increasing energy thresholds however is not always ideal.

- Limits the ability to study physics processes with low energy taus.

QCD jet rejection of the L1 tau trigger may be increased by

- Increasing E_T threshold in specific regions of the detector
- This limits the efficiency of the trigger for low energy taus
- E_T isolation at L1 in the EM and/or HAD calorimeters
- This limits the efficiency of the trigger for high energy taus

References

[1] Observation of $W \rightarrow \tau \nu$ Decays with the ATLAS Experiment, *The ATLAS collaboration*, ATLAS-CONF-2010-097, Nov 2010

Other Tau Trigger Posters

https://twiki.cern.ch/twiki/bin/viewauth/Atlas/TapmTau#Presentations_in_Conferences