MICRONUCLEI INDUCTION IN *VICIA FABA* ROOTS

PART II: BIOLOGICAL EFFECTS OF NEUTRONS BELOW 1 cGy

I. Marshall*)

Kernforschungszentrum Karlsruhe,
Institut für Kernphysik, Karlsruhe, Fed. Rep. Germany

M. Bianchi
CERN, Geneva, Switzerland

ABSTRACT

A dose-effect relationship has been established for high-energy neutrons (maximum energy 600 MeV) within a dose range of 0.2 to 80 cGy and for low-energy neutrons produced by a ^{252}Cf source (mean energy 2.35 MeV) for doses between 0.2 and 5 cGy. The number of micronuclei produced by neutron radiations was found to increase linearly with dose. The Relative Biological Effectiveness (RBE) values calculated using ^{60}Co radiation as a reference were, in the high-dose region, 4.7 ± 0.4 and 11.8 ± 1.3 for the high- and low-energy neutrons, respectively. At doses below 1 cGy constant values of 25.4 ± 4.4 and 63.7 ± 12 were reached for the respective neutron energies.

(Submitted to Int. J. Radiat. Biol.)

*) Present address: Rad. Environ. Lab., Brookhaven National Laboratory, Upton, NY, USA.

RESUME

Nous avons établi des courbes d'effet de dose pour des neutrons de haute énergie (énergie max.: 600 MeV) pour des doses comprises entre 0,2 et 80 cGy, et pour des neutrons de basse énergie, provenant d'une source de ^{252}Cf (énergie moyenne: 2,35 MeV), pour des doses entre 0,2 et 5 cGy. Le nombre de micronoyaux augmente linéairement avec la dose. Aux hautes énergies, les valeurs de l'efficacité biologique relative (EBR) calculée en utilisant les rayons γ du ^{60}Co comme radiations de référence étaient de $4,7 \pm 0,4$ et $11,8 \pm 1,3$ respectivement pour les neutrons de haute et de basse énergie. A des doses inférieures à 1 cGy, nous avons trouvé, pour les deux énergies, des valeurs constantes de l'EBR: $25,4 \pm 4,4$ et $63,7 \pm 12$.

ABRISS

Eine Dosis-Effekt-Beziehung wurde aufgestellt für hochenergetische Neutronen (max. Energie 600 MeV) bei Bestrahlungsdosen von 0,2 bis 80 cGy und für Neutronen niederer Energien (mittlere Energie 2,35 MeV), die aus einer ^{252}Cf Quelle produziert wurden, mit Bestrahlungsdosen zwischen 0,2 und 5 cGy. Es zeigte sich, dass die Zahl der durch die Neutronenbestrahlung produzierten Mikrokerne linear mit der Dosis ansteigt. Die Werte für Relative Biologische Wirksamkeit (RBW) wurden unter Bezugnahme auf Resultate mit ^{60}Co Gammastrahlen berechnet und betrugen bei hohen Bestrahlungsdosen $4,7 \pm 0,4$ für hochenergetische Neutronen und $11,8 \pm 1,3$ für Neutronen niederer Energien. Bei Bestrahlungsdosen unter 1 cGy ergaben sich für beiden Neutronenenergien konstante Werte von $25,4 \pm 4,4$ bzw. $63,7 \pm 12$.

1. INTRODUCTION

The Relative Biological Effectiveness (RBE) of neutrons varies both with neutron energy and with dose, the maximum effect being observed for neutrons of 0.43 MeV (Broerse and Barendsen 1973, Hall et al., 1973, Sparrow et al. 1972, Field and Hornsey 1979). Owing to the difference in shape of the dose-effect curves of neutrons and gamma-rays, the RBE value is highest in the low-dose region.

The data reported in the literature for various biological systems using both plant and animal cells suggest RBE values as high as 100 (Sparrow et al. 1972, Bateman and Snead 1970, Vogel 1969, Shellabarger et al. 1980). In view of the high values observed in the dose region of particular interest for radiation protection, it is necessary to have more experimental data and careful evaluation of the damage at doses below 10 cGy in order to reduce the extent of extrapolation in the estimate of the shape of the dose-response curve. From recent investigations on the effects of low doses of neutrons of various energies on lens opacification, similar RBE values were obtained at a dose of 1 cGy (Di Paola et al. 1981). Therefore we decided to extend the investigations into the dose region below 1 cGy, using micronuclei induction in *Vicia faba* roots as a biological indicator.

This system has been used to establish a dose-response curve for ^{60}Co γ -rays over a wide range of doses, and damage produced by low doses of neutrons can easily be detected (Diehl-Marshall and Bianchi 1981, Marshall 1981, Marshall and Bianchi 1982). The neutrons studied in the present experiment were high-energy neutrons produced by 600 MeV protons and neutrons of low energy from a ^{252}Cf source.

2. IRRADIATIONS AND DOSIMETRY

The high-energy neutrons were produced at the CERN Synchro-cyclotron by 600 MeV protons incident on a Be target and extracted in the forward direction. The energy spectra of the secondary neutrons is expected to be rather broad, with a maximum energy equal to the initial 600 MeV proton energy and a peak maximum at about 50 MeV below the incident energy (Dunn 1975, Fischer et al. 1976). The contribution to the dose from γ contamination was about 10% of the total dose (Baarli et al. 1978).

The bean meristems were irradiated in aerated water at a depth of 0.25 g/cm²; this position was chosen since the contribution to the dose from nuclear star interactions is dominant. The dose rate at the position of irradiation was about 20 cGy/h. The doses given were 0.2, 0.5, 1, 5, 15, 40, and 80 cGy.

The ²⁵²Cf neutron source emits neutrons and γ -rays by spontaneous fission. The mean energy of the neutrons is 2.35 MeV, whereas most of the γ -rays are produced with energies up to 1 MeV. The dose measurements were carried out using a 0.1 cm³ tissue-equivalent ionization chamber. The dose rate at the position of irradiation was about 7 cGy/h. The γ contribution to the total absorbed dose, as determined with a Geiger-Müller counter, was 2.9 cGy/h, whereas the neutron dose-rate was 4.1 cGy/h. From measurements with indium activation the neutron flux was calculated to be 4.5×10^5 n/cm²/s. The irradiation position was located 60 mm from the central axis of the source at a water depth of 6 mm inside a circular container made of plexiglas, where aerated water was circulating. The meristems of the beans were positioned at a height of ± 10 mm from the central axis of the source. The doses given were 0.2, 0.5, 1, and 5 cGy.

Details of the cytological technique used are given in Part I.

3. RESULTS

The data obtained for the high-energy neutrons have been compiled over the course of several experiments and most of the data at doses above 1 cGy have already been published (Diehl-Marshall and Bianchi 1981). As mentioned in Part I, the mean number of micronuclei observed in the period between 24 and 48 h after irradiation has been used for a quantitative assessment of the radio-induced damage. The fixations were made at 24, 30, 36, 42, and 48 h after irradiation. The mean number of micronuclei (MN), the mean number of cells with micronuclei, and the mean value of mitotic index (MI) determined for neutrons produced by 600 MeV protons is reported in table 1. When the number of micronuclei was plotted against dose a departure from linearity was observed for the two highest doses, as illustrated in figure 1. Correspondingly, the average number of cells under division dropped with increasing

dose, reaching a value close to 50% for a dose of 80 cGy. At doses below 1 cGy, the percentage of cells undergoing division at the period of time comprised between 24 and 48 h after irradiation remains unaltered.

After applying a correction which takes into account the reduction in mitotic index (see Part I), the data were best fitted by a linear equation as illustrated in figure 2. The α value was 0.48 ± 0.02 and the χ^2/f was 1.4. The fit was made with and without the control value and the intercept was within 1 standard deviation of this value. The mean value of micronuclei observed after irradiation with the dose of 0.2 cGy was 0.19 ± 0.07 , which is close to the value of 0.13 ± 0.03 obtained for controls. However, when the frequency of beans having 0, 1, or more micronuclei was considered, a difference in the distribution was evident, as illustrated in figure 3.

The data obtained after irradiation with low-energy neutrons are reported in table 2 and figure 4. The dose value given in the table corresponds to the total absorbed dose and no subtraction of the γ contamination was attempted. The data obtained for the ^{252}Cf source were best fitted by a linear equation ($E = C + \alpha D$), the α value was 1.2 ± 0.1 , and the χ^2/f was 0.1.

In the case of the high-energy neutrons a split dose of 20 + 20 cGy was also given, using a 1 h interval between the two fractions. The results are reported in table 2 together with the corresponding results for the single dose given during the same experiment. The data indicate the absence of recovery, although a sensitization effect cannot be excluded.

The RBE value, which is the ratio of the doses ($D_{\text{ref.rad.}}/D_{\text{test rad.}}$) needed to produce the same effect, becomes constant at low doses, owing to the linearity displayed by both the γ and neutron radiations at low doses. This value was 25.4 ± 4.4 for the high-energy neutrons and 63.7 ± 12.0 for the neutrons of the ^{252}Cf source, with about 40% γ contamination. For a neutron dose of about 5 cGy the RBE values were, respectively, 9.2 and 21.0 for the two neutron energies. At higher doses the RBE values decrease even more, and for a degree of damage equivalent to 190 cGy of ^{60}Co γ -rays are 4.7 ± 0.4 and 11.8 ± 1.8 .

4. DISCUSSION

The dependence of RBE on neutron energy, on dose, and on the biological system is not a special feature of neutrons, since it is also found with other types of radiations. Neutrons are produced in accelerators and reactors with a broad range of energies. Because of the difficulties involved in measuring accurately a neutron spectrum and because of the large spread in RBE values observed between them, high- and low-energy neutrons pose a special problem in radiation protection. Furthermore, our knowledge of the dependence of the biological effects on the biological system at low doses needs to be broadened with information at high energies.

By comparing the results reported here with data obtained from this laboratory using the same high-energy beam and a 10-day growth in bean roots as an assay, a good correlation is found. The RBE value was for a level of damage equivalent to 190 cGy of ^{60}Co γ rays 3.9 ± 0.3 (Bergoglio 1981).

However, when the data obtained for low-energy neutrons are compared with those reported in the literature a discrepancy is found. Hall and Fairchild (1970) quoted, for a 10-day growth in *Vicia faba*, RBE values varying from 6.5 to 5.3 at doses between 16 and 82 cGy of neutrons from a ^{252}Cf source with a similar contribution of γ -rays to the absorbed dose. For micronuclei induction in *Vicia faba* roots after irradiation with fission neutrons (average energy of 3 MeV and about 8% γ contamination), an RBE value of 10.9 was reported when the effect of a dose of 17 cGy of neutrons was compared at the various fixation times with that of a dose of 160 cGy of γ -rays (Evans et al. 1959). In the same experiment, when the mean number of micronuclei observed in the period between 3½ and 24 h after irradiation is considered, both for neutrons and γ -rays a linear dependence on the dose-effect response is obtained, which leads to an RBE value of 10.29 ± 2.04 . Furthermore, RBE values approaching 100 have been obtained for the same fission neutrons from the analysis of the induction of chromosome aberrations in *Tradescantia* microspores (Neary et al. 1963). The values we have found are higher than those reported by Hall and Fairchild (1970) but conform to previous results obtained in our laboratory using a PuBe neutron source which had an average energy of 6 MeV (Baarli et

al. 1976), where an RBE value of 7.5 for a dose of 20 cGy was obtained for 10 day-growth inhibition in bean roots.

The dependence of the RBE values on dose for the induction of micronuclei has been calculated from the fitted curves and is shown in figs. 5 and 6, together with data reported in the literature; the broken part of the line in figure 5 represents the region where extrapolation has to be made owing to the lack of experimental data. A good correspondence is found between the data obtained for the induction of micronuclei in bean roots and that of lens opacification in mice, determined after irradiation in the same high-energy neutron beam. Furthermore, the present data give an RBE dependence on dose which is very close to that reported in the literature for the induction of chromosome aberrations in *Tradescantia* stamen hairs (Sparrow et al. 1972), lens opacification in mice (Bateman et al. 1972), induction of mammary tumours in rats (Vogel 1969), chromosome aberrations in human lymphocytes (Biola et al. 1971), growth reduction in *Vicia faba* (Hall et al. 1973), and skin damage in humans, rats, and pigs (Field 1969). However, some caution has to be taken when making a comparison with data reported in the literature; in fact in some of the curves shown in figure 6, X-rays have been used as the reference radiation instead of γ -rays. Experimental evidence exists that at low doses, between 220 kV X-rays and ^{60}Co γ -rays, an RBE value of about 2 is found both in plant and mammalian cells (Sparrow et al. 1972, Lloyd et al. 1975, Hall and Miller 1981, Barile et al. 1982). For example, the curve which is shown in the figure for the induction of chromosome aberrations in *Tradescantia* is at doses below 10 cGy lower by a factor of about 2.

The absence of recovery or the eventual effect of sensitization observed for the dose of high-energy neutrons, which was fractionated, agrees with previous observations obtained for a 10-day growth in *Vicia faba* using the 600 MeV neutron beam (Bianchi and Diehl-Marshall 1981). A sensitization effect by fractionation with low-energy neutrons has been reported for the induction of mammary tumours in mice (Vogel and Dickson, 1982) and for the survival of the protoplasts of *Nicotiana* (Magnien et al. 1981).

The results obtained in the present experiment seem to indicate that a constant value of RBE is reached at low doses and this is independent of dose rate; however, still a range of RBE values between 20 and 100 can be obtained for various energies at low doses.

Owing to the presence of low-energy neutrons in most neutron radiation fields it might be advisable to consider a value as high as possible in the case of protection.

Acknowledgements

We should like to thank J. Baarli, A.H. Sullivan, J.W.N. Tuyn and C. Raffnsøe for having made the dosimetry measurements in the high-energy neutron beam. We are specially indebted to C.R. Raffnsøe who made the dosimetry of the Cf source. Furthermore, we should like to thank R. Paris, K. Panman and D. Williams for their help with the preparation of the biological samples.

REFERENCES

- Baarli, J., Bakke, O., Sullivan, A.H., and Tuyn, J.W.N., 1978, Proceedings 3rd Symposium on Neutron Dosimetry in Biology and Medicine, Munich, 1977, p. 243. EURATOM, Luxembourg.
- Baarli, J., Bianchi, M., Sullivan, A.H., and Di Paola, M., 1976, Proceedings IAEA Symposium on Biological Effects of Low-Level Radiation Pertinent to Protection of Man and His Environment, Chicago, 1976, p. 195. IAEA, Vienna.
- Barile, G., Bertoncello, G., Capuano, J., Coppola, M., and Quintiliani, M., 1982, ARR Winter Meeting, Book of Abstracts, Univ. Manchester.
- Bateman, J.L., and Snead, M.R., 1970, Symposium on Neutrons in Radiobiology. Oak Ridge, Division of Technical Information Extension.
- Bateman, J.L., Rossi, H.H., Kellerer, A.M., Robinson, C.V., and Pond, V.P., 1972, Radiat. Res., 51, 381.
- Bergoglio, F., 1981, PhD. Thesis, Scuola di Perfezionamento in Fisica Sanitaria e Ospedaliera, University of Milan.
- Bianchi, M., and Diehl-Marshall, I., 1981, Brit. J. Radiol. 54, 532.
- Biola, M.T., Le Go, R., Ducatez, G., Dacher, J., and Bourguignon, M., 1971, Proceedings Symposium on New Developments in Physical and Biological Radiation Detectors, IAEA, Vienna, 1970, p. 633. IAEA, Vienna.
- Broerse, J.J., and Barendsen, G.W., 1973, Curr. Top. Radiat. Res. Quarterly, 8, 305.
- Diehl-Marshall, I., and Bianchi, M., 1981, Brit. J. Radiol., 54, 530.
- Di Paola, M., Coppola, M., Baarli, J., Bianchi, M., and Sullivan, A.H., 1981, Radiat. Res., 84, 453.
- Dunn, E., 1975, Los Alamos Report LA-5845-PR, p. 51.
- Evans, H.J., Neary, G.J., and Williamson, F.S., 1959, Int. J. Radiat. Biol., 3, 216.
- Field, S.B., 1969, Radiology 93, 915.
- Field, S.B., and Hornsey, S., 1979, Adv. Radiat. Biol., 8, 1.
- Fischer, T., Fischer, U., Hagedorn, W., Hammel, G., Huerste, W., Kern, K., Kettle, R., Kleinschmidt, M., Lehmann, L., Moser, K., Roessle, E., and Schmitt, H., 1976, SIN Newsletter 7, 14.

- Hall, E.J., and Fairchild, R.G., 1970, Brit. J. Radiol., 43, 263.
- Hall, E.J., and Miller, R.C., 1981, Radiat. Res., 87, 208.
- Hall, E.J., Rossi, H.H., Kellerer, A.M., Goodman, L.J., and Marino, S., 1973, Radiat. Res., 54, 431.
- Lloyd, D.C., Purrott, R.J., Dolphi, G.W., Bolton, D., Edwards, A.A., and Carp, M.J., 1975, Int. J. Radiat. Biol., 25, 75.
- Magnien, E., Dalschaert, X., and Coppola, M., 1981, Int. J. Radiat. Biol., 40, 463.
- Marshall, I., 1981, PhD Thesis, University of Karlsruhe, Kernforschungszentrum publication, KfK 3265.
- Marshall, I., and Bianchi M., 1982, Int. J. Radiat. Biol., Part I.
- Neary, G.J., Savage, J.R.K., Evans, H.J., and Whittle, J., 1963, Int. J. Radiat. Biol., 6, 127.
- Shellabarger, C.J., Chmelevsky, D., and Kellerer, A.M., 1980, J. Nat. Cancer Inst., 64, 821.
- Sparrow, A.H., Underbrink, A.G., and Rossi, H.H., 1972, Science, 176, 916.
- Vogel, H.H., 1969, Nature, 222, 1279.
- Vogel, H.H., and Dickson, H., 1982. Abstract for the 6th Annual Meeting of the European Society for Radiation Biology, Cracow, p. 166.

Table 1

The experimental data of the high-energy neutron experiments

Dose (cGy)	Mitotic index (MI)	No. of MN/ 100 cells	No. of cells with MN/ 100 cells
0 ¹⁾	9.8 ± 0.5	0.21 ± 0.05	0.21 ± 0.05
0 ²⁾	11.2 ± 0.6	0.15 ± 0.05	0.14 ± 0.04
0.2 ²⁾	10.9 ± 0.4	0.19 ± 0.07	0.17 ± 0.06
0.5 ²⁾	10.9 ± 0.3	0.41 ± 0.11	0.38 ± 0.11
1.0 ¹⁾	10.3 ± 0.4	1.0 ± 0.2	0.9 ± 0.2
5.0 ¹⁾	9.5 ± 0.4	2.0 ± 0.2	1.8 ± 0.2
15.0 ¹⁾	8.3 ± 0.4	7.1 ± 0.5	6.4 ± 0.4
40.0 ¹⁾	6.6 ± 0.4	14.3 ± 0.6	12.6 ± 0.5
40.0 ²⁾	8.3 ± 0.7	13.1 ± 1.0	11.2 ± 0.8
80.0 ¹⁾	6.1 ± 0.5	22.0 ± 0.9	18.2 ± 0.6
40	8.3 ± 0.4	13.1 ± 0.6	11.2 ± 0.5
(20 + 20)	8.0 ± 0.4	15.6 ± 0.6	13.3 ± 0.5
(1 h split)			

The indices 1 and 2 refer to two separate experiments.

Table 2

The experimental data of the ²⁵²Cf-neutron experiment

Dose (cGy)	Mitotic index (MI)	No. of MN/ 100 cells	No. of cells with MN/ 100 cells
0.0	9.7 ± 0.4	0.09 ± 0.05	0.09 ± 0.05
0.2	8.8 ± 0.5	0.37 ± 0.15	0.31 ± 0.09
0.5	9.4 ± 0.5	0.62 ± 0.15	0.57 ± 0.12
1.0	9.7 ± 0.6	1.14 ± 0.18	1.08 ± 0.17
5.0	8.9 ± 0.5	5.7 ± 0.4	5.2 ± 0.3

Figure captions

- Fig. 1 : The mean number of micronuclei between 24 and 48 h after radiation plotted against neutron dose (uncorrected data).
- Fig. 2 : The number of micronuclei as a function of neutron dose, after correction for the mitotic index is applied.
- Fig. 3 : Distribution of the frequency of micronuclei per bean.
- Fig. 4 : The number of micronuclei as a function of ^{252}Cf -neutron dose, after correction for the mitotic index is applied.
- Fig. 5 : A comparison of the curves derived from the RBE values found for micronuclei induction is made with the values found for lens opacification in mice.
- Fig. 6 : The RBE curves found for micronuclei induction (dashed lines) compared with the RBE curves found for different end points. 1) Opacification of lens in mice (430 keV neutrons). 2) Opacification of the lens in mice (1.8 MeV neutrons). 3) Mutations of *Tradescantia* stamen hairs (430 keV neutrons). 4) Mammary neoplasm in the Sprague-Dawley rat (fission neutrons). 5) Chromosome aberrations in human lymphocytes (fission neutrons). 6) Growth reduction of *Vicia faba* roots (6.0 MeV neutrons). 7) Skin damage (human, rat, mouse, pig) (6.0 MeV neutrons).

Fig. 1

Fig. 2

Fig. 3

Fig. 4

Fig. 5

Fig. 6