CERN Data Handling Division

DD/82/10

2 July 1982

K.S. Kölbig

ON THE INTEGRAL $\int_0^\infty e^{-ut} t^{\nu-1} \log^m t dt$

ABSTRACT

A recurrence relation is given for the integral in the title. Formulae which allow easy evaluation by formula manipulation are given for integer and half-integer values of ν . Explicit expressions for these integrals for small values of m and $\nu = n$, $\nu = n + \frac{1}{2}$, are also presented.

Geneva
1982

1. INTRODUCTION

The purpose of this note is to give expressions for the Laplace (or Mellin) integral

$$R_m(\mu, \nu) = \int_0^{\infty} e^{-\mu t} t^{\nu-1} \log^m t \, dt \quad (\operatorname{Re} \mu > 0, \operatorname{Re} \nu > 0), \quad (1)$$

in particular for integer and half-integer values of ν . This integral, with integer values of ν , occurs when establishing an asymptotic expansion for the Landau density function [2,6], as $x \rightarrow \infty$,

$$\phi(x) = \frac{1}{\pi} \int_0^{\infty} e^{-xt} t^{-t} \sin \pi t \, dt. \quad (2)$$

For $m \leq 3$, $R_m(\mu, \nu)$ can be found, for example, in [3, No. 4.3521, 4.3582, 4.3583]:

$$R_1(\mu, \nu) = \frac{\Gamma(\nu)}{\mu^\nu} [\psi(\nu) - \log \mu] \quad (3)$$

$$R_2(\mu, \nu) = \frac{\Gamma(\nu)}{\mu^\nu} \left\{ [\psi(\nu) - \log \mu]^2 + \zeta(2, \nu) \right\} \quad (4)$$

$$R_3(\mu, \nu) = \frac{\Gamma(\nu)}{\mu^\nu} \left\{ [\psi(\nu) - \log \mu]^3 + 3[\psi(\nu) - \log \mu] \zeta(2, \nu) - 2\zeta(3, \nu) \right\}^* \quad (5)$$

where

$$\psi(x) = \frac{d}{dx} \log \Gamma(x) = \frac{\Gamma'(x)}{\Gamma(x)}, \quad (6)$$

and where

$$\zeta(r, q) = \sum_{j=0}^{\infty} \frac{1}{(q+j)^r} \quad (7)$$

is the generalized zeta function, related to $\psi(x)$ by [3, No. 8.3638]

$$\psi^{(k)}(x) = (-1)^{k+1} k! \zeta(k+1, x). \quad (8)$$

*) Note that the corresponding formula in [3], copied from McLachlan et al. [8], is incorrect, as has been pointed out in [7]. In addition, $\zeta(2, \nu)$ and $\zeta(3, \nu)$ must be replaced by $\zeta(2, \nu+1)$ and $\zeta(3, \nu+1)$ respectively in the formulae corresponding to $m=2$ and $m=3$ in [8].

2. A RECURRENCE FORMULA FOR $R_m(\mu, \nu)$

In order to give a recurrence formula for $R_m(\mu, \nu)$, we need the following

Lemma:

Let $f(x)$ be m times differentiable. Then

$$\left(\frac{d}{dx}\right)^m e^{f(x)} = e^{f(x)} F_m(x) \quad (9)$$

where $F_0(x) = 1$, $F_1(x) = f'(x)$, and

$$F_k(x) = F'_{k-1}(x) + F_1(x)F_{k-1}(x), \quad (k = 2, 3, \dots, m) \quad (10)$$

The proof by induction is easy.

We apply this lemma to $R_m(\mu, \nu)$ and obtain

$$\begin{aligned} R_m(\mu, \nu) &= \int_0^\infty e^{-\mu t} t^{\nu-1} \log^m t \, dt \\ &= \left(\frac{d}{d\nu}\right)^m \int_0^\infty e^{-\mu t} t^{\nu-1} \, dt \\ &= \left(\frac{d}{d\nu}\right)^m \frac{\Gamma(\nu)}{\mu^\nu} = \frac{\Gamma(\nu)}{\mu^\nu} G_m(\mu, \nu), \end{aligned} \quad (11)$$

where $G_0(\mu, \nu) = 1$, $G_1(\mu, \nu) = \psi(\nu) - \log \mu$, and

$$G_k(\mu, \nu) = \frac{d}{d\nu} G_{k-1}(\mu, \nu) + G_1(\mu, \nu) G_{k-1}(\mu, \nu). \quad (12)$$

With a formula manipulation system, such as REDUCE [4], it is easy to compute $R_m(\mu, \nu)$ from (8) and (12) in terms of $\psi(\nu) - \log \mu$ and $\zeta(k, \nu)$ ($2 \leq k \leq m$).

The resulting expressions for $R_m(\mu, \nu)$ for $m = 1(1)9$ are given in Table 1.

3. A FORMULA FOR $R_m(\mu, n+1)$, n INTEGER

Expressions for $R_m(\mu, n+1)$, where n is integer, can be obtained from (12) by setting $\nu = n+1$ in $G_m(\mu, \nu)$ and using

$$\zeta(k, n+1) = \zeta(k) - \sigma(n, k) \quad (13)$$

$$\psi(n+1) = -\gamma + \sigma(n, 1), \quad (14)$$

where

$$\sigma(n, k) = \sum_{j=1}^n \frac{1}{j^k}, \quad (15)$$

γ is Euler's constant and $\zeta(k)$ is the Riemann zeta function.

In order to derive another formula for this case, we set $\tilde{R}_m(\mu, n) = R_m(\mu, n+1)$ and write

$$\begin{aligned} \tilde{R}_m(\mu, n) &= \int_0^\infty e^{-\mu t} t^n \log^m t \, dt \quad (n \geq 0) \\ &= \left(\frac{d}{d\alpha} \right)^m \int_0^\infty e^{-\mu t} t^{n+\alpha} \, dt \Big|_{\alpha=0} \\ &= \frac{1}{\mu^{n+1}} \left(\frac{d}{d\alpha} \right)^m \left[\frac{1}{\mu^\alpha} \Gamma(1+n+\alpha) \right]_{\alpha=0} \\ &= \frac{1}{\mu^{n+1}} \left(\frac{d}{d\alpha} \right)^m \left[\frac{1}{\alpha} (\alpha)_{n+1} \frac{1}{\mu^\alpha} \Gamma(1+\alpha) \right]_{\alpha=0}, \end{aligned} \quad (16)$$

where [1, No. 24.1.3]

$$(\alpha)_k = \alpha(\alpha+1) \dots (\alpha+k-1) = \sum_{j=1}^k (-1)^{j+k} S_k^{(j)} \alpha^j \quad (17)$$

is the Pochhammer symbol. The $S_k^{(j)}$ are the Stirling numbers of the 1st kind, defined by

$$\begin{aligned} S_{k+1}^{(j)} &= S_k^{(j-1)} - k S_k^{(j)} \quad (k \geq j \geq 1) \\ S_k^{(1)} &= (-1)^{k-1} (k-1)! \\ S_k^{(k)} &= 1, \quad S_k^{(0)} = \delta_{0k} \\ S_k^{(j)} &= 0 \quad \text{for } k < j. \end{aligned} \quad (18)$$

Applying the Leibniz formula, we have

$$R_m(\mu, n) = \frac{1}{\mu^{n+1}} \sum_{\rho=0}^m \binom{m}{\rho} \left[\frac{1}{\alpha} (\alpha)_{n+1} \right]^{(\rho)} \left[\frac{1}{\mu^\alpha} \Gamma(1+\alpha) \right]^{(m-\rho)} \Big|_{\alpha=0}. \quad (19)$$

We see from (17) that

$$\left(\frac{d}{d\alpha} \right)^\rho \frac{1}{\alpha} (\alpha)_{n+1} \Big|_{\alpha=0} = \begin{cases} (-1)^{\rho+n} \rho! S_{n+1}^{(\rho+1)} & (\rho \leq n) \\ 0 & (\rho > n) \end{cases} \quad (20)$$

We note that [3, No. 8.3421]

$$\log \mu^{-\alpha} \Gamma(1 + \alpha) = -(\gamma + \log \mu) \alpha + \sum_{k=2}^{\infty} (-1)^k \zeta(k) \alpha^k / k \quad (21)$$

(|α| < 1) ,

and therefore, by exponentiation (see, for example, Henrici [5, §1.6]),

$$\mu^{-\alpha} \Gamma(1 + \alpha) = \sum_{k=0}^{\infty} b_k \alpha^k \quad (|x| < 1) \quad (22)$$

where $b_0 = 1$,

$$b_k = \frac{1}{k} \sum_{\kappa=1}^k (-1)^\kappa \tilde{\zeta}(\kappa) b_{k-\kappa} \quad (23)$$

and $\tilde{\zeta}(1) = \gamma + \log \mu$, $\tilde{\zeta}(\kappa) = \zeta(\kappa)$ for $\kappa > 1$.

Substituting (20) and (22) into (19), we finally have

$$\tilde{R}_m(\mu, n) = \int_0^\infty e^{-\mu t} t^n \log^m t \, dt = \frac{(-1)^n n!}{\mu^{n+1}} \sum_{\rho=0}^{\min(m, n)} (-1)^\rho S_{n+1}^{(\rho+1)} b_{m-\rho} \quad (24)$$

(n ≥ 0) .

We consider a few special cases. For $n = 0$ we obtain

$$\tilde{R}_m(\mu, 0) = \frac{m!}{\mu} b_m . \quad (25)$$

For $m = 1$ we have

$$\begin{aligned} \tilde{R}_1(\mu, n) &= \frac{(-1)^n}{\mu^{n+1}} \left(S_{n+1}^{(1)} b_1 - S_{n+1}^{(2)} \right) \\ &= \frac{n!}{\mu^{n+1}} [\sigma(n, 1) - \gamma - \log \mu] \end{aligned} \quad (26)$$

in accordance with (3), since from (18),

$$S_{n+1}^{(1)} = (-1)^n n! , \quad S_{n+1}^{(2)} = (-1)^{n+1} n! \sigma(n, 1) . \quad (27)$$

For $m = 2$ it follows from (24) and (4) that

$$\begin{aligned}\tilde{R}_2(\mu, n) &= 2 \frac{(-1)^n}{\mu^{n+1}} \left(S_{n+1}^{(1)} b_2 - S_{n+1}^{(2)} b_1 + S_{n+1}^{(3)} \right) \\ &= \frac{n!}{\mu^{n+1}} \left\{ (\gamma + \log \mu)^2 - 2(\gamma + \log \mu) \sigma(n, 1) + \frac{\pi^2}{6} + 2 \frac{(-1)^n}{n!} S_{n+1}^{(3)} \right\} \quad (28) \\ &= \frac{n!}{\mu^{n+1}} \left\{ (\gamma + \log \mu)^2 - 2(\gamma + \log \mu) \sigma(n, 1) + \sigma^2(n, 1) - \sigma(n, 2) + \frac{\pi^2}{6} \right\},\end{aligned}$$

where we have used $\zeta(2) = \pi^2/6$. Note that $S_{n+1}^{(2)} = 0$ for $n = 0$ and $S_{n+1}^{(3)} = 0$ for $n = 0, 1$. From (28), we obtain the relation

$$S_{n+1}^{(3)} = \frac{1}{2} (-1)^n n! [\sigma^2(n, 1) - \sigma(n, 2)], \quad (29)$$

which is not immediately obvious from (18). In general, by comparing the result obtained from the recurrence relation (12) with that given by (24), we find that

$$S_{n+1}^{(k+1)} = (-1)^{n+k} \frac{n!}{k!} \omega[\sigma(n, \kappa)] \quad (30)$$

with $\kappa = 1, 2, \dots, k$. The function

$$\omega[\sigma(n, \kappa)] = \sum_{j=1}^k \sum_{\{j_i\}} c_{\{j_i\}} \sigma(n, j_1) \dots \sigma(n, j_k) \quad (31)$$

where the inner sum runs over all ordered sets of integers $\{j_i\}$ with

$$j_i \geq 1, \quad \sum_{i=1}^k j_i = k,$$

is a "homogeneous" polynomial in $\sigma(n, \kappa)$, with integer coefficients. We give two more examples:

$$S_{n+1}^{(4)} = \frac{1}{6} (-1)^{n+1} n! [\sigma^3(n, 1) - 3\sigma(n, 1)\sigma(n, 2) + 2\sigma(n, 3)] \quad (32)$$

$$S_{n+1}^{(5)} = \frac{1}{24} (-1)^n n! [\sigma^4(n, 1) + 3\sigma^2(n, 2) - 6\sigma^2(n, 1)\sigma(n, 2) + 8\sigma(n, 1)\sigma(n, 3) - 6\sigma(n, 4)].$$

Note that $\omega[\sigma(n, \kappa)] \equiv 0$ for $0 \leq n < k$. This fact is not apparent when one derives $R_m(\mu, \nu)$ by means of the recurrence (12), replacing ν by $n + 1$ in $R_m(\mu, \nu)$.

In general, formula (24) can easily be evaluated by formula-manipulation. Expressions for $\tilde{R}_m(\mu, n)$, $m = 0(1)5$, $n = 0(1)5$, are given in Table 2. For special values of m and n (e.g. $n > 0$, $m = 1$, or $n = 0$, $m \leq 3$), these expressions can also be found in the relevant handbooks or tables of integral transforms.

4. A FORMULA FOR $R_m(\mu, n + \frac{1}{2})$, n INTEGER

Setting $\nu = n + \frac{1}{2}$ and [1, No. 6.3.4, 23.2.20]

$$\zeta(k, n + \frac{1}{2}) = (2^k - 1)\zeta(k) - 2^k \hat{\sigma}(n, k) \quad (33)$$

$$\psi(n + \frac{1}{2}) = -\gamma - \log 4 + 2\hat{\sigma}(n, 1) , \quad (34)$$

where

$$\hat{\sigma}(n, k) = \sum_{j=1}^n \frac{1}{(2j-1)^k} , \quad (35)$$

in the expression for $R_m(\mu, \nu)$ obtained from (12) leads to expressions for $R_m^*(\mu, n) = R_m(\mu, n + \frac{1}{2})$. In the same way as in Section 3, we may also derive another formula in this case. We start with

$$\begin{aligned} R_m^*(\mu, n) &= \int_0^\infty e^{-\mu t} t^{n-1/2} \log^m t \, dt \quad (n \geq 0) \\ &= \left(\frac{d}{d\alpha} \right)^m \int_0^\infty e^{-\mu t} t^{n-1/2+\alpha} \, dt \Big|_{\alpha=0} \\ &= \frac{1}{\mu^{n+1/2}} \left(\frac{d}{d\alpha} \right)^m \left[\frac{1}{\mu^\alpha} \Gamma(n + \frac{1}{2} + \alpha) \right]_{\alpha=0} . \end{aligned} \quad (36)$$

Using

$$\begin{aligned} \Gamma\left(\frac{1}{2} + n + \alpha\right) &= \left(\alpha + \frac{1}{2}\right)_n \Gamma\left(\frac{1}{2} + \alpha\right) \\ &= \sqrt{\pi} \left(\alpha + \frac{1}{2}\right)_n 2^{-2\alpha} \frac{\Gamma(1 + 2\alpha)}{\Gamma(1 + \alpha)} , \end{aligned} \quad (37)$$

we obtain

$$R_m^*(\mu, \nu) = \frac{\sqrt{\pi}}{\mu^{n+1/2}} \left(\frac{d}{d\alpha} \right)^m \left[\left(\alpha + \frac{1}{2}\right)_n H(\alpha, \mu) \right]_{\alpha=0} \quad (38)$$

where

$$H(\alpha, \mu) = \frac{1}{(4\mu)^\alpha} \frac{\Gamma(1 + 2\alpha)}{\Gamma(1 + \alpha)} . \quad (39)$$

As for the series (22), we can write

$$H(\alpha, \mu) = \sum_{k=0}^{\infty} b_k^* \alpha^k \quad \left(|\alpha| < \frac{1}{2} \right) . \quad (40)$$

where $b_0^* = 1$

$$b_k^* = \frac{1}{k} \sum_{\kappa=1}^k (-1)^\kappa \zeta^*(\kappa) b_{k-\kappa}^* \quad (41)$$

and

$$\zeta^*(\kappa) = \begin{cases} \gamma + \log 4\mu & (\kappa = 1) \\ (2^\kappa - 1)\zeta(\kappa) & (\kappa > 1) \end{cases} \quad (42)$$

Applying the Leibniz formula to (38), we use (17) and write

$$\begin{aligned} \left(\frac{d}{d\alpha}\right)^\rho \left(\alpha + \frac{1}{2}\right)_n \Big|_{\alpha=0} &= \left(\frac{d}{d\alpha}\right)^\rho \sum_{k=1}^n (-1)^{k+n} S_n^{(k)} \left(\alpha + \frac{1}{2}\right)^k \\ &= \begin{cases} \rho! \sum_{k=\rho}^n (-1)^{k+n} S_n^{(k)} \binom{k}{\rho} 2^{\rho-k} & (\rho \leq n) \\ 0 & (\rho > n) \end{cases} \end{aligned} \quad (43)$$

and therefore finally

$$\begin{aligned} R_m^*(\mu, n) &= \int_0^\infty e^{-\mu t} t^{n-1/2} \log^m t \, dt \quad (n \geq 0) \\ &= \sqrt{\frac{\pi}{\mu}} \frac{(-1)^n n!}{\mu^n} \sum_{\rho=0}^{\min(m, n)} b_{m-\rho}^* \sum_{k=\rho}^n (-1)^k S_n^{(k)} \binom{k}{\rho} 2^{\rho-k} \end{aligned} \quad (44)$$

We consider two special cases. For $n = 0$, we have

$$R_m^*(\mu, 0) = \sqrt{\frac{\pi}{\mu}} m! b_m^* \quad (45)$$

For $m = 1$ we obtain from (3) and (44) the expression [3, No. 4.3523]

$$\begin{aligned} R_1^*(\mu, n) &= \frac{\Gamma\left(n + \frac{1}{2}\right)}{\mu^{n+1/2}} [-\gamma - \log 4\mu + 2\hat{\sigma}(n, 1)] \\ &= \sqrt{\frac{\pi}{\mu}} \frac{(2n-1)!!}{(2\mu)^n} [-\gamma - \log 4\mu + 2\hat{\sigma}(n, 1)] \\ &= \sqrt{\frac{\pi}{\mu}} \frac{1}{\mu^n} \left[-\left(\frac{1}{2}\right)_n (\gamma + \log 4\mu) + 2(-1)^n \sum_{k=1}^n (-1)^k S_n^{(k)} k 2^{-k} \right] \end{aligned} \quad (46)$$

Thus, by comparison,

$$\sum_{k=1}^n (-1)^k S_n^{(k)} k 2^{-k} = (-1)^n \frac{(2n-1)!!}{2^n} \hat{\sigma}(n, 1) \quad (47)$$

Similarly, for $m = 2$, one finds

$$\sum_{k=2}^n (-1)^k S_n^{(k)} k(k-1) 2^{-k} = (-1)^n \frac{(2n-1)!!}{2^n} [\hat{\sigma}^2(n,1) - \hat{\sigma}(n,2)] . \quad (48)$$

In general, as in the case of $\tilde{R}_m(\mu, n)$, a comparison of the result obtained from (12) with expression (44) shows that

$$\sum_{k=\rho}^n (-1)^k S_n^{(k)} \frac{k!}{(k-\rho)!} 2^{-k} = (-1)^n \frac{(2n-1)!!}{2^n} \omega[\hat{\sigma}(n, \rho')] , \quad (49)$$

where ω is defined by (31), and $\omega[\hat{\sigma}(n, \rho')] \equiv 0$ for $0 \leq n < \rho$.

Formula (44) can be evaluated easily by formula manipulation. Expressions for $R_m^*(\mu, n)$, $m = 0(1)5$, $n = 0(1)5$, are given in Table 3. $R_1^*(\mu, n)$ can also be found in [3, No. 4.3523].

5. TWO RELATED INTEGRALS

Substituting $t = T^2$ in $\tilde{R}_m(\mu, n)$, we find that

$$\int_0^\infty e^{-\mu T^2} T^{2n+1} \log^m T \, dT = \frac{1}{2^{m+1}} \tilde{R}_m(\mu, n) , \quad (50)$$

and, by the same substitution in $R_m^*(\mu, n)$,

$$\int_0^\infty e^{-\mu T^2} T^{2n} \log^m T \, dT = \frac{1}{2^{m+1}} R_m^*(\mu, n) . \quad (51)$$

Three particular combinations of these integrals for $m = 1$ are given in [3, No. 4.3551, 4.3553, 4.3554].

REFERENCES

- [1] M. Abramowitz and I.A. Stegun (Editors), Handbook of Mathematical Functions, with Formulas, Graphs, and Mathematical Tables, 5th printing with corrections, Nat. Bur. Standards Appl. Math. Series 55, US Government Printing Office, Washington DC, 1966.
- [2] W. Börsch-Supan, On the evaluation of the function $\phi(\lambda) = 1/2\pi i \int_{\sigma-i\infty}^{\sigma+i\infty} e^{u \ln u + \lambda u} du$ for real values of λ , J. Res. Nat. Bur. Standards, v. 65B, 1961, pp. 245-250.
- [3] I.S. Gradshteyn and I.M. Ryzhik, Table of Integrals, Series, and Products, Academic Press (New York, London, Toronto, Sydney, San Francisco), 1980.
- [4] A.C. Hearn, REDUCE Users Manual, 2nd Edition, Univ. of Utah, Salt Lake City, Utah, USA, 1973.
- [5] P. Henrici, Applied and computational complex analysis, v. 1, John Wiley & Sons (New York, London, Sydney, Toronto), 1974.
- [6] L. Landau, On the energy loss of fast particles by ionization, J. Physics, v. 8, 1944, pp. 201-205.
- [7] Y.L. , Review of [3], Math. Comp., v. 36, 1981, pp. 310-314.
- [8] N.W. McLachlan, P. Humbert and L. Poli, Supplément au Formulaire pour le Calcul symbolique, Mémor. Sci. Math., Fasc. 113, Gauthier-Villars, Paris, 1950.

Table 1

Let

$$Y_m = \frac{\mu^\nu}{\Gamma(\nu)} \int_0^\infty e^{-\mu t} t^{\nu-1} \log^m t \, dt = \frac{\mu^\nu}{\Gamma(\nu)} R_m(\mu, \nu), \quad (\operatorname{Re} \mu > 0, \operatorname{Re} \nu > 0)$$

$$\phi = \psi(\nu) - \log \mu,$$

and

$$\beta_2 = \zeta(2, \nu)$$

$$\beta_3 = \zeta(3, \nu)$$

$$\beta_4 = \zeta^2(2, \nu) + 2\zeta(4, \nu)$$

$$\beta_5 = 5\zeta(2, \nu)\zeta(3, \nu) + 6\zeta(5, \nu)$$

$$\beta_6 = 3\zeta^3(2, \nu) + 18\zeta(2, \nu)\zeta(4, \nu) + 8\zeta^2(3, \nu) + 24\zeta(6, \nu)$$

$$\beta_7 = 35\zeta^2(2, \nu)\zeta(3, \nu) + 84\zeta(2, \nu)\zeta(5, \nu) + 70\zeta(3, \nu)\zeta(4, \nu) + 120\zeta(7, \nu)$$

$$\begin{aligned} \beta_8 = & 15\zeta^4(2, \nu) + 180\zeta^2(2, \nu)\zeta(4, \nu) + 160\zeta(2, \nu)\zeta^2(3, \nu) + 480\zeta(2, \nu)\zeta(6, \nu) \\ & + 384\zeta(3, \nu)\zeta(5, \nu) + 180\zeta^2(4, \nu) + 720\zeta(8, \nu) \end{aligned}$$

$$\begin{aligned} \beta_9 = & 315\zeta^3(2, \nu)\zeta(3, \nu) + 1134\zeta^2(2, \nu)\zeta(5, \nu) + 1890\zeta(2, \nu)\zeta(3, \nu)\zeta(4, \nu) \\ & + 3240\zeta(2, \nu)\zeta(7, \nu) + 280\zeta^3(3, \nu) + 2520\zeta(3, \nu)\zeta(6, \nu) \\ & + 2268\zeta(4, \nu)\zeta(5, \nu) + 5040\zeta(9, \nu). \end{aligned}$$

Then

$$Y_0 = 1$$

$$Y_1 = \phi$$

$$Y_2 = \phi^2 + \beta_2$$

$$Y_3 = \phi^3 + 3\beta_2\phi - 2\beta_3$$

$$Y_4 = \phi^4 + 6\beta_2\phi^2 - 8\beta_3\phi + 3\beta_4$$

$$Y_5 = \phi^5 + 10\beta_2\phi^3 - 20\beta_3\phi^2 + 15\beta_4\phi - 4\beta_5$$

$$Y_6 = \phi^6 + 15\beta_2\phi^4 - 40\beta_3\phi^3 + 45\beta_4\phi^2 - 24\beta_5\phi + 5\beta_6$$

$$Y_7 = \phi^7 + 21\beta_2\phi^5 - 70\beta_3\phi^4 + 105\beta_4\phi^3 - 84\beta_5\phi^2 + 35\beta_6\phi - 6\beta_7$$

$$Y_8 = \phi^8 + 28\beta_2\phi^6 - 112\beta_3\phi^5 + 210\beta_4\phi^4 - 224\beta_5\phi^3 + 140\beta_6\phi^2 - 48\beta_7\phi + 7\beta_8$$

$$Y_9 = \phi^9 + 36\beta_2\phi^7 - 168\beta_3\phi^6 + 378\beta_4\phi^5 - 504\beta_5\phi^4 + 420\beta_6\phi^3 - 216\beta_7\phi^2 \\ + 63\beta_8\phi - 8\beta_9 .$$

Thus

$$Y_m = \phi^m + \sum_{j=2}^m c_{m,j} \beta_j \phi^{m-j} , \quad (m = 2, \dots, 9)$$

$$c_{m,j} = (-1)^j (j-1) \sum_{k=j}^m \binom{k-1}{j-1} .$$

Table 2

Let

$$I_{mn} = \mu^{n+1} \int_0^{\infty} e^{-\mu t} t^n \log^m t \, dt = \mu^{n+1} \tilde{R}_m(\mu, n) ; \quad C = \gamma + \log \mu .$$

(Re $\mu > 0$)

Then

$$I_{0n} = n! \quad (n \geq 0)$$

$$I_{10} = -C$$

$$I_{1n} = n! \left[\sum_{j=1}^n j^{-1} - C \right] \quad (n > 0)$$

$$I_{20} = C^2 + \frac{\pi^2}{6}$$

$$I_{21} = C^2 - 2C + \frac{\pi^2}{6}$$

$$I_{22} = 2C^2 - 6C + \frac{\pi^2}{3} + 2$$

$$I_{23} = 6C^2 - 22C + \pi^2 + 12$$

$$I_{24} = 24C^2 - 100C + 4\pi^2 + 70$$

$$I_{25} = 120C^2 - 548C + 20\pi^2 + 450$$

$$I_{30} = -C^3 - \frac{\pi^2}{2} C - 2\zeta(3)$$

$$I_{31} = -C^3 + 3C^2 - \frac{\pi^2}{2} C - 2\zeta(3) + \frac{\pi^2}{2}$$

$$I_{32} = -2C^3 + 9C^2 - (\pi^2 + 6)C - 4\zeta(3) + \frac{3}{2} \pi^2$$

$$I_{33} = -6C^3 + 33C^2 - 3(\pi^2 + 12)C - 12\zeta(3) + \frac{11}{2} \pi^2 + 6$$

$$I_{34} = -24C^3 + 150C^2 - 6(2\pi^2 + 35)C - 48\zeta(3) + 25\pi^2 + 60$$

$$I_{35} = -120C^3 + 822C^2 - 30(2\pi^2 + 45)C - 240\zeta(3) + 137\pi^2 + 510$$

$$I_{40} = C^4 + C^2\pi^2 + 8\zeta(3)C + \frac{3}{20} \pi^4$$

$$I_{41} = C^4 - 4C^3 + \pi^2 C^2 + 2(4\zeta(3) - \pi^2)C - 8\zeta(3) + \frac{3}{20} \pi^4$$

$$I_{42} = 2C^4 - 12C^3 + 2(\pi^2 + 6)C^2 + 2(8\zeta(3) - 3\pi^2)C - 24\zeta(3) + \frac{3}{10} \pi^4 + 2\pi^2$$

$$I_{43} = 6C^4 - 44C^3 + 6(\pi^2 + 12)C^2 + 2(24\zeta(3) - 11\pi^2 - 12)C - 88\zeta(3) + \frac{9}{10} \pi^4 + 12\pi^2$$

$$I_{44} = 24C^4 - 200C^3 + 12(2\pi^2 + 35)C^2 + 4(48\zeta(3) - 25\pi^2 - 60)C - 400\zeta(3) \\ + \frac{18}{5}\pi^4 + 70\pi^2 + 24$$

$$I_{45} = 120C^4 - 1096C^3 + 60(2\pi^2 + 45)C^2 + 4(240\zeta(3) - 137\pi^2 - 510)C \\ - 2192\zeta(3) + 18\pi^4 + 450\pi^2 + 360$$

$$I_{50} = -C^5 - \frac{5}{3}\pi^2C^3 - 20\zeta(3)C^2 - \frac{3}{4}\pi^4C - \frac{10}{3}\zeta(3)\pi^2 - 24\zeta(5)$$

$$I_{51} = -C^5 + 5C^4 - \frac{5}{3}\pi^2C^3 - 5(4\zeta(3) - \pi^2)C^2 + \left(40\zeta(3) - \frac{3}{4}\pi^4\right)C - \frac{10}{3}\zeta(3)\pi^2 \\ - 24\zeta(5) + \frac{3}{4}\pi^4$$

$$I_{52} = -2C^5 + 15C^4 - 10\left(\frac{1}{3}\pi^2 + 2\right)C^3 - 5(8\zeta(3) - 3\pi^2)C^2 + \left(120\zeta(3) - \frac{3}{2}\pi^4 - 10\pi^2\right)C \\ - \frac{20}{3}\zeta(3)\pi^2 - 40\zeta(3) - 48\zeta(5) + \frac{9}{4}\pi^4$$

$$I_{53} = -6C^5 + 55C^4 - 10(\pi^2 + 12)C^3 - 5(24\zeta(3) - 11\pi^2 - 12)C^2 \\ + \left(440\zeta(3) - \frac{9}{2}\pi^4 - 60\pi^2\right)C - 20\zeta(3)\pi^2 - 240\zeta(3) - 144\zeta(5) \\ + \frac{33}{4}\pi^4 + 10\pi^2$$

$$I_{54} = -24C^5 + 250C^4 - 20(2\pi^2 + 35)C^3 - 10(48\zeta(3) - 25\pi^2 - 60)C^2 \\ + 2(1000\zeta(3) - 9\pi^4 - 175\pi^2 - 60)C - 80\zeta(3)\pi^2 - 1400\zeta(3) - 576\zeta(5) \\ + \frac{75}{2}\pi^4 + 100\pi^2$$

$$I_{55} = -120C^5 + 1370C^4 - 100(2\pi^2 + 45)C^3 - 10(240\zeta(3) - 137\pi^2 - 510)C^2 \\ + 10(1096\zeta(3) - 9\pi^4 - 225\pi^2 - 180)C - 400\zeta(3)\pi^2 - 9000\zeta(3) - 2880\zeta(5) \\ + \frac{411}{2}\pi^4 + 850\pi^2 + 120$$

Table 3

Let

$$J_{mn} = \sqrt{\frac{\mu}{\pi}} \mu^n \int_0^{\infty} e^{-\mu t} t^{n-1/2} \log^m t \, dt = \sqrt{\frac{\mu}{\pi}} \mu^n R_m^*(\mu, n) \quad (\operatorname{Re} \mu > 0)$$

and

$$C = \gamma + \log 4\mu.$$

Then

$$J_{00} = 1$$

$$J_{0n} = \frac{(2n-1)!!}{2^n} \quad (n > 0)$$

$$J_{10} = -C$$

$$J_{1n} = \frac{(2n-1)!!}{2^n} \left(-C + 2 \sum_{j=1}^n \frac{1}{2j-1} \right) \quad (n > 0)$$

$$J_{20} = C^2 + \frac{\pi^2}{2}$$

$$J_{21} = \frac{1}{2} C^2 - 2C + \frac{\pi^2}{4}$$

$$J_{22} = \frac{3}{4} C^2 - 4C + \frac{3}{8} \pi^2 + 2$$

$$J_{23} = \frac{15}{8} C^2 - \frac{23}{2} C + \frac{15}{16} \pi^2 + 9$$

$$J_{24} = \frac{105}{16} C^2 - 44C + \frac{105}{32} \pi^2 + 43$$

$$J_{25} = \frac{945}{32} C^2 - \frac{1689}{8} C + \frac{945}{64} \pi^2 + \frac{475}{2}$$

$$J_{30} = -C^3 - \frac{3}{2} \pi^2 C - 14\zeta(3)$$

$$J_{31} = -\frac{1}{2} C^3 + 3C^2 - \frac{3}{4} \pi^2 C - 7\zeta(3) + \frac{3}{2} \pi^2$$

$$J_{32} = -\frac{3}{4} C^3 + 6C^2 - 3\left(\frac{3}{8} \pi^2 + 2\right)C - \frac{21}{2} \zeta(3) + 3\pi^2$$

$$J_{33} = -\frac{15}{8} C^3 + \frac{69}{4} C^2 - 9\left(\frac{5}{16} \pi^2 + 3\right)C - \frac{105}{4} \zeta(3) + \frac{69}{8} \pi^2 + 6$$

$$J_{34} = -\frac{105}{16} C^3 + 66C^2 - 3\left(\frac{105}{32} \pi^2 + 43\right)C - \frac{735}{8} \zeta(3) + 33\pi^2 + 48$$

$$J_{35} = -\frac{945}{32} C^3 + \frac{5067}{16} C^2 - \frac{15}{2}\left(\frac{189}{32} \pi^2 + 95\right)C - \frac{6615}{16} \zeta(3) + \frac{5067}{32} \pi^2 + 345$$

$$J_{40} = C^4 + 3\pi^2 C^2 + 56\zeta(3)C + \frac{7}{4}\pi^4$$

$$J_{41} = \frac{1}{2}C^4 - 4C^3 + \frac{3}{2}\pi^2 C^2 + 2(14\zeta(3) - 3\pi^2)C - 56\zeta(3) + \frac{7}{8}\pi^4$$

$$J_{42} = \frac{3}{4}C^4 - 8C^3 + 3\left(\frac{3}{4}\pi^2 + 4\right)C^2 + 6(7\zeta(3) - 2\pi^2)C - 112\zeta(3) + \frac{21}{16}\pi^4 + 6\pi^2$$

$$J_{43} = \frac{15}{8}C^4 - 23C^3 + 9\left(\frac{5}{8}\pi^2 + 6\right)C^2 + 3\left(35\zeta(3) - \frac{23}{2}\pi^2 - 8\right)C - 322\zeta(3) + \frac{105}{32}\pi^4 + 27\pi^2$$

$$J_{44} = \frac{105}{16}C^4 - 88C^3 + 3\left(\frac{105}{16}\pi^2 + 86\right)C^2 + 3\left(\frac{245}{2}\zeta(3) - 44\pi^2 - 64\right)C - 1232\zeta(3) + \frac{735}{64}\pi^4 + 129\pi^2 + 24$$

$$J_{45} = \frac{945}{32}C^4 - \frac{1689}{4}C^3 + 15\left(\frac{189}{32}\pi^2 + 95\right)C^2 + 3\left(\frac{2205}{4}\zeta(3) - \frac{1689}{8}\pi^2 - 460\right)C - \frac{11823}{2}\zeta(3) + \frac{6615}{128}\pi^4 + \frac{1425}{2}\pi^2 + 300$$

$$J_{50} = -C^5 - 5\pi^2 C^3 - 140\zeta(3)C^2 - \frac{35}{4}\pi^4 C - 70\zeta(3)\pi^2 - 744\zeta(5)$$

$$J_{51} = -\frac{1}{2}C^5 + 5C^4 - \frac{5}{2}\pi^2 C^3 + 5(3\pi^2 - 14\zeta(3))C^2 + 35\left(8\zeta(3) - \frac{\pi^4}{8}\right)C - 35\zeta(3)\pi^2 - 372\zeta(5) + \frac{35}{4}\pi^4$$

$$J_{52} = -\frac{3}{4}C^5 + 10C^4 - 5\left(\frac{3}{4}\pi^2 + 4\right)C^3 + 15(2\pi^2 - 7\zeta(3))C^2 + 5\left(112\zeta(3) - \frac{21}{16}\pi^4 - 6\pi^2\right)C - \frac{105}{2}\zeta(3)\pi^2 - 280\zeta(3) - 558\zeta(5) + \frac{35}{2}\pi^4$$

$$J_{53} = -\frac{15}{8}C^5 + \frac{115}{4}C^4 - 15\left(\frac{5}{8}\pi^2 + 6\right)C^3 + 15\left(\frac{23}{4}\pi^2 - \frac{35}{2}\zeta(3) + 4\right)C^2 + 5\left(322\zeta(3) - \frac{105}{32}\pi^4 - 27\pi^2\right)C - \frac{525}{4}\zeta(3)\pi^2 - 1260\zeta(3) - 1395\zeta(5) + \frac{805}{16}\pi^4 + 30\pi^2$$

$$J_{54} = -\frac{105}{16}C^5 + 110C^4 - 5\left(\frac{105}{16}\pi^2 + 86\right)C^3 + 15\left(22\pi^2 - \frac{245}{4}\zeta(3) + 32\right)C^2 + 5\left(1232\zeta(3) - \frac{735}{64}\pi^4 - 129\pi^2 - 24\right)C - \frac{3675}{8}\zeta(3)\pi^2 - 6020\zeta(3) - \frac{9765}{2}\zeta(5) + \frac{385}{2}\pi^4 + 240\pi^2$$

$$J_{55} = -\frac{945}{32}C^5 + \frac{8445}{16}C^4 - 25\left(\frac{189}{32}\pi^2 + 95\right)C^3 + 15\left(\frac{1689}{16}\pi^2 - \frac{2205}{8}\zeta(3) + 230\right)C^2 + 15\left(\frac{3941}{2}\zeta(3) - \frac{2205}{128}\pi^4 - \frac{475}{2}\pi^2 - 100\right)C - \frac{33075}{16}\zeta(3)\pi^2 - 33250\zeta(3) - \frac{87885}{4}\zeta(5) + \frac{59115}{64}\pi^4 + 1725\pi^2 + 120$$