EUROPEAN ORGANIZATION FOR NUCLEAR RESEARCH

CERN-EP/83-44
16 March 1983

STUDIES OF TWO-PARTICLE CORRELATION IN RAPIDITY SPACE

IN (pp) COLLISIONS AT $(\sqrt{s})_{pp} = 30, 44, \text{ AND } 62 \text{ GeV}$

J. Berbiers, G. Bonvicini, G. Cara Romeo, L. Cifarelli, A. Contin

M. Curatolo, G. D'Ali, C. Del Papa, B. Esposito, P. Giusti,

T. Massam, R. Nania, G. Natale, F. Palmonari, G. Sartorelli,

G. Susinno, L. Votano and A. Zichichi

CERN, Geneva, Switzerland

Istituto di Fisica dell'Università di Bologna, Italy

Istituto Nazionale di Fisica Nucleare, Laboratori Nazionali di Frascati, Italy

Istituto Nazionale di Fisica Nucleare, Sezione di Bologna, Italy

ABSTRACT

Two particle correlations in rapidity space in (pp) collisions are reported, using the leading proton subtraction method. The study is performed at different values of the nominal c.m. (pp) total energy $(\sqrt{s})_{pp} = 30, 44, \text{ and } 62 \text{ GeV}$. The dependence of the correlation function $R(y, y')$ at $y = y' = 0$, versus the effective hadron energy available for particle production, is reported.

(Submitted to Nuovo Cimento Letters)

We have already reported¹⁾ evidence for the same two-particle correlation in rapidity space in (e^+e^-) annihilation and in (pp) collisions, when the leading proton effects are taken into account. The method of subtracting the leading particle effects has already produced striking similarities between the properties of multiparticle systems produced in (e^+e^-) annihilation and in soft (pp) collisions²⁻¹⁷⁾. The aim of the present work is to go further in the studies of two-particle correlations and to show that this important feature of the hadronization mechanism is related to the effective total hadronic energy available for particle production, $\sqrt{(q_{\text{tot}}^{\text{had}})^2}$, not to the nominal c.m. total energy of the two colliding protons $(\sqrt{s})_{\text{pp}}$. The definition of $q_{\text{tot}}^{\text{had}}$ is:

$$q_{\text{tot}}^{\text{had}} = q_1^{\text{inc}} + q_2^{\text{inc}} - q_1^{\text{leading}} - q_2^{\text{leading}}$$

where $q_{1,2}^{\text{inc}}$ are the four-vectors of the incident protons and $q_{1,2}^{\text{leading}}$ the four-vectors of the leading protons. Following the (e^+e^-) method, the rapidity is calculated with respect to the sphericity axis, in the reference system where the space-like part of the four momentum $q_{\text{tot}}^{\text{had}}$ is zero. For details on the analysis we refer the reader to our previous paper¹⁾.

Only events with two identified protons and at least four charged tracks, besides the protons, are retained for the analysis.

The number of events, for each $\sqrt{(q_{\text{tot}}^{\text{had}})^2}$ interval obtained at three different values of $(\sqrt{s})_{\text{pp}}$ is shown in Table 1.

Table 1

Number of events for different intervals of $(\sqrt{s})_{\text{pp}}$
and $\sqrt{(q_{\text{tot}}^{\text{had}})^2}$ (both in GeV)

$(\sqrt{s})_{\text{pp}}$ $\sqrt{(q_{\text{tot}}^{\text{had}})^2}$	30	44	62
5-15	817	1543	247
15-25	71	1890	1226
25-36	-	24	1264

The correlation function $R(y, y')$ is defined as:

$$R(y, y') = \frac{\rho_2(y, y')}{f \rho_1(y) \rho_1(y')} - 1$$

where $\rho_1(y)$ is the rapidity single-particle density

$$\rho_1(y) = \frac{1}{\sigma_{in}} \frac{d\sigma(y)}{dy}$$

and $\rho_2(y, y')$ is the rapidity two-particle density

$$\rho_2(y, y') = \frac{1}{\sigma_{in}} \frac{d\sigma(y, y')}{dy dy'}$$

The correct normalization is granted by the parameter $f = \langle n_{ch}(n_{ch} - 1) \rangle / \langle n_{ch} \rangle^2$.

In Fig. 1 the comparison of two-particle rapidity correlations for two different values of $(\sqrt{s})_{pp}$ 30 and 44 GeV but with the same $\sqrt{(q_{tot}^{had})^2}$ in the range 5-15 GeV is reported.

The same comparison in the $\sqrt{(q_{tot}^{had})^2}$ range 15-25 GeV at two different values of $(\sqrt{s})_{pp}$ 44 and 62 GeV is shown in Fig. 2.

The agreement between two-particle correlations measured at different $(\sqrt{s})_{pp}$, but with the same $\sqrt{(q_{tot}^{had})^2}$ is very satisfactory.

These results show that we can add all the contributions from different $(\sqrt{s})_{pp}$ and evaluate the two-particle correlations in rapidity space for three different $\sqrt{(q_{tot}^{had})^2}$ intervals. The results are reported in Figs. 3a,b,c.

An interesting result of these measurements is that the maximum of $R(y, y')$ at $y = y' = 0$ is increasing with increasing $\sqrt{(q_{tot}^{had})^2}$. How $R(0,0)$ varies with $\sqrt{(q_{tot}^{had})^2}$ is shown in Fig. 4. For comparison, in the same Fig. 4, we have reported the values of $R(0,0)$ measured for final states of (pp) collisions without the subtraction of the leading proton effects. In this case the energy variable is $(\sqrt{s})_{pp}$. The results show no variation of $R(0,0)$ versus $(\sqrt{s})_{pp}$.

These studies are a further confirmation of the fact that the relevant quantity in describing high-energy and soft (pp) interactions is the effective total hadronic energy $\sqrt{(q_{\text{tot}}^{\text{had}})^2}$ available for multiparticle production, and not the nominal c.m. total energy $(\sqrt{s})_{\text{pp}}$ of the colliding protons.

REFERENCES

- 1) G. Bonvicini, G. Cara Romeo, L. Cifarelli, A. Contin, M. Curatolo, G. D'Alì, C. Del Papa, B. Esposito, P. Giusti, T. Massam, R. Nania, G. Natale, F. Palmonari, G. Sartorelli, G. Susinno, L. Votano and A. Zichichi, Evidence for the same two-particle correlations in rapidity space in (pp) collisions and (e^+e^-) annihilation, CERN-EP/83-29 (1983), submitted to Nuovo Cimento Letters.
- 2) M. Basile, G. Cara Romeo, L. Cifarelli, A. Contin, G. D'Alì, P. Di Cesare, B. Esposito, P. Giusti, T. Massam, F. Palmonari, G. Sartorelli, G. Valenti and A. Zichichi, Phys. Lett. 92B, 367 (1980).
- 3) M. Basile, G. Cara Romeo, L. Cifarelli, A. Contin, G. D'Alì, P. Di Cesare, B. Esposito, P. Giusti, T. Massam, R. Nania, F. Palmonari, G. Sartorelli, G. Valenti and A. Zichichi, Nuovo Cimento 58A, 193 (1980).
- 4) M. Basile, G. Cara Romeo, L. Cifarelli, A. Contin, G. D'Alì, P. Di Cesare, B. Esposito, P. Giusti, T. Massam, R. Nania, F. Palmonari, G. Sartorelli, G. Valenti and A. Zichichi, Phys. Lett. 95B, 311 (1980).
- 5) M. Basile, G. Cara Romeo, L. Cifarelli, A. Contin, G. D'Alì, P. Di Cesare, B. Esposito, P. Giusti, T. Massam, R. Nania, F. Palmonari, G. Sartorelli, G. Valenti and A. Zichichi, Nuovo Cimento Lett. 29, 491 (1980).
- 6) M. Basile, G. Cara Romeo, L. Cifarelli, A. Contin, G. D'Alì, P. Di Cesare, B. Esposito, P. Giusti, T. Massam, R. Nania, F. Palmonari, G. Sartorelli, M. Spinetti, G. Susinno, G. Valenti and A. Zichichi, Phys. Lett. 99B, 247 (1981).
- 7) M. Basile, G. Cara Romeo, L. Cifarelli, A. Contin, G. D'Alì, P. Di Cesare, B. Esposito, P. Giusti, T. Massam, R. Nania, F. Palmonari, G. Sartorelli, M. Spinetti, G. Susinno, G. Valenti and A. Zichichi, Nuovo Cimento Lett. 30, 389 (1981).

- 8) M. Basile, G. Cara Romeo, L. Cifarelli, A. Contin, G. D'Alì, P. Di Cesare, B. Esposito, P. Giusti, T. Massam, R. Nania, F. Palmonari, G. Sartorelli, M. Spinetti, G. Susinno, G. Valenti, L. Votano and A. Zichichi, Nuovo Cimento Lett. 31, 273 (1981).
- 9) M. Basile, G. Cara Romeo, L. Cifarelli, A. Contin, G. D'Alì, P. Di Cesare, B. Esposito, P. Giusti, T. Massam, R. Nania, F. Palmonari, V. Rossi, G. Sartorelli, M. Spinetti, G. Susinno, G. Valenti, L. Votano and A. Zichichi, Nuovo Cimento 65A, 400 (1981).
- 10) M. Basile, G. Cara Romeo, L. Cifarelli, A. Contin, G. D'Alì, P. Di Cesare, B. Esposito, P. Giusti, T. Massam, R. Nania, F. Palmonari, V. Rossi, G. Sartorelli, M. Spinetti, G. Susinno, G. Valenti, L. Votano and A. Zichichi, Nuovo Cimento 65A, 414 (1981).
- 11) M. Basile, G. Cara Romeo, L. Cifarelli, A. Contin, G. D'Alì, P. Di Cesare, B. Esposito, P. Giusti, T. Massam, R. Nania, F. Palmonari, V. Rossi, G. Sartorelli, M. Spinetti, G. Susinno, G. Valenti, L. Votano and A. Zichichi, Nuovo Cimento Lett. 32, 210 (1981).
- 12) M. Basile, G. Cara Romeo, L. Cifarelli, A. Contin, G. D'Alì, P. Di Cesare, B. Esposito, P. Giusti, T. Massam, R. Nania, F. Palmonari, V. Rossi, G. Sartorelli, M. Spinetti, G. Susinno, G. Valenti, L. Votano and A. Zichichi, Nuovo Cimento 66A, 129 (1981).
- 13) M. Basile, G. Cara Romeo, L. Cifarelli, A. Contin, G. D'Alì, P. Di Cesare, B. Esposito, P. Giusti, T. Massam, R. Nania, F. Palmonari, V. Rossi, G. Sartorelli, M. Spinetti, G. Susinno, G. Valenti, L. Votano and A. Zichichi, Nuovo Cimento Lett. 32, 321 (1981).
- 14) M. Basile, G. Cara Romeo, L. Cifarelli, A. Contin, G. D'Alì, P. Di Cesare, B. Esposito, P. Giusti, T. Massam, R. Nania, F. Palmonari, V. Rossi, F. Rohrbach, G. Sartorelli, M. Spinetti, G. Susinno, G. Valenti, L. Votano and A. Zichichi, Nuovo Cimento 67A, 53 (1982).

- 15) M. Basile, G. Bonvicini, G. Cara Romeo, L. Cifarelli, A. Contin, M. Curatolo, G. D'Alì, P. Di Cesare, B. Esposito, P. Giusti, T. Massam, R. Nania, F. Palmonari, A. Petrosino, V. Rossi, G. Sartorelli, M. Spinetti, G. Susinno, G. Valenti, L. Votano and A. Zichichi, Nuovo Cimento 67A, 244 (1982).
- 16) M. Basile, G. Cara Romeo, L. Cifarelli, A. Contin, M. Curatolo, G. D'Alì, P. Di Cesare, B. Esposito, P. Giusti, T. Massam, R. Nania, F. Palmonari, A. Petrosino, V. Rossi, G. Sartorelli, M. Spinetti, G. Susinno, G. Valenti, L. Votano and A. Zichichi, preprint CERN-EP/81-147 (1981).
- 17) M. Basile, G. Bonvicini, G. Cara Romeo, L. Cifarelli, A. Contin, M. Curatolo, G. D'Alì, C. Del Papa, B. Esposito, P. Giusti, T. Massam, R. Nania, F. Palmonari, G. Sartorelli, M. Spinetti, G. Susinno, L. Votano and A. Zichichi, Experimental proof that the leading protons are not correlated, preprint CERN-EP/82-182 (1982), submitted to Nuovo Cimento.

Figure captions

Fig. 1 : Correlation function $R(y, y')$ measured in the $\sqrt{(q_{\text{tot}}^{\text{had}})^2}$ range 5-15 GeV for two different values of $(\sqrt{s})_{\text{pp}}$. The results for $(\sqrt{s})_{\text{pp}} = 44$ GeV are plotted as open circles; for $(\sqrt{s})_{\text{pp}} = 30$ GeV as black points.

Fig. 2 : Correlation function $R(y, y')$ measured in the $\sqrt{(q_{\text{tot}}^{\text{had}})^2}$ range 15-25 GeV for two different values of $(\sqrt{s})_{\text{pp}}$. The results for $(\sqrt{s})_{\text{pp}} = 62$ GeV are plotted as open squares; for $(\sqrt{s})_{\text{pp}} = 44$ GeV as black points.

Fig. 3 : Correlation function $R(y, y')$ measured for three different $\sqrt{(q_{\text{tot}}^{\text{had}})^2}$ ranges, adding the contributions from the data taken at $(\sqrt{s})_{\text{pp}} = 30, 44, \text{ and } 62$ GeV.

a) $5 \leq \sqrt{(q_{\text{tot}}^{\text{had}})^2} \leq 15$ GeV

b) $15 \leq \sqrt{(q_{\text{tot}}^{\text{had}})^2} \leq 25$ GeV

c) $25 \leq \sqrt{(q_{\text{tot}}^{\text{had}})^2} \leq 36$ GeV

Fig. 4 : Correlation function $R(y, y')$ measured at $y = y' = 0$, versus $\sqrt{(q_{\text{tot}}^{\text{had}})^2}$. The data are indicated as black points. The broken line is the best fit.

For comparison, the open triangles show $R(0,0)$ versus $(\sqrt{s})_{\text{pp}}$. In this case the analysis of the final state is made without subtracting the leading proton effects. The dash-dotted line is the best fit to these data.

Notice that the abscissa for the black points is $\sqrt{(q_{\text{tot}}^{\text{had}})^2}$; for the open triangles, it is $(\sqrt{s})_{\text{pp}}$.

Fig. 1

Fig. 2

$\sqrt{(q_{\text{tot}}^{\text{had}})^2} = 5-15 \text{ GeV}$

Fig. 3a

$\sqrt{(q_{\text{tot}}^{\text{had}})^2} = 15\text{-}25 \text{ GeV}$

Fig. 3b

$\sqrt{(q_{\text{tot}}^{\text{had}})^2} = 25-36 \text{ GeV}$

Fig. 3c

Fig. 4