Ref.TH.3581-CERN

TESTING THE ORIGIN OF THE $\Delta I = \frac{1}{2}$ RULE THROUGH K_L DALITZ DECAYS

L. Bergström, E. Massó ^{*)} and P. Singer ^{**)}

CERN - Geneva

A B S T R A C T

It is pointed out that the process $K_L \rightarrow \ell^+ \ell^- \gamma$ has some specific properties making it a useful tool to study the structure of the weak non-leptonic interaction. In particular, it is demonstrated how the scheme using penguin diagrams to enhance the $K-\pi$ transition may experimentally be distinguished from, e.g., Sakurai's phenomenological Hamiltonian model through a measurement of the ratio $\Gamma(K_L \rightarrow e^+ e^- \gamma) / \Gamma(K_L \rightarrow \mu^+ \mu^- \gamma)$.

^{*)} On leave of absence from Departament de Física Teòrica, Universitat Autònoma de Barcelona, Bellaterra, Spain.

^{**)} On leave from Technion-Israel Institute of Technology, Haifa, Israel.

Non-leptonic decays of hadrons constitute a framework where the interplay between the $SU(2) \times U(1)$ and $SU(3)_c$ gauge models, describing electroweak and strong interactions respectively, is present. Indeed, in order to analyze those decays one must consider the rôle played by QCD in the structure of the weak transitions [see, e.g., Ref. 1) for a review]. To lowest order in weak interactions, the modifications induced by QCD in the non-leptonic weak Hamiltonian are expressed through the Wilson operator expansion ^{2),3)}

$$H_w^{NL} = \frac{G_F}{\sqrt{2}} \sum_i c_i O_i, \quad (1)$$

where c_i are c number coefficients which can be calculated by using renormalization group techniques, and O_i are the lowest (non-trivial) dimension operators that form the expansion. Since quarks are not degenerate in mass, Eq. (1) receives contributions not only from the diagrams shown in Fig. 1a ²⁾, but also from the "penguin" diagrams of Fig. 1b ³⁾.

This approach has been applied in particular to the study of the $\Delta I = \frac{1}{2}$ rule in kaonic decays, i.e., to explain the experimental ratio $r_{\text{exp}} \equiv A(\Delta I = \frac{1}{2}) / A(\Delta I = \frac{3}{2}) \approx 20$. The diagrams of the type shown in Fig. 1a give only a contribution $|r_a| \approx 4-5$. The penguin diagrams 1b which are purely $\Delta I = \frac{1}{2}$ also give a sizeable contribution, since although their coefficients c_i are quite small the structure of the penguin operators induces a compensating enhancement factor $O(m_\pi^2/m_u m_s)$ when calculating the matrix element for the $K-\pi$ transition ³⁾. Hence the penguin diagrams are obviously relevant for the description of the non-leptonic decays. Although their evaluation is plagued by some uncertainties ⁴⁾, it has nevertheless been claimed that they lead to a quantitative understanding of kaon non-leptonic decays ³⁾.

To elucidate further the rôle of the penguin contributions, one should consider processes where penguins are either believed to be clearly dominating or where their absence distinguishes them from other explanations of the $\Delta I = \frac{1}{2}$ rule. The purpose of this paper is to show that the Dalitz decay $K_L^0 \rightarrow \ell^+ \ell^- \gamma$ could shed some light on the question of the penguins through the latter approach. The CP violating K^0 decays ⁵⁾ and the Cabibbo suppressed decays of naked bottom mesons ⁶⁾ are, on the other hand, processes where penguins are believed to dominate. Unfortunately, the predictions for these decays are for the time being difficult to test, in contrast to the process we suggest.

In order to discuss the $K_L \rightarrow \ell^+ \ell^- \gamma$ transition we must refer to the $K_L \rightarrow \gamma\gamma$ decay and, naturally, a comparison with the similar decays of the pseudo-scalar mesons, π^0, η, η' is in order. The two-photon mediated decays of neutral non-strange pseudoscalar mesons (P), i.e., $P \rightarrow \gamma\gamma, P \rightarrow \ell^+ \ell^- \gamma$ and $P \rightarrow \ell^+ \ell^-$ have been discussed extensively in the literature and a fair understanding of these processes has been achieved. In particular, Ametller et al. ⁷⁾ have recently presented closed-form expressions for the two-photon form factor $\langle 0 | T \{ j_\mu^{\text{em}}(x) j_\nu^{\text{em}}(y) \} | P \rangle$ in a quark triangle model with pointlike γ_5 couplings between quarks and mesons, and used them for analyzing the above-mentioned processes. It turns out that a comparison of the quark-model calculation with the vector meson dominance approach ⁸⁾ reveals their equivalence in a local sense, both for $P \rightarrow \ell^+ \ell^- \gamma$ ⁹⁾ and $P \rightarrow \ell^+ \ell^-$ ¹⁰⁾.

Although the processes $K_L \rightarrow \gamma\gamma$ and $\ell^+ \ell^- \gamma$ are superficially similar to the non-strange P decays, the weak interaction now introduces a new element, the physical quantity involved being $\langle 0 | T \{ j_\mu^{\text{em}}(x) j_\nu^{\text{em}}(y) H_W^{\text{NL}}(0) \} | K_L \rangle$. As the non-leptonic weak Hamiltonian H_W^{NL} generates several types of diagrams, the structure of the $K_L \rightarrow \ell^+ \ell^- \gamma$ matrix element may be quite different from the otherwise analogous electromagnetic processes $P \rightarrow \ell^+ \ell^- \gamma$ [for a somewhat related approach to the structure of the $K^+ \rightarrow \pi^+ e^+ e^-$ matrix element, see Ref. 11)]. Indeed, there are two general classes of tree diagrams for $K_L \rightarrow \gamma^* \gamma$, with H_W^{NL} acting on the meson or a (virtual) photon leg, as illustrated in Figs. 1c and 1d, respectively. Diagram 1d does not contribute to the $K_L \rightarrow \gamma\gamma$ decay, since the general structure of the $s\bar{d} \rightarrow \gamma^*$ transition requires it to vanish for a real photon ^{12), 13)}. On the other hand, diagram 1c is directly related to the $K_L \rightarrow \gamma\gamma$ decay. For an off-shell photon, both diagrams 1c and 1d contribute; there is obviously no similar situation in the purely electromagnetic transitions of π^0, η, η' .

It is generally recognized ¹²⁾⁻¹⁵⁾ that the strength of the $K_L \rightarrow \gamma\gamma$ transition is determined by the long-range (pole) contributions, the quark loop calculations yielding only some 10% of the observed strength. Our calculation of $K_L \rightarrow \ell^+ \ell^- \gamma$ will thus involve an estimation of the long-range contributions yielding diagram 1c. For diagram 1d, we take the $K^* K \gamma$ coupling from the experimental value for $K^* \rightarrow K \gamma$ ¹⁶⁾ and present two alternatives for calculating the $K^* \rightarrow \rho, \omega, \phi$ transition: a quark model calculation and a calculation based on the same phenomenological Hamiltonian that accounts for the observed $K_L \rightarrow \gamma\gamma$. The sizeably different results to which the two approaches lead is what makes $K_L \rightarrow \ell^+ \ell^- \gamma$ the sensitive probe alluded to beforehand.

For the long-range contribution we use the phenomenological current-current Hamiltonian of Sakurai¹⁷⁾, which is known to account remarkably well for a large number of weak baryonic and K meson decays¹⁷⁾⁻¹⁹⁾. Its explicit form is

$$H_W^{NL(s)}(\Delta I = \frac{1}{2}) = \sqrt{2} G_{NL} d_{6ab} \bar{J}_\mu^a(x) J^{\mu b}(x), \quad (2)$$

where $\bar{J}_\mu^a(x)$ are the hadronic SU(3) currents and d_{6ab} is an SU(3) constant. The effective coupling G_{NL} turns out to be practically equal to the Fermi constant, this enhancement being achieved in this approach by simply disregarding the Cabibbo angle in (2). The hadronic currents are saturated by the low-lying particles with the appropriate quantum numbers. In the calculation of diagram 1c we retain the contribution of the pseudoscalar meson poles. There is then the question of how to supplement (2) to include the η' contribution. We have considered three possibilities :

i) extension of Eq. (2) to SU(4) leading to¹⁵⁾

$$f_{K_L \eta} \equiv \langle \eta | H_W^{NL} | K_L \rangle = \frac{1}{\sqrt{6}} \langle \pi^0 | H_W^{NL} | K_L \rangle \equiv \frac{1}{\sqrt{6}} f_{K_L \pi^0} ;$$

ii) nonet symmetry for the coupling of pseudoscalar mesons, giving²⁰⁾

$$f_{K_L \eta} = 2\sqrt{\frac{2}{3}} f_{K_L \pi^0} ;$$

iii) assuming the effective neutral current in Eq. (2) to behave like $(u\bar{u}-c\bar{c})$ which is the dominant term in the effective non-leptonic Hamiltonian¹²⁾; this assumption gives

$$f_{K_L \eta} = \sqrt{\frac{2}{3}} f_{K_L \pi^0} .$$

To calculate the $K_L \rightarrow \gamma\gamma$ amplitude we use a mixing angle for η, η' of $\theta = -8.4^\circ$, which we derive from the reported 2γ widths of π^0, η, η' . We also consider it appropriate to use $F_K = F_\eta = F_{\eta'} = F_\pi$ for the pseudoscalar couplings of PCAC, in view of the uncertainties in these constants. For the $P \rightarrow \gamma\gamma$ form factor $F_p(q^2)$ we use the value at $q^2 = m_K^2$, as calculated in Ref. 7). We thus obtain for the effective amplitude of $K_L \rightarrow \gamma\gamma$, defined by

$$A(K_L \rightarrow \gamma\gamma) = A_{\gamma\gamma} \epsilon_{\mu\nu\sigma\tau} \epsilon_1^\mu \epsilon_2^\nu k_1^\sigma k_2^\tau, \quad (3)$$

in the three models described

$$\begin{aligned} A_{\gamma\gamma} &= 3.0 \times 10^{-12} \text{ MeV}^{-1} & (i) \\ &= 6.7 \times 10^{-12} \text{ MeV}^{-1} & (ii) \\ &= 4.2 \times 10^{-12} \text{ MeV}^{-1} & (iii) \end{aligned}$$

versus an experimental value of

$$|A_{\gamma\gamma}|_{\text{exp}} = (3.19 \pm 0.33) \times 10^{-12} \text{ MeV}^{-1}.$$

Thus, both models (i) and (iii) account well for the $K_L \rightarrow \gamma\gamma$ rate and one has thus acquired confidence that we have a suitable effective Hamiltonian to estimate the long-range contributions also to $K_L \rightarrow \ell^+ \ell^- \gamma$.

The $K_L \rightarrow \ell^+ \ell^- \gamma$ amplitude is a sum of lc and ld. For the former, we use the measured $|A_{\gamma\gamma}|_{\text{exp}}$ of the $K_L \rightarrow \gamma\gamma$ transition, in combination with vector meson dominance of the virtual photon. Using Eq. (2), the form factor corresponding to diagram ld is given by ¹⁹⁾

$$A_{K^*}(s) = \sqrt{2} e G_{NL} f_{K^*} K_Y \frac{m_\rho^2}{f_{K^*} f_\rho^2} \left(\frac{1}{1 - s/m_{K^*}^2} \right) \left\{ \frac{4}{3} - \frac{1}{1 - s/m_\rho^2} - \frac{1}{9} \left(\frac{1}{1 - s/m_\omega^2} + \frac{2}{1 - s/m_\phi^2} \right) \right\}, \quad (4)$$

which vanishes for a real photon ($s=0$), as required by gauge invariance ^{12),13),19)}. Here the coupling strength $\sim G_{NL}$ (without Cabibbo suppression) of the $K^* \rightarrow \rho, \omega, \phi$ transition is a direct reflection of using (2) universally for both pseudoscalar and vector mesons. In the general case, the form factor A_D for the Dalitz decay $K_L \rightarrow \ell^+ \ell^- \gamma$ is given by

$$A_D(s) = \frac{|A_{\gamma\gamma}|_{\text{exp}}}{1 - s/m_\rho^2} + \alpha A_{K^*}(s) \quad (5)$$

where the first term is a good representation of diagram lc with one virtual photon, and where we have introduced a parameter α to parametrize the relative strength between diagrams lc and ld. The model just considered gives $|\alpha| = 1$, whereas we get a significantly smaller value for $|\alpha|$ in the quark model calculation, to which we now turn.

In the standard model, the non-leptonic effective Hamiltonian contains the Cabibbo suppression factor $\sin\theta_c \cos\theta_c$ (or $\sin\theta_1 \cos\theta_1 \cos\theta_3$ in the Kobayashi-Maskawa scheme). Gluonic (QCD) corrections to the bare Hamiltonian (Fig. 1a) may enhance some four-quark $(V-A) \times (V-A)$ operators, whereas others are suppressed ²⁾. In particular, as we have mentioned, the $\Delta I = \frac{1}{2}$ operators are enhanced over the $\Delta I = \frac{3}{2}$. The innovation brought in by Shifman, Vainshtein and Zakharov (SVZ) ³⁾

was the realization that the penguin diagrams lb in addition give rise to terms with a $(V-A) \times (V+A)$ structure. These can be Fierz transformed to $(SS-PP)$, i.e., with the right quantum numbers for scalar or pseudoscalar particles. This is the physical content of the penguin enhancement of, e.g., the pseudoscalar-pseudoscalar $K-\pi$ transition. The point is now that, accordingly, this particular enhancement will be absent for the vector-vector $K^* \rightarrow \rho, \omega, \phi$ transition we are interested in.

For simplicity of the meson quark contents, we consider the $K^{*+}-\rho^+$ transition. The renormalization group improved effective SVZ Hamiltonian (including penguins) is ³⁾

$$H_w^{NL}(\Delta S=1) = \sqrt{2} G_F \sin\theta_c \cos\theta_c \left\{ -2.5 O_1 + 0.08 O_2 + 0.08 O_3 + \right. \\ \left. + 0.4 O_4 - (0.06 \div 0.14) O_5 - (0.01 \div 0.05) O_6 \right\}, \quad (6)$$

where O_1-O_6 are four-quark operators with definite isospin and $SU(3)_F$ transformation properties ; e.g.,

$$O_1 = (\bar{s}_i \gamma_\mu d^i)_L (\bar{u}_j \gamma^\mu u^j)_L - (\bar{s}_i \gamma_\mu u^i)_L (\bar{u}_j \gamma^\mu d^j)_L \quad (\{8_F\}, \Delta I = \frac{1}{2}),$$

where $i = 1, 2, 3$ is a colour index [for a complete list of the O_i , see Ref. 3)]. The (small) effects of the $(V-A) \times (V-A)$ parts of the penguin diagrams are included in the coefficients of O_1-O_4 , whereas O_5 and O_6 are the additional $(V-A) \times (V+A)$ operators. Following SVZ, we use the valence quark model with vacuum insertion for the $K^{*+}-\rho^+$ transition amplitude induced by Eq. (6). A straightforward calculation gives

$$\langle K^{*+} | H_w^{NL}(\Delta S=1) | \rho^+ \rangle \simeq 1.2 \sqrt{2} G_F \frac{m_{K^*}^2 m_\rho^2}{f_{K^*} f_\rho} \sin\theta_c \cos\theta_c \epsilon_\mu^*(k^*) \epsilon^\mu(\rho), \quad (7)$$

where we have defined

$$\langle 0 | (\bar{d}_i \gamma^\mu u^i)_L | \rho^+ \rangle = \frac{1}{3} \delta_i^j \frac{1}{\sqrt{2}} \frac{m_\rho^2}{f_\rho} \epsilon^\mu(\rho) \quad (8)$$

and a similar expression for the K^{*+} matrix element. The result in Eq. (7) is only a 20% enhancement over the bare value without QCD corrections and not at all enough to overcome the Cabibbo suppression. It is smaller by $\sim 1.2 \sin\theta_c$ than the result for this transition calculated with (2) and used in Eq. (4). The value in Eq. (7) would correspond to $|\alpha| \simeq 0.2-0.3$ in Eq. (5) including all vector mesons.

We can thus draw the conclusion that in the SVZ scheme there is no significant enhancement of the vector-vector non-leptonic transitions. This is in contrast to the case of the phenomenological Hamiltonian in Eq. (2) where pseudoscalar-pseudoscalar and vector-vector transitions are of the same, universal strength.

Turning now to the numerical results, the decay spectrum for $K_L \rightarrow \ell^+ \ell^- \gamma$, normalized to the $\gamma\gamma$ width is given by

$$\frac{1}{\Gamma_{\gamma\gamma}} \frac{d\Gamma}{ds} = \frac{e^2}{6\pi^2} \frac{1}{s} (1-s/m_K^2)^3 (1+2m_\ell^2/s) \sqrt{1-4m_\ell^2/s} |F(s,0;m_K^2)|^2, \quad (9)$$

where $F(s,0;m_K^2)$ is the two-photon form factor with one photon off-shell, normalized such that $F(0,0;m_K^2) = 1$. In our case, $F(s,0;m_K^2) = A_D(s)/|A_{\gamma\gamma}|_{\text{exp}}$, with $A_D(s)$ given by Eq. (5). Using this form factor, with α kept as a free parameter, we can integrate (9) to get the branching ratios for $K_L \rightarrow \mu^+ \mu^- \gamma$ and $K_L \rightarrow e^+ e^- \gamma$ as a function of α . This is shown in Fig. 2, where also the ratio $R(\alpha) \equiv \Gamma(K_L \rightarrow e^+ e^- \gamma) / \Gamma(K_L \rightarrow \mu^+ \mu^- \gamma)$ is displayed. As can be seen, $R(\alpha)$ is a quite rapidly varying function of α . The value for $\alpha = 0$ (no K^* contribution) is $R(0) = 29.2$ which coincides with an old result of Seghal²¹⁾, who did not consider contributions of the type $1d$. The Sakurai model corresponds to $|\alpha| = 1$, giving $R(-1) = 14.2$, $R(+1) = 87.8$. With the SVZ type non-leptonic Hamiltonian, Eq. (6), we obtained $|\alpha| \approx 0.2-0.3$, giving, e.g., $R(|\alpha|) \approx 37$ or 24 for $|\alpha| = 0.25$, depending on the sign of α (about which we a priori have nothing to say).

It is clear that an experiment with reasonably good precision and statistics should be able to distinguish between these possibilities. We note that there already exists data for both decay modes²²⁾ but with only a handful of events. In Fig. 2, we also present the branching ratios into $\mu^+ \mu^- \gamma$ and $e^+ e^- \gamma$ separately as a function of α . As can be seen, almost the entire variation of $R(\alpha)$ is caused by the sensitivity of the $\mu^+ \mu^- \gamma$ branching ratio to α , whereas the $e^+ e^- \gamma$ rate is almost independent of α . This is easily understood from the $1/s$ dependence of the spectrum [Eq. (9)]. The $e^+ e^- \gamma$ decay mode is strongly peaked near the threshold $s = 4m_e^2$. For such low values of s , the contribution from the K^{*0} term is small since it necessarily vanishes on the photon mass shell $s = 0$. For the $\mu^+ \mu^- \gamma$ decay, on the other hand, the threshold is much higher ($s = 4m_\mu^2$) and there the K^{*0} contribution is already significant. In Fig. 3, we have displayed the influence of the K^{*0} term on the differential invariant mass spectrum [Eq. (9)] for $K_L \rightarrow \mu^+ \mu^- \gamma$. It is clear that also a measurement of this invariant mass distribution itself could give an indication of the magnitude and sign of α , although the ratio $R(\alpha)$ probably is preferable experimentally since some systematic uncertainties may cancel in the ratio.

Our conclusions are thus the following.

- A) $K_L \rightarrow \gamma\gamma$ is well understood in a pole model.
- B) $K_L \rightarrow \ell^+\ell^-\gamma$ adds a new non-trivial feature related to the structure of the weak non-leptonic interaction. We therefore expect deviations from the old predictions of Seghal²¹⁾.
- C) A dedicated experiment provides a new test of models for the weak non-leptonic transitions. In particular, it would easily enable a distinction between the model of Sakurai¹⁷⁾⁻¹⁹⁾ and that of Shifman, Vainshtein and Zakharov³⁾. In fact, the measurement of $R(\alpha)$ provides a unique opportunity for a direct determination of a weak non-leptonic vector-vector transition, which is of obvious interest in itself.

ACKNOWLEDGEMENTS

We thank D. Nanopoulos and D. Stech for useful discussions and for a careful reading of the manuscript. One of us (E.M.) is grateful to "Caixa de Barcelona" for financial support.

REFERENCES

- 1) R.D. Peccei - Proc. XVII Karpacz Winter School, Poland, 1980 (Harwood Academic Publishers, New York, 1981).
- 2) G. Altarelli and L. Maiani - Phys.Lett 52B (1974) 351 ;
M.K. Gaillard and B.W. Lee - Phys.Rev.Lett. 33 (1974) 108.
- 3) M.A. Shifman, A.I. Vainshtein and V.I. Zakharov - Nucl.Phys. B120 (1977) 316 ;
Soviet Phys. JETP 45 (1977) 670.
- 4) J.F. Donoghue et al. - Phys.Rev. D21 (1980) 186.
- 5) F. Gilman and M. Wise - Phys.Rev. D20 (1979) 2392 ;
B. Guberina and R.D. Peccei - Nucl.Phys. B163 (1980) 289.
- 6) B. Guberina, R.D. Peccei and R. Rückl - Phys.Lett. 90B (1980) 169.
- 7) Ll. Ametller et al. - CERN Preprint TH. 3501 (1983), to be published.
- 8) C. Quigg and J.D. Jackson - UCRL Report No 18487 (1968), unpublished.
- 9) A. Bramon and E. Massó - Phys.Lett. 104B (1981) 311.
- 10) L. Bergström et al. - Phys.Lett. B, in press.
- 11) A.I. Vainshtein et al. - Soviet J.Nucl.Phys. 24 (1976) 427.
- 12) M.K. Gaillard and B.W. Lee - Phys.Rev. D10 (1974) 897.
- 13) E. Witten - Nucl.Phys. B122 (1977) 109.
- 14) D.V. Nanopoulos and G.G. Ross - Phys.Lett. 56B (1975) 279.
- 15) E. Ma and A. Pramudita - Phys.Rev. D24 (1981) 2476.
- 16) W.C. Carithers et al. - Phys.Rev.Lett. 35 (1975) 349.
- 17) J.J. Sakurai - Phys.Rev. 156 (1967) 1508.
- 18) M. Moshe and P. Singer - Phys.Rev.Lett. 22 (1971) 1685 ; Phys.Rev. D6 (1972) 1379.
- 19) R.F. Sarraga and H.J. Munczek - Phys.Rev. D4 (1971) 2884.
- 20) Y. Dupont and T.N. Pham - Ecole Polytechnique Preprint, unpublished.
- 21) L.M. Seghal - Phys.Rev. D11 (1973) 3303.
- 22) A.S. Carroll et al. - Phys.Rev.Lett. 44 (1980) 525.

FIGURE CAPTIONS

Figure 1 a) Conventional four-quark diagram for the weak non-leptonic interaction.
b) Penguin diagram contribution.
c) Pole model for $K_L \rightarrow \gamma\gamma$.
d) Additional contribution through $K^*K\gamma$ coupling and $K^* \rightarrow \rho, \omega, \phi$ transition. This contribution is non-vanishing only if the photon originating from the K^* leg is virtual.

Figure 2 Branching ratios for $K_L \rightarrow e^+e^-\gamma$ and $K_L \rightarrow \mu^+\mu^-\gamma$ as a function of α , which measures the admixture of contributions of the type in Fig. 1d. Also shown is $R(\alpha) = \Gamma(K_L \rightarrow e^+e^-\gamma) / \Gamma(K_L \rightarrow \mu^+\mu^-\gamma)$.

Figure 3 The decay distribution for $K_L \rightarrow \mu^+\mu^-\gamma$ as a function of $x = s/m_K^2$, where s is the invariant mass of the lepton pair. The shape for $\alpha = 0$ (no K^{*0} contribution) coincides with that of Seghal²¹⁾. The values $\alpha = \pm 1$ correspond to the Sakurai model discussed in the text.

- Figure 1 -

- Figure 2 -

- Figure 3 -