EUROPEAN ORGANIZATION FOR NUCLEAR RESEARCH

CERN-EP/83-77

9 June 1983

NEUTRINO-ELECTRON SCATTERING

Klaus Winter

CERN, Geneva, Switzerland

Invited paper presented at
the Europhysics Study Conference on
Electroweak Effects at High Energies
Erice, Sicily, 1-10 February 1983
(To be published by Plenum Publishing Corporation)

NEUTRINO-ELECTRON SCATTERING

Klaus Winter

CERN, Geneva, Switzerland

1. INTRODUCTION

The first experimental observation of a weak neutral-current phenomenon was on the scattering of $\bar{\nu}_\mu$ on electrons ten years ago by the Gargamelle Collaboration at CERN¹⁾. Further observations, in particular on semileptonic neutral-current neutrino interactions²⁾ and on the scattering of polarized electrons on deuterium³⁾, have by now lent strong support to a unified gauge theory of the weak and the electromagnetic interaction as proposed by Weinberg and Salam. The main goal of pursuing the study of neutrino scattering on electrons is to determine the coupling constants of the leptonic weak neutral current, thus avoiding the ambiguities inherent in the use of hadronic targets. Until now, however, the extremely low cross-section has drastically limited the number of observed events, as illustrated in Tables 1 and 2. Taking all experiments together, about 100 events have now been observed in each channel.

At this conference we have heard reports on recent results on the ratio of $\sigma(\nu_\mu e)/\sigma(\bar{\nu}_\mu e)$, obtained by the CHARM collaboration at CERN⁴⁾, on progress from a beautiful new experiment⁵⁾ at the AGS in Brookhaven and on preliminary results from the flash-tube detector at Fermilab.

The axial-vector coupling constant g_A^e and the vector coupling constant g_V^e can be determined by a phenomenological, model-independent analysis. Assuming that leptons are point-like and that the outgoing neutrino is identical to the incoming neutrino, we obtain for the differential cross-section,

Table 1. Summary of past activities on $\sigma(\nu_\mu e)$

Experiment		$\nu_\mu e$	Background	$10^{-42} \frac{\sigma}{\text{cm}^2/\text{GeV}}$
1973 GGM PS		1	0.3 ± 0.1	$< 3(90\% \text{ c.l.})$
Aachen/Pad		11	3	1.1 ± 0.6
GGM SPS	64K	9	0.5 ± 0.2	$2.4 \pm \begin{smallmatrix} 1.2 \\ 0.9 \end{smallmatrix}$
Col-BNL 15' BC	84K	8	0.5 ± 0.5	1.8 ± 0.8
LM O		40	12	1.4 ± 0.3
CHARM	10^6	46 ± 12	64 ± 10	$2.1 \pm 0.55(\text{stat}) \pm 0.49$
World average $\sin^2\theta = 0.24 \pm \begin{smallmatrix} 0.06 \\ 0.04 \end{smallmatrix}$				1.55 ± 0.26

Table 2. Summary of past activities on $\sigma(\bar{\nu}_\mu e)$

GGM PS		3	0.4 ± 0.1	$1.0 \pm \begin{smallmatrix} 2.1 \\ 0.9 \end{smallmatrix}$
A-P		8	1.7	2.2 ± 1.0
CHARM SPS	10^6	77	146	$1.6 \pm 0.35(\text{stat}) \pm 0.36$
World average $\sin^2\theta = 0.37 \pm 0.05$				1.58 ± 0.30

$$\frac{d\sigma}{dy}(\nu_\mu e) = \frac{\sigma_0}{2} E_\nu \left[\left(g_V^e + g_A^e \right)^2 + \left(g_V^e - g_A^e \right)^2 (1-y)^2 + \frac{m_e}{E_\nu} \left(g_V^{e^2} - g_A^{e^2} \right) \right] \rho^2 \quad (1a)$$

and

$$\frac{d\sigma}{dy}(\bar{\nu}_\mu e) = \frac{\sigma_0}{2} E_\nu \left[\left(g_V^e - g_A^e \right)^2 + \left(g_V^e + g_A^e \right)^2 (1-y)^2 - \frac{m_e}{E_\nu} \left(g_V^{e^2} - g_A^{e^2} \right) \right] \rho^2 \quad (1b)$$

where $\sigma_0 = G^2 m_e / \pi = 8.6 \times 10^{-42} \text{ cm}^2$. ρ is an overall factor

related to the coupling at the $\nu_\mu Z^0 \nu_\mu$ vertex. If $\rho = 1$, $M_W = M_Z \cos \theta$ and the coupling of the neutral and charged current is of universal strength, $G_F^{NC} = G_F^{CC}$.

In the standard model $\rho = 1$ if one doublet of Higgs bosons is sufficient to describe the spontaneous symmetry breaking. Recently, C. Jarlskog and F. Yndurain⁷⁾ have estimated deviations of ρ from one due to the possible existence of lepton doublets with large mass differences between the neutral (neutrino) and charged component. They find

$$\epsilon = |\rho^2 - 1| \geq \frac{G_F}{8\sqrt{2}\pi^2} \sum_{\ell=1}^{N_e} m_\ell^2$$

for N_ℓ new families beyond the (τ, ν_τ) . Inserting the present experimental limit of

$$\epsilon < 3\%,$$

derived from a recent analysis of semileptonic neutral current data⁸⁾, and assuming $m_\ell > 20$ GeV, as suggested by searches for new heavy leptons at PETRA⁹⁾, a limit on the number of new lepton generations can be derived, giving

$$N_\nu < 737.$$

Combining this analysis with the present limit on modifications to the photon propagator in the reaction $e^+e^- \rightarrow \mu^+\mu^-$ due to heavy leptons

$$\sigma = \sigma^{(0)} (1 + \delta).$$

With

$$\delta = \frac{2\alpha s}{15\pi} \sum \frac{1}{m_\ell^2} < 3\% \text{ at } s = 1600 \text{ GeV}^2$$

the authors conclude that $N_\nu < 134$. At the large e^+e^- colliders a measurement of the cross-section of the reaction

$$e^+e^- \rightarrow \gamma + \text{nothing}$$

10 GeV above the Z^0 pole can give a sensitivity of $\Delta N_\nu \sim 0.2$.

2. Electron-Neutrino-Electron Scattering

The cross-section of the scattering of antielectron-neutrinos on electrons has been measured by Reines et al.¹⁰⁾. It gives important constraints on the neutral current coupling of the electron, as well as a sensitive test of their generation universality. Both neutral and charged currents contribute to the reaction cross-section. Averaging over the unobserved electron helicities λ and λ'

$$\sigma = \frac{1}{2} \sum_{\lambda, \lambda'} \left| \begin{array}{c} e_{\lambda} \\ \bar{\nu}_e \end{array} \begin{array}{c} W^+ \\ \text{---} \end{array} \begin{array}{c} \bar{\nu}_e \\ e_{\lambda} \end{array} + \begin{array}{c} \bar{\nu}_{\text{out}} \\ \bar{\nu}_e \end{array} \begin{array}{c} W^+ \\ \text{---} \end{array} \begin{array}{c} e_{\lambda'} \\ e_{\lambda} \end{array} \right|^2,$$

where the interference term is given, in the context of the Weinberg-Salam model, by

$$\left[T_L^3(e) + \sin^2 \theta \right] \cong -\frac{1}{2} + 0.23,$$

and should therefore be negative. Table 3 shows the predicted values in terms of σ_{V-A}^{CC} for two energy regions and for three assumed cases, destructive, constructive and no interference, the latter corresponding to a situation in which $\nu_{\text{in}} \perp \nu_{\text{out}}$. The experimental results of Reines et al.¹⁰⁾ favour the destructive interference case but cannot reject, in a compelling way, the other two cases.

It seems, therefore, that a new experiment, using modern technology would be required.

Observation of this interference term would prove that the outgoing and incoming neutrinos are identical; that the neutral current couples to left-handed electrons, solving the ambiguity under exchange of g_V^e and g_A^e in Eq. (1); and that the neutral current is helicity-conserving in the same way as the charged current¹¹⁾, and hence that its space-time structure is given by V and A contributions only.

Table 3. Cross-section for $\bar{\nu}_e e \rightarrow \bar{\nu}_e e$ for three cases

Case	$1.5 \leq T_e \leq 3 \text{ MeV}$	$3.0 \leq T_e \leq 4.5 \text{ MeV}$
Destructive	$0.83 \sigma_{V-A}^{CC}$	$1.2 \sigma_{V-A}^{CC}$
Constructive	2.2	2.8
No interference ($\nu_{\text{in}} \perp \nu_{\text{out}}$)	1.5	2.0
Exp (Reines)	0.87 ± 0.25	1.7 ± 0.44

3. Coupling of the weak neutral electron current

The effective Lagrangian of neutrino-electron interactions and of electron-hadron interactions involves the weak neutral electron current

$$J_{\mu}^e = \bar{e} \gamma_{\mu} (g_V^e + g_A^e \gamma_5) e. \quad (2)$$

It is determined by two coupling constants, g_A^e and g_V^e . The study of neutrino-electron scattering at high energy leaves a four-fold ambiguity, one sign ambiguity and an ambiguity under the exchange of g_V and g_A . Combining these results with those from $\bar{\nu}_e e \rightarrow \bar{\nu}_e e$ scattering¹⁰⁾ and from $e^+e^- \rightarrow \mu^+\mu^-$ at PETRA¹²⁾ a unique solution emerges, the one predicted by the standard model of electroweak interactions, with

$$\begin{aligned} g_A^e &= -0.523 \pm 0.035 \\ \text{and}^{13)} \quad \sin^2\theta &= 0.215 \pm 0.043. \end{aligned}$$

Figure 1 summarizes the present situation. Using the value of g_A^e determined from neutrino-electron scattering one can deduce, from measurements of the angular asymmetry in $e^+e^- \rightarrow f^+f^-$, which is proportional to the product $g_A^e \cdot g_A^f$, the axial vector coupling of the other leptonic neutral currents. Table 4 summarizes these results.

Table 4. Neutral current coupling of the leptonic currents

Current	Process	g_A^f	I_3^R
ee	νe	-0.523 ± 0.035	0.02 ± 0.03
$\mu\mu$	$e\bar{e} \rightarrow \mu\bar{\mu}$	-0.50 ± 0.08	0.01 ± 0.08
$\tau\tau$	$e\bar{e} \rightarrow \tau\bar{\tau}$	-0.39 ± 0.12	-0.11 ± 0.12

The SU(2) structure is clearly confirmed by these measurements, the left-handed leptons transform as doublets and the right-handed leptons as singlets. An independent way of resolving the ambiguities has been achieved before¹⁴⁾, using results of an experiment performed at SLAC on the helicity asymmetry of electron-deuteron scattering and of experiments on neutrino-hadron scattering, under the assumption of factorization.

The overall coefficient ρ^2 , multiplying the theoretical expressions of the cross sections of the neutrino-electron scattering, determines the relative strength of the neutral and of the charged current coupling. In the standard model $\rho = 1$, provided the Higgs fields form an isospin doublet. This simplest version of the theory is already confirmed by the present measurement¹⁵⁾ giving

$$\rho = 1.12 \pm 0.16 .$$

At a higher level of accuracy small deviations of ρ from one are expected as discussed in the introduction. It would therefore be important to determine $|\rho^2 - 1|$ to $\sim \pm 1\%$.

The most precise determination of the value of $\sin^2\theta$ in the leptonic sector has been obtained by the CHARM Collaboration¹⁵⁾, making use of the direct relation between the ratio of $\sigma(\nu_e e)$ and $\sigma(\bar{\nu}_\mu e)$ and $\sin^2\theta$.¹⁶⁾

$$R = \frac{\sigma(\nu_e e)}{\sigma(\bar{\nu}_\mu e)} = 3 \frac{1 - 4\sin^2\theta + (16/3)\sin^4\theta}{1 - 4\sin^2\theta + 16\sin^4\theta} .$$

In the vicinity of $\sin^2\theta = \frac{1}{4}$ this relation gives $\Delta\sin^2\theta \sim \frac{1}{8} \Delta R$ and, hence, a very precise determination of the mixing angle. The detection efficiency cancels in the ratio and many systematic uncertainties are reduced. Using their upgraded detector, incorporating now measurements of the shower barycentre in two orthogonal projections behind each target plate, the CHARM Collaboration will further improve the accuracy of this result. The plans for further measurements at CERN, FNAL and BNL are summarized in Table 5.

Are further, improved measurements of neutrino-electron scattering required? The electroweak theory predicts the Born term ($Q^2 \sim 0$) of the mass of the Z^0 as a function of $\sin^2\theta$

$$M_{Z^0} = \frac{37.4 \text{ GeV}}{\sin\theta\cos\theta}$$

Second order corrections shift M_{Z^0} by 5 GeV¹⁷⁾, e.g. for $\sin^2\theta = 0.220 \pm 0.015$, as determined by the CHARM Collaboration from semi-leptonic neutral current interactions¹⁸⁾. Wheater and Llewellyn-Smith¹⁷⁾ predict

$$M_{Z^0} (\text{Born}) = 90 \pm 2.3 \text{ GeV}$$

$$M_{Z^0} (\text{physical}) = 94.6 \pm 2.3 \text{ GeV}$$

and hence a shift of ~ 5 GeV. A precise experimental determination of this shift would constitute a decisive test of the underlying electroweak gauge theory. A significant test could be claimed if the present error on M would be reduced to 0.7 GeV corresponding to an error on $\sin^2\theta$ of ± 0.005 ¹⁹⁾.

Table 5. Summary of present and future activities on $\nu_\mu e$ scattering

Group	$\sigma(\theta)$ E = 15 GeV	S/B	Events	$\Delta \sin^2 \theta$ (stat)
Flashtube det. FNAL 200 tons	~ 8 mrad	1/1	$\sim 10 \nu_\mu e$	
CHARM 1982 70 tons	~ 12 mrad	1/2	77 $\bar{\nu}_\mu e$ 46 $\nu_\mu e$	± 0.04
CHARM 1983	~ 8 mrad	1/1	$\sim 60 \nu_\mu e$ $\sim 60 \bar{\nu}_\mu e$	Total $\sim \pm 0.02$
BNL 1982	$\frac{12 \text{ mrad}}{\sqrt{E}}$	1/0.25	80 $\nu_\mu e$ 25 $\bar{\nu}_\mu e$	Total ± 0.017
BNL 1982+1983 80 tons			Total 140 $\nu_\mu e$ 90 $\bar{\nu}_\mu e$	

Can this accuracy be achieved in experiments? Llewellyn-Smith²⁰⁾ has discussed the theoretical uncertainties of the quark model picture of the nucleon and its effects on attempts to measure $\sin^2 \theta$ in semi-leptonic neutral current interactions. These uncertainties do not exist in the leptonic reactions of neutrino-electron scattering. Here only the experimental problems have to be solved, namely:

- 1) event rate requiring a large fiducial tonnage and high selection efficiency over a wide window of electron energies;
- 2) background, dominantly due to quasi-elastic electron-neutrino scattering and to coherent π^0 production¹⁵⁾ has to be reduced by efficient e/π discrimination and precise measurements of the shower direction;
- 3) monitoring of the relative flux of the different beam components ν_μ , $\bar{\nu}_\mu$, ν_e and $\bar{\nu}_e$ is required to determine the ratio of $\sigma(\nu_\mu e)/\sigma(\bar{\nu}_\mu e)$.
4. Study of a new detector for neutrino-electron scattering

A tentative design of a new dedicated ($\nu_\mu e$) detector has been discussed by the CHARM Collaboration. It is based on the principle of a fine-grain target-calorimeter and on the accumulated experience

of the CHARM Collaboration with instrumentation of this type. The main concern of this study was the question: can the present technique used for $\nu_\mu e$ studies be sufficiently improved to match the aim of $\Delta \sin^2 \theta = 0.005$ or is a new technique required?

The accuracy of shower direction measurements depends mainly on three contributions

$$\sigma^2(\theta) \sim (\Delta\theta_{\sqrt{N}})^2 + (\Delta\theta_V)^2 + (\Delta\theta_{\text{SMEAR}})^2$$

statistics vertex smearing

- on the sampling frequency and on the method used to count the number of shower particles N ;
- on the plate thickness and grain size of the calorimeter near the vertex;
- on the lateral shower sampling used to determine the barycentre of the shower.

An optimization is required if limited space is available.

The limiting accuracy is given by the Z number of the target material,

$$\sigma(\theta) \sim \frac{Z}{\sqrt{E}} \text{ const.}$$

The mean Z of the present marble target is 13, with MgCO_3 (dolomite) a value of ~ 8 could be achieved. In these light targets with a nuclear absorption length of $\Lambda_{\text{abs}} \sim 4$ radiation lengths, electromagnetic and hadronic showers have very similar longitudinal profiles²¹⁾. The CHARM Collaboration has developed a new method to discriminate between electromagnetic and hadronic showers in low Z materials based on the characteristic difference of their lateral profiles (see figure 2). A discrimination by a factor of ~ 100 has been achieved²¹⁾. Figure 3 shows the resolution in shower direction measurements achieved with the present CHARM detector and with the upgraded version.

The structure of the new, dedicated detector is sketched in figure 4. It consists of 330 modules of $3.5 \times 3.5 \text{ m}^2$ surface area, each composed of a 4 cm thick target plate (marble) and of a plane of streamer tubes with 1 cm wire spacing, read out by crossed cathode strips of 2 cm spacing in two orthogonal projections. Using analog electronics the centroid position of a track or a shower can be reconstructed with $\pm 2 \text{ mm}$ accuracy. Simulating this structure using Monte Carlo methods (EGS) we find an angular resolution of $\sigma(\theta) \sim 16 \text{ mrad}/\sqrt{E/\text{GeV}}$ corresponding to $\sim 4 \text{ mrad}$ at 15 GeV and an energy resolution of $\sigma(E)/E \sim 20\%/\sqrt{E/\text{GeV}}$. Hence, we expect a reduction of background proportional to $\sigma^2(\theta)$ by a factor of ~ 9 .

Figure 3 shows this estimate of $\sigma(\theta)$ compared to the present CHARM I detector. The choices made of analog versus digital readout of strips and of the strip width are illustrated in figures 5a and 5b. This configuration has a fiducial target weight of ~ 320 tons, whereas the CHARM I detector has 70 tons.

Monitoring of the beam composition has been studied in detail for the CHARM I experiments using quasielastic $\nu_\mu(\bar{\nu}_\mu)$ events. This method can be extended to lower neutrino energies by restricting the value of Q^2 to less than 0.01 GeV^2 to equalize the cross sections for the neutrino and the antineutrino induced reactions. Figure 6 shows the measurements for quasielastic reactions²²⁾ at low Q^2 . At $E_\nu > 10.8 \text{ GeV}$ a peak appears at small Q^2 because of the inverse μ decay reaction, $\nu_\mu e \rightarrow \mu^- \nu_e$. The CHARM II detector will have good angular resolution for measurement of the muon direction and, combined with a muon spectrometer, will therefore directly monitor the flux ratio $\phi(\nu_\mu)/\phi(\bar{\nu}_\mu)$ to $\pm 2\%$, required to evaluate the ratio R and the neutrino spectra.

Electron neutrino-electron scattering has a ten time higher cross section than $\nu_\mu e$ scattering. The beam contains approximately 2% electron neutrinos and antineutrinos. Aiming at a 5% measurement of R requires a knowledge of the $(\nu_e + \bar{\nu}_e)$ flux to better than $\pm 10\%$. I think that this can be achieved by making dE/dx measurements in the plane following the vertex, at angles outside the forward peak. The CHARM Collaboration has also developed a method to recognize inclusive $\nu_e N \rightarrow eX$ events with $y < 0.621$).

I summarize this study in Table 6 which gives the selection criteria, the angular resolution, the rate and background for $\bar{\nu}_\mu e$ scattering which would be obtained for 10^{19} protons on target.

Table 6. Rates and background of $\bar{\nu}_\mu e$ scattering in CHARM I and in CHARM II for 10^{19} protons

Selection	CHARM I	CHARM II
Rate for $7.5 < E_e < 30 \text{ GeV}$	154 ± 27	909 ± 33
ϵ_e (1 hit)	61%	90%
$\sigma(\theta)$ at $\bar{E}_e = 15 \text{ GeV}$	12 mrad	4 mrad
Signal/background	1/2	1/0.22
Rate for $2.5 < E_e < 30 \text{ GeV}$		1364 ± 40

The same statistics for $\nu_\mu e$ would be obtained with $5 \cdot 10^{18}$ protons. Hence, we expect to measure

$$R = \frac{N(\nu_\mu e)}{N(\bar{\nu}_\mu e)} \times F = \frac{1364 \pm 40}{1364 \pm 40} (F \pm 2\%) = \bar{R} \pm 5\%,$$

corresponding to $\Delta \sin^2 \theta = \pm 0.005$.

An additional physics result may be obtained on the value of ρ , derived from a measurement of the ratio of cross sections for $\nu_\mu e \rightarrow \nu_\mu e$ and for $\nu_\mu e \rightarrow \mu^- \nu_e$. The flux monitoring is now internal and, apart from systematic uncertainties due to background subtraction, electron detection efficiency and spectrum determination, the statistical error of ρ would be $\pm 2\%$.

The main conclusion that we draw from this study is that the experiment seems to be feasible using the presently developed fine-grain calorimeter technique. Of course, many details have to be investigated further and it is not the aim of this study to be at the level of a proposal.

5. Concluding remarks

With the recent advances in experiments showing helicity-asymmetry in electron-deuteron scattering and forward-backward angular asymmetry in $e^+e^- \rightarrow \mu^+\mu^-$ our understanding of the weak neutral current has progressed.

Neutrino-electron scattering is a field that is expected to progress rapidly, following the demonstration that experiments at high energy, using fine-grain electronic calorimeters, can detect these rare events and separate them reliably from the dominant semileptonic reactions.

Future results at the Z^0 pole and on neutrino-electron scattering together will provide an important test of the electroweak standard model.

Acknowledgements

It is a great pleasure to thank the organizers of the Study Conference on Electroweak Interactions for their kind and warming hospitality and for having created a stimulating atmosphere of discussions.

In preparing these notes I have enjoyed many discussions with my colleagues of the CHARM Collaboration whom I wish to thank sincerely.

References

- 1) F.J. Hasert et al., Phys. Letters 46B (1973) 121.
J. Blietschau et al., Nucl. Phys. B114 (1976) 189.
- 2) For a recent review see, for example, F. Niebergall in Proceedings of the International Conference Neutrino '82, Balatonfüred, Hungary 1982 (Budapest, 1982) Volume II, p.62.
- 3) C.Y. Prescott et al., Phys. Letters 77B (1978) 347 and 84B (1979) 524.
- 4) Report by B. Borgia (CHARM Collaboration), these proceedings.
- 5) Report by R.E. Lanou, these proceedings.
- 6) Report by J. Morfin, these proceedings.
- 7) C. Jarlskog and F.J. Yndurain, Phys. Letters 120B (1983) 361.
- 8) M. Jonker et al., (CHARM Collaboration), Phys. Letters 99B (1981) 265.
- 9) PLUTO Collaboration, Ch. Berger et al., Phys. Letters 99B (1981) 489
MARK-J Collaboration, D.P. Barber et al., Phys. Rev. Letters 45 (1980) 1904
TASSO Collaboration, R. Brandelik et al., Phys. Letters 99B (1981) 163
JADE Collaboration, W. Bartel et al., Phys. Letters 123B (1983) 353.
- 10) F. Reines et al., Phys. Rev. Letters 37 (1976) 315.
- 11) M. Jonker et al., (CHARM Collaboration), CERN-EP/82-207 and Z. f. Physik C. (in print).
- 12) See e.g. A. Böhm, DESY 82-084, December 1982, and M. Davier, LAL 82/93, Oct. 1982.
- 13) F. Bergsma et al., (CHARM Collaboration), Phys. Letters 117B (1982) 272.
- 14) See e.g. review of K. Winter, in Proc. of the 1979 Int. Symposium on Lepton and Photon Interactions at High Energies, (FNAL, Batavia, 1979) p. 258.
- 15) See reference 13).
- 16) A precision measurement of $\sin^2\theta$ by the ratio of $\sigma(\nu_\mu e)/\sigma(\bar{\nu}_\mu e)$ has been discussed by K. Winter in New Aspects of Subnuclear Physics (Plenum Press, New York and London 1980) p.218.
- 17) J. Wheeler, C.H. Llewellyn-Smith, Phys. Letters 105B (1981) 486.
- 18) See reference 8).
- 19) K. Winter, in Workshop on Weak Interactions, Javea (Spain) September 1982.
- 20) Ch. Llewellyn-Smith, contribution to the SPS fixed target Physics Workshop at CERN, Dec. 1982 and Nucl. Phys. (to be published)
J. Panman, Contribution to the SPS Fixed Target Physics Workshop at CERN, Dec. 1982 (to be published).
- 21) A.N. Diddens et al. (CHARM Collaboration) Nucl. Instr. Methods 178 (1980) 27
M. Jonker et al. (CHARM Collaboration) Nucl. Instr. Methods 200 (1982) 183.
- 22) F. Bergsma et al., (CHARM Collaboration), Phys. Letters 122B (1983) 465.

Fig. 1 Results of various leptonic reactions determining g_A^e and g_V^e

Fig. 2 Distribution of the width Γ of electron and pion induced showers as measured by the lateral energy deposition profile in the scintillator planes of the CHARM I detector

Fig. 3 Angular resolution of electron shower direction as a function of electron energy. In the CHARM I detector close and far refer to the projection following the target plate in which the event started (close) and the next plate (far). In the upgraded CHARM I detector additional planes of streamer tubes have been installed and the resolution is now the same in both projections

Fig. 4 Schematic sketch of a new dedicated CHARM II detector

Fig. 5a Shower profile sampling by detectors with analog and with digital measurements

Fig. 5b Resolution of the shower direction as a function of the strip width

Fig. 6 Observed Q^2 dependence of quasielastic $\nu_\mu(\mu^-)$ and $\bar{\nu}_\mu(\mu^+)$ events