Ref.TH.3830-CERN

DETERMINATION OF THE ELEMENTS V_{us} AND V_{ud}
OF THE KOBAYASHI-MASKAWA MATRIX^{*)}

H. Leutwyler⁺⁾
CERN -- Geneva

and

M. Roos
CERN -- Geneva
and
Department of High Energy Physics
University of Helsinki

A B S T R A C T

We discuss the theoretical problems encountered in the standard evaluations of the Cabibbo angle and show that the data on $K_{\ell 3}$ and hyperon decays require $|V_{us}| = 0.221 \pm 0.002$. With this value, unitarity of the Kobayashi-Maskawa matrix predicts $|V_{ud}| = 0.9753 \pm 0.0005$. The ratio of pseudoscalar decay constants is found to be $F_K/F_\pi = 1.22 \pm 0.01$.

^{*)} Work supported in part by Schweizerischer Nationalfonds.

⁺⁾ On leave of absence from Institut für Theoretische Physik, Sidlerstrasse 5, CH-3012 Bern.

1. Introduction

The information on the quark mass matrix is gradually accumulating. For a review of the status of our knowledge about the eigenvalues m_u , m_d , m_s , m_c and m_b see ref. [1]; a review of the information concerning the Kobayashi-Maskawa matrix

$$\begin{pmatrix} V_{ud} & V_{us} & V_{ub} \\ V_{cd} & V_{cs} & V_{cb} \\ V_{td} & V_{ts} & V_{tb} \end{pmatrix} = \begin{pmatrix} c_1 & s_1 c_3 & s_1 s_3 \\ -s_1 c_2 & c_1 c_2 c_3 - s_2 s_3 e^{i\delta} & c_1 c_2 s_3 + s_2 c_3 e^{i\delta} \\ -s_1 s_2 & c_1 s_2 c_3 + c_2 s_3 e^{i\delta} & c_1 s_2 s_3 - c_2 c_3 e^{i\delta} \end{pmatrix}$$

which relates the eigenvectors of the weak charge operator to the eigenvectors of the quark mass matrix is given in ref. [2]. In the present paper we focus on the elements $V_{us} = \sin \theta_1 \cos \theta_3$ and $V_{ud} = \cos \theta_1$ of the K-M matrix. (Our interest in an accurate value for V_{us} stems from chiral perturbation theory. In this framework the ratio F_K/F_π plays an important role. To extract a reliable value for this ratio from data on $K_{\mu 2}$ and $\pi_{\mu 2}$ decays we need to know the ratio V_{us}/V_{ud} rather precisely).

Any determination of the elements of the mixing matrix contains two theoretical ingredients: a calculation of weak vector or axial current matrix elements between hadronic states (QCD) and a calculation of the radiative corrections to the process in question (QED). In the case of V_{ud} and V_{us} the QCD part of the analysis concerns current matrix elements involving the light quark fields u , d and s . In the theoretical limit $m_u = m_d = m_s$ the vector current is conserved and the matrix elements of this current at zero momentum transfer are given by Clebsch-Gordan coefficients. Even in this limit the axial current matrix elements cannot be calculated reliably with presently available methods; the symmetry properties of the theory allow one to express these matrix elements for transitions within say the baryon octet in terms of two constants (F and D coupling, "Cabibbo-fit") which may then be determined phenomenologically.

In the real world, the matrix elements of the vector current differ from the Clebsch-Gordan coefficients mentioned above and the matrix elements of the axial current cannot be described by only two coupling constants. In the case of the vector current matrix elements the Ademollo-Gatto theorem [3] asserts that the corrections are of second order in the symmetry breaking; there is no such suppression of symmetry breaking effects in the case of the axial current. For this

reason, determinations of V_{us} based on transitions to which the axial current makes a significant contribution are contaminated by theoretical uncertainties, unless one determines the size of these contributions directly from experiment and avoids the use of a Cabibbo-fit. We will present an analysis of the hyperon data obtained by the WA2 collaboration [4] in section 5.

In Section 2 we discuss the experimental situation concerning the $K_{\ell 3}$ decay rates [5]. These decays are not plagued by the above-mentioned theoretical uncertainty, because now only the vector current contributes. The QCD part of the theoretical input then reduces to a calculation of the symmetry breaking effects in the vector matrix element, which we will discuss in detail in section 3. We use the results of this calculation in section 4, where we determine the value of $|V_{us}|$ required by the observed rates of the decays $K^+ \rightarrow \pi^0 e^+ \nu$ and $K^0 \rightarrow \pi^- e^+ \nu$.

In section 6 we show that the values of the elements in the first row of the Kobayashi-Maskawa matrix are consistent with unitarity.

Section 7 contains our conclusions.

2. $K_{\ell 3}$ decay

The rates of the four decays

$$K^+ \rightarrow \pi^0 e^+ \nu_e$$

$$K^+ \rightarrow \pi^0 \mu^+ \nu_\mu$$

$$K_L^0 \rightarrow \pi^- e^+ \nu_e$$

$$K_L^0 \rightarrow \pi^- \mu^+ \nu_\mu$$

may be written in the form

$$\Gamma = \frac{G_F^2}{192\pi^3} M_K^5 \cdot |V_{us}|^2 \cdot C^2 \cdot |f_+(0)|^2 \cdot I \cdot (1+\delta)$$

$$I = \frac{1}{M_K^8} \int_{M_\ell^2}^{(M_K - M_\pi)^2} dt \lambda^{\frac{3}{2}} \left(1 + \frac{M_\ell^2}{2t}\right) \left(1 - \frac{M_\ell^2}{t}\right)^2 F(t)$$

$$F(t) = \bar{f}_+(t)^2 + \frac{3M_\ell^2(M_K^2 - M_\pi^2)^2}{(2t + M_\ell^2) \cdot \lambda} \bar{f}_0(t)^2$$

$$\lambda = t^2 + M_K^4 + M_\pi^4 - 2tM_K^2 - 2tM_\pi^2 - 2M_K^2M_\pi^2 \quad (2.1)$$

where $f_+(t)$, $f_0(t)$ are the form factors of the strangeness changing vector current in pure QCD ($e = 0$):

$$\langle K, p' | \bar{u} \gamma_\mu s | \pi, p \rangle = C \{ (p'_\mu + p_\mu) f_+(t) + (p'_\mu - p_\mu) f_-(t) \}$$

$$f_0(t) = f_+(t) + \frac{t}{M_K^2 - M_\pi^2} f_-(t) \quad (2.2)$$

$$\bar{f}_+(t) = f_+(t) / f_+(0)$$

$$\bar{f}_0(t) = f_0(t) / f_0(0)$$

The factor $(1 + \delta)$ stands for the radiative corrections which we normalize by

$$G_F^2 = G_\mu^2 \left\{ 1 + \frac{2\alpha}{\pi} \ln \frac{M_Z}{M_p} \right\} \quad (2.3)$$

where M_p is the mass of the proton and G_μ determines the lifetime of the muon according to the formula [6]

$$\frac{1}{\tau_\mu} = \frac{G_\mu^2}{192\pi^3} M_\mu^5 \left(1 - 8 \frac{M_e^2}{M_\mu^2} \right) \left\{ 1 - \frac{\alpha}{2\pi} \left(\pi^2 - \frac{25}{4} \right) \right\} \quad (2.4)$$

Numerically, using the muon lifetime quoted by the Particle Data Group [7], this becomes

$$G_\mu = 1.16632 \pm 0.00002 \cdot 10^{-11} \text{ MeV}^{-2}$$

$$G_F^2 = 1.021 \cdot G_\mu^2 \quad (2.5)$$

Note that the masses as well as the form factors depend on the channel considered. For the decay $K_L^0 \rightarrow \pi^- \mu^+ \nu$ e.g. the masses M_K, M_π and M_ℓ are to be identified

with M_{K^0} , M_{π^-} and M_{μ} respectively. The four decay channels are described by two different sets of form factors, which we denote more explicitly by $\{f_+^{K^+\pi^0}(t), f_0^{K^+\pi^0}(t)\}$ and $\{f_+^{K^0\pi^-}(t), f_0^{K^0\pi^-}(t)\}$ respectively. The quantity C is the Clebsch-Gordan coefficient referred to in the introduction

$$C^2 = \begin{cases} \frac{1}{2} & K^+ \rightarrow \pi^0 \\ 1 & K^0 \rightarrow \pi^- \end{cases}$$

(In this normalization both $f_+^{K^+\pi^0}(0)$ and $f_+^{K^0\pi^-}(0)$ become equal to 1 if the quark mass differences are neglected.)

The reduced form factors $\bar{f}_+(t)$ and $\bar{f}_0(t)$, which are normalized to 1 at $t=0$, can be measured directly. The data are consistent with a linear t -dependence in the decay region

$$\begin{aligned} \bar{f}_+(t) &= 1 + \lambda_+ t \\ \bar{f}_0(t) &= 1 + \lambda_0 t \end{aligned} \tag{2.6}$$

The overall fit to the K_{e3}^0 data on the slope parameter λ_+ is [7]

$$\lambda_+ = 0.0300 \pm 0.0016 \tag{2.7}$$

in units of M_{π^+} . (The λ_+ value obtained from K_{e3}^+ agrees with this number; sign and order of magnitude are consistent with pole dominance, which predicts $\lambda_+ \simeq (M_{\pi}/M_{K^*})^2 = 0.024$.)

Since the contribution from the scalar form factor is proportional to the square of the lepton mass, the value of λ_0 does not significantly affect the electronic decay rates. Evaluating the phase space integrals with (2.7) one finds

$$\begin{aligned} I_{K_{e3}^+} &= 0.1605 \{ 1 + 3.4 \Delta\lambda_+ \} \\ I_{K_{e3}^0} &= 0.1561 \{ 1 + 3.4 \Delta\lambda_+ \} \end{aligned} \tag{2.8}$$

where $\Delta\lambda_+$ stands for the deviation of λ_+ from 0.03. With the error bar for λ_+ given above the uncertainty in the phase space integrals thus affects the value of V_{us} by 0.3%.

Concerning λ_0 the experimental situation is not clear. The value $\lambda_0 = 0.019 \pm 0.004$ obtained in a high statistics experiment in 1974 [8] confirmed the theoretical expectations. (For $f_0(t)$ to be approximately linear and to pass through the Callan-Treiman prediction $f_0(M_K^2 - M_\pi^2) = F_K/F_\pi$, the value of λ_0 must be close to 0.02; pole dominance with a scalar meson in the range 1 to 1.4 GeV also suggests $\lambda_0 = 0.01$ to 0.02.) More recent measurements of this parameter however disagree with the above value. The muonic phase space integrals are quite sensitive to λ_0 . (If λ_0 is increased from 0.019 to 0.046 [9] the phase space integrals for $K_{\mu 3}^+$ and $K_{\mu 3}^0$ increase by 6%.) For this reason, and also because the values for the $K_{\mu 3}$ branching ratios quoted in ref. [7] are not completely independent from the data on $K_{e 3}$, we do not make use of the experimental information on muonic decays (electron-muon universality of course requires the values of V_{us} extracted from $K_{\mu 3}^+$ and $K_{\mu 3}^0$ to coincide with those found in $K_{e 3}^+$ and $K_{e 3}^0$ respectively.)

The radiative correction δ may be split into two parts, $\delta = \delta_1 + \delta_2$, where δ_1 is sensitive to the lepton kinematics and δ_2 is not. The quantity δ_1 may be identified with the radiative correction in the old, nonrenormalizable V-A model. To split this piece off in a unique manner one needs to specify the cutoff Λ_W in the logarithmic divergence which one encounters in this model. In accordance with the prescriptions [10] used in the analysis of the data, it is convenient to set $\Lambda_W = M_p$. The order of magnitude of δ_1 then is - 2%, - 1/2%, + 1/2% and + 2% for the rates of the decays $K_{e 3}^+$, $K_{\mu 3}^+$, $K_{e 3}^0$ and $K_{\mu 3}^0$ respectively. (The first two numbers were calculated on the basis of Sirlin's prescriptions - the values agree with the results of Becherrawy [10]. The last two are from ref. [11]). The remainder δ_2 does not involve any large logarithms and can therefore not be calculated from first principles at this time. (In the notation of Sirlin [6], δ_2 stands for $\frac{\alpha}{\pi} (\frac{1}{2} \ln \frac{M_p}{M_A} + C)$. The large logarithm which stems from the fact that the Glashow-Salam-Weinberg theory leads to an effective cutoff which is much larger than M_p , viz. $\Lambda_W = M_Z$, is accounted for by the 2.1% difference between G_F^2 and G_μ^2). The quantity δ_2 is expected to be small, of order $\alpha/\pi = 0.2\%$.

Some of the large experiments have applied radiative corrections to their data on the Dalitz plot, according to the prescriptions of Ginsberg [10]. The rates which follow from the semileptonic branching ratios given by the Particle Data Group [7]

$$\Gamma_{K_{e3}^+} = 2.565 \cdot (1 \pm 0.011) \cdot 10^{-15} \text{ MeV} \quad (2.9)$$

$$\Gamma_{K_{e3}^0} = 4.915 \cdot (1 \pm 0.015) \cdot 10^{-15} \text{ MeV}$$

however also involve data which are not corrected for radiative effects. In view of this and in view of the theoretical uncertainties in δ_2 we allow for an overall error due to radiative corrections of

$$\delta = \pm 1\% \quad (2.10)$$

With the values (2.8) of the phase space integrals and the value (2.5) of the Fermi constant the rates quoted above imply

$$K_{e3}^+ : \quad f_+^{K^+\pi^0}(0) \cdot |V_{us}| = 0.2161 \pm 0.0017 \quad (2.11)$$

$$K_{e3}^0 : \quad f_+^{K^0\pi^-}(0) \cdot |V_{us}| = 0.2103 \pm 0.0020 \quad (2.12)$$

where the errors include the uncertainties in δ and in λ_+ .

3. Symmetry breaking in $f_+(0)$

As mentioned in the introduction, any determination of the elements in the Kobayashi-Maskawa matrix requires theoretical input concerning the hadronic matrix elements of the weak currents. In the case of $K_{\ell 3}$ decay theory is needed to pin down the value of the constant $f_+(0)$ (matrix element of the weak charge between K and π at infinite momentum). The matrix element $f_+(0)$ is a function of the renormalization group invariant quark masses and of the renormalization group invariant scale Λ of QCD. We treat the mass term associated with the light quarks as a perturbation and study the expansion of $f_+(0)$ in powers of m_u , m_d and m_s at fixed ratio $m_u : m_d : m_s$

$$f_+(0) = f_0 + f_1 + f_2 + \dots \quad (3.1)$$

where f_0 , f_1 and f_2 are of order 1, m_q and m_q^2 respectively. Consider first the lowest order term f_0 . In the limit $m_u = m_d = m_s = 0$ the weak vector current is

conserved. The weak charge operator therefore only generates transitions between states of the same energy and momentum, i.e. acts within the multiplets of SU(3) (which in the massless theory are degenerate). The pseudoscalar mesons form a degenerate multiplet at $M_\pi = M_K = M_\eta = 0$ (Goldstone bosons of the spontaneously broken chiral symmetry). The matrix element $f_0^{K^0\pi^-}$ is therefore a Clebsch-Gordan coefficient which in our normalization is equal to 1:

$$f_0^{K^0\pi^-} = 1 \quad (3.2)$$

In the case of the matrix element $f_0^{K^+\pi^0}$ there is a complication: in the massless theory there is no physical difference between π^0 and η (the other states of the octet are unambiguously specified by their electric charge and strangeness). To identify the states $|\pi^0\rangle_0$, $|\eta\rangle_0$ in the massless theory we need to look at the perturbation generated by the quark mass term $\bar{q}Mq = m_u\bar{u}u + m_d\bar{d}d + m_s\bar{s}s$. First order perturbation theory shows that the physical states diagonalize the perturbation: ${}_0\langle\eta|\bar{q}Mq|\pi^0\rangle_0 = 0$. In terms of the standard octet basis $|1\rangle$, $|2\rangle$, ..., $|8\rangle$ the physical states of the massless theory are

$$\begin{aligned} |\pi^0\rangle_0 &= \cos \epsilon |3\rangle + \sin \epsilon |8\rangle \\ |\eta\rangle_0 &= -\sin \epsilon |3\rangle + \cos \epsilon |8\rangle \end{aligned} \quad (3.3)$$

$$\tan 2\epsilon = \frac{\sqrt{3}}{2} \frac{m_d - m_u}{m_s - \hat{m}} \quad ; \quad \hat{m} = \frac{1}{2} (m_u + m_d)$$

The crucial point here is that although the $\pi^0\eta$ mixing angle ϵ is proportional to the small mass difference $m_u - m_d$, it counts as an effect of order zero in our expansion, because what matters is not the size of $m_u - m_d$ in comparison to the scale Λ of the theory but the size of $m_u - m_d$ in comparison to the mass difference $m_s - \hat{m}$. It is a simple matter to determine the corresponding isospin breaking effect in $f_0^{K^+\pi^0}$: In the octet basis the matrix elements of f_0 are Clebsch-Gordan coefficients. To first order in ϵ one finds

$$\begin{aligned} f_0^{K^+\pi^0} &= 1 + \sqrt{3} \epsilon \\ &= 1 + \frac{3}{4} \frac{m_d - m_u}{m_s - \hat{m}} \end{aligned} \quad (3.4)$$

The ratio of quark mass differences is known rather accurately [1]:

$$\frac{m_d - m_u}{m_s - \hat{m}} = \frac{1}{43.5 \pm 2.2} \quad (3.5)$$

Inserting this value in (3.4) one finds that the lowest order isospin breaking effect enhances the rate of the decay $K^+ \rightarrow \pi^0 \bar{\ell} \nu$ in comparison to $K^0 \rightarrow \pi \bar{\ell} \nu$ by 3.4%, a contribution which is substantially larger than the radiative corrections.

Before we proceed to discuss the higher order terms in the expansion (3.1) let us consider the commutation relation of the null plane charges

$$[Q^{\bar{u}s}, Q^{\bar{s}u}] = Q^{\bar{u}u - \bar{s}s} \quad (3.6)$$

which implies the sum rule [12]

$$\sum_n |(K^0 | Q^{\bar{u}s} | n)|^2 - \sum_n |(K^0 | Q^{\bar{s}u} | n)|^2 = 1 \quad (3.7)$$

The contribution of the π^- intermediate state to the first term on the l.h.s. of this sum rule is the quantity we are looking for:

$$(K^0 | Q^{\bar{u}s} | \pi^-) = f_+^{K^0 \pi^-}(0) \quad (3.8)$$

All other matrix elements vanish in the limit $m_s = m_u$ and are therefore of order $m_s - m_u$. This implies that the deviation of $f_+^{K^0 \pi^-}(0)$ from 1 is of order $(m_s - m_u)^2$ (Ademollo-Gatto theorem) [3]. Furthermore, the second term on the left hand side of (3.7) involves exotic states with strangeness 2. Neglecting these contributions we obtain

$$|f_+^{K^0 \pi^-}(0)|^2 = 1 - \sum_{n \neq \pi^-} |(K^0 | Q^{\bar{u}s} | n)|^2 \quad (3.9)$$

i.e. $f_+^{K^0 \pi^-}(0)$ is predicted to be smaller than one [12].

Note that the quantity $f_+^{K^+ \pi^0}(0)$ is not protected from receiving corrections of first order in $m_u - m_d$; the sum rule analogous to (3.7), obtained by sandwiching the commutator between K^+ states receives contributions both from π^0

and η intermediate states. Only the sum $\frac{1}{4} |f_+^{K^+\pi^0}|^2 + \frac{3}{4} |f_+^{K^+\eta}|^2$ obeys an Ademollo-Gatto theorem.

We now return to the expansion (3.1). First order perturbation theory in the quark mass term leads to an expression for f_1 which is logarithmically divergent [13]. The divergence stems from the fact that the unperturbed system ($m_u = m_d = m_s = 0$) contains massless physical states (π, K, η). The divergence implies that the quantity $f_+(0)$ does not admit a simple Taylor series expansion in powers of the quark masses. The first order contribution f_1 is not of order m_q , it is of order $m_q \log m_q$. Explicitly, the expression for $f_1^{K^0\pi^-}$ reads [14]

$$f_1^{K^0\pi^-} = H_{K^0\pi} + \frac{1}{2} H_{K^+\pi} + \frac{3}{2} H_{K^+\eta} + \epsilon\sqrt{3} (H_{K\pi} - H_{K\eta})$$

$$H_{12} = - \frac{1}{128\pi^2 F_\pi^2} \left\{ M_1^2 + M_2^2 + \frac{2M_1^2 M_2^2}{M_1^2 - M_2^2} \ln \frac{M_2^2}{M_1^2} \right\} \quad (3.10)$$

where $F_\pi = 93.3$ MeV is the pion decay constant and where we have retained isospin breaking contributions to first order in ϵ . (In the isospin symmetry limit (3.10) reduces to the expression given by Wada [15]). Numerically, the correction (3.10) lowers the value of $f_+^{K^0\pi^-}(0)$ from 1 to 0.977.

It is instructive to confront the above explicit expression for $f_1^{K^0\pi^-}$ with the general properties of $f_+^{K^0\pi^-}(0)$ derived from the charge commutator. We first note that $f_1^{K^0\pi^-}$ is indeed negative, as expected: the function H_{12} is negative for any value of the ratio M_1^2/M_2^2 . In the vicinity of the symmetry limit $M_1^2 = M_2^2$ the function H_{12} is of the form

$$H_{12} = - \frac{1}{384\pi^2 F_\pi^2} \frac{(M_1^2 - M_2^2)^2}{M_1^2} \left\{ 1 + O\left(\frac{M_1^2 - M_2^2}{M_1^2}\right) \right\} \quad (3.11)$$

In agreement with the Ademollo-Gatto theorem, H_{12} is of second order in the symmetry breaking $M_1^2 - M_2^2 \sim m_s - \hat{m}$, however with a coefficient which blows up if the quark masses are sent to zero. This is how f_1 manages to be of first order in the quark masses, despite being proportional to $(m_s - \hat{m})^2$: it picks up a small energy denominator of order m_s which transforms $(m_s - \hat{m})^2$ into $(m_s - \hat{m})^2/m_s$. In fact, this is the reason why the first order contribution to $f_+(0)$ can be calculated

without solving QCD; one only needs to identify the dangerous intermediate states which can give rise to small energy denominators and to calculate the contribution generated by these intermediate states [14].

For the quantity $f_1^{K^+\pi^0}$ an expression analogous to (3.10) can be given. It is convenient to instead consider the ratio $R = f_+^{K^+\pi^0}(0)/f_+^{K^0\pi^-}(0)$. In the isospin symmetry limit $m_u = m_d$, this ratio equals 1. To first order in ϵ it is of the form $R = 1 + \epsilon k + O(\epsilon^2)$ where k may be expanded in powers of $\hat{m}$ and m_s . The relations (3.2) and (3.4) imply that the expansion starts with $R = 1 + \epsilon\sqrt{3} + O(\epsilon m_q)$. Including terms of order ϵm_q the ratio is given by [14]

$$\frac{f_+^{K^+\pi^0}(0)}{f_+^{K^0\pi^-}(0)} = 1 + \sqrt{3} \bar{E} + O(\epsilon m_q^2)$$

$$\bar{E} = \frac{\sqrt{3}}{4} \frac{(M_{K^0}^2 - M_{K^+}^2 - M_{\pi^0}^2 + M_{\pi^+}^2)}{M_K^2 - M_\pi^2} \{ 1 + \Delta_{GMO} + C \}$$

$$\Delta_{GMO} = (4M_K^2 - M_\pi^2 - 3M_{\eta'}^2) / (M_{\eta'}^2 - M_\pi^2)$$

$$C = \frac{1}{64\pi^2 F_\pi^2} \left[4M_K^2 + (M_K^2 - M_\pi^2)^{-1} \{ 9M_{\eta'}^4 \ln \frac{M_K^2}{M_{\eta'}^2} - M_\pi^4 \ln \frac{M_K^2}{M_\pi^2} \} \right] \quad (13.12)$$

Numerically, this becomes

$$\frac{f_+^{K^+\pi^0}(0)}{f_+^{K^0\pi^-}(0)} = 1.022 \quad (13.13)$$

The corrections of order ϵm_q thus increase the zero order isospin breaking effect from 3.4% to 4.4% in the rates.

Finally, we consider the effects of order m_q^2 (including $m_q^2 \log^2 m_q$) described by f_2 . Concerning the ratio $f_+^{K^+\pi^0}/f_+^{K^0\pi^-}$ the above calculation accounts for contributions of order ϵ and of order ϵm_q . There are no other contributions of order ϵm_q to this ratio - the contribution from f_2 is of order ϵm_q^2 . To illustrate this we consider the contribution from $\pi^0 \eta'$ mixing [16]. If the π^0 mixes with the η , do we have to expect a corresponding sizeable contribution from the η' ? There are two reasons why the contribution from $\pi^0 \eta'$ mixing to the matrix

element $f_+^{K^+\pi^0}(0)$ is suppressed. First, the $\pi^0\eta'$ mixing angle is not of order $(m_u - m_d)/(m_s - \hat{m})$, but only of order $(m_u - m_d)/\Lambda$ where Λ is the scale of QCD. (Note that this is an algebraic statement; in this language the mass of the proton, the pion decay constant, and the average transverse momentum of light quarks in a bound state are all of order Λ . It is misleading to identify Λ with the renormalization group invariant scale in numerical estimates: the scale to compare symmetry breaking effects with must be large in comparison with $m_s - \hat{m}$, as there would otherwise be no reason for SU(3) to be an approximate symmetry of the theory.) We should therefore expect the π^0 to mix more with the η than with the η' . The second reason is connected with the fact that in the SU(3) symmetry limit $m_u = m_d = m_s$ the weak charge is conserved. Since the mass of the η' does not coincide with the mass of the K in this limit, the matrix element $\langle K^+ | Q^{\bar{u}s} | \eta' \rangle$ must tend to zero if the quark masses take a common value. This implies that the contribution from $\pi^0\eta'$ mixing to $f_+^{K^+\pi^0}$ is of order $(m_u - m_d) \cdot (m_s - \hat{m}) \sim \epsilon(m_s - \hat{m})^2$. In comparison to the contribution from the η the contribution from the η' is suppressed by two powers of the SU(3) breaking parameter $(m_s - \hat{m})/\Lambda$, confirming the above claim that the presence of the η' can only show up in f_2 .

We expect the contribution from f_2 to the ratio $f_+^{K^+\pi^0}/f_+^{K^0\pi^-}$ to be tiny; the numerical value given in (3.13) should be correct within an error of the order of 2 % (note that the correction from f_1 only amounts to 5 %).

The contribution from f_2 to the quantity $f_+^{K^0\pi^-}(0)$ is not a priori of this irrelevant order of magnitude. On the basis of a model-independent analysis given in the Appendix, we conclude that the contribution of order m_q^2 is expected to slightly decrease the value of $f_+^{K^0\pi^-}(0)$. Our estimate for this contribution is $f_2 = -0.016 \pm 0.008$ (see the Appendix), which leads to our final theoretical number

$$f_+^{K^0\pi^-}(0) = 0.961 \pm 0.008 \quad (3.14)$$

4. V_{us} from K_{e3} decays

With the theoretical value (3.13) for the ratio $f_+^{K^+\pi^0}/f_+^{K^0\pi^-}$ we can convert the experimental information on the semileptonic decay of the charged kaon into information about the quantity $f_+^{K^0\pi^-}|V_{us}|$. Dividing (2.11) by 1.022 we obtain

$$K_{e3}^+ : \quad f_+^{K^0\pi^-}(0) |V_{us}| = 0.2114 \pm 0.0017 \quad (4.1)$$

to be compared with the value

$$K_{e3}^0 : \quad f_+^{K^0\pi^-}(0) |V_{us}| = 0.2103 \pm 0.0020 \quad (4.2)$$

extracted from the decay of the neutral kaon. The fact that the two numbers agree within the errors shows that the data confirm the large isospin violation required by chiral perturbation theory. (In the square of the matrix element the predicted asymmetry is 4.4%. The observed rates correspond to an asymmetry of $5.7 \pm 1.9\%$. Incidentally, the agreement of the two values obtained from K_{e3}^+ and K_{e3}^0 decays respectively is a good check on the radiative corrections which are quite different in the two cases.)

Combining the two independent measurements we get

$$f_+^{K^0\pi^-}(0) |V_{us}| = 0.2110 \pm 0.0013 \quad (4.3)$$

Using the theoretical number (3.14) for $f_+^{K^0\pi^-}(0)$ we finally arrive at

$$|V_{us}| = 0.2196 \pm 0.0023 \quad (4.4)$$

5. Hyperon semileptonic decays

The hyperon β decays have traditionally furnished rather precise information on $|V_{us}|$. With the exception of a small number of enigmatic results, all experiments have agreed remarkably well [17] with the Cabibbo theory [18] without any sign of SU(3)-symmetry breaking.

The recent results of the WA2 experiment [4] at CERN confirm this picture, at the same time as it removes the enigmatic contradictions.

At the level of precision on $|V_{us}|$ now obtainable, it is certainly advisable not to combine different experiments (as in the past [17]), but to treat each experiment consistently, including experimental cuts in the phase space integrals, and applying radiative corrections. This is what the WA2 collaboration has done, and we shall in the following make use of their hyperon semileptonic decay data alone.

Neglecting contributions of order $q = p - p'$ the transition matrix element of the strangeness changing weak current is determined by two constants g_V, g_A :

$$\langle B' | \bar{u} \gamma_\mu (1 - \gamma_5) s | B \rangle = \bar{u}(p') \gamma_\mu (g_V - g_A \gamma_5) u(p) \quad (5.1)$$

The terms neglected contain the momentum dependence of the form factors $g_V(q^2)$, $g_A(q^2)$, the magnetic form factors and terms proportional to q_μ (which have a negligible effect on the rate because their contributions are proportional to m_e^2). The effect of these higher order terms is small; to a good approximation the radiatively corrected rate is given by [4]

$$\Gamma = \frac{G_F^2}{60\pi^3} |V_{us}|^2 (\Delta M)^5 (1 - 3\delta) (g_V^2 + 3g_A^2) \quad (5.2)$$

$$\Delta M = M - M' ; \quad \delta = (M - M') / (M + M')$$

where M, M' are the masses of the parent and daughter baryons respectively.

The WA2 experiment furnishes information on the four $|\Delta S| = 1$ decay rates $\Gamma(\Lambda \rightarrow p e \nu)$, $\Gamma(\Sigma^- \rightarrow n e \nu)$, $\Gamma(\Xi^- \rightarrow \Lambda e \nu)$, $\Gamma(\Xi^- \rightarrow \Sigma^0 e \nu)$, as well as on the ratios $\hat{g}_A = g_A/g_V$ for three of these transitions ($\hat{g}_A^{\Lambda p}$, $\hat{g}_A^{\Sigma^- n}$, $\hat{g}_A^{\Xi^- \Lambda}$).

In ref. [4] the data are analyzed under the hypothesis that the constants g_V, g_A are SU(3)-symmetric (momentum dependence of the form factors, magnetic contributions and radiative corrections are taken into account). Since the vector charges are generators of SU(3), the constants g_V are given by Clebsch-Gordan coefficients. Furthermore, the symmetry fixes the axial constants g_A up to two parameters F and D ("Cabibbo fit"). It also determines the strangeness conserving

transitions $n \rightarrow p e \nu$ and $\Sigma^- \rightarrow \Lambda e \nu$ in terms of the same parameters. The data are consistent with this parametrization, provided $|V_{us}| = 0.231 \pm 0.003$.

Unfortunately, the quality of the theoretical input (exact SU(3)) does not match the quality of the data. The mass difference $m_s - \hat{m}$ generates first order symmetry breaking in g_A ; the relations expressing the constants g_A in terms of F and D may well be off by 20%. Little is known theoretically about the size or even about the sign of these asymmetries [19, 20].

The symmetry breaking in g_V is analogous to the symmetry breaking in $f_+(0)$ discussed in section 3. At leading order in the quark mass expansion there is an isospin breaking effect [20] of order $\epsilon = \frac{1}{4} \sqrt{3} (m_d - m_u)/(m_s - \hat{m}) = 0.010$ in the g_V matrix elements involving Σ^0 or Λ (in the limit $m_u = m_d = m$ there is no physical distinction between these two states). The quantities $g_V^{\Xi^- \Sigma^0 s}$, $g_V^{\Sigma^0 p}$ are larger than the Clebsch-Gordan coefficient by the factor $1 + \sqrt{3} \epsilon = 1.017$ whereas $g_V^{\Lambda p}$ and $g_V^{\Xi^- \Lambda}$ are smaller by the factor $1 - \epsilon/\sqrt{3} = 0.994$. Higher order effects presumably amplify these asymmetries somewhat (in the case of $f_+(0)$ the higher order contributions increase the factor $1 + \sqrt{3} \epsilon$ from 1.017 to 1.022).

The asymmetries in g_V due to SU(3) breaking are suppressed by the Ademollo-Gatto theorem. The asymmetry is of order $(m_s - \hat{m})^2$, however with a coefficient which blows up in the chiral limit. As was shown by Langacker and Pagels [13] the infrared singularities in the baryon matrix elements are worse than in the meson matrix elements: square root singularities rather than logarithms. A formal expansion in powers of square roots of quark masses does not lead to meaningful results. A coherent method to deal with the problem is described in Appendix C of ref. [1]. The method has however not been applied to a calculation of g_V .

The absence of exotic states in the sum rule obtained from the expectation value of the charge commutator in the neutron implies that the physical value of $g_V^{\Sigma^- n}$ is smaller than the Clebsch-Gordan coefficient. There is no such statement for the constants $g_V^{\Lambda p}$, $g_V^{\Xi^- \Lambda}$ and $g_V^{\Xi^- \Sigma^0}$ which may be either smaller or larger than in the SU(3) limit. More theoretical work is needed to pin these asymmetries down.

In the following we take the symmetry breaking in g_A into account, but neglect the asymmetries in g_V which we expect to be substantially smaller. To obtain information which does not rely on the assumption that the g_A values are SU(3)-symmetric we use the experimental values of $\hat{g}_A = g_A/g_V$ measured in the

Dalitz plot distribution [4]:

$$\begin{aligned}
 \hat{g}_A^{\Lambda p} &= +0.70 \pm 0.03 \\
 \hat{g}_A^{\Sigma^- n} &= -0.34 \pm 0.05 \\
 \hat{g}_A^{\Xi^- \Lambda} &= +0.25 \pm 0.05
 \end{aligned} \tag{5.3}$$

According to the approximate formula (5.2) the rate is proportional to

$$\gamma = |V_{us}|^2 (1 + 3 \hat{g}_A^2) \tag{5.4}$$

To get rid of the kinematical factors we convert the measured rates [4] into information about the factor γ (using the SU(3) values for g_V) with the result

$$\begin{aligned}
 \gamma^{\Lambda p} &= (13.1 \pm 0.57) \cdot 10^{-2} \\
 \gamma^{\Sigma^- n} &= (6.7 \pm 0.37) \cdot 10^{-2} \\
 \gamma^{\Xi^- \Lambda} &= (6.7 \pm 0.45) \cdot 10^{-2}
 \end{aligned} \tag{5.5}$$

With the experimental values (5.3) we thus obtain three independent measurements of V_{us} :

$$|V_{us}| = \begin{cases} 0.230 \pm 0.008 & \Lambda \rightarrow p e \nu \\ 0.223 \pm 0.010 & \Sigma^- \rightarrow n e \nu \\ 0.238 \pm 0.011 & \Xi^- \rightarrow \Lambda e \nu \end{cases} \tag{5.6}$$

As pointed out above the physical value of $g_V^{\Sigma^- n}$ is smaller than the Clebsch-Gordan coefficient used in this determination of $|V_{us}|$. Symmetry breaking effects will therefore increase the lowest one of the three values for $|V_{us}|$ somewhat, whereas the other two may be shifted up or down.

A simultaneous fit to all three decays which accounts for the corrections to the approximate formula (5.2) leads to

$$|V_{us}| = 0.231 \pm 0.005 \quad (5.7)$$

with $\chi^2 = 1$ for 2 degrees of freedom. The mean value thus remains the same as for the "Cabibbo fit", but the error bar is larger, because the fit now involves more parameters (the values of the constants g_A are treated as unknowns, not as given in terms of the two parameters F and D, constrained to also fit the strangeness conserving decays.)

If one combines the hyperon data with the value (4.4) obtained from K_{e3} decays one obtains

$$|V_{us}| = 0.221 \pm 0.002 \quad (5.8)$$

with a χ^2 of 4.2 for 3 degrees of freedom. The fit is still acceptable, but the g_A values are forced to deviate quite substantially from the values predicted by the "Cabibbo fit". (For detailed information on the values of the axial coupling constants obtained in this manner, we refer the reader to a forthcoming paper by Gaillard and Sauvage.)

On general grounds, sizeable asymmetries in the axial coupling constants are to be expected (compare the observed deviations from the SU(3) predictions for the magnetic moments); both the sign and the magnitude of the deviations from SU(3) however remain to be understood.

6. Unitarity

If there are only three generations of quarks (more generally, if the weak transitions between the first generation and the fourth, fifth, ... generations are strongly suppressed) unitarity of the Kobayashi-Maskawa matrix requires

$$|V_{ud}|^2 + |V_{us}|^2 + |V_{ub}|^2 = 1 \quad (6.1)$$

The recent B lifetime measurements [21], together with the observation that the b quark preferentially decays into c leads to [2, 22]

$$|V_{ub}| < 0.008 \quad (6.2)$$

This bound implies that the contribution from V_{ub} in (6.1) can safely be neglected in comparison with the accuracy to which V_{ud} can be determined. A measurement of the quantity $|V_{ud}|$ thus provides us with a measurement of $|V_{us}|$

$$|V_{us}| = (1 - |V_{ud}|^2)^{\frac{1}{2}} \quad (6.3)$$

and vice versa. Recently, Wilkinson [23] has reanalyzed the determination of the quantity $G_F |V_{ud}|$ from superallowed Fermi transitions. As a "tentatively final value" he gives

$$\begin{aligned} G_F^R |V_{ud}| &= (1.41232 \pm 0.00046) \cdot 10^{-49} \text{ erg cm}^3 \\ &= (1.14723 \pm 0.00037) \cdot 10^{-11} \text{ MeV}^{-2} \end{aligned} \quad (6.4)$$

In the notation used above the quantity G_F^R is related to the constant G_F defined in (2.3) by

$$(G_F^R)^2 = G_F^2 (1 + \delta_2) \quad (6.5)$$

where δ_2 is the piece of the radiative correction which is independent of the decay kinematics and does not contain any large logarithms. (In Sirlin's notation, $\delta_2 = \frac{\alpha}{\pi} (\frac{1}{2} \ln \frac{M_p}{M_A} + C)$).

Several rule of thumb estimates for δ_2 may be found in the literature. We quote the value for V_{ud} given in ref. [4]

$$|V_{ud}| = 0.9735 \pm 0.0015 \quad (6.6)$$

which is based on Wilkinson's new Ft -value (with the estimate $\delta_2 \simeq \alpha/\pi \simeq 0.2\%$ added linearly to the experimental error, scaled up by $S = 1.8$). With (6.6) the unitarity relation (6.3) then implies

$$|V_{us}| = 0.229 \pm 0.006 \quad (6.7)$$

The error bar is very sensitive to the estimate used for δ_2 . Shrock and Wang [17] and Paschos and Türke [17] use a more conservative estimate for these theoretical uncertainties and hence end up with a larger error bar for V_{ud} (± 0.0025). The unitarity prediction for V_{us} based on their numbers is 0.229 ± 0.011 .

7. Summary and conclusion

(i) Determinations of V_{ud} based on K_{e3} decays used to give two different results, depending on whether the decay rate of the K^+ or of the K^0 was used. (Shrock and Wang [17] e.g. obtain $|V_{us}| = 0.221 \pm 0.003$ and $|V_{us}| = 0.212 \pm 0.005$ from K_{e3}^+ and K_{e3}^0 respectively.) The source of this discrepancy was identified in ref. [14]: the quark mass difference $m_u - m_d$ generates substantial isospin breaking in the form factors ($f_+^{K^+\pi^0}(0) > f_+^{K^+\pi^-}(0)$). The asymmetry can be calculated in a parameter free manner up to and including contributions of order ϵm_q . The correction brings the two experimental values for $f_+(0)|V_{us}|$ into good agreement, at the 1% level of accuracy.

(ii) To extract the matrix element $|V_{us}|$ from these data one needs a theoretical value for $f_+(0)$. We have analyzed the symmetry breaking effects due to $m_s - \bar{m}$. The infrared singular contributions of order $(m_s - \bar{m})^2/m_s$ can be calculated in a parameter free manner [13, 14, 15]; they decrease the value of $f_+^{K^0\pi^-}(0)$ by 2.3% from the symmetry limit $f_+^{K^0\pi^-} = 1$. The absence of exotic states in the sum rule obtained from the commutation relation of the strangeness changing weak charge implies that the remaining, infrared stable, contributions decrease the value of $f_+^{K^0\pi^-}(0)$ further. We have estimated the size of these residual effects at $1.6 \pm 0.8\%$ which leads to $f_+^{K^0\pi^-}(0) = 0.961 \pm 0.008$. The value of $|V_{us}|$ which follows from the K_{e3} data with this theoretical input is

$$|V_{us}| = 0.220 \pm 0.002 \quad (7.1)$$

(iii) The value of the quantity $|V_{us}|$ extracted from hyperon decays is traditionally somewhat higher. The analysis of the data is complicated by the fact that in this case the axial vector current also contributes to the transition. In contrast to the vector current matrix elements g_V , in which the asymmetries are suppressed by a nonrenormalization theorem, the symmetry breaking in the axial coupling constant g_A is of first order in the quark mass difference

$m_s - \bar{m}$ and may well be sizeable. We have reanalyzed the data of the WA2 collaboration by exploiting the experimental information on the ratio g_A/g_V derived from the Dalitz plot distribution. Neglecting symmetry breaking effects in g_V we obtain $|V_{us}| = 0.231 \pm 0.005$. The difference with the K_{e3} value (7.1) persists, a simultaneous fit of the two sets of data however shows that the difference is not statistically significant. The fit leads to

$$|V_{us}| = 0.221 \pm 0.002 \quad (7.2)$$

with $\chi^2 = 4.2$ for 3 degrees of freedom. (The difference may also be due to asymmetries in the vector coupling constants g_V which could be of the same size as in $f_+(0)$, but need not have the same sign.)

(iv) Since the matrix element V_{ub} is known to be very small, unitarity of the three generation Kobayashi-Maskawa matrix requires $|V_{ud}|^2 + |V_{us}|^2 = 1$ to a good approximation. With the value (7.2) we therefore predict

$$|V_{ud}| = 0.9753 \pm 0.0005 \quad (7.3)$$

to be compared with the experimental value 0.9735 ± 0.0015 [4, 23]. We conclude that the value of V_{us} given above is consistent with the unitarity condition for three generations.

(v) It is interesting to compare the elements of the mixing matrix with the ratio of quark masses [1]

$$\frac{m_s}{m_d} = 19.6 \pm 1.6 \quad (7.4)$$

The values are consistent with the familiar formula [24, 25]

$$m_s |V_{us}|^2 = m_d |V_{ud}|^2 \quad (7.5)$$

(With (7.2) and (7.3) this relation predicts $m_s/m_d = 19.5 \pm 0.4$.)

(vi) Finally, we determine the ratio of decay constants F_K/F_π . The ratio of $K_{\mu 2}$ to $\pi_{\mu 2}$ decay rates is given by

$$\frac{\Gamma_{K \rightarrow \mu\nu}}{\Gamma_{\pi \rightarrow \mu\nu}} = \frac{|V_{us}|^2}{|V_{ud}|^2} \cdot \frac{F_K^2}{F_\pi^2} \cdot \frac{M_{K^+}}{M_{\pi^+}} \cdot \frac{\{1 - (M_\mu/M_{K^+})^2\}^2}{\{1 - (M_\mu/M_{\pi^+})^2\}^2} \cdot (1 + \delta_K - \delta_\pi) \quad (7.6)$$

We have calculated the radiative corrections δ_K , δ_π on the basis of the formulae given by Sirlin [26] with the result $\delta_K - \delta_\pi = -0.006$. Using the rates given by the Particle Data Group [7] this leads to

$$\frac{|V_{us}|}{|V_{ud}|} \frac{F_K}{F_\pi} = 0.2758 \pm 0.0009 \quad (7.7)$$

Inserting the values (7.1) or (7.2) and (7.3) for V_{us} , V_{ud} we obtain

$$\frac{F_K}{F_\pi} = 1.22 \pm 0.01 \quad (7.8)$$

Acknowledgements

We thank the WA2 group, and in particular D. Froidevaux, J.-M. Gaillard and G. Sauvage for performing the analysis described in section 5 with their data. We profited from discussions with G. Karl and J. Trampetić concerning symmetry breaking effects in constituent models, and we thank J. Gasser for help in our struggle with the monstrosities of chiral perturbation theory. One of us (M.R.) is indebted to INFN, Sezione di Roma, Rome, Italy, and in particular to L. Maiani for hospitality extended to him in Rome, where part of this analysis was done. We also thank P. Schlein for letting us use his computing facilities.

Appendix: Estimates of second order effects in $f_+(0)$

In the language of the sum rule (3.9) the contribution f_1 generated by the infrared logarithms of chiral perturbation theory [given in (3.10)] is due to the threshold region of three-meson intermediate states. States containing more than three Goldstone bosons or massive states such as A_1 or π' contribute at order $(m_q)^2$. To estimate the order of magnitude of these residual, infrared stable contributions we describe the pion and the kaon in terms of constituent wave functions and relate the quantity $f_+(0)$ to an overlap integral of these wave functions. Since the masses of the pseudoscalars are very small and very different, it does not make sense to analyze the symmetry breaking effects in terms of rest frame wave functions (there is no rest frame in the chiral limit). Instead we use the infinite momentum frame [27] which has the further advantage that the matrix elements of the charge operator carry zero invariant momentum transfer t , i.e. measure the form factor $f_+(t)$ at $t = 0$. (The charge operator conserves three-momentum. In the π to K matrix element the charge operator connects a state of energy $E = (M_\pi^2 + \vec{p}^2)^{1/2}$ to a state of energy $E' = (M_K^2 + \vec{p}^2)^{1/2}$. The invariant momentum transfer is $t = (E' - E)^2$. In the rest frame, t is equal to $(M_K - M_\pi)^2 \neq 0$; for large momenta, t is of order $(M_K^2 - M_\pi^2)^2 / \vec{p}^2$ and hence vanishes in the infinite momentum limit.)

It is somewhat cumbersome to look at the system at time $x^0 = 0$ and then to take the infinite momentum limit of the relevant matrix elements. It is more economic to instead work at finite momentum and to look at the system at time $x^0 = x^3$ (null plane) [28]. In this language the three kinematical components of the momentum of a state are $p_\mu = (p_1, p_2, p_+)$ where p_+ stands for $\frac{1}{2}(p_0 + p_3)$. The normalization of a covariant one particle state is independent of its mass:

$$\langle p' | p \rangle = (2\pi)^3 \cdot 2p_+ \cdot \delta^3(p' - p) \quad (A.1)$$

We describe the bound states of d and $\bar{s}$ in terms of a set of basis vectors labelled by the three-momenta of the two constituents:

$$|K^0, p\rangle = (2\pi)^{3/2} \sqrt{2} \int d^3p^d d^3p^s \delta^3(p - p^d - p^s) \cdot a_d^\dagger(p^d) b_s^\dagger(p^s) |0\rangle \cdot \phi_K(p^d, p^s) \quad (A.2)$$

The quantity ϕ_K is the (null-plane) wave function of the kaon [29]. For simplicity we disregard the spin of the quarks (the following analysis carries through without any essential modification if spin is taken into account [28]).

Except for colour the constituent basis vectors carry the same quantum numbers as a $q\bar{q}$ state in the Fock space of QCD. We are however not talking about this sector of Fock space. The sectors containing additional gluons and quark pairs are equally important for an adequate description of the kaon within QCD. To discuss constituent wave functions does not mean to suppose that these higher sectors of Fock space can be neglected. What matters is that the main properties of the bound states, in particular, the symmetry properties of current matrix elements can be understood in terms of the degrees of freedom of the two quarks which drag these gluons and pairs along.

Normalizing the basis vectors by

$$\{a_q(p), a_{q'}^\dagger(p')\} = \delta^3(p-p') \quad (A.3)$$

the probability for finding a d-quark with momentum in the interval d^3p^d becomes

$$dp_K = \frac{d^3p^d}{p_+} [\phi_K(p^d, p^s)]^2 \quad (A.4)$$

(Time reversal invariance implies that in a suitable phase convention the wave function is real.)

The transformation properties under Lorentz transformations which leave the plane $x^0 = x^3$ invariant imply that the wave function depends on the momenta p^d , p^s of the two constituents only through the combinations

$$\begin{aligned} k_T &= (p_+^d p_-^s - p_+^s p_-^d) / (p_+^s + p_+^d) \\ k_L &= (p_+^s - p_+^d) / 2(p_+^s + p_+^d) \end{aligned} \quad (A.5)$$

with $k_T = (k_1, k_2)$. This is analogous to the fact that the fixed time wave functions only depend on the relative momentum $\vec{k}$ of the constituents; in fact, rotational invariance implies that the fixed time wave function of a pseudoscalar bound state only depends on the single variable $\vec{k}^2$. The null plane wave function

instead depends on two variables $0 \leq k_T^2 < \infty$ and $-\frac{1}{2} \leq k_L \leq \frac{1}{2}$, in terms of which the probability distribution reads

$$dp_K = d^2k_T dk_L \varphi_K(k_T^2, k_L)^2; \int dp_K = 1 \quad (A.6)$$

In the nonrelativistic limit (m_d, m_s heavy) the null plane description reduces to the standard framework: the three components of the nonrelativistic momentum vector are given by [28]

$$\vec{k} = \{k_1, k_2, m_s(\frac{1}{2} - k_L) - m_d(\frac{1}{2} + k_L)\} \quad (A.7)$$

With these preliminaries out of the way we can now discuss the matrix elements of the strangeness changing charge operator

$$Q = \int_{x^0=x^3} d\sigma_\mu \bar{u} \gamma^\mu s \quad (A.8)$$

between K^0 and π^- :

$$\langle K^0, p' | Q | \pi^-, p \rangle = (2\pi)^3 \cdot 2p_+ \delta^3(p' - p) f_+(0) \quad (A.9)$$

The discussion is based on the assumption that the constituents transform in a simple manner under this charge:

$$[Q, b_u^+(p)] = -b_s^+(p) \quad (A.10)$$

This implies that the quantity $f_+(0)$ is given by the overlap integral

$$f_+(0) = \int d^2k_T dk_L \varphi_K(k_T^2, k_L) \varphi_\pi(k_T^2, k_L) \quad (A.11)$$

Clearly, if the pion and kaon wave functions coincide, we obtain $f_+(0) = 1$. Defining the function $\delta(k_T^2, k_L)$ by

$$\varphi_K = (1 + \delta) \varphi_\pi \quad (A.12)$$

we have

$$\int d^2 k_T dk_L \delta^2 \phi_\pi^2 = 2 \{1 - f_+(0)\} \quad (\text{A.13})$$

which shows that the symmetry breaking lowers the value of $f_+(0)$ by an amount which is of second order in the symmetry breaking:

$$f_+(0) = 1 - \frac{1}{2} \langle \delta^2 \rangle_\pi \quad (\text{A.14})$$

In this relation the symbol $\langle F \rangle_\pi$ stands for the mean value of the function F in the probability distribution associated with the pion wave function:

$$\langle F \rangle_\pi = \int dp_\pi F \quad (\text{A.15})$$

We parametrize the asymmetry δ by three constants:

$$\delta(k_T^2, k_L) = a + b k_T^2 + c k_L \quad (\text{A.16})$$

The constant b accounts for a difference in the distribution of the transverse momentum (for $b > 0$ the mean k_T^2 is larger in the K than in the π). The constant c generates an asymmetry in the longitudinal momentum distribution of the K (we disregard isospin breaking and hence assume that ϕ_π is symmetric with respect to the operation $k_L \rightarrow -k_L$ which interchanges the two quarks.)

To first order in δ the normalization $\int dp_\pi = \int dp_K = 1$ requires

$$\langle \delta \rangle_\pi = 0$$

This determines the constant a :

$$a = -b \langle k_T^2 \rangle_\pi \quad (\text{A.17})$$

Next we compare the mean values of k_T^2 in the two states. The expectation value of k_T^2 in the K wave function is given by

$$\langle k_T^2 \rangle_K = \int dp_K k_T^2 = \langle (1 + 2\delta) k_T^2 \rangle_\pi$$

Since the term proportional to c drops out by virtue of the symmetry of the pion probability distribution with respect to $k_L \rightarrow -k_L$ we obtain

$$\langle k_T^2 \rangle_K = \langle k_T^2 \rangle_\pi + 2b\sigma_T \quad (A.18)$$

where σ_T is the dispersion in the transverse momentum distribution

$$\sigma_T^2 = \langle (k_T^2 - \langle k_T^2 \rangle_\pi)^2 \rangle_\pi \quad (A.19)$$

Analogously, one shows that the mean value of k_L in the K is determined by the constant c:

$$\langle k_L \rangle_K = 2c \langle k_L^2 \rangle_\pi \quad (A.20)$$

(note that the symmetry of the pion wave function implies $\langle k_L \rangle_\pi = 0$).

It is convenient to introduce two dimensionless parameters ϵ_T, ϵ_L which measure the asymmetries of the wave functions:

$$\begin{aligned} \epsilon_T &= \frac{1}{\sigma_T} \{ \langle k_T^2 \rangle_K - \langle k_T^2 \rangle_\pi \} \\ \epsilon_L &= \frac{1}{\langle k_L^2 \rangle_\pi} \{ \langle k_L \rangle_K - \langle k_L \rangle_\pi \} \end{aligned} \quad (A.21)$$

The function δ may be expressed in terms of these parameters as

$$\delta(k_T^2, k_L) = \frac{1}{2} \epsilon_T \frac{k_T^2 - \langle k_T^2 \rangle_\pi}{\sigma_T} + \frac{1}{2} \epsilon_L \frac{k_L}{\langle k_L^2 \rangle_\pi} \quad (A.22)$$

Inserting this result in the expression (A.14) for $f_+(0)$ one obtains the simple formula

$$f_+(0) = 1 - \frac{1}{8} (\epsilon_T^2 + \epsilon_L^2) \quad (A.23)$$

expressing the deviation of $f_+(0)$ from 1 in terms of the asymmetries of the wave functions.

What remains to be done is to estimate the asymmetry parameters ϵ_T, ϵ_L . In order for the wave functions to be approximately SU(3) symmetric, we must have $\epsilon_T, \epsilon_L \ll 1$. To quantify this condition we first note that the symmetry breaking

effects in $f_+(0)$ due to the presence of infrared singularities in chiral perturbation theory are accounted for in the value $f_+(0) = 0.977$ quoted in section 3. Here we are discussing the remaining, infrared stable contributions of second order in $m_s - \hat{m}$. Recent work on chiral symmetry breaking [14] confirms that the asymmetries generated by the mass difference $m_s - \hat{m}$ in infrared stable quantities are small. We assume that the residual asymmetries in the probability distributions $dp_K/dp_\pi \simeq 1 + 2\delta$ do not exceed 40% in the central region where the probabilities are not small. (More specifically, we demand $2|\delta| < 0.4$ for $k_T = 0$, $k_L^2 = \langle k_L^2 \rangle$ and assume that the transverse momentum distribution is approximately Gaussian: $\sigma_T \simeq \langle k_T^2 \rangle$). Expressed in terms of the asymmetry parameters $\varepsilon_T, \varepsilon_L$ this condition implies

$$|\varepsilon_T| + |\varepsilon_L| < 0.4 \quad (\text{A.24})$$

Inserting this constraint in (A.23) one finds that the asymmetries lower the value of $f_+(0)$ by at most 2% (the upper limit is reached for $\varepsilon_T = 0.4$, $\varepsilon_L = 0$ or vice versa.) The main point here is that even considerable wave function asymmetries only produce a very small effect in $f_+(0)$. A more realistic estimate is $\varepsilon_T \simeq \varepsilon_L \simeq 0.2$ in which case we obtain a 1% contribution to $f_+(0)$. On the basis of these considerations we estimate the size of the second order contributions to $f_+(0)$ at

$$f_2 = -0.016 \pm 0.008 \quad (\text{A.25})$$

where the error bar is chosen such that unexpectedly large wave function asymmetries like ($\varepsilon_T = 0.4$, $\varepsilon_L = 0.3$) and unexpectedly small asymmetries ($\varepsilon_T \simeq \varepsilon_L \simeq 0$) amount to two standard deviations from the mean value.

To see whether these estimates are consistent with the observed difference between F_K and F_π we assume that the decay constants are proportional to the null plane wave function at the origin [30]

$$\begin{aligned} F_\pi &= N \int d^3k_T dk_L \phi_\pi \\ F_K &= N \int d^3k_T dk_L \phi_K \end{aligned} \quad (\text{A.26})$$

Inserting the above representation for ϕ_K this leads to

$$\frac{F_K}{F_\pi} = 1 + \frac{1}{2} \kappa \epsilon_T \quad (A.27)$$

$$\kappa = \frac{1}{\sigma_T} \int d^2 k_T dk_L (k_T^2 - \langle k_T^2 \rangle) \varphi_\pi / \int d^2 k_T dk_L \varphi_\pi$$

The value of κ depends on the shape of the transverse momentum distribution. For a Gaussian distribution we have $\kappa = 1$. With $\epsilon_T = 0.2$ and $\epsilon_T = 0.4$ we therefore get $F_K/F_\pi = 1.1$ and 1.2 respectively. Note that the ratio F_K/F_π also receives a nonanalytic contribution from long range phenomena which must be treated separately, like the corresponding contribution to $f_+(0)$. The size of the nonanalytic piece depends on the scale μ at which one normalizes the chiral logarithms. For $\mu = 1$ GeV the correction due to the infrared singularities alone amounts to $F_K/F_\pi = 1.13$ [14]. We conclude that the above estimates of the asymmetries in the wave functions are consistent with the observed asymmetry in the decay constants ($F_K/F_\pi \simeq 1.2$).

An independent, very crude estimate for $f_+(0)$ may be obtained from the model discussed in ref. [30]. Taking the "Melosh angles" $\beta_u = \beta_d$, β_s introduced in that paper to be independent of k_T^2 and k_L one finds

$$F_\pi / F_\rho = \cos 2\beta_u \quad ; \quad F_K / F_\rho = \cos (\beta_u + \beta_s)$$

$$f_+(0) = \cos (\beta_u - \beta_s)$$

The observed values $F_\rho/F_\pi \simeq 1.5$, $F_K/F_\pi \simeq 1.2$ allow one to determine the angles: $\beta_u \simeq 0.42$, $\beta_s \simeq 0.22$. Inserting this result in the expression for $f_+(0)$ one obtains $f_+(0) \simeq 0.98$. This simple model confirms that sizeable asymmetries in the wave functions, measured here in terms of a different relative weight of the S and P wave components of the wave function, indeed have a small effect on $f_+(0)$.

The quantity $f_+(0)$ may also be calculated within specific models such as the nonrelativistic oscillator model or the bag model [31]. The numbers one finds in general confirm the model independent estimate given above - the value of the deviation $1 - f_+(0)$ however depends on the model. It is notoriously difficult within these models to account for $M_\pi \ll M_K$ in a coherent manner; the problem manifests itself e.g. in the fact that some of these model calculations fail to produce the proper sign ($1 - f_+(0) > 0$).

References

- 1 J. Gasser and H. Leutwyler, Phys.Rep. 87 (1982) 77.
- 2 L.-L. Chau, Phys.Rep. 95 (1983) 1;
K. Kleinknecht and B. Renk, Z.Phys. C - Particles and Fields 20 (1983) 71;
R. Rückl, "Weak decays of heavy flavours", to be published in Phys.Rep.
- 3 R.E. Behrends and A. Sirlin, Phys.Rev.Lett. 4 (1960) 186;
M. Ademollo and R. Gatto, Phys.Rev.Lett. 13 (1964) 264.
- 4 M. Bourquin et al., Z.Phys. C - Particles and Fields 12 (1982) 307;
21 (1983) 1, 17, 27.
- 5 For an excellent review see L.M. Chounet, J.M. Gaillard and M.K. Gaillard,
Phys.Rep. 4C (1972) 199.
- 6 A. Sirlin, Nucl.Phys. B196 (1982) 83; B71 (1974) 29; Phys.Rev. D22 (1980)
971; D5 (1972) 436; 164 (1967) 1767; Rev.Mod.Phys. 50 (1978) 573;
W.J. Marciano and A. Sirlin, Phys.Rev. D22 (1980) 2695; Phys.Rev.Lett. 36
(1976) 1425;
- 7 Review of particle properties, Phys.Lett. 111B (1982).
- 8 G. Donaldson et al., Phys.Rev. D9 (1974) 2960.
- 9 Y. Cho et al., Phys.Rev. D22 (1980) 2688.
- 10 E.S. Ginsberg, Phys.Rev. D1 (1970) 229; D4 (1971) 2893;
E.S. Ginsberg and J. Smith, Phys.Rev. D8 (1973) 3887.
T. Becherrawy, Phys.Rev. D1 (1970) 1452.
- 11 H.H. Williams et al., Phys.Rev.Lett. 33 (1974) 240.
- 12 The pioneering paper on this sum rule is G. Furlan, F.G. Lannoy,
C. Rossetti and G. Segrè, Nuovo Cim. 38 (1965) 1747.
- 13 P. Langacker and H. Pagels, Phys.Rev.Lett. 30 (1973) 630;
M. Bačé and D.T. Cornwell, Phys.Rev. D10 (1974) 2694;
for a review see H. Pagels, Phys.Rep. 16C (1975) 219.
- 14 J. Gasser and H. Leutwyler, CERN TH.3798, 3828, 3829, to be published.
- 15 S. Wada, Phys.Lett. 49B (1974) 175
- 16 N. Isgur, H.R. Rubinstein, A. Schwimmer and H. Lipkin, Phys.Lett. 89B
(1979) 79.

- 17 M. Roos, Nucl.Phys. B77 (1974) 420;
R.E. Shrock and L.L. Wang, Phys.Rev.Lett. 41 (1978) 1692;
E.A. Paschos and U. Türke, Phys.Lett. 116B (1982) 360.
- 18 N. Cabibbo, Phys.Rev.Lett. 10 (1963) 531.
- 19 See e.g. A. Garcia, Phys.Rev. D3 (1971) 2638;
B.H. Kellet, Phys.Rev. D10 (1974) 2269;
T. de Grand, R.L. Jaffe, K. Johnson and I. Kiskis, Phys.Rev. D12 (1975) 2060;
J.F. Donoghue and B.R. Holstein, Phys.Rev. D25 (1982) 206;
J. Trampetić, "Fermi motions and static properties of baryons", preprint Max-Planck-Institut, MPI-PAE/PTH 60/83, München (1983).
- 20 P. Bracken, A. Frenkel and G. Karl, Phys.Rev. D24 (1981) 2984.
- 21 E. Fernandez et al., Phys.Rev.Lett. 51 (1983) 1022;
N.S. Lockyer et al., Phys.Rev.Lett. 51 (1983) 1316.
- 22 C. Jarlskog (rapporteur) in Proc. HEP83, Brighton (UK), Eds. J. Guy and C. Costain, Rutherford Appleton Lab. (1983), p. 768.
- 23 D.H. Wilkinson, in "Nuclear Physics with Heavy Ions and Mesons", Eds. R.Balian, M. Rho and G. Ripka, North-Holland, Amsterdam (1978).
- 24 R. Gatto, G. Sartori and M. Tonin, Phys.Lett. 28B (1968) 128;
N. Cabibbo and L. Maiani, Phys.Lett. 28B (1968) 131.
R.J. Oakes, Phys.Lett. 29B (1969) 683.
- 25 For a recent proposal concerning the link between quark masses and mixing angles see B. Stech, Phys.Lett. 130B (1983) 189.
- 26 A. Sirlin, Phys.Rev. D5 (1972) 436.
- 27 S. Fubini and G. Furlan, Physics 1 (1965) 229.
- 28 For a detailed discussion of this framework and references to the literature see H. Leutwyler and J. Stern, Ann.Phys. (N.Y.) 112 (1978) 94.
- 29 Null plane wave functions of the pseudoscalar mesons were analyzed by V.L. Chernyak and A.R. Zhitnisky, Preprints Novosibirsk 83-103, 104, 105, 106, 107, 108 (1983)
- 30 H. Leutwyler, Nucl.Phys. B76 (1974) 413.
- 31 M. Böhm and D. Rein, Nucl.Phys. B11 (1969) 61;
D. Rein, Z.Phys. 259 (1973) 205;
N. Isgur, Phys.Rev. D12 (1975) 3661;
G. Karl, private communication;
J. Trampetić, private communication.