$\eta \rightarrow 3\pi$ TO ONE LOOP^{*)}

J. Gasser

Institut für Theoretische Physik der Universität Bern

and

H. Leutwyler⁺⁾

CERN - Geneva

A B S T R A C T

We calculate the amplitude of the decays $\eta \rightarrow \pi^0 \pi^+ \pi^-$ and $\eta \rightarrow \pi^0 \pi^0 \pi^0$ to first non-leading order in the low energy expansion. It is shown that the infra-red singularities of chiral perturbation theory strongly enhance the perturbation generated by the quark mass term. The corrections of order m_s increase the current algebra prediction for the rate by more than a factor of two and remove the well-known discrepancy between the low energy theorems of current algebra and experiment.

^{*)} Work supported in part by Schweizerischer Nationalfonds.

⁺⁾ On leave of absence from Universität Bern; permanent address: Institut für Theoretische Physik, Sidlerstrasse 5, 3012 Bern.

1. Introduction

At low energies, up to effects of order E^2/M_W^2 , the standard model reduces to the gauge theory of $SU(3)_C \times U(1)_{e.m.}$. If the mass difference $m_u - m_d$ and the electromagnetic interaction are switched off, this theory conserves isospin and the transition $\eta \rightarrow 3\pi$ is forbidden by Bose symmetry. Accordingly, the amplitude

$$\langle \pi^0 \pi^+ \pi^- \text{ out} | \eta \rangle = i(2\pi)^4 \delta^4(P_f - P_i) A(s, t, u)$$

$$s = (p_\eta - p_{\pi^0})^2, \quad t = (p_\eta - p_{\pi^+})^2, \quad u = (p_\eta - p_{\pi^-})^2 \quad (1.1)$$

only receives a contribution proportional to the quark mass difference

$A_{\text{QCD}} = O(m_u - m_d)$ and a contribution of electromagnetic origin, $A_{\text{QED}} = O(e^2)$ (Sutherland 1966; Bose and Zimerman 1966; Bell and Sutherland 1968; Wilson 1969).

The transformation properties of the isospin violating piece of the interaction imply that the three pions emerge in a configuration with $I = 1$. This is evident for the QCD part: the isospin breaking piece of the QCD Hamiltonian is the $\Delta I = 1$ operator

$$\mathcal{H}_{\text{QCD}}(x) = \frac{1}{2} (m_u - m_d) (\bar{u}u - \bar{d}d) \quad (1.2)$$

In the case of the electromagnetic contribution, described by the operator

$$\mathcal{H}_{\text{QED}}(x) = -\frac{1}{2} e^2 \int dy D^{\mu\nu}(x-y) T j_\mu(x) j_\nu(y) \quad (1.3.)$$

(where $D^{\mu\nu}(x-y)$ is the photon propagator), the selection rule arises because only the interference term between the isovector and isoscalar components of the electromagnetic current $j_\mu = j_\mu^3 + \frac{1}{2} j_\mu^Y$ carries the proper G-parity.

The selection rule $\Delta I = 1$ implies that the function $A(s, t, u)$ also determines the decay into neutral pions

$$\langle \pi^0 \pi^0 \pi^0 \text{ out} | \eta \rangle = i(2\pi)^4 \delta^4(P_f - P_i) \bar{A}(s, t, u)$$

$$\bar{A}(s, t, u) = A(s, t, u) + A(t, u, s) + A(u, s, t) \quad (1.4)$$

(Invariance under charge conjugation requires $A(s,t,u)$ to be symmetric under $t \leftrightarrow u$.)

The QCD part of the amplitude is determined by the scale Λ of the theory and by the renormalization group invariant quark masses

$$A_{\text{QCD}}(s,t,u) = (m_u - m_d) f(p^2, \Lambda, m_u, m_d, m_s, m_c, \dots) \quad (1.5)$$

where p^2 stands for the invariant momentum variables, say for t and u . Our analysis of the function f is based on the assumption that the light quark masses m_u , m_d , m_s may be treated as perturbations.

In the chiral limit $m_u = m_d = m_s = 0$ both the pion and the eta are massless. The square of the mass is of first order in the light quark masses

$$\begin{aligned} \overset{\circ}{M}_\pi^2 &= (m_u + m_d) B_0 \\ \overset{\circ}{M}_\eta^2 &= \frac{1}{3} (4m_s + m_u + m_d) B_0 \end{aligned} \quad (1.6)$$

(In these expressions as throughout this paper terms of order $(m_u - m_d)^2/m_s$ are neglected. The index $\circ$ in $\overset{\circ}{M}_\pi$, $\overset{\circ}{M}_\eta$ indicates that these formulae are valid up to corrections of order M^2 where M is the mass matrix of the light quarks u, d, s .) Kinematics restricts the variables s, t, u to be small, of order M_η^2 , i.e. of first order in the quark masses. We consider a simultaneous expansion of f in powers of the light quark masses and of the momenta, at fixed ratios $p^2 : m_u : m_d : m_s$

$$f = f_0 + f_1 + \dots \quad (1.7)$$

The expansion starts with a term of order 1: f_0 depends on p^2 , m_u , m_d and m_s only through their ratios. The next term f_1 is of order p^2 , i.e. may be written as m_s times as function of the ratios, etc. (this only holds up to logarithmic contributions, see below).

The lowest order term f_0 is well known (Osborn and Wallace 1970; this paper also contains references to the earlier literature). The structure of this contribution is determined by chiral symmetry

$$f_0 = \frac{B_0}{3\sqrt{3} F_\pi^2} \left\{ 1 + \frac{3(S-S_0)}{M_\eta^2 - M_\pi^2} \right\}$$

$$S_0 = \frac{1}{3} (s+t+u) = M_\pi^2 + \frac{1}{3} M_\eta^2 \quad (1.8)$$

In this formula F_π is the pion decay constant [in our normalization $F_\pi = 93.3$ MeV; strictly speaking, the lowest order term in the expansion (1.7) is obtained from (1.8) by replacing F_π by its value F_0 in the chiral limit and by using the first order expressions (1.6) for the meson masses].

The electromagnetic contribution to the transition amplitude may be represented in an analogous fashion

$$A_{QED}(s,t,u) = e^2 g(p^2, \Lambda, m_u, m_d, m_s, \dots)$$

$$g = g_0 + g_1 + \dots \quad (1.9)$$

where g_0 is of order 1, g_1 is of order p^2 etc. (always expanding at fixed ratios $p^2 : m_u : m_d : m_s$). As was shown by Sutherland (1966), chiral symmetry implies that the lowest order term g_0 vanishes: in contrast to the mass differences $M_{\pi^+}^2 - M_{\pi^0}^2$ and $M_{K^0}^2 - M_{K^+}^2$, which do receive an electromagnetic contribution of order $e^2 \Lambda^2$, chiral symmetry suppresses the matrix element $\langle \pi^0 | \mathcal{H}_{QED} | \eta \rangle$ as well as the amplitude $\langle 3\pi | \mathcal{H}_{QED} | \eta \rangle$ at leading order in the low energy expansion (for a derivation of these results within the effective Lagrangian approach see ref. IV).

The structure of the effective electromagnetic interaction at leading order in the low energy expansion also allows us to relate the quantity $(m_u - m_d) \cdot B_0$, which determines the size of the amplitude $\eta \rightarrow 3\pi$, to measurable mass differences. The reasoning is as follows. In the absence of the electromagnetic interaction, the mass difference $M_{K^0}^2 - M_{K^+}^2$ is of first order in $m_u - m_d$

$$(M_{K^0}^2 - M_{K^+}^2)_{QCD} = (m_d - m_u) \cdot B_0 \left\{ 1 + O(m) \right\} \quad (1.10)$$

whereas the mass difference $(M_{\pi^+}^2 - M_{\pi^0}^2)_{QCD}$ is negligibly small, being of second order in $m_u - m_d$. According to Dashen's theorem, at leading order in the low

energy expansion, the electromagnetic contribution to these mass differences is the same (Dashen 1969)

$$(M_{K^0}^2 - M_{K^+}^2 - M_{\pi^0}^2 + M_{\pi^+}^2)_{QED} = O(e^2 m) \quad (1.11)$$

This implies the low energy theorem

$$M_1^2 = (m_d - m_u) \cdot B_0 \{ 1 + O(m) \} + O(e^2 m) \quad (1.12)$$

where M_1^2 stands for the differences of squares of physical masses

$$M_1^2 \equiv M_{K^0}^2 - M_{K^+}^2 - M_{\pi^0}^2 + M_{\pi^+}^2 \quad (1.13)$$

At leading order in the low energy expansion chiral symmetry therefore fully determines the η -decay amplitude in terms of measurable quantities:

$$A(s, t, u) = -\frac{M_1^2}{3\sqrt{3}F_\pi^2} \left\{ 1 + 3 \frac{s - s_0}{M_\eta^2 - M_\pi^2} \right\} \left\{ 1 + O(p^2) + O\left(e^2 \frac{m_s - \hat{m}}{m_d - m_u}\right) \right\} \quad (1.14)$$

The symbol $O(p^2)$ stands for the higher order terms in the low energy expansion. It includes contributions of order s, t, u as well as contributions of order m_u, m_d, m_s . The last term in (1.14) stems from possible violations of Dashen's theorem. [In (Gasser and Leutwyler 1982) we have estimated the uncertainty in the $K^0 - K^+$ mass difference in pure QCD due to these violations at ± 0.33 MeV. This estimate implies that the contributions of order $e^2 m_s / (m_u - m_d)$ to the amplitude $A(s, t, u)$ generate an uncertainty of $0.33/5.3 \sim 6\%$.]

The differential rate for the decay $\eta \rightarrow \pi^0 \pi^+ \pi^-$ is given by

$$d\Gamma_{\eta \rightarrow \pi^0 \pi^+ \pi^-} = \frac{1}{2M_\eta} (2\pi)^4 \delta^4(P_f - P_i) d\mu_0 d\mu_+ d\mu_- |A(s, t, u)|^2$$

$$d\mu = \frac{d^3 p}{(2\pi)^3 2p^0} \quad (1.15)$$

Evaluating the phase space integral with (1.14) and using the value $F_\pi = 93.3$ MeV one obtains [note that phase space is so small that the mass difference $M_{\pi^+} - M_{\pi^0}$ plays a significant role in the numerical prediction for the rate]:

$$\Gamma_{\eta \rightarrow \pi^0 \pi^+ \pi^-} = 66 \text{ eV} \quad (1.16)$$

to be compared with the experimental value (Particle Data Group 1982)

$$\Gamma_{\eta \rightarrow \pi^0 \pi^+ \pi^-} = 197 \pm 29 \text{ eV} \quad (1.17)$$

The prediction (1.16) is off by a factor of three. (Note that the uncertainty in the current algebra prediction for the rate due to violations of Dashen's theorem as estimated above only amounts to ± 8 eV). The flat disagreement between the current algebra prediction and experiment has prompted many authors to investigate the problem. (See, e.g., Osborn and Wallace 1970; Cantor 1971; Schechter and Ueda 1971; Dittner, Dondi and Eliezer 1973; Cicogna, Strocchi and Vergara Caffarelli 1974; Hudnall and Schechter 1974; Langacker and Pagels 1974; Weinberg 1975; Kogut and Susskind 1975; Crewther 1975; Raby 1976; Matsubara and Tameike 1977; Dominguez and Zepeda 1978; Becherrawy and Nasrallah 1978; Milton, Palmer and Prinsky 1980; Kawarabayashi and Ohta 1980; Roiesnel and Truong 1981, 1982; Minkowski 1982; Tarrach 1982; Truong 1983).

To compare the corresponding low energy theorem for the decay into neutral pions with the data, we consider the branching ratio

$$r = \Gamma_{\eta \rightarrow 3\pi^0} / \Gamma_{\eta \rightarrow \pi^0 \pi^+ \pi^-} \quad (1.18)$$

With (1.4) the low energy theorem (1.16) predicts

$$r = 1.51 \quad (1.19)$$

whereas the Particle Data Group quotes

$$r = 1.344 \pm 0.053 \quad (1.20)$$

in disagreement with (1.19) by three standard deviations.

The low energy theorems of current algebra are sensitive tests of the chiral properties of the theory; the flat disagreement with observation manifesting itself in the η -decay rates poses a serious problem. One may of course blame the discrepancy on the fact that, after all, the mass of the strange quark is not

that small and that an expansion in powers of m_s may therefore fail to converge rapidly. Since the mass m_s however practically coincides with the SU(3) breaking parameter $m_s - \hat{m}$, this "resolution" of the puzzle would imply that SU(3) is not a reliable approximate symmetry of the strong interactions. How can it happen that the Gell-Mann-Okubo formula for the meson masses holds to a remarkable degree of accuracy and that F_K is approximately equal to F_π , if at the same time the low energy theorem for η decay fails by a factor of three ?

The origin of the discrepancy was identified by Roiesnel and Truong (1981), who have shown that the unitarity corrections generated by $\pi\pi$ final state interactions are large. In the systematic expansion discussed above the unitarity corrections are corrections of order p^2 , contained in the quantity f_1 . The purpose of the present paper is to show that f_1 can be calculated explicitly in terms of measurable quantities. The calculation shows that the large deviation from the low energy theorem of current algebra is due to the fact that the expansion in powers of the quark masses is plagued by infrared singularities, which enhance the effect of the perturbation generated by the quark mass term. It so happens that the low energy theorem for the decay $\eta \rightarrow 3\pi$ is very unstable with respect to these perturbations, while the Gell-Mann-Okubo formula and the lowest order prediction $F_K : F_\pi = 1$ are stable.

Before we proceed to show how f_1 can be calculated, we briefly comment on the electromagnetic contribution to the decay rate. It is instructive to compare the effect of the electromagnetic interaction on the decay $\eta \rightarrow 3\pi$ with its effect on the mass difference $M_{K^0} - M_{K^+}$. As was discussed above, this mass difference also receives a contribution both from $m_u - m_d$ and from electromagnetism. The electromagnetic contribution can be calculated from the observed value of $M_{\pi^+} - M_{\pi^0}$ with the result $(M_{K^0} - M_{K^+})_{\text{QED}} = -1.27 \pm 0.3 \text{ MeV}$ (see ref. I). The measured mass difference shows that the contribution from $m_u - m_d$ is of opposite sign: $(M_{K^0} - M_{K^+})_{\text{QCD}} \simeq 5.3 \text{ MeV}$. Despite the fact that chiral symmetry does not suppress the electromagnetic contribution in this case, the effect of $m_u - m_d$ on the mass difference is substantially larger than the effect of the electromagnetic interaction.

In the case of the decay $\eta \rightarrow 3\pi$ the electromagnetic interaction gives rise to two effects: (i) it increases the mass of the charged pion from $(M_{\pi^\pm})_{\text{QCD}} \simeq M_{\pi^0}$ to the observed value. This effect modifies the rate of the decay $\eta \rightarrow \pi^0 \pi^+ \pi^-$ considerably, as it reduces phase space by 11%. (ii) it generates an electromagnetic con-

tribution A_{QED} to the transition amplitude. We expect this effect to be very small for the following reason. As was mentioned above, chiral symmetry implies that, in contrast to the mass differences $M_{K^0} - M_{K^+}$, $M_{\pi^+} - M_{\pi^0}$ the transition amplitude does not receive an electromagnetic contribution at leading order in the low energy expansion. In fact it is shown in Appendix A that chiral symmetry suppresses the electromagnetic contribution even at first nonleading order (Bell and Sutherland 1968; Dittner, Dondi and Eliezer 1973; and ref. IV): in the physical region of the decay, A_{QED} is of order $e^2(m_u + m_d)$, rather than of order $e^2 m_s$. At this order of magnitude ordinary radiative corrections, proportional to $e^2(m_u - m_d)$, come into play. We therefore disregard the electromagnetic interaction, except for using the physical pion masses in the evaluation of the phase space integrals. (Even if carried to first nonleading order, an expansion in powers of m_s does not reach the accuracy at which these electromagnetic contributions become relevant. Note, however, that the Dalitz plot distribution and the ratio $\Gamma_{\eta \rightarrow 3\pi^0} / \Gamma_{\eta \rightarrow \pi^0 \pi^+ \pi^-}$ are controlled by $SU(2) \times SU(2)$ - these relative quantities only show small deviations from the current algebra predictions. In the analysis of these deviations, electromagnetic effects are not a priori negligible.)

2. Transition amplitude to one loop

The technology required to calculate the function f_1 defined in (1.5), (1.7) is described in detail in ref. III. It suffices to evaluate the four-point function of the axial current from the formulae given in that reference; the amplitude $A(s,t,u)$ we are looking for is the residue of the poles which occur in this Green's function. (The calculation of the on-shell amplitude simplifies considerably if use is made of the method described in the Appendix). We write the result in the form

$$A(s,t,u) = -\frac{M_1^2}{3\sqrt{3} F_\pi^2} \left\{ T(s) + U(s,t,u) + V(s) + W(s,t,u) \right\} \quad (2.1)$$

The term proportional to $T(s)$

$$T(s) = 1 + 3 \frac{s - s_0}{\Delta} ; \quad s_0 \equiv M_\pi^2 + \frac{1}{3} M_\eta^2 ; \quad \Delta \equiv M_\eta^2 - M_\pi^2 \quad (2.2)$$

is the leading order contribution given in section 1. The quantity M_1^2 is defined in (1.13) in terms of the physical masses and F_π denotes the physical pion decay constant, $F_\pi = 93.3$ MeV.

The contribution $U(s,t,u)$ denotes the unitarity correction generated by the one loop graphs which appear at this order in the low energy expansion. The main labor concerns the isospin gymnastics in the evaluation of this contribution: one needs to expand the general expression for the unitarity correction given in ref. III to first order in the mass difference $m_u - m_d$. This mass difference occurs in three places : (i) The lowest order effective Lagrangian explicitly contains the quark mass matrix and hence contains vertices proportional to $m_u - m_d$ (σ -terms). (ii) The explicit expression for the one loop integrals refers to a meson field basis in which the meson mass term is diagonal. Because of $\pi^0 \eta$ mixing this basis does not coincide with the octet basis:

$$\lambda_{\pi^0} = \lambda^3 + \varepsilon \lambda^8 ; \quad \lambda_\eta = \lambda^8 - \varepsilon \lambda^3$$

$$\varepsilon = \frac{\sqrt{3}}{4} \frac{m_d - m_u}{m_s - \hat{m}} \quad (2.3)$$

(iii) Finally, the masses of the mesons which run around the loop implicitly depend on the quark masses. Since M_K^2 receives a contribution from $m_u - m_d$, K-meson loops produce a unitarity correction due to a mass insertion in the propagators (there is no such contribution from loops involving pions or etas, because the masses of these particles do not contain a term linear in $m_u - m_d$).

Explicitly, the unitarity correction is of the form

$$U(s,t,u) = \frac{1}{3} \Delta_0(s) \{ 3T(s) + T(t) + T(u) \}$$

$$+ \frac{1}{2} \Delta_1(t) \{ T(s) - T(u) \} + \frac{1}{2} \Delta_1(u) \{ T(s) - T(t) \}$$

$$+ \frac{1}{2} \Delta_2(t) \{ T(s) + T(u) \} + \frac{1}{2} \Delta_2(u) \{ T(s) + T(t) \}$$

$$+ \Delta_3(s) \quad (2.4)$$

The functions $\Delta_i(s)$ may be expressed in terms of the renormalized loop integrals $J^r(s)$ and $M^r(s)$ defined in ref. III:

$$\begin{aligned}
 \Delta_0(s) &= \frac{1}{2F_\pi^2} (2s - M_\pi^2) J_{\pi\pi}^r(s) \\
 \Delta_1(s) &= \frac{2s}{F_\pi^2} \left\{ M_{\pi\pi}^r(s) + \frac{1}{2} M_{KK}^r(s) \right\} \\
 \Delta_2(s) &= -\frac{1}{2F_\pi^2} (s - 2M_\pi^2) J_{\pi\pi}^r(s) + \frac{1}{4F_\pi^2} (3s - 4M_K^2) J_{KK}^r(s) \\
 &\quad + \frac{M_\pi^2}{3F_\pi^2} J_{\pi\eta}^r(s) \\
 \Delta_3(s) &= -\frac{1}{6F_\pi^2 \Delta} (s - 2M_\pi^2)(3s - 4M_K^2) J_{\pi\pi}^r(s) \\
 &\quad - \frac{s(3s - 4M_\pi^2)}{4F_\pi^2 \Delta} J_{KK}^r(s) + \frac{M_\pi^2}{3F_\pi^2 \Delta} (3s - 4M_\pi^2) J_{\pi\eta}^r(s) \\
 &\quad - \frac{M_\pi^2}{2F_\pi^2} J_{\eta\eta}^r(s) - \frac{3s}{8F_\pi^2} \frac{(3s - 4M_K^2)}{(s - 4M_K^2)} \left\{ J_{KK}^r(s) - J_{KK}^r(0) - \frac{1}{8\pi^2} \right\}
 \end{aligned} \tag{2.5}$$

The imaginary part of $\Delta_0(s)$ is the $I = 0$, S-wave $\pi\pi$ phase shift

$$\Delta_0(s) = \frac{s^2}{\pi} \int_{4M_\pi^2}^{\infty} \frac{ds' \delta_0^0(s')}{s'^2(s' - s - i\epsilon)} + C_0 + C_1 s \tag{2.6}$$

To leading order in the low energy expansion, this phase shift is determined by F_π

$$\delta_0^0(s) = \frac{1}{32\pi F_\pi^2} (2s - M_\pi^2) \left\{ 1 - 4M_\pi^2/s \right\}^{1/2} \quad (2.7)$$

The imaginary parts of $\Delta_1(s)$ and $\Delta_2(s)$ contain the $I = 1$, P-wave and $I = 2$, S-wave $\pi\pi$ phase shifts respectively

$$\begin{aligned} \delta_1^1(s) &= \frac{1}{96\pi F_\pi^2} (s - 4M_\pi^2) \left\{ 1 - 4M_\pi^2/s \right\}^{1/2} \\ \delta_0^2(s) &= -\frac{1}{32\pi F_\pi^2} (s - 2M_\pi^2) \left\{ 1 - 4M_\pi^2/s \right\}^{1/2} \end{aligned} \quad (2.8)$$

as well as contributions from $K\bar{K}$ and $\pi\eta$ intermediate states. The function $\Delta_3(s)$ collects the remainder of the unitarity correction; in particular, this term also contains a contribution

$$\Delta_3^{\pi\pi}(s) = -\frac{1}{3} \left\{ T(t) + T(u) \right\} \Delta_2^{\pi\pi}(s)$$

from the $\pi\pi$ final state interaction.

Note that the functions $J^r(s)$ and $M^r(s)$ depend on the scale μ at which the loops are renormalized; this ambiguity reflects the fact that the dispersion relation (2.6) involves subtractions. The scale μ drops out in the expression (2.1) for the full amplitude - only the splitting of this amplitude into a unitarity correction U and a contribution $V + W$ from tree graphs and tadpoles requires the introduction of such a scale.

The functions V and W are polynomials in s, t, u , obtained by expanding the general expression for the tadpole and tree graph contributions given in III to first order in $m_u - m_d$. The expression involves the constants F_0 and B_0 as well as the low energy constants $L_3, L_4, \dots, L_8$, which occur in the effective Lagrangian at order p^4 . The quantity F_0 may be expressed in terms of the physical pion decay constant F_π and L_4, L_5 . Similarly, the product $(m_d - m_u)B_0$ is given by M_1^2 , up to a correction involving L_4, L_5, L_6 and L_8 . Finally, we may eliminate two of the low energy constants in favour of the two measurable quantities

$$d_1 = \frac{1}{\Delta} (4M_K^2 - M_\pi^2 - 3M_\eta^2)$$

$$d_2 = \frac{F_K}{F_\pi} - 1 \quad (2.9)$$

With these substitutions the expressions for V and W take the form

$$V(s) = T(s) \{a_1 + 3a_2\Delta + a_3(9M_\eta^2 - M_\pi^2)\} + a_4$$

$$W(s,t,u) = \frac{2}{3} \left\{ d_1 + \frac{4M_\pi^2}{\Delta} d_2 \right\} T(s) - \frac{8}{3} \frac{M_\pi^2}{\Delta} d_1$$

$$+ \frac{4L_3}{F_\pi^2 \Delta} \{s^2 + 2tu - 3s_0 s\} \quad (2.10)$$

The constants a_i only depend on the meson masses, the pion decay constant and the scale μ :

$$a_1 = \frac{2}{3\Delta} M_\pi^2 (-\mu_\pi - 2\mu_K + 3\mu_\eta)$$

$$a_2 = \frac{2}{3\Delta^2} (-M_\pi^2 \mu_\pi + 4M_K^2 \mu_K - 3M_\eta^2 \mu_\eta)$$

$$a_3 = \frac{1}{128\pi^2 F_\pi^2} \left(\ln \frac{M_K^2}{\mu^2} + 1 \right)$$

$$a_4 = -8M_\pi^2 a_2 - 12M_K^2 a_3 + \frac{3M_\pi^2}{32\pi^2 F_\pi^2} \left\{ 1 - \frac{M_\pi^2}{M_K^2 - M_\pi^2} \ln \frac{M_K^2}{M_\pi^2} \right\}$$

$$\mu_P = \frac{1}{32\pi^2 F_\pi^2} M_P^2 \ln \frac{M_P^2}{\mu^2} \quad (2.11)$$

Since the low energy constant L_3 is scale independent ($\Gamma_3 = 0$), the amplitude W is independent of μ . The scale dependence of the constants a_i must therefore counterbalance the scale dependence of the unitarity correction in such a manner that $U + V$ is independent of μ . We have checked that this is indeed the case - a rather thorough consistency check on our calculation. (Note that the functions U and V are accurate only to leading order in the quark mass expansion; in lead-

ing order the meson masses satisfy the Gell-Mann-Okubo formula).

In the physical region, the imaginary part of the amplitude is exclusively generated by the pion loop integrals contained in $U(s,t,u)$. Analytically continued, the same amplitude also describes the scattering process $\eta\pi \rightarrow \pi\pi$. We have verified that, to order p^4 , the absorptive part of the functions $J_{\pi\pi}^r, M_{\pi\pi}^r$ contained in $U(s,t,u)$ indeed reproduces the contributions of $\pi\pi$ intermediate states to the unitarity condition for this scattering process. In fact, unitarity fully determines the amplitude at order p^4 up to a real second order polynomial in s, t, u (in the notation used above, this polynomial is denoted by $V+W$). The heavy machinery developed in III is needed only to pin down this polynomial in terms of $M_\pi, M_K, M_\eta, F_\pi, F_K$ and L_3 .

As a final check, one easily verifies that the corrections preserve the Adler zero, which manifests itself in the current algebra prediction: for $s, u = O(M_\pi^2)$, $t = M_\eta^2 + O(M_\pi^2)$ the amplitude is of order M_π^2 .

In addition to the constants d_1, d_2 defined in (2.9), which are well determined experimentally [$d_1 = 0.21$; the recent determination of F_K/F_π (Leutwyler and Roos 1984) implies $d_2 = 0.22 \pm 0.01$], the amplitude depends on the low energy constant L_3 , which is not known with the same accuracy. (This constant could presumably be measured directly in $K_{\ell 4}$ decay or in $K\pi$ scattering, but this yet remains to be done). In III we have shown that a theoretical estimate for L_3 may be derived from the large N_c -expansion (Okubo-Iizuka-Zweig rule). The theoretical prediction is

$$L_3 = \frac{5}{2} \pi F_\pi^4 (-2 a_2^0 + 5 a_2^2) - \frac{1}{1536 \pi^2} \quad (2.12)$$

where a_2^0, a_2^2 are the D-wave $\pi\pi$ scattering lengths. The observed values of these quantities as reported by Petersen in the compilation of coupling constants and low energy parameters (Nagels et al. 1979) imply that L_3 is small:

$$L_3 = (-4.4 \pm 2.5) \cdot 10^{-3} \quad (2.13)$$

With this information the above explicit expressions for U, V and W provide us with a parameter free representation of the decay amplitude. Note that this representation not only accounts for π_η^0 mixing, but also automatically includes

mixing with the η' : in the decay amplitude the η' induces symmetry breaking effects of order $(m_u - m_d) \cdot (m_s - \hat{m})$, which the low energy representation given above takes care of. [In the language used here, the η' affects the low energy constant L_7 , which generates a substantial contribution to the quantity d_1 , defined in (2.9). See ref. III, sections 10 and 12].

3. Results

a. Decay rate for the charged mode

Evaluating the decay rate $\Gamma_{\eta \rightarrow \pi^0 \pi^+ \pi^-}$ with the representation of the amplitude given in the last section we find

$$\Gamma_{\eta \rightarrow \pi^0 \pi^+ \pi^-} = 160 \pm 50 \text{ eV} \quad (3.1)$$

(the error bar represents an estimate of the theoretical uncertainties due to higher order contributions in the low energy expansion, see below.) The result (3.1) is to be compared with the experimental value $197 \pm 29 \text{ eV}$.

We thus confirm the conclusion of Roiesnel and Truong (1981, 1982): the current algebra prediction (66 eV) receives a large correction from effects of order m_s ; which increase the rate by a factor 2.4 ± 0.8 and remove the discrepancy with the measured rate.

Note that the recent data (Weinstein et al. 1983) on the decay $\eta \rightarrow 2\gamma$ imply $\Gamma_{\eta \rightarrow 2\gamma} = 0.56 \pm 0.16 \text{ keV}$. In view of the large error bar this result is not inconsistent with the value $\Gamma_{\eta \rightarrow 2\gamma} = 0.324 \pm 0.046 \text{ keV}$ (Browman et al. 1974), quoted by the Particle Data Group. (With the known branching ratios, the new value implies $\Gamma_{\eta \rightarrow \pi^0 \pi^+ \pi^-} = 340 \pm 100 \text{ eV}$). If the new value should however be confirmed, the calculated rate (3.1) would be off by a factor of two. In our opinion this is beyond the uncertainties involved in the calculation described here and would resurrect the puzzle we claim to solve.

To understand why the corrections of order m_s are large, it suffices to consider the value of the amplitude in the center of the Dalitz plot ($s = t = u = s_0$). With $T(s_0) = 1$ the correction factor is given by

$$C = 1 + U + V + W$$

Numerically, $C = 1.50 + i 0.18$, $|C|^2 = 2.3$. An important contribution stems from the $\pi\pi$ final state interaction, contained in the unitarity correction $U(s,t,u)$. In the center of the Dalitz plot the P-waves described by $\Delta_1(s)$ do not contribute. The $I = 0$, S-wave $\pi\pi$ phase shift generates the function $\Delta_0(s)$, given explicitly by

$$\Delta_0(s) = \frac{1}{32\pi^2 F_\pi^2} (2s - M_\pi^2) \left\{ \sigma \ln \frac{1-\sigma}{1+\sigma} + 1 - \ln \frac{M_\pi^2}{\mu^2} + i\pi\sigma \right\}$$

$$\sigma = (1 - 4M_\pi^2/s)^{1/2} \quad (3.2)$$

The contribution from the $I = 0$ final state interaction to the correction factor C is given by $\frac{5}{3} \Delta_0(s_0)$ with $s_0 = M_\pi^2 + \frac{1}{3} M_\eta^2$. The $I = 2$, S-wave phase shift generates contributions both to $\Delta_2(s)$ and to $\Delta_3(s)$. In the center of the Dalitz plot these contributions may also be expressed in terms of $\Delta_0(s_0)$; in the correction factor C the $I = 2$ piece of the final state interaction amounts to $\frac{4}{3} \Delta_2^{\pi\pi}(s_0) \simeq -\frac{1}{2} \Delta_0(s_0)$ (the sign is opposite to the contribution from $I = 0$: in the isoscalar channel the interaction is attractive and enhances the decay rate, in the $I = 2$ channel the interaction is repulsive).

In the center of the Dalitz plot the net effect of the final state interaction is a unitarity correction given by $C^{\pi\pi} = \frac{7}{6} \Delta_0(s_0)$. This contribution is responsible, in particular, for the imaginary part of C (the contributions from $K\bar{K}$, $\pi\eta$ and $\eta\eta$ loops and the polynomials V and W are real). The $I = 0$ and $I = 2$ channels contribute $\text{Im } C_{I=0}^{\pi\pi} = 0.26$ and $\text{Im } C_{I=2}^{\pi\pi} = -0.08$ respectively. In the square of the amplitude, the contribution from the imaginary part is negligibly small.

As can be seen from (3.2), the real part of Δ_0 depends on the scale μ at which the unitarity correction is normalized (i.e. on the subtraction point in the dispersion relation for $\Delta_0(s)$). For $\mu = 500$ MeV and $\mu = 1$ GeV we obtain $\text{Re } C^{\pi\pi} = 0.25$ and $\text{Re } C^{\pi\pi} = 0.38$ respectively. The final state interaction among the pions therefore increases the square of the amplitude in the center of the Dalitz plot by $|1 + C^{\pi\pi}|^2 = 1.6$ or by $|1 + C^{\pi\pi}|^2 = 1.9$ (depending on the manner in which the unitarity correction is split off from the rest).

The final state interaction is thus responsible for about half of the difference between the lowest order current algebra prediction for the decay rate and the prediction (3.1) which accounts for all corrections of order m_S .

Fixing the scale at $\mu = 750$ MeV, the loops containing $K\bar{K}$, $\pi\eta$ and $\eta\eta$ intermediate states contribute $C^{K\bar{K}} = 0.05$, $C^{\pi\eta} = 0.008$ and $C^{\eta\eta} = 0.002$ respectively. The chiral logarithms collected in the amplitude V amount to $V = -0.13$ and $V = +0.05$ for $\mu = 500$ MeV and $\mu = 1$ GeV respectively. The remainder, given by $W = 0.14$ mainly stems from the quantity d_1 which measures the deviation from the Gell-Mann-Okubo formula.

Fixing the scale at $\mu = 750$ MeV we thus have the following decomposition of the correction factor: $\text{Re}C = 1 + 0.33 + 0.06 - 0.03 + 0.14 = 1.50$, where the first correction is due to $\pi\pi$ final state interactions, the second term comes from the analogous contributions involving virtual K or η mesons, the third piece is generated by the chiral logarithms collected in V and the remainder comes from the mass correction in W .

In the language used here the unitarity correction considered by Roiesnel and Truong (1981, 1982) concerns the contribution $C^{\pi\pi}$ from $\pi\pi$ final state interactions to the function $U(s,t,u)$. These authors estimate $\text{Re}C^{\pi\pi} = 0.83$ and hence predict a correction in the decay rate by the factor $(1.83)^2 = 3.3$ from final state interactions alone. This estimate for $\text{Re}C^{\pi\pi}$ is substantially larger than the numbers given above. There are several reasons for this difference. In their work

(i) the two body unitarity correction is applied only to a part of the amplitude. If an analogous correction is carried out also in the remainder, the contribution due to $I = 0$, S -wave final state interactions is reduced by the factor $\{3T(s) + T(t) + T(u)\}/9$. In the center of the Dalitz plot, this contribution is therefore overestimated by a factor $9/5$. (In part, the subsequent unitarization performed by these authors compensates for the missing factor.)

(ii) the quantity $\Delta_0(s)$ is evaluated phenomenologically, with $a_0^0 = 0.25$, whereas the lowest order current algebra formula (3.2) is equivalent to $a_0^0 = 0.16$.

(iii) the final state interaction is treated in exponential form, whereas our low energy representation only accounts for the first two terms in the Taylor series, $\exp \Delta_0 = 1 + \Delta_0 + \dots$

(iv) the repulsive $I = 2$ interaction, which reduces the unitarity correction, is neglected.

As was shown in ref. II, higher order corrections indeed increase the current algebra prediction for a_0^0 , to $a_0^0 = 0.20 \pm 0.01$ (the analogous corrections to a_0^2

are very small). To include these particular higher order corrections in the low energy representation for the decay amplitude, it suffices to reduce the value of F_π in the expression (3.2) for $\Delta_0(s)$ by the factor $(0.16/0.20)^{1/2} = 0.89$. With this modification, and fixing the scale at $\mu = 750$ MeV, the function $\exp \Delta_0(s)$ reproduces the phenomenological representation for the quantity $D^{-1}(s)$ given by Roiesnel and Truong quite well ($a_0^0 = 0.20$; the value $a_0^0 = 0.25$ is inconsistent with chiral symmetry.) If one calculates the correction factor in the center of the Dalitz plot with this modified representation for the $I = 0$, S-wave contribution $\Delta_0(s)$ and, furthermore, replaces $\Delta_0(s)$ by $\exp \Delta_0(s) - 1$, one finds $\text{Re}C^{\pi\pi} = 0.52$, instead of the value $\text{Re}C^{\pi\pi} = 0.33$ which results from the lowest order formula with $\mu = 750$ MeV. The exercise shows that higher order effects presumably increase the unitarity correction contained in our low energy representation. We consider it however very unlikely that these higher order effects more than double the leading correction. With the simple model for higher order corrections just described, the prediction for the rate is increased by 50eV, the theoretical uncertainty quoted in (3.1).

b. Dalitz plot

We discuss the distribution over the Dalitz plot in terms of the standard variables x and y (Particle Data Group 1982), defined by

$$\begin{aligned} x &= \sqrt{3} \frac{T_+ - T_-}{Q} = \frac{\sqrt{3}}{2M_\eta Q} (u - t) \\ y &= \frac{3T_0}{Q} - 1 = \frac{3}{2M_\eta Q} \{ (M_\eta - M_{\pi^0})^2 - s \} - 1 \\ Q &= M_\eta - 2M_{\pi^+} - M_{\pi^0} \end{aligned} \quad (3.3)$$

[Note that the scale of the variables x, y is rather sensitive to the mass difference $M_{\pi^+} - M_{\pi^0}$, because the Q -value is small. To compare our amplitude with the data we therefore need to distinguish M_{π^0} from M_{π^+} . The difference between these two masses is of electromagnetic origin. As discussed in section 1, the electromagnetic contributions to the amplitude $A(s, t, u)$ itself are expected to be very small. We therefore simply use the relations (3.3), together with $s + t + u = M_\eta^2 + 2M_{\pi^+}^2 + M_{\pi^0}^2$, to calculate the values of s, t, u which correspond to a given point x, y on the Dalitz plot and insert these values in the calculated amplitude. The difference between $M_\eta^2 + 2M_{\pi^+}^2 + M_{\pi^0}^2$ and $3s_0 = M_\eta^2 + 3M_{\pi^0}^2$ amounts

to less than 1%, which is the order of magnitude of the neglected electromagnetic contributions to $A(s,t,u)$.]

We analyze the Dalitz plot distribution in terms of the mean value of $|A(s,t,u)|^2$ over the variable x

$$\hat{M}^2(y) = \frac{1}{2x_1(y)} \int_{-x_1(y)}^{x_1(y)} dx |A(s,t,u)|^2 \quad (3.4)$$

(where $x = \pm x_1(y)$ is the boundary of the physical region) and define the normalized distribution $M^2(y)$ as the ratio of $\hat{M}^2(y)$ to $\hat{M}^2(-1)$:

$$M^2(y) = \hat{M}^2(y) / \hat{M}^2(-1) ; -1 \leq y \leq y_1 \quad (3.5)$$

The function $M^2(y)$ is plotted in Fig. 1. Curve 1 describes the current algebra prediction. Since the current algebra formula implies that the amplitude is independent of x and is linear in y , the corresponding representation for the normalized distribution is of the form

$$M^2(y) = (1 + \frac{a}{2}y)^2 / (1 - \frac{a}{2})^2 \quad (\text{current algebra}) \quad (3.6)$$

$$a = -1.01$$

More generally, the distribution (3.6) follows if (i) $A(s,t,u)$ is approximately linear in s, t, u and (ii) $A(s,t,u)$ vanishes at $s \simeq M_\pi^2$ (Adler zero).

The curve labelled 2 displays the distribution which follows from the low energy representation of the amplitude to second order, given in section 2. Curves L and G are the experimental distributions of Layter et al. (1973) and Gormley et al. (1970) respectively.

The figure shows that, although the overall corrections to the amplitude are large, these corrections do not modify the Dalitz plot distribution strongly and do therefore not ruin the agreement between the current algebra prediction and the observed distribution. The main feature of the calculated distribution is the presence of an Adler zero near $s \simeq M_\pi^2$ which is clearly confirmed by the data.

A closer look at these distributions shows that the shape of the experimental curves exhibits less curvature than what is predicted by the current algebra

formula [in Layter et al. (1973), it is pointed out that an amplitude linear in s - as required by current algebra - provides a very poor fit to the experimental distribution]. The corrections of order p^2 do not eliminate this difference between the calculated and the observed distributions, but instead even enhance the discrepancy somewhat (to a good approximation, the amplitude remains linear in s ; the zero of the linear approximation occurs at $s \simeq 2.5 M_\pi^2$, instead of occurring at $s = \frac{4}{3} M_\pi^2$ as for the current algebra prediction).

If the distribution $M^2(y)$ is parametrized by a second order polynomial

$$M^2(y) = N \{ 1 + ay + by^2 \}$$

we find $a = -1.30$, $b = 0.38$. The current algebra prediction is $a = -1.01$, $b = \frac{1}{4} a^2 = 0.26$, the data of Layter et al. are well fit by $a = -1.07 \pm 0.013$, $b = 0$, the data of Gormley et al. yield $a = -1.18 \pm 0.02$, $b = 0.20 \pm 0.03$.

Representing the square of the amplitude by a polynomial of the form $N(1 + ay + by^2 + cy^3)$, we obtain $a = -1.33$, $b = 0.42$, $c = 0.08$, to be compared with the values found by Layter et al.: $a = -1.08 \pm 0.014$, $b = 0.034 \pm 0.027$, $c = 0.046 \pm 0.031$ or by Gormley et al.: $a = -1.17 \pm 0.02$, $b = 0.21 \pm 0.03$, $c = 0.06 \pm 0.04$.

In the central region $y \simeq 0$, the calculated value of $M^2(y)$ [which roughly measures the ratio of the square of the amplitude in the center of the Dalitz plot to the square of the amplitude at the boundary $s = (M_\eta - M_{\pi^0})^2$] underestimates the observed value of this quantity by 10 to 20 %.

The discrepancy may be due to one of the following reasons:

(i) Since the overall corrections of order p^2 are large, higher order contributions may affect the distribution significantly, if they are not uniform over the Dalitz plot. As pointed out by Roiesnel and Truong, one may estimate the higher order effects on the basis of the three particle unitarity conditions given by Khuri and Treiman (1960). Roiesnel and Truong indeed obtain considerable modifications of the Dalitz plot distribution in this manner. We did not perform such a calculation, because it is not free of ambiguities: unitarity alone does not determine the amplitude.

(ii) Part of the curvature in the calculated distribution is generated by the constant L_3 . As discussed in section 2, we determine the value of this constant from the D-wave $\pi\pi$ scattering lengths by using large N_c arguments. Since neither the experimental information concerning the D-waves, nor the relevance of the large N_c limit are beyond doubt, it is possible that we have underestimated the uncertainties in the value of the constant L_3 obtained in this manner. (In the prediction for the rate the uncertainties due to L_3 are taken care of by the theoretical error bar given in (3.1). The value of L_3 does however affect the Dalitz plot distribution significantly: if L_3 is set equal to zero, the calculated value of $M^2(0)$ increases by 7%.)

We conclude that within the accuracy to which we trust our low energy representation, the predicted Dalitz plot distribution is consistent with the measured distribution.

c. Branching ratio $\Gamma_{\eta \rightarrow 3\pi^0} / \Gamma_{\eta \rightarrow \pi^0 \pi^+ \pi^-}$

We discuss the information on the neutral decay mode in terms of the branching ratio

$$r = \frac{\Gamma_{\eta \rightarrow 3\pi^0}}{\Gamma_{\eta \rightarrow \pi^0 \pi^+ \pi^-}} \quad (3.7)$$

The current algebra prediction is

$$r = 1.51 \quad (3.8)$$

[This value follows if the amplitude $A(s,t,u)$ is linear in s, t, u and vanishes at $s = \frac{4}{3} M_\pi^2$. If the amplitude is linear, with a slope that differs from the current algebra value $a = -1.01$ by Δa , then the branching ratio is modified by $\Delta r = 0.3 \Delta a$.]

With the low energy representation for the amplitude given in section 2, we find

$$r = 1.43 \quad (3.9)$$

It is not a simple matter to pin down the uncertainties in the prediction (3.9). They are due to our neglect of higher order terms in the low energy expansion and of the electromagnetic contributions (see Bell and Sutherland 1968 and Dittner, Dondi and Eliezer 1973 for an analysis of the latter). Simple estimates for the size of these contributions suggest that the value given should be accurate to within about 2%.

The experimental data concerning the branching ratio r are not very precise. The overall fit to the data, provided by the Particle Data Group (1982), is $r = 1.344 \pm 0.053$. The bulk of the information underlying this value stems from combinations of other branching ratios. These indirect measurements by themselves constrain r to

$$r = 1.36 \pm 0.06 \quad (3.10)$$

The direct measurements of r are internally inconsistent, some of the older data implying a value of r below 1 with a rather small error bar. We therefore disregard the direct measurements [depending on whether one includes or excludes the older data the average of the direct measurements is 1.28 ± 0.14 or 1.46 ± 0.13 respectively. The overall fit mentioned above results from the combination of the first one of these two numbers with the indirect measurements].

The prediction (3.9) agrees with the experimental value (3.10) at the one standard deviation level, whereas the current algebra prediction (3.8) disagrees with this value by $2\sqrt{2} \sigma$.

d. Connection between Dalitz plot distribution and branching ratio

The isospin selection rule, which implies that the processes $\eta \rightarrow \pi^0 \pi^+ \pi^-$ and $\eta \rightarrow 3\pi^0$ are governed by the same function $A(s,t,u)$, leads to a relation between the branching ratio r and the Dalitz plot distribution $M^2(y)$ of the charged decay mode. To establish this relation we assume that the momentum dependence predicted by current algebra provides us with a good first approximation to the actual behaviour of the amplitude. More precisely, we write the full amplitude in the form of a correction to the current algebra prediction

$$A(s, t, u) = N T(s) \{ 1 + \Theta(x, y) \}$$

$$T(s) = 1 + \frac{3(s-s_0)}{M_\eta^2 - M_\pi^2} \quad (3.11)$$

and assume that with a suitable normalization of the constant N (such as, e.g., $N = A(s_0, s_0, s_0)$) the function $\theta(x, y)$ satisfies $|\theta(x, y)|^2 \ll 1$ over the entire Dalitz plot [the corrections of order p^2 to the current algebra prediction indeed have this property, with $|\theta(x, y)|^2 < 0.07$].

Dropping terms of order $|\theta|^2$, the normalized distribution $M^2(y)$ defined in section 3b becomes

$$M^2(y) = \left(\frac{1 + \frac{1}{2}ay}{1 - \frac{1}{2}a} \right)^2 \{ 1 + 2\bar{\Theta}(y) \} \quad (3.12)$$

with

$$\begin{aligned} \bar{\Theta}(y) &= \bar{\Theta}(y) - \bar{\Theta}(-1) \\ \bar{\Theta}(y) &= \frac{1}{2x_1(y)} \int_{-x_1(y)}^{x_1(y)} dx \operatorname{Re} \Theta(x, y) \end{aligned} \quad (3.13)$$

One easily verifies that the projection $\bar{\Theta}(y)$ of the function $\theta(x, y)$ also fixes the branching ratio r (again dropping terms of order $|\theta|^2$):

$$\begin{aligned} r &= 1.51 \left\{ 1 - \int_{-1}^{y_1} dy f_1(y) \bar{\Theta}(y) + \int_{-1}^{y_2} dy f_2(y) \bar{\Theta}(y) \right\} \\ f_1(y) &= 2x_1(y) T^2(s) / \int_{-1}^{y_1} dy' x_1(y') T^2(s') \\ f_2(y) &= 2x_2(y) T(s) / \int_{-1}^{y_2} dy' x_2(y') \end{aligned} \quad (3.14)$$

In this relation we have kept track of the unequal size of phase space in the two channels: $x = \pm x_1(y)$ denotes the boundary of the Dalitz plot for the decay $\eta \rightarrow \pi^0 \pi^+ \pi^-$, whereas $x = \pm x_2(y)$ is the boundary of the physical region for the neutral decay mode. [Strictly speaking, the integrand of the second term in the curly brackets involves the mean of the function $\theta(x, y)$ from $-x_2(y)$ to $+x_2(y)$;

since the amplitude only very weakly depends on x , this difference is however numerically not significant. Note that the functions $f_1(y)$ and $f_2(y)$ satisfy

$$\int_{-1}^{y_1} dy f_1(y) = \int_{-1}^{y_2} dy f_2(y) = 2 \quad (3.15)$$

[We have checked that the approximate formula (3.14) correctly reproduces the value $r = 1.43$, if $\bar{\theta}(y)$ is calculated with the amplitude (2.1).] A measurement of the Dalitz plot distribution $M^2(y)$ therefore implies a specific value of the branching ratio r . In particular, if the distribution agrees with the one predicted by current algebra ($\bar{\theta}(y) = 0$), then the value of r must coincide with the current algebra prediction $r = 1.51$. Conversely, since the experimental information on r implies $r < 1.51$, the Dalitz plot distribution cannot coincide with the current algebra prediction.

Using the experimental distribution given by Layter et al. or by Gormley et al. to calculate the quantity $\bar{\theta}(y)$ from (3.12) and inserting the result in (3.14) one obtains $r = 1.44 \pm 0.02$, in good agreement with the value $r = 1.43$ required by our low energy representation. (One would have to distort the Dalitz plot distribution beyond recognition to obtain a value of r as low as 1.36).

4. Summary and conclusions

(i) We have shown that the corrections of order p^2 to the low energy theorems of current algebra for the decays $\eta \rightarrow \pi^0 \pi^+ \pi^-$ and $\eta \rightarrow 3\pi^0$ can unambiguously be calculated in terms of measured low energy observables. The calculation includes $\pi^0 \eta$ mixing as well as mixing with the η' (the η' contribution is contained in the term proportional to $4 M_K^2 - 3 M_\eta^2 - M_\pi^2$, which occurs in W).

(ii) The corrections of order p^2 turn out to be large; they increase the current algebra prediction (66 eV) for the rate $\Gamma_{\eta \rightarrow \pi^0 \pi^+ \pi^-}$ to 160 eV, i.e. by a factor 2.4. A substantial part of the correction is due to $I = 0$, S-wave $\pi\pi$ final state interactions, which current algebra predicts to be attractive and which therefore enhance the rate.

(iii) The final state interaction theorem implies that, in the case of two particle final states, the unitarity corrections exponentiate. Correspondingly, we expect the contributions of order p^4 , which we have neglected, to amplify the calculated corrections. We estimate the theoretical uncertainties due to these

higher order contributions at 50 eV in the rate. The overall fit to the data, provided by the Particle Data Group, $\Gamma_{\eta \rightarrow \pi^0 \pi^+ \pi^-} = 197 \pm 29$ eV, is in good agreement with our result.

(iv) Instead of using Dashen's theorem to determine the quantity $(M_{K^0}^2 - M_{K^+}^2)_{\text{QCD}}$, which normalizes the amplitude, one may use the relation established in III

$$(M_{K^0}^2 - M_{K^+}^2)_{\text{QCD}} = \mathcal{H} \frac{M_K^2}{M_\pi^2} (M_K^2 - M_\pi^2) \{ 1 + O(M^2) \}$$

$$\mathcal{H} \equiv \frac{m_d - m_u}{m_s - \hat{m}} \cdot \frac{2\hat{m}}{m_s + \hat{m}}$$

[Note that mass formulae like $(M_{K^0}^2 - M_{K^+}^2)_{\text{QCD}} = M_\pi^2(m_d - m_u)/(m_d + m_u) = (M_K^2 - M_\pi^2)(m_d - m_u)/(m_s - \hat{m})$ only hold at leading order. At the accuracy at which we are working here, these relations are not valid.] Comparing the prediction $\Gamma_{\eta \rightarrow \pi^0 \pi^+ \pi^-} = 160 \pm 50$ eV with the measured rate, we obtain

$$\mathcal{H} = (1.9 \pm 0.3) \cdot 10^{-3}$$

to be compared with the value $\kappa = (1.77 \pm 0.19) \cdot 10^{-3}$, which follows from the ratios $(m_s - \hat{m})/(m_d - m_u) = 43.5 \pm 2.2$, $m_s/\hat{m} = 25.0 \pm 2.5$ given in I. This shows that η decay does offer a nontrivial check on the validity of Dashen's theorem.

(v) A recent experiment (Weinstein et al. 1983) yields $\Gamma_{\eta \rightarrow 2\gamma} = 0.56 \pm 0.16$ keV, a result which is considerably higher than the rate $\Gamma_{\eta \rightarrow 2\gamma} = 0.324 \pm 0.046$ keV (Browman et al. 1974) quoted by the Particle Data Group (the rate of the decay $\eta \rightarrow 2\gamma$ is of crucial importance, because it normalizes the decay rates of all other channels). With the known branching ratios the new value implies $\Gamma_{\eta \rightarrow \pi^0 \pi^+ \pi^-} = 340 \pm 100$ eV. In view of the large error bar this result is not inconsistent with the value quoted above. Should the new value however be confirmed by more accurate measurements, the old problem which we claim to solve would emerge again: the result of our analysis, $\Gamma = 160 \pm 50$ eV, is not consistent with $\Gamma = 340$ eV.

(vi) Despite the fact that the overall corrections are large, the corrections to the normalized Dalitz plot distribution are rather small, because the main feature of this distribution is an $SU(2) \times SU(2)$ Adler zero near $s \sim M_\pi^2$, whose

presence is clearly borne out by experiment. Since the corrections are consistent with chiral symmetry, they do not affect this property of the distribution.

(vii) We have established a relation between the Dalitz plot distribution of the charged decay mode and the branching ratio

$$r = \frac{\Gamma_{\eta \rightarrow 3\pi^0}}{\Gamma_{\eta \rightarrow \pi^0 \pi^+ \pi^-}}$$

With the observed distribution, this relation predicts $r = 1.44 \pm 0.02$, to be compared with the value $r = 1.36 \pm 0.06$ which follows from the experimental information on the various branching ratios.

(viii) The current algebra prediction is $r = 1.51$. [In contrast to the current algebra prediction for the rate, which involves $SU(3) \times SU(3)$, the prediction for the branching ratio r follows if $A(s,t,u)$ is linear in s, t, u and has an $SU(2) \times SU(2)$ Adler zero near $s \simeq M_\pi^2$.] The corrections of order p^2 decrease the branching ratio to $r = 1.43$, in good agreement with the experimental information mentioned in the preceding paragraph.

(ix) The decay $\eta \rightarrow 3\pi$ is a model case of an amplitude for which the low energy theorems of current algebra are not stable with respect to the perturbations generated by the quark masses. We have shown that these perturbations are responsible for the flat disagreement between current algebra and experiment and conclude that the experimental information on this decay confirms the low energy predictions which follow from broken chiral symmetry.

Appendix: Extracting the scattering matrix from the generating functional

In this appendix we give a convenient recipe for extracting the scattering amplitudes from the generating functional. For definiteness we consider the T matrix elements involving four on-shell mesons (this covers quasi-elastic scattering as well as the decay $\eta \rightarrow 3\pi$).

To calculate the scattering matrix we determine the on-shell residue of the corresponding singularity in the four-point function $\langle 0 | T A_{\mu 1} \dots A_{\mu 4} | 0 \rangle$. This Green's function is the coefficient of $a_{\mu 1} \dots a_{\mu 4}$ in the expansion of the generating functional in powers of the external field $a_{\mu}(x)$. We make use of the fact that contributions which do not contain a pole in each one of the four legs do not contribute on-shell - as we will show below this simplifies the calculation considerably (if one wants to check the Ward identities directly, the information contained in the on-shell scattering amplitude does however not suffice - in the Ward identities the scattering amplitude is balanced against various contact terms).

We switch the external fields $v_{\mu}(x)$ and $p(x)$ off and set the scalar field $s(x)$ equal to the quark mass matrix - only the external field $a_{\mu}(x)$ is retained. The generating functional depends on this field both explicitly and implicitly, through the meson field $U = \exp i\phi$, which is fixed by a_{μ} through the equation of motion

$$Z = Z(a, \varphi) ; \quad \varphi = \varphi[a] \quad (A.1)$$

As was discussed in section 10 of ref. III we may replace the equations of motion associated with the leading order effective Lagrangian by the condition

$$\frac{\delta Z}{\delta \varphi^a(x)} = 0 \quad (A.2)$$

We expand Z in powers of a_{μ} and ϕ

$$Z(a, \varphi) = Z^{(0)} + Z^{(2)} + Z^{(4)} + \dots \quad (A.3)$$

where $Z^{(0)}$ is an irrelevant constant. $Z^{(2)}$ is the quadratic form

$$Z^{(2)}(a, \varphi) = \frac{1}{2} \int d^4x \sum_{a,b} \{ A^{ab} (\partial_\mu \varphi^a - a_\mu^a) (\partial^\mu \varphi^b - a^{\mu b}) - B^{ab} \varphi^a \varphi^b \} \quad (A.4)$$

and $Z^{(4)}$ is a (nonlocal) fourth order polynomial. More explicitly, the equation of motion becomes

$$A^{ab} \square \varphi^a + B^{ab} \varphi^b = A^{ab} \partial_\mu a^{\mu b} + \frac{\delta Z^{(4)}}{\delta \varphi^a} + \dots \quad (A.5)$$

This equation can be solved by iteration,

$$\varphi = \varphi^{(1)} + \varphi^{(3)} + \dots \quad (A.6)$$

where $\varphi^{(1)}$ is of order a_μ , $\varphi^{(3)}$ is of order $[a_\mu]^3$ etc. Explicitly, $\varphi^{(1)}$ is given by

$$\varphi^{(1)} = (A \square + B)^{-1} A \partial_\mu a^\mu \quad (A.7)$$

In fact, the correction $\varphi^{(3)}$ is not needed. Since we are evaluating the generating functional at its extremum, the difference $Z(a, \varphi) - Z(a, \varphi^{(1)})$ is of order $[\varphi^{(3)}]^2 \sim [a_\mu]^6$ and does therefore not contribute to the four-point function $[\varphi^{(3)}]$ e.g. generates a one particle reducible contribution to the scattering amplitude involving six mesons].

To calculate the four-point function, it thus suffices to expand $Z(a, \varphi^{(1)})$ to fourth order in a_μ . Since $\varphi^{(1)}$ is a linear functional of a_μ , the quadratic piece $Z^{(2)}$ of the generating functional does not contribute - the relevant piece is $Z^{(4)}(a, \varphi)$ where φ is to be replaced by $\varphi^{(1)}$.

Next, we observe that the poles in the square of the external four momenta are generated by the propagator $(A \square + B)^{-1}$, which determines the solution $\varphi^{(1)}$ in terms of the external field: only terms in $Z^{(4)}(a, \varphi)$ which involve four meson fields contribute to the T matrix.

As was shown in III the propagator is diagonalized by the matrix F which collects the pseudoscalar decay constants

$$\langle 0 | A_\mu^a | P, p \rangle = i p_\mu F^P_a$$

$$A \square + B = F^+ (\square + M^2) F \quad (A.8)$$

(The eigenvalues of the diagonal matrix M are the physical masses.) The solution (A.7) therefore takes the form

$$\varphi^{(1)} = F^{-1} (\square + M^2)^{-1} F \partial_\mu a^\mu \quad (A.9)$$

The quantity $F \partial_\mu a^\mu$ is the factor $\langle 0 | A_\mu^a | P, p \rangle a^\mu$ which occurs with each one of the four legs of the four-point function. To obtain the T matrix element we simply have to drop this factor, together with the pole term $(\square + M^2)^{-1}$ in (A.9). In other words, the wave function renormalization of the external legs is given by the physical decay constants.

This amounts to the following simple recipe:

(i) Switch all external fields off, including the field $a_\mu(x)$, and consider the generating functional as an effective action for the meson field :
 $Z(0, \phi) = Z(\phi)$.

(ii) The normalization of the meson field is fixed by the quadratic terms in $Z(\phi)$: determine the matrix F such that the transformation

$$\varphi(x) = F^{-1} \pi(x) \quad (A.10)$$

brings the quadratic terms in $Z(\phi)$ to normal form

$$Z(\phi) = \frac{1}{2} \int d^4x \sum_P \left\{ \partial_\mu \pi^P(x) \partial^\mu \pi^P(x) - M_P^2 \pi^P(x) \pi^P(x) \right\} + O(\pi^4) \quad (A.11)$$

(If the masses and decay constants have not been worked out before, they are determined in this step.)

(iii) The T matrix may now be extracted from the terms of order π^4 in the standard manner, treating the functional Z as the renormalized effective action of the system.

We add a remark concerning the fact that, even within pure QCD, the η is not a stable particle. The two-point function of the current A_μ^8 , e.g., does not contain an isolated pole on the real axis, but contains a continuum starting at $s = 9 M_\pi^2$, which develops a pole on the second sheet at $s = (M_\eta - \frac{i}{2} \Gamma_{\eta \rightarrow 3\pi})^2$. The above recipe concerns the residue of a pole at $s = M_\eta^2$, when in reality there is no pole there - how can this recipe be right ?

The point is that the instability of the η manifests itself in the analytic structure of the Green's functions only if the low energy expansion is carried beyond one loop. The pole is driven away from the real axis by the virtual transition $\eta \rightarrow 3\pi \rightarrow \eta$; in the low energy expansion, the leading contribution to this transition is given by the two loop self energy graph depicted in Fig. 2. Two loop graphs contribute to the generating functional at order p^6 . Since our calculation only includes the contributions of order p^4 , it is not sensitive to the precise structure of the η singularity; at this order in the low energy expansion the η propagator is adequately represented by a pole on the real axis.

This can also be seen directly, by looking at the expansion of $M_\eta - \frac{i}{2} \Gamma_{\eta \rightarrow 3\pi}$ in powers of the quark masses. Modulo logarithms the expansion of M_η is of the form $M^{1/2} + M^{3/2} + M^{5/2} + \dots$. One loop graphs contribute at order $M^{3/2}$, two loop graphs affect the expansion at order $M^{5/2}$. The leading contribution to the decay width (current algebra) is of order $(m_u - m_d)^2 m_s^{1/2}$ (times a function of Λ and of quark mass ratios; the factor $m_s^{1/2} \sim M_\eta$ arises from phase space). The imaginary part of the complex mass is therefore of the same order as the two loop contributions to the real part. To extract the decay amplitude from the expression for the Green's function we need the position of the pole only to one loop accuracy.

Bibliography

- Bell, J.S. and Sutherland, D.G., Nucl.Phys. B4 (1968) 315.
- Becherrawy, T. and Nasrallah, N.F., Nuovo Cim. 46A (1978) 17.
- Bose, S.K. and Zimerman, A.H., Nuovo Cim. 43A (1966) 1165; Errata: Nuovo Cim. 45A (1966) 1056.
- Browman, A. et al., Phys.Rev.Lett. 32 (1974) 1067.
- Cantor, A.J., Phys.Rev. D3 (1971) 3195, 3205.
- Cicogna, G., Strocchi, F. and Vergara Caffarelli, R., Lett. Nuovo Cim. 10 (1974) 25.
- Crewther, R.J., Asymptotic behaviour in quantum field theory, in: Weak and Electromagnetic Interactions at High Energies (Cargèse 1975) Part A, eds. M. Lévy, J.L. Basdevant, D. Speiser and R. Gastmans, NATO Advanced Study Institute Series, Vol 13a (Plenum, New York, 1976) pp. 345-457.
- Dashen, R., Phys.Rev. 183 (1969) 1245.
- Dittner, P., Dondi, P.H. and Eliezer, S., Phys.Rev. D8 (1973) 2253.
- Dominguez, C.A. and Zepeda, A., Phys.Rev. D18 (1978) 884.
- Gasser, J. and Leutwyler, H., Phys.Rep. C87 (1982) 77 (this reference is quoted as I);
Phys.Lett. 125B (1983) 321, 325; "Chiral perturbation theory to one loop", Preprint CERN TH-3689 (1983), to be published in Ann.of Phys. (these references are quoted as II);
"Chiral perturbation theory: Expansion in the mass of the strange quark", Preprint CERN TH-3798 (1983) (this reference is quoted as III);
"Electromagnetic and weak interactions in chiral perturbation theory", in preparation (this reference is quoted as IV).
- Gormley, M. et al., Phys.Rev. D2 (1970) 501.
- Hudnall, W. and Schechter, J., Phys.Rev. D9 (1974) 2111.
- Kawarabayashi, K. and Ohta, N., Nucl.Phys. B175 (1980) 477.
- Khuri, N.N. and Treiman, S.B., Phys.Rev. 119 (1960) 1115.

Kogut, J. and Susskind, L., Phys.Rev. D11 (1975) 3594.

Langacker, P. and Pagels, H., Phys.Rev. D10 (1974) 2904.

Layter, J.G. et al., Phys.Rev. D7 (1973) 2565.

Leutwyler, H. and Roos, M., "On the elements V_{us} and V_{ud} of the Kobayashi-Maskawa matrix", Preprint CERN TH-3830 (1983), submitted to Z.Phys.C.

Llanta, E. and Tarrach, R., Ann.Phys. 141 (1982) 116.

Matsubara, Y. and Tameike, S., Progr.Theor.Phys. 57 (1977) 1679.

Milton, K.A., Palmer, W.F. and Pinsky, S.S., Phys.Rev. D22 (1980) 1124, 1647.

Minkowski, P., Phys.Lett. 116B (1982) 373.

Nagels, M.M. et al., Nucl.Phys. B147 (1979) 189.

Osborn, H. and Wallace, D.J., Nucl.Phys. B20 (1970) 23.

Particle Data Group, Review of particle properties, Phys.Lett. 111B (1982)

Raby, S., Phys.Rev. D13 (1976) 2594.

Roiesnel, C. and Truong, T.N., Nucl.Phys. B187 (1981) 293; CPT, Ecole Polytechnique Preprint A515-0982 (Sept. 1982).

Schechter, J. and Ueda, Y., Phys.Rev. D3 (1971) 2874; D4 (1971) 733.

Sutherland, D.G., Phys.Lett. 23 (1966) 384.

Truong, T.N., "Departure from chiral symmetry: Role of unitarity and analyticity", Lecture given at the XVIII Cracow School of Theoretical Physics, Zakopane, June 1983, and at the Asia Pacific Conference, Singapore, June 12-18, 1983.

Weinberg, S., Phys.Rev. D11 (1975) 3583.

Weinstein, A. et al., Phys.Rev. D28 (1983) 2896.

Wilson, K., Phys.Rev. 179 (1969) 1499.

Figure captions

Fig. 1: Normalized Dalitz plot distribution, defined in (3.4), (3.5). Curve 1 is the current algebra prediction (first term in the low energy expansion). The curve labelled 2 displays the distribution resulting from the low energy expansion to second order. L and G are the experimental distributions of Layter et al. and Gormley et al. respectively.

Fig. 2: The leading contribution to the virtual transition $\eta \rightarrow 3\pi \rightarrow \eta$. This transition drives the pole in the two-point function of the current A_μ^8 away from the real axis onto the second sheet.

Fig.1

Fig.2