LAPP-TH-108
May, 1984

Missing energy sources at large p_T from standard QCD processes

P. Aurenche
LAPP, Annecy-le-Vieux, France

and

R. Kinnunen *)
LAPP, Annecy-le-Vieux, France
CERN, Geneva, Switzerland

A B S T R A C T

A detailed analysis is given of processes involving the production of heavy vector bosons as sources of events with a single hadronic jet balanced by a large missing energy or possibly a charged large p_T lepton and missing energy. In particular, the hadronic channels of the τ lepton produced in $W^\pm$ decays provide an appreciable rate of monojet events, with energetic narrow jets of low charged multiplicity. Furthermore it is shown that higher order QCD corrections to weak boson production, in particular the Compton like process $qg \rightarrow Zq$ or $qg \rightarrow Wq$ are an appreciable source of the types of events mentioned above.

*) On leave from the Department of High Energy Physics, University of Helsinki, Helsinki, Finland.

Events with large missing transverse energy at the SPS collider have recently raised much interest both experimentally and theoretically. The primary source of "missing energy" is of course the production of the weak boson $W^\pm$ and its subsequent decay into the channels $W \rightarrow e \nu_e$ or $W \rightarrow \mu \nu_\mu$ ^{1,2)}. However several events have been reported which could not, in an obvious way, be attributed to known processes in the standard theory. The UA1 collaboration has found seven events with a missing transverse energy in excess of 40 GeV associated either to a narrow hadronic jet (five events) or to a neutral electromagnetic cluster³⁾ and the UA2 collaboration has reported about three events with a missing transverse energy larger than 50 GeV associated to a large p_T electron and hadronic jets⁴⁾. On the theoretical side, models based on supersymmetry and/or supergravity⁵⁾ offer a great variety of ways to generate events with a large missing energy via the production of the heavy partners of the known particles which subsequently decay with the emission of an undetectable photino for example. In some cases, such supersymmetric events could produce the experimentally observed topologies. In this paper we adopt a more pragmatical attitude and analyse in detail the standard model processes which lead to final state neutrinos as a source of missing energy. We do not consider the production and semi-leptonic decays of heavy flavours: they tend to give a small missing transverse energy and have a rich jet structure which would set them in a different category. Our main concern will be the process

$$\begin{aligned}
 p \bar{p} &\rightarrow W X \\
 &\searrow \tau \nu_\tau \\
 &\quad \downarrow \\
 &\quad \left\{ \begin{array}{l} e \nu_e \nu_\tau \\ \text{hadrons} + \nu \end{array} \right.
 \end{aligned}
 \tag{1}$$

The production of τ is identical to that of the direct electron or muon from the weak boson decay. The calculation for the latter process has already been presented⁶⁾ and we use, for τ production, the same mechanism which takes into account higher order QCD corrections. Various decay channels of the τ lepton which make up 80% of the total width will be studied in detail:

$$\begin{array}{ll}
 \tau \rightarrow e \nu_e \nu_\tau & B_r^e = 17 \% \\
 \tau \rightarrow \pi \nu_\tau & B_r^\pi = 11 \% \\
 \tau \rightarrow \pi \bar{\pi} \nu_\tau & B_r^{2\pi} = 22 \% \\
 \tau \rightarrow \pi \pi \pi \nu_\tau & B_r^{3\pi} = 14 \%
 \end{array} \quad (2)$$

Polarization effects arising from the V-A structure of the weak current will be kept throughout as it was shown that they have significant effects on the final particle spectra⁷⁾. For each channel, transverse momentum and angular distributions will be given. Experimentally related distributions summing up over all considered channels will also be given and comparison to the events with large missing transverse energy will be made: the possible identification of the jets observed in such events³⁾ with that made up of the hadronic debris of the τ lepton will be discussed.

The other source of missing p_T is naturally the process

$$\begin{array}{c}
 p \bar{p} \rightarrow Z_0 X \\
 \quad \quad \quad \searrow \\
 \quad \quad \quad \nu \bar{\nu}
 \end{array} \quad (3)$$

The higher order QCD corrections, by the emission of a large p_T parton, recoiling from the weak boson will obviously give rise to "monojet" events³⁾. Quantitative estimates for this process will be presented. Related to this mechanism is the reaction

$$\begin{array}{c}
 p \bar{p} \rightarrow W \text{ jet } X \\
 \quad \quad \quad \searrow \\
 \quad \quad \quad e \nu_e
 \end{array} \quad (4)$$

which leads to events involving a lepton and a hard jet associated to a large missing transverse energy such as those observed by the UA2 collaboration⁴⁾ and it will also briefly be considered.

I. The process $p\bar{p} \rightarrow WX$, $W \rightarrow \tau\nu_\tau$

The direct production of a lepton in hadronic collisions has already been described extensively in previous studies^{6,7)} and we shall not repeat the discussion here. It suffices to say that besides the basic Drell-Yan mechanism we consider the QCD corrections associated to soft gluon emission, using the well known impact parameter space formalism⁸⁾, and the hard QCD corrections which are treated perturbatively to first order. The hard corrections contain two types of terms: the annihilation diagrams, $q\bar{q} \rightarrow Wg$ and the Compton like diagrams, $qg \rightarrow Wq$. Although the Compton process contributes only about 8% of the total cross-section it dominates the production of the weak boson at p_T values above ~ 25 GeV.

A massless lepton is produced from W decay in a pure helicity state (left handed for leptons, right handed for antileptons) owing to the V-A nature of the weak current. Taking the lepton mass into account will induce a small admixture of the unfavoured helicity component. For the Drell-Yan process alone the ratio of the unfavoured component to the favoured component has been estimated to be⁷⁾ $1/2 (m_\tau/m_W)^2$, which is very small ($\approx 2.5 \cdot 10^{-4}$) in our case. This figure also holds true for the full calculation as has been checked by the numerical evaluation of the complete formulae. All properties of the τ production mechanism (e.g. the transverse momentum and the angular distribution of the lepton) are very similar to direct e or μ production. In the following study of τ decays, we shall neglect, for simplicity, the contribution of the unfavoured component. All our predictions for the proton-antiproton collisions at $\sqrt{s} = 540$ GeV are calculated using the Glück-Hoffmann-Reya parametrization of the parton densities.

II. The leptonic decay of the τ lepton

For definiteness we consider the channel

$$\tau^+ \rightarrow e^+ \nu_e \bar{\nu}_\tau$$

with the experimental branching ratio $B_\tau^e = 17\%$. All final state particles are taken to be massless. We recall here the expression for the exclusive decay of a heavy polarized antilepton in its rest frame⁷⁾:

$$|D_\lambda|^2 = C E_\nu (m - 2E_\nu) (1 - \lambda \cos \theta_\nu) \quad (5)$$

The electron neutrino has an energy E_ν and its polar angle is defined with respect

to the direction given by the τ momentum in the laboratory. The quantities m and λ are respectively the mass and the helicity ($\lambda = \pm 1$) of the τ^+ . The constant factor appearing in Eq.(5) is left arbitrary since the normalization of our prediction is obtained from the value of the branching ratio. In the case considered here, λ is equal to $+1$ and consequently the neutrino ν_e will be distributed dominantly in the direction opposite to that of the τ lepton. Because of kinematical constraints the positron tends to be somewhat aligned along the τ momentum: an analytic calculation of the inclusive positron spectrum yields the distribution $(1 + \frac{1}{3} \cos \theta_e)$ in the τ rest frame¹⁰⁾. In Fig. 1 we show the transverse momentum and angular distributions of the positron, the decay product of the τ^+ . We also show the corresponding spectra for the τ^+ itself which are in fact almost identical to that of a directly produced e^+ or μ^+ . The p_T -spectra are understandably quite different ($\langle p_T^e \rangle_\tau \sim \langle p_T^{\text{miss}} \rangle_\tau \sim 12$ GeV compared to $\langle p_T^e \rangle_{\text{direct}} \sim \langle p_T^{\text{miss}} \rangle_{\text{direct}} \sim 30$ GeV) whereas the angular distributions appear to be identical because of the collimation, in the τ decay, of the positron with the τ direction, which is due to the small mass of the unstable lepton compared to its momentum. Distinguishing between the two processes on an event to event basis is very difficult since they lead to the same topology: isolated electron at large p_T and a large missing transverse momentum sometimes associated to a QCD jet. The differences between the two processes appear best in the two-dimensional plot of the transverse momentum versus the polar angle of the lepton shown in Fig. 2. The striking feature is the strong asymmetry observed in the case of the directly produced lepton at the lower p_T values. It may be interesting to compare the probability of occurrence of a positron produced at a given p_T and a given rapidity for the two processes. The result of such an estimate is shown in Fig. 3 where the lines correspond to constant ratio of probabilities: the most probable domain for τ mediated processes is the area delimited by the curves and the horizontal rapidity axis. On the basis of the 52 $W \rightarrow e\nu$ events recorded by the UA1 collaboration¹⁾ one expects to observe, after taking account of the experimental cuts ($p_T^{\text{miss}} > 15$ GeV, no hadronic jet recoiling from the lepton within an angle of 30°), about 3 τ events where the τ decays leptonically.

III. Hadronic decays of τ : $\tau^+ \rightarrow \pi^+ \nu$

The matrix element for this process, the simplest of the hadronic decay channel of the τ is written as¹¹⁾

$$D^\pi = \bar{V}(l) \gamma_\mu (1 - \gamma_5) V(p_\nu) p_\pi^\mu \quad (6)$$

where ℓ , p_ν and p_π are the momenta of the τ^+ , $\bar{\nu}_\tau$ and π^+ respectively. As before we ignore possible normalization factors in our analytical expressions. Squaring Eq.(6) and projecting out states of definite τ polarization defined by the four-vector s one finds

$$|D^\pi|^2 = 2 p_\pi \cdot p_\nu p_\tau \cdot (\ell + m s) - m_\pi^2 p_\nu \cdot (\ell + m s) \quad (7)$$

In the τ rest frame and choosing the helicity basis where $s = \lambda(0,0,0,1)$, Eq.(7) can be written

$$\begin{aligned} |D_\lambda^\pi|^2 &= m^2 \left[(m^2 - m_\pi^2 - m_\nu^2) (E_\pi - \lambda |\vec{p}_\pi| \cos \theta_\pi) - m_\pi^2 (E_\nu + \lambda |\vec{p}_\pi| \cos \theta_\pi) \right] \\ &= \frac{m^4}{2} (1 - \lambda \cos \theta_\pi) \end{aligned} \quad (8)$$

where the last line, which is valid when all masses are neglected compared to the τ mass, could have been obtained directly using elementary helicity conservation arguments. Equation (8) shows that the pion tends to be produced in the direction opposite to that of the τ momentum in the laboratory. Hence a rather soft p_T spectrum is expected for the pion. On the contrary the $\bar{\nu}_\tau$ from τ^+ decay will tend to be quite hard and produced in the τ -direction thereby partially compensating the transverse momentum of the ν_τ pair-produced with τ^+ in the W -decay: quite a small missing p_T will be observed in such an event. We display in Figs 4 and 5 the π transverse momentum and the total missing transverse momentum spectra of the process.

IV. Hadronic decays of τ : $\tau^+ \rightarrow \rho^+ \bar{\nu}_\tau$, $\tau^+ \rightarrow A_1^+ \bar{\nu}_\tau$

Following previous studies¹¹⁾ we assume that the 2π and 3π decay channels of the τ decay are resonance dominated i.e.

$$\begin{aligned} \tau^+ &\rightarrow \rho^+ \bar{\nu}_\tau, \quad \rho^+ \rightarrow \pi^+ \pi^0 \\ \tau^+ &\rightarrow A_1^+ \bar{\nu}_\tau, \quad A_1^+ \rightarrow \pi^+ \pi^0 \pi^0 \text{ or } \pi^+ \pi^+ \pi^- \end{aligned} \quad (9)$$

Considering now the production of spin-one resonances we can treat the ρ and A_1 cases together. The matrix element is defined as:

$$D = \bar{v}(\ell) \gamma_\mu (1 - \gamma_5) v(p_\nu) V^\mu \quad (10)$$

where V^μ describes the polarization state of the vector resonance. The square of the decay matrix element for a τ lepton in a given helicity state can be expressed as

$$|D_\lambda^V|^2 = 4 \left[p_\nu \cdot V^\dagger (\ell + m_s) \cdot V + p_\nu \cdot V (\ell + m_s) \cdot V^\dagger + p_\nu \cdot (\ell + m_s) + i \epsilon_{\alpha\beta\gamma\delta} (\ell + m_s)^\alpha V^\dagger^\beta V^\gamma p_\nu^\delta \right] \quad (11)$$

After some algebra, the following results are obtained in the τ rest-frame, when we explicitly write out the contribution to various meson helicity states:

$$\begin{aligned} |D_\lambda^V(\lambda_V=0)|^2 &= \frac{m |\vec{p}_V|}{m_V^2} \left[2m(|\vec{p}_V| - \lambda E_V \cos \theta_V) + m_V^2 (1 + \lambda \cos \theta_V) \right] \\ |D_\lambda^V(\lambda_V=1)|^2 &= 2m |\vec{p}_V| (1 + \lambda \cos \theta_V) \\ |D_\lambda^V(\lambda_V=-1)|^2 &= 0 \end{aligned} \quad (12)$$

In Eqs (12) the τ -neutrino mass is neglected. The result of the last line is simply a consequence of spin conservation: only a particle of spin 3/2 or higher can decay into particles of spin 1/2 and spin 1 with opposite helicities. One observes that the vector meson in the helicity state $\lambda_V = +1$ tends to be harder than in the $\lambda_V = 0$ case because of the $(1 + \cos \theta_V)$ factor which collimates the resonance in the direction of the τ momentum in the former case. The rate of production in the $\lambda_V = 1$ state can easily be estimated to be

$$\frac{\Gamma(\lambda_V=+1)}{\sum_{\lambda_V} \Gamma(\lambda_V)} = \frac{2 m_V^2}{m^2 + 2 m_V^2} \quad (13)$$

It increases with the mass of the vector meson with the consequence that the

A_1 p_T -spectrum will be harder than the corresponding ρ -spectrum because of this dynamical effect. This is further enhanced by the well known kinematical effect whereby in a two-body decay with asymmetric masses the heavier particle carries more momentum. These effects are shown in Fig. 4. To illustrate quantitatively the spin and mass effects discussed above, we give below the average p_T -values of the hadronic "jets" from the decay of the τ :

$$\langle p_T^\pi \rangle = 12.1 \text{ GeV}$$

$$\langle p_T^S \rangle = 16.7 \text{ GeV}$$

$$\langle p_T^{A_1} \rangle = 22.8 \text{ GeV}$$

The hardness of the hadronic jets will reflect itself in the hardness of the missing energy spectrum: the mismatch between the jet and the missing transverse momentum is, in our formalism, due to the higher order QCD corrections. Therefore, the " A_1 jet" will be associated to a larger missing p_T than a " ρ -jet" and a fortiori a " π -jet" (see Fig. 5). The angular distribution, not shown, exhibits a roughly similar behaviour for all cases: the shape of the curves as well as the mean values are almost identical to the angular distribution of the τ itself reflecting the strong collimation of the hadronic debris with the τ momentum.

The properties of the τ -generated jets can further be studied by letting the resonances decay into stable particles. The $\rho \rightarrow 2\pi$ decay, described by the usual matrix element $V(\lambda_\rho) \cdot (p_1 - p_2)$, can easily be folded in Eq.(11) to give for the full process $\tau^+ \rightarrow \pi^+(p_1)\pi^0(p_2)\bar{\nu}_\tau(p_\nu)$:

$$\begin{aligned} |D_\lambda^{2\pi}|^2 &= \frac{m_\rho^2 \Gamma_\rho^2}{((p_1+p_2)^2 - m_\rho^2) + \Gamma_\rho^2 m_\rho^2} \left\{ 2((p_1-p_2) \cdot \ell)^2 + 2(p_1-p_2) \cdot p_\nu m_S \cdot (p_1-p_2) \right. \\ &\quad \left. - (p_1-p_2)^2 (m^2 - (p_1+p_2) \cdot \ell + m_S \cdot (p_1+p_2)) \right\} \\ &= \frac{m^2 m_\rho^2 \Gamma_\rho^2}{((p_1+p_2)^2 - m_\rho^2) + \Gamma_\rho^2 m_\rho^2} \left\{ 2(E_1 - E_2)^2 + E_\nu (m - 2E_\nu) \right. \\ &\quad \left. - \lambda (E_1 (m - 4E_2) \cos\theta_1 + E_2 (m - 4E_1) \cos\theta_2) \right\} \end{aligned} \quad (14)$$

The last line is valid in the τ rest frame, with the usual definition of the 4-vector s , assuming all stable particles massless. The pion p_T -spectrum is shown in Fig. 6 and it is only slightly softer than that of the single π produced from τ decay.

The A_1 decay into three pions is more complicated since two channels are possible with equal probability: $A_1 \rightarrow \rho^+ \pi^0(p_3)$, $\rho^+ \rightarrow \pi^+(p_1) \pi^0(p_2)$ and $A_1 \rightarrow \rho^0 \pi^+(p_3)$, $\rho^0 \rightarrow \pi^+(p_1) \pi^-(p_2)$. The usual decay matrix element is

$$\begin{aligned} D^{A_1 \rightarrow 3\pi} &= \epsilon_{\alpha\beta\gamma\delta} A_1^\alpha f^{\beta\gamma} p_1^\delta p_2^\delta p_3^\delta \\ &= \epsilon_{\alpha\beta\gamma\delta} A_1^\alpha p_1^\beta p_2^\gamma p_3^\delta \end{aligned} \quad (15)$$

when only stable particles are considered. The final expression $\tau \rightarrow 3\pi\nu$ decay is not very illuminating and will not be shown here. The resulting single charged π spectrum is compared in Fig. 6 with the predictions for the other channels.

V. Phenomenology of hadronic τ decays

After these lengthy theoretical premises we present, in this section, our results in a form more appropriate for comparison to experimental data. All the processes discussed above lead to a well collimated low multiplicity jet, the transverse momentum of which being, as we saw before, roughly balanced by the neutrino momenta. Following the recent presentation of the UA1 collaboration³⁾ we shall distinguish jets without charged tracks ("neutral electromagnetic clusters") from those with at least one charged track: in the latter case we shall also give predictions according to the charged multiplicity. All our plots are presented with the following cuts: $p_T^{\text{miss}} > 15 \text{ GeV}$ and $p_T^{\text{QCDjet}} < 15 \text{ GeV}$ where QCD jet denotes the jet recoiling from the weak boson in the higher order QCD production processes.

We show in Fig. 7, the transverse momentum spectrum of the hadronic jet, from τ decay, independent of its charged content: it has a broad shape extending to rather large p_T values. The hardest jets tend to have a higher charged multiplicity because they originate from A_1 decay (see Fig. 4). We find the following figures for the integrated cross-sections

$$\sigma(p_T^{\text{jet}} > 30 \text{ GeV}) = 42 \text{ pb}$$

$$\sigma(p_T^{\text{jet}} > 40 \text{ GeV}) = 5.1 \text{ pb}$$

For an integrated luminosity of $.113 \text{ pb}^{-1}$ this corresponds to about 5 events and .6 event respectively. These numbers are lower bounds on the jet production cross-section since we have left out the τ decay modes into four pions or more. Although these modes represent about 20% of the total τ width, they dominate the large p_T -tail of the distribution for the same kinematical reasons why the 3π system dominates over the 2π system. A rough estimate of their contribution (assuming a four pion mode dominated by the ρ' resonance) leads to a considerable increase of the expected number of events: we find

$$\sigma(p_T^{\text{jet}} > 30 \text{ GeV}) = 93 \text{ pb} \quad (10 \text{ events})$$

$$\sigma(p_T^{\text{jet}} > 40 \text{ GeV}) = 16 \text{ pb} \quad (1.8 \text{ event})$$

For comparison, the UAl collaboration³⁾ reports five to seven events at large p_T (excluded from the comparison are all events which would appear to contain a lepton, like event A of Ref. 3) for example). In the production mechanism discussed here, there is a rather strong correlation between the jet transverse momentum and the missing transverse momentum, a trend which is not in disagreement with the data.

A more exclusive analysis of the predictions can be made by considering the differential distribution in p_T for a given topology: an event is defined as a n -track jet when n charged pions carry each a transverse momentum above 1 GeV/c. Thus the π and ρ dominated τ decay will contribute only to one track jet for example whereas the $A_1^+ \rightarrow \pi^+\pi^+\pi^-$ channel will be seen as a one track event if two of the pions have a $p_T < 1 \text{ GeV/c}$. (In the absence of dynamical models for high multiplicity τ decays we consider in the following only the contribution of three pions or less.) The corresponding distributions are shown in Fig. 8. The high p_T tail ($p_T > 35 \text{ GeV}$) of one track events is dominated by π mediated τ -decay and therefore in this region the charged track should appear unaccompanied by neutral energy. We estimate the following cross-sections

$$\sigma(p_T^{1\text{ track}} > 30 \text{ GeV}) = 6.5 \text{ pb} \quad (.7 \text{ event})$$

$$\sigma(p_T^{1\text{ track}} > 40 \text{ GeV}) = .5 \text{ pb} \quad (.06 \text{ event})$$

where the figures in brackets are obtained for a luminosity $\mathcal{L} = .113 \text{ pb}^{-1}$. In Fig. 9 the correlation between the charged p_T and the total p_T is displayed (one pion events are not included): the accumulation of points along the diagonal is due to the channel where A_1 decays into three charged particles.

We consider now the possibility that only electromagnetic energy (i.e. π^0 's) is produced at large p_T . To estimate the cross-section of such a process we impose that no charged tracks has a transverse momentum greater than 1 GeV. The corresponding p_T spectrum is shown in Fig. 11 yielding the following cross-section estimates

$$\sigma(p_T^{\text{emg}} > 20 \text{ GeV}) = 4 \text{ pb} \quad (.5 \text{ event})$$

$$\sigma(p_T^{\text{emg}} > 30 \text{ GeV}) = 1.5 \text{ pb} \quad (.2 \text{ event})$$

$$\sigma(p_T^{\text{emg}} > 40 \text{ GeV}) = .2 \text{ pb} \quad (.02 \text{ event})$$

These are to be compared with the one or two "isolated photon" events seen by the UA1 collaboration, where in both cases the "photon" transverse energy is greater than 40 GeV. As can be seen the rate for the τ decay is too small to explain these events. However, it should be recalled that not all hadronic decay modes of the τ have been included in our predictions: in particular the channel $\tau \rightarrow 3(\pi^\pm) + \text{neutrals} + \nu$, which has a branching ratio of about 20%, and which we have not considered here, will increase the production rates given above.

We have not tried in this work to perform a detailed comparison with the available data. Instead we wanted to show that the hadronic decay modes of the τ lepton produced in W decays should be considered as serious candidates to explain some or all the reported "monojet" events. Folding in the acceptance of the detector will probably decrease the estimates reported here. On the other hand, the neglected hadronic channels will definitely have the opposite effect (specially in what concern the high p_T tail of the jet distributions).

VI. Other processes as source of high p_T jets

We discuss in this section alternate sources of generating events with a large missing energy together with a large p_T hadronic jet. The most obvious one is of course the process

$$p\bar{p} \rightarrow Z_0 \text{ jet } X$$

$$\quad \quad \quad \searrow \nu \bar{\nu}$$
(16)

The crucial observation is that the branching ratio of Z_0 into neutrinos is much greater than that into e^+e^- for example: the vector part of the neutral current coupling to charged leptons almost vanishes whereas it is equal to the axial vector part for neutrinos. Furthermore at least three neutrino types can be produced in Z_0 decay so that one expects a factor 6 for the decay $Z_0 \rightarrow \nu\bar{\nu}$ compared to the channel $Z_0 \rightarrow e^+e^-$. The jet produced in association with the weak boson arises from the first order QCD subprocesses

$$q\bar{q} \rightarrow Z_0 q$$

$$qg \rightarrow Z_0 q$$
(17)

We have estimated the following cross-sections for the production of a jet at large p_T in reaction Eq.(16)

$$\sigma(p_T^{\text{jet}} > 20 \text{ GeV}) = 21 \text{ pb} \quad (.24 \text{ event})$$

$$\sigma(p_T^{\text{jet}} > 30 \text{ GeV}) = 7.6 \text{ pb} \quad (.9 \text{ event})$$

$$\sigma(p_T^{\text{jet}} > 40 \text{ GeV}) = 3.2 \text{ pb} \quad (.4 \text{ event})$$

$$\sigma(p_T^{\text{jet}} > 50 \text{ GeV}) = 1.5 \text{ pb} \quad (.2 \text{ event})$$

These figures do not depend on the particular way the soft gluon emission is dealt with (it would affect only the low p_T jet spectrum) but it results from the straightforward evaluation of the subprocesses in Eq.(17). Because of its much harder p_T dependence the Compton process dominates entirely the cross-section

for p_T^{jet} above 30 GeV.

Related to this mechanism is the process

$$p \bar{p} \rightarrow W^\pm \text{jet } X \quad (18)$$

which leads to events with an electron, a hadronic jet and missing momentum all at large p_T . Three such events have been observed by the UA2 collaboration⁴⁾. One notes that the p_T values of the electron in these events are 18, 22 and 34 GeV/c respectively compatible with $W \rightarrow e\nu$ decay: we calculate the average lepton transverse momentum to be $\langle p_T^l \rangle = 30$ GeV/c. More striking are the large p_T values of the jets: 39 ± 4 , 67 ± 7 and 66 ± 6 GeV/c. Our estimates for the probability of such events are as follows

$$\sigma(p_T^{\text{jet}} > 30 \text{ GeV}) = 8.3 \text{ pb} \quad (.96 \text{ event})$$

$$\sigma(p_T^{\text{jet}} > 40 \text{ GeV}) = 2.8 \text{ pb} \quad (.33 \text{ event})$$

$$\sigma(p_T^{\text{jet}} > 50 \text{ GeV}) = 1.1 \text{ pb} \quad (.13 \text{ event})$$

$$\sigma(p_T^{\text{jet}} > 60 \text{ GeV}) = .35 \text{ pb} \quad (.04 \text{ event})$$

The event numbers are based on an integrated luminosity $\int d\mathcal{L} = .116 \text{ pb}^{-1}$. No acceptance cut is folded in our predictions. The remarks concerning the production of Z^0 at large p_T apply equally well here. We would like to stress that the figures presented above are on underestimate of the QCD predictions calculated to all order. It is by now well known that the higher order corrections to particle production at large p_T increase the predictions obtained at the Born level¹²⁾: here the diagrams we have considered correspond to the Born terms for the production of the weak boson at large p_T and the higher order corrections would be related to the production of two or more hadronic jets. An indication of the importance of such processes can be seen in the rather rich jet structure of the events reported by the UA2 collaboration⁴⁾.

Conclusions

We have shown that the weak boson decays, sometimes associated to a hard QCD jet, can produce the topologies of the "monojet" events observed at the SPS collider. In particular the hadronic final states of the τ lepton from W decay lead to narrow energetic jets with few particles and a substantial neutral component approximately balanced by a large missing transverse energy. Applying a simple cut $p_T^{\text{jet}} > 30 \text{ GeV}$ we find about 10 "monojet" events corresponding to the integrated luminosity of 113 nb^{-1} . Such effects like triggering and detection efficiencies would of course somewhat reduce the calculated rate. Particularly important for the jet production at large p_T are the τ decay modes into three pions or more. It was also found that the τ final states with π^0 's, especially the ρ and A_1 dominated channels, $\pi^\pm \pi^0 \nu$ and $\pi^\pm \pi^0 \pi^0 \nu$, can simulate the "isolated photon" events in the case the charged pion is too slow for observation. However we find only a small contribution, about 0.2 events above $p_T^{\text{jet}} > 30 \text{ GeV}$, for the electromagnetic jets from this mechanism.

An important source for the "monojet" events extending beyond the kinematical range of the weak boson decays are the Z^0 decays to neutrinos in the presence of a hard QCD jet. We find about half of event for $p_T^{\text{jet}} > 40 \text{ GeV}$ based on the well defined first order QCD processes for Z^0 production. Again the predicted number is small to be confronted with the two events of the UA1 collaboration at very large transverse momenta. Our prediction, however, is underestimated by the neglect of the higher order (> 1) corrections. Such processes could produce weak bosons associated to more than one jet, namely the type of events found by UA2 collaboration.

Acknowledgements

We thank J. Colas, M.N. Minard, S. Narison and C. Roiesnel for discussions.

REFERENCES

- 1) UA1 Collaboration, G. Arnison et al., Phys. Lett. 122B (1983) 103; 126B (1983) 398; CERN-EP/83-111.
- 2) UA2 Collaboration, G. Banner et al., Phys. Lett. 122B (1983) 476; P. Bagnaia et al., Phys. Lett. 129B (1983) 130; CERN-EP/84-39.
- 3) UA1 Collaboration, G. Arnison et al., CERN-EP/84-42.
- 4) UA2 Collaboration, P. Bagnaia et al., CERN-EP/84-40.
- 5) M.J. Herrero, L.E. Ibanez, C. Lopez and F.J. Yndurain, Phys. Lett. 132B (1983) 199;
R.M. Barnett, K.S. Lackner and H.E. Haber, Phys. Lett. 126B (1983) 64;
J. Ellis, J.S. Hagelin, D.V. Nanopoulos and M. Srednicki, CERN-TH-3572 (1983);
G. Altarelli, B. Mele and S. Petrarca, CERN-TH-3822 (1984).
- 6) P. Aurenche and R. Kinnunen, Phys. Lett. 135B (1984) 493.
- 7) P. Aurenche and R. Kinnunen, Annecy preprint LAPP-TH-104 (1984).
- 8) G. Parisi and R. Petronzio, Nucl. Phys. B154 (1979) 427;
P. Chiapetta and M. Greco, Nucl. Phys. B199 (1982) 77; B221 (1983) 269.
- 9) M. Glück, E. Hoffmann and E. Reya, Z. Phys. C13 (1982) 119.
- 10) P. Aurenche, R. Kinnunen and K. Mursula, Annecy preprint LAPP-TH-106 (1984).
- 11) S. Narison, Z. Physik C2 (1979) 11.
- 12) See for example, R.K. Ellis, M.A. Furman, H.E. Haber and I. Hinchliffe, Nucl. Phys. B173 (1980) 397;
P. Aurenche, R. Baier, A. Douiri, M. Fontannaz and D. Schiff, Annecy preprint LAPP-TH-96 (1983). This calculation concerns higher order corrections to direct photon at large p_T . The diagrams involved are similar to those involved in the calculation of the corrections to the weak boson production at large p_T . The K-factor was rather larger (~ 1.5) at $\sqrt{s} = 540$ GeV.

FIGURE CAPTIONS

- Fig. 1 a) Transverse momentum distribution of the positron from the decay of τ^+ lepton compared to that of a directly produced positron, in $\bar{p}p$ collisions at $\sqrt{s} = 540$ GeV.
- b) The polar angle distribution for a directly produced positron or a sequential positron coming from τ^+ decay.
- Fig. 2 a) Event density plot as a function of the transverse momentum and the polar angle for a direct positron in $\bar{p}p$ collisions at $\sqrt{s} = 540$ GeV.
- b) The same as a) for the positron, the decay product of τ^+ .
- Fig. 3 Comparison of the probability rates for the directly and sequentially produced positron. R is defined as the ratio of the cross-section for the sequential process over the cross-section for the direct process at given p_T and rapidity.
- Fig. 4 Transverse momentum distributions of the hadronic resonance π , ρ or A_1 produced in the decay of τ in the reaction $\bar{p}p \rightarrow WX$, $W \rightarrow \tau\nu$ at $\sqrt{s} = 540$ GeV.
- Fig. 5 Missing transverse momentum distributions associated to the production of π , ρ and A_1 decaying from τ^+ in the process $\bar{p}p \rightarrow WX$, $W \rightarrow \tau\nu_\tau$ at $\sqrt{s} = 540$ GeV.
- Fig. 6 The charged pion transverse momentum spectra for the various hadronic decay channels of the τ produced in $\bar{p}p \rightarrow WX$, $W \rightarrow \tau\nu_\tau$ at $\sqrt{s} = 540$ GeV.
- Fig. 7 The jet transverse momentum distribution in the sequential process $\bar{p}p \rightarrow WX$, $W \rightarrow \tau\nu_\tau$ followed by $\tau \rightarrow \text{jet} + \nu_\tau$ at $\sqrt{s} = 540$ GeV. The cuts $p_T^{\text{miss}} > 15$ GeV and $p_T^{\text{QCDjet}} < 15$ GeV are enforced.
- Fig. 8 Transverse momentum distributions for one, two or three charged tracks in the process $\bar{p}p \rightarrow WX$, $W \rightarrow \tau\nu_\tau$ and $\tau \rightarrow n \text{ tracks} + \nu_\tau$ at $\sqrt{s} = 540$ GeV.
- Fig. 9 Correlation plot between the charged and the total transverse energy in the hadronic decay modes of the τ . The channel $\tau \rightarrow \pi \nu_\tau$ is not included.
- Fig. 10 "Isolated" π^0 transverse momentum distribution in the hadronic decay modes of the τ .

FIG. 1

FIG. 1

FIG. 2

FIG. 2

FIG. 3

FIG. 4

FIG. 5

FIG. 6

FIG. 7

FIG. 8

FIG. 9

FIG. 10