EFFECTS OF HEAVY UNCONVENTIONAL SCALARS IN LEPTONIC INTERACTIONS

John N. Ng

TRIUMF and Physics Department, University of British Columbia
 4004 Wesbrook Mall, Vancouver, B.C., Canada V6T 2A3

Abstract

We derived the contributions of virtual scalar in $e^+e^- + \mu^+\mu^-$, $e^+e^- + e^+e^-$, muon decays and $(\bar{\nu}_e + e^- + \bar{\nu}_e)e^-$ scattering to obtain constraints on Yukawa couplings and masses of these scalars. Implications for anomalous $e^+e^- \gamma$ events in the $p\bar{p}$ collider are also discussed.

(Submitted to Physics Letters B)

CERN LIBRARIES, GENEVA

CM-P00067327

Recent investigations in supersymmetric gauge models have suggested that there could be more than one Higgs doublet in electroweak theories [1]. The coupling of the physical scalars to fermions need not be simply proportional to the fermions' masses. For two Higgs doublets there exist two physical charged Higgs and three neutral ones. Production of these non-standard Higgs in e^+e^- collisions have been considered by various authors [2]. Suggestions have also been made that the anomalous $e^+e^- \gamma$ events recently observed in the $p\bar{p}$ collider at CERN could be due to the decay of either Z^0 [3] or toponium [4] into a scalar or an unconventional Higgs plus a photon. Being unconventional, the neutral Higgs subsequently decays into e^+e^- pair with a large branching ratio.

In this paper we investigate the constraints placed on the Yukawa couplings of both unconventional charged and neutral scalars as a function of their masses. We concentrate on flavor conserving scalars. We use the reactions (a) $e^+e^- + \mu^+\mu^-$, (b) $e^+e^- + e^+e^-$, (c) muon decays and (d) $\bar{\nu}_e e + \bar{\nu}_e e$ scattering. The constraints are nontrivial for scalar masses around 50 GeV/c² and become loose as the masses increase. We shall take a general phenomenological approach and parameterize the effective coupling of the charged scalar, $\chi^\pm$ coupling to $\bar{\nu}_\ell \nu_\ell$ by the effective Lagrangian

$$\mathcal{L} = y_\ell \bar{\nu}_\ell \chi^\pm + \text{h.c.} \quad (1)$$

and similarly the neutral scalar ϕ^0 $\ell\bar{\ell}$ effective Lagrangian is taken to be

$$\mathcal{L} = y_\ell \bar{\ell} \ell \phi^0. \quad (2)$$

Since we are not considering CP violating effects we can take y_ℓ and y_ℓ^i to be real. In a gauge model where ϕ^0 and $\chi^\pm$ belong to the same multiplet $y_\ell = \sqrt{2} y_\ell^i$. In general the lowest mass physical scalars could come from

different multiplets and hence the Yukawa couplings need not be simply related. In the standard model $y_e^i = 0$ and $y_e = 2^{1/4} G_F^{1/2} m_\ell$ where m_ℓ is the mass of the lepton ℓ . The subscript ℓ for each lepton is kept for these Yukawa couplings since they need not be universal in their strengths.

In the standard model they are not. However, some models with discrete symmetries [5] impose on the three families has $e - \mu - \tau$ universality holds; and thus

$$y_e = y_\mu = y_\tau. \quad (3)$$

We shall assume that the masses of $\chi^\pm$ and ϕ^0 , denoted by M_χ and M_ϕ respectively, are larger than 45 GeV/c² otherwise they would have already been produced at PETRA. Alternatively, if scalars are light they must have $y^2/(4\pi) \ll \alpha$ in order to escape detection. Then they behave like standard Higgs boson and we are not considering them. For massive scalars only virtual effects can be discerned in the reactions of (a) - (d) at present energies. We study them systematically below:

a) $e^+e^- \rightarrow \mu^+\mu^-$

To the lowest order only, ϕ^0 , contributes in the annihilation channel. The Feynman diagrams are depicted in fig. 1. The cross section is given by [3]

$$\sigma = \frac{4\pi\alpha^2}{3} s \left[1 + \frac{(g_V^2 + g_A^2)^2}{16 s_W^4 c_W^4} \left(1 - \frac{M_Z^2}{s} \right) \right] + \frac{g_V^2}{2 s_W^2 c_W^2} \frac{1}{\left(1 - \frac{M_Z^2}{s} \right)} + \frac{w_e^2 w_\mu^2}{\left(1 - \frac{M_\phi^2}{s} \right)^2} \quad (4)$$

where $w_\ell = \frac{y_\ell}{e}$, $s_W^2 = \sin^2\theta_W$, $c_W^2 = \cos^2\theta_W$ (5a)

and $g_V^2 = \frac{1}{4} (1 - 4 s_W^2)^2$, $g_A^2 = \frac{1}{4}$ (5b)

We assume the validity of the standard electroweak model and use the

average values of $M_Z = 92$ GeV/c² and $\sin^2\theta_W = 0.23$ given by the UA1 and UA2 experiments. Using the experimental result of MARKJ [6] we obtain the allowed values in $y_\mu y_e - M_\phi^2$ space. This is the area below the solid line plotted in fig. 2. We have used the data [6] from PETRA at $\sqrt{s} = 41.3$ GeV and allowed for a 5 percent error in the data to obtain this constraint. For $M_\phi = 50$ GeV/c² we get $y_\mu y_e < 0.010$ and for $M_\phi = 1$ TeV/c² the limit is $y_\mu y_e < 13$

b) $e^+e^- \rightarrow e^+e^-$

In this case the t-channel scalar exchange scattering process can interfere with the s-channel annihilation process of γ and Z^0 exchanges (see fig. 3a) and vice versa (fig. 3b). The interference between s and t-channel ϕ exchange also takes place. In general this is a more sensitive test for virtual scalar effects than the previous case. The differential cross section is given explicitly as

$$\frac{d\sigma}{d\cos\theta} = \frac{\pi\alpha^2}{s} \left[\frac{1}{2} \frac{(3 + \cos^2\theta)^2}{(1 - \cos\theta)} + \frac{1}{2s_W^2 c_W^2} \frac{(g_V^2 + g_A^2)u^2 + (g_V^2 - g_A^2)t^2}{s(s - M_Z^2)} \right] + \frac{1}{16s_W^4 c_W^4} \frac{\{(g_V^2 + g_A^2)^2 + 4g_V^2 g_A^2\} u^2 + (g_V^2 + g_A^2)^2 t^2}{(s - M_\phi^2)^2} + \frac{1}{16s_W^4 c_W^4} \frac{\{(g_V^2 + g_A^2)^2 + 4g_V^2 g_A^2\} u^2 + (g_V^2 + g_A^2)^2 s^2}{(t - M_\phi^2)^2} + \frac{1}{2s_W^2 c_W^2} \frac{(g_V^2 + g_A^2)u^2 + (g_V^2 - g_A^2)s^2}{t(t - M_Z^2)} + \frac{1}{2s_W^2 c_W^2} (g_V^2 + g_A^2)u^2 \left\{ \frac{1}{s(t - M_Z^2)} + \frac{1}{t(s - M_Z^2)} \right\} + \frac{(g_V^2 + g_A^2)^2 + 4g_V^2 g_A^2}{16s_W^4 c_W^4} \frac{u^2}{(s - M_\phi^2)(t - M_\phi^2)} + w_e^4 \left\{ \frac{u^2 - s^2 - t^2}{4(s - M_\phi^2)(t - M_\phi^2)} + \frac{s^2}{2(s - M_\phi^2)^2} + \frac{t^2}{2(t - M_\phi^2)^2} \right\}$$

$$-w_e^2 \left\{ \frac{t^2}{s(t-M_\phi^2)} + \frac{s^2}{t(s-M_\phi^2)} + \frac{(g_V^2 - g_A^2)t^2}{4s_W^2 c_W^2 (t-M_\phi^2)(s-M_Z^2)} \right\} + \frac{(g_V^2 - g_A^2)s^2}{4s_W^2 c_W^2 (t-M_Z^2)(s-M_\phi^2)} \left\{ \right\},$$

where s , t , and u are the Mandelstam variables. The first term is the lowest order QED result and the terms not involving w_e are the standard electroweak model contribution [7]. We have written them in a form where one can identify the interference terms. Using the MARKJ data [8] at $\sqrt{s} = 34.6$ we performed a fit to find the maximum allowed value for w_e^2 for a given M_ϕ . Notice that in this reaction a constraint on y_e^2 alone can be obtained. Fig. 4 depicts the allowed region for y_e^2 and M_ϕ^2 to be right of the dashed curve. For low masses of $m_\phi \approx 50$ GeV/ c^2 , y_e^2 is constrained to be less than 0.003; and for large masses, e.g. $m_\phi = 1$ TeV/ c^2 , we obtain $y_e^2 < 0.91$. We mention in passing that since the lowest order QED completely dominates at small angles the better place to look for the effects of these currents is at large angles. However, the data is poorer there.

$$c) \mu^+ + e^+ \nu_\mu \nu_e$$

It is well known that the Michel spectrum of the e^+ for muon decays is sensitive to the presence of charged scalar currents. The precision measurement of the η -parameter gives us knowledge of $y_e^2 y_\mu^2$ and is crucial for testing the universality of the y_e 's which could arise from discrete or family symmetries. The e^+ -spectrum is [9]

$$\frac{dR}{dx} = \frac{G_F^2 m_\mu^5}{96\pi^3} \left\{ \left[1 + \left(\frac{M_W}{M_\chi} \right)^4 \left(\frac{y_e^2 y_\mu^2}{g^2} \right)^2 \right] x^2 (3-2x) + 12 \left(\frac{M_W}{M_\chi} \right)^2 \left(\frac{y_e^2 y_\mu^2}{g^2} \right) \frac{m_e}{m_\mu} x(1-x) \right\}$$

with $x \equiv (2E_e)/(m_\mu)$, where g is the SU(2) gauge coupling. As expected the charged scalar and W exchange interference term has a

coefficient of m_e/m_μ and vanishes in the limit $m_e \rightarrow 0$. Using the current value of η measurement [10] where

$$\eta \equiv \frac{2 \left(\frac{M_W s_W}{M_\chi} \right)^2 \frac{y_e^2 y_\mu^2}{e^2}}{1 + \left(\frac{M_W s_W}{M_\chi} \right)^4 \frac{y_e^2 y_\mu^2}{e^2}} < 0.05$$

we rule out the area to the left of the dashed line in fig. 2. At $M_\chi = 1$ TeV/ c^2 we find $y_e^2 y_\mu^2 < 16$. The dot-dash line indicated the region $y_e^2 y_\mu^2 - M_\chi^2$ space that will be ruled out by a two percent measurement of η if no effects are found. This accuracy is within reach of current experiments [11].

$$d) \left(\frac{-}{\nu_e} + e^- + \left(\frac{-}{\nu_e} + e^- \right) \right)$$

In this process an interference between the exchange of a charged scalar which flips helicity and the neutral current exchange process can occur [see fig. 5]. The effect arises in general for any neutral weak boson whose coupling to the right-handed electrons is not zero. This is the case for the Z^0 in the standard model. This is in addition to the known interference between the W and Z^0 terms. Since we assume that scalar particles do not couple to neutrinos then $\nu_\mu e + \nu_\mu e$ plays no role in detecting the effects of these particles. The differential cross sections we obtain are (neglecting electron mass)

$$\begin{aligned} \frac{d\sigma}{dE'} (\nu_e e + \nu_e e) &= \frac{2G_F^2 m_e}{\pi} \left[\frac{1}{4} + s_W^2 + s_W^4 \right] \left\{ 1 + \left(1 - \frac{E'}{E_\nu} \right)^2 \right\} \\ &\quad - 2 \left(\frac{y_e^2 M_W}{g M_\chi} s_W \right)^2 - 4 \left(\frac{y_e^2 M_W}{g M_\chi} \right)^4 \left\{ 1 - \frac{E'}{E_\nu} \right\}^2 \\ \text{and} \quad \frac{d\sigma}{dE'} (\bar{\nu}_e e + \bar{\nu}_e e) &= \frac{2G_F^2 m_e}{\pi} \left[\frac{1}{4} + s_W^2 + s_W^4 \right] \left\{ 1 - \frac{E'}{E_\nu} \right\}^2 + s_W^4 \\ &\quad - 2 \left(\frac{y_e^2 M_W s_W}{g M_\chi} \right)^2 + 4 \left(\frac{y_e^2 M_W}{g M_\chi} \right)^4 \end{aligned}$$

where E' is the energy of the scattered electron and $E_{\nu(\bar{\nu})}$ is the incident neutrino (antineutrino) energy. The first three terms contain the usual W and Z^0 interference [12], and next terms are the χ - Z^0 interference, and the contribution from χ itself.

These reactions now play a similar role to Bhabha scattering in that they are sensitive only to charged scalar Yukawa coupling, y_e' . Using the reactor data [13] and the standard model parameters given above, the allowed values for $y_e'^2$ and M_χ^2 is to the right of the solid line in fig. 4. For $M_\chi = 1 \text{ TeV}/c^2$ we get $y_e'^2 < 6.8$.

We see from the above that the most stringent limit currently available on the Yukawa couplings is given by Bhabha scattering. The value of $m_\chi = 50 \text{ GeV}/c^2$ is of interest to the anomalous $e^+e^- \gamma$ event from $p\bar{p}$ collider [14,15]. Even allowing for the uncertainty in $\sin^2\theta_W$ and M_Z we get at best $y_e^2 < 0.01$. This rules out unconventional Higgs coupling of strength determined by the top quark mass, i.e. $y_e^2 \approx 0.04$. Applying the above values we assess the possibility that the anomalous events originate from mixing of Z^0 and toponium which decays into unconventional Higgs plus a photon [4]. The only solution allowed by Bhabha scattering is maximal mixing with

$$y_{eBR}^2(\phi + e^+e^-) = 0.002.$$

This leads to a branching of ϕ into e^+e^- about 0.2. Although permissible phenomenologically it is highly unlikely that any reasonable model can give such a large branching ratio for $\phi + e^+e^-$. Furthermore, the invariant mass plot of the e^+e^- pairs in the preliminary data does not support such a resonance interpretation.

Finally the role of muon $g-2$ plays in the study of unconventional scalars has to be discussed. In $g-2$ calculations additional theoretical

issues concerning effective $\phi^0 Z^0 \gamma$ and $\phi^0 \gamma \gamma$ terms enter. These are highly model dependent. If we leave them out or if they are small, then the constraints on the Yukawa coupling are [14],

$$y_\mu^2 < 5 \times 10^{-4} \frac{m_\mu^2}{m_\chi^2}$$

for charged scalars and a factor of two better neutral scalars. This is certainly not competitive with Bhabha scattering constraints.

We note in passing that similar considerations can be extended to the quark sector. Precision measurements of pion decays such as the branching ratio of $\pi_{e\nu}/\pi_{\mu\nu}$ can be used to limit the Yukawa couplings of charged scalars to quarks and leptons. However, the uncertainty in dealing with the radiative corrections involving hadrons which cannot be ignored hampers the accurate extraction of information on new weak currents from these measurements.

In conclusion, we have demonstrated the usefulness of leptonic interactions to constrain the Yukawa couplings of scalars. What we have considered can be carried through for pseudoscalars by inserting a $i\gamma_5$ for L_{eff} and can be extended verbatim to include the τ lepton. Optimistically if effects are found, one can use Bhabha scattering and $(\bar{\nu})e$ scattering to test whether y_e and y_e' are simply related. This is important in general and can be used to pin down the group structure of the scalar sector. Then analysing $e^+e^- + \mu^+\mu^-$ and the η -parameter in muon decays one can test whether Yukawa couplings for these unconventional scalars or pseudoscalars to e and μ obey universality as discussed by some models.

Acknowledgements

The author would like to thank Dr. M. Jacob for the kind hospitality at CERN where this work was completed. Discussions with Dr. H. Chen, Dr. J.E. Kim and members of the CERN theory division are gratefully acknowledged.

References

- [1] For reviews see H.P. Mlles, UCVA-DPT (1983) and A. Arnouitt, A.H. Chameddine and P. Nath, Northeastern Univ. preprint (1984).
- [2] N.G. Deshpande, X. Tata and D.A. Dicus, Phys. Rev. D29 (1984) 1527.
For earlier work see C. Albright, J. Smith, and H. Tye, Phys. Rev. D21 (1980) 711.
For a review see R.A. Flores and M. Sher, Ann. Phys. 148 (1983) 95.
- [3] R.D. Peccei, Phys. Lett. 136B (1984) 121.
U. Bauer, H. Fritzsch, and H. Faisner, Phys. Lett. 135B (1984) 313.
- [4] J. Lindfors, S. Chu and B.R. Desai, Univ. of California, Riverside preprint (1984).
- [5] J.E. Kim and G. Segre, Phys. Lett. 78B (1978) 75.
B. McWilliams and L.F. Li, Nucl. Phys. B179 (1981) 62.
- [6] S. Naroska, Proceedings of the 1983 International Symposium on Lepton and Photon Interactions at High Energies, Cornell University.
- [7] S. Budny, Phys. Lett. 55B (1975) 227.
- [8] For results of Mark J collaboration see M.I.T. report 131 (1983).
- [9] F. Scheck, Phys. Rept. 44C (1978) 187.
- [10] R. Bossingham, Submitted to Xth Int. Conf. on Particles and Nuclei, Heidelberg, (1984). TRIUMF EXP E134.
- [11] H.L. Anderson, LAMPF Experiment in progress.
- [12] B. Kayser, E. Fishbach, S.P. Rosen and H. Spitzvak, Phys. Rev. D20 (1979) 87.
- [13] E. Pasierb, et al., Phys. Rev. Lett. 43 (1979) 96.
- [14] UA1 Collaboration, Phys. Lett. 126B (1983) 398.
- [15] UA2 Collaboration, Phys. Lett. 129B (1983) 130.
- [16] J.P. Leveille, Nucl. Phys. 137B (1978) 73.

Figure Captions

- [1] Feynman diagrams contributing to $e^+e^- \rightarrow \mu^+\mu^-$ scattering including virtual scalar, ϕ^0 , exchange.
- [2] Constraints on the Yukawa couplings $y_\mu y_e$ and $y'_\mu y'_e$ of the neutral and charged scalars ϕ and χ as functions of the masses squared. The solid line is from $e^+e^- \rightarrow \mu^+\mu^-$, the dashed line from muon decay with 5 percent measurement of η , and dash-dot line is for 2 percent of measurement. Allowed parameters are to the right of the curves.
- [3] Feynman diagrams for Bhabha scattering.
- [4] Constrained Yukawa couplings y_e and y'_e as a function of M_ϕ^2 and M_χ^2 . The solid line is given by $\bar{\nu}e \rightarrow \bar{\nu}e$ scattering and the dashed line by Bhabha scattering. Allowed parameters are to the right of the lines.
- [5] Feynman diagrams for $\nu e^+ \nu e^-$ scattering including virtual charged scalar, χ , exchange.

Fig. 1

Fig. 2

Fig. 3

Fig. 4

Fig. 5