EUROPEAN ORGANIZATION FOR NUCLEAR RESEARCH

CERN/EP 84-90
25 July 1984

LOW MOMENTUM TRANSFER ELASTIC SCATTERING

AT THE CERN PROTON-ANTIPROTON COLLIDER

UA4 Collaboration

Amsterdam¹-CERN²-Genova³-Napoli⁴-Pisa⁵

M. Bozzo³, P.L. Braccini^{5(*)}, F. Carbonara⁴, R. Castaldi⁵,
F. Cervelli⁵, G. Chiefari⁴, E. Drago⁴, M. Haguenauer², V. Innocente⁴,
B. Koene¹, S. Lanzano⁴, G. Matthiae⁴, L. Merola⁴, M. Napolitano⁴,
V. Palladino⁴, G. Sanguinetti⁵, S. Scapellato^{5(**)}, G. Sciacca⁴,
G. Sette³, R. van Swol¹, J. Timmermans¹, C. Vannini⁵,
J. Velasco^{2(***)}, P.G. Verdini⁵ and F. Visco⁴

- ¹ NIKHEF-H, Amsterdam, The Netherlands
- ² CERN, European Organization for Nuclear Research, Geneva, Switzerland
- ³ Istituto di Fisica dell'Università e Sezione INFN, Genoa, Italy
- ⁴ Istituto di Fisica Sperimentale dell'Università e Sezione INFN, Naples, Italy
- ⁵ Istituto di Fisica dell'Università e Sezione INFN, Pisa, Italy

ABSTRACT

Proton-antiproton elastic scattering was measured at the centre-of-mass energy $\sqrt{s} = 546$ GeV in the four-momentum-transfer range $0.03 < -t < 0.32$ GeV². For $-t < 0.15$ GeV² the data are well described by a simple exponential form $\exp(bt)$ with slope parameter $b = 15.2 \pm 0.2$ GeV⁻². This value is significantly larger than the one measured in the region $0.21 < -t < 0.50$ GeV².

Submitted to Physics Letters B

(*) Also at Istituto di Fisica dell'Università, Perugia, Italy.
(**) Also at Scuola Normale Superiore, Pisa, Italy.
(***) Permanent address: IFIC, University of Valencia (Spain).

In this letter we present the results of the measurement of low momentum transfer proton-antiproton elastic scattering at the CERN SPS Collider at a centre-of-mass energy $\sqrt{s} = 546$ GeV. Since the experimental method is the same as in our earlier measurement [1], we only recall briefly a few relevant points of the technique employed. Elastic events are observed in a system of four telescopes placed symmetrically above and below the machine plane at a distance of about 40 m from the beam crossing point. A telescope is composed of two movable sections of the vacuum chamber (pots) which are 6 m apart. Each pot contains a scintillation counter hodoscope and a wire chamber; the chamber has four drift planes measuring the vertical coordinate with 0.13 mm resolution per plane and a proportional plane measuring the horizontal coordinate by the charge division method with 0.4 mm resolution [2].

The present data were collected in a dedicated Collider run with the machine optics at the intersection region in the so-called high- β mode, the betatron function β at the crossing point being 100 m in both the horizontal and vertical plane. This optics was especially designed to allow measuring low- t elastic scattering, and is nearly symmetric on the proton and antiproton sides. At the crossing point the transverse beam size and the beam angular divergence had r.m.s. values of ~ 1.4 mm and ~ 0.014 mrad respectively. The average luminosity was about $4 \cdot 10^{26} \text{ cm}^{-2} \text{ s}^{-1}$.

The particles scattered in the crossing region traverse quadrupoles of the machine before reaching the detectors. Particle deflection through the quadrupoles was calculated with the standard formalism of transfer matrices using the accurately known strength and effective length of the quadrupoles. As a consequence, each track seen in a telescope is reconstructed back to the crossing region with errors depending only on the resolution of the detectors and in an about equal amount on the multiple scattering in the pot windows and counters. The "effective distance" of the detectors from the crossing point (i.e. the ratio of the particle displacement at the detector plane to the production angle) was about 35 m and 40 m in the horizontal and vertical plane respectively, very close to the geometric distance.

Calibration and detection efficiency of the wire chambers were studied using the data collected with the elastic trigger itself. In the track reconstruction procedure a hit was required in at least five of the eight drift planes and in at least one of the two proportional planes in a telescope. Due to the redundancy in the number of planes a fully efficient detection could be ensured over the complete sensitive area of the chambers.

After track reconstruction in the individual telescopes further analysis was carried out according to the following steps:

- (a) Search for two roughly collinear tracks in two opposite telescopes.
- (b) Rejection of the events where the number of tracks in a telescope is larger than two (this allows including elastic events with an accompanying δ -ray).
- (c) Cut on the vertical coordinate of the proton and antiproton trajectories at the centre of the crossing region.
- (d) Cut on the collinearity distributions of the proton and antiproton trajectories in both the vertical and horizontal plane.

The cuts (c) and (d) were done at the three-standard-deviation level. Most of the rejected events had many tracks in several chambers and can be attributed to random coincidences of beam-wall and beam-gas interactions. About 10^5 events, corresponding to 54% of the triggers, survived all cuts and were taken to represent our final sample of elastic scattering data.

The distributions of the vertical displacement of the antiproton and proton trajectories at the crossing region and the collinearity distributions are shown in fig. 1. The widths of these distributions, after unfolding the experimental resolution and the multiple scattering contribution, agree well with the measured beam emittances. The off-set in the horizontal collinearity distribution is due to the displacement of the beam in this plane. By taking the average of the scattering angles on the proton and antiproton side, the effect of this off-set is canceled.

In order to calculate the four-momentum transfer t given by $-t = p^2 \theta^2$ where θ is the measured scattering angle and p the beam momentum, one must know the absolute calibration of the machine energy. The nominal beam momentum is 270 GeV/c. However, recent measurements have shown that the actual beam momentum is 273.0 ± 1.4 GeV/c, always the same since the start of Collider operation [3]. The quoted error is a conservative upper limit of the systematic uncertainty. As a consequence, the t -scale of all earlier elastic scattering data from the SPS Collider [1,4,5] should be corrected upwards by $(2.2 \pm 1.0)\%$ and the values of the slope parameter should be lowered by the same amount.

The observed angular distributions for the two detection arms (up-left $\times$ down-right) and (down-left $\times$ up-right) were fitted simultaneously leaving the angle of the incoming beam as a free parameter. The beam angle was found to be very small, less than 0.01 mrad in both the horizontal and vertical planes, and has been used to correct the measured scattering angle. The geometrical acceptance was calculated by the Monte-Carlo method using the known values of the beam parameters.

As already noted, scattered particles traverse quadrupoles of the machine before reaching the detectors. The transverse section of the vacuum chamber in the quadrupoles follows the shape of the iron poles. As a consequence the accepted range of the horizontal component θ_H of the scattering angle becomes narrower as the vertical component θ_V increases. The range in θ_H was suitably defined to ensure sufficient clearance inside the vacuum chamber, taking into account the transverse beam dimension and also a small misalignment and deformation of the vacuum chamber itself. It was checked that limiting θ_H to even smaller values did not significantly change the results for the parameters in the fits of the angular distributions.

The measured differential cross section as a function of t in the range $0.03 < -t < 0.32$ GeV² is presented in Table 1. The absolute normalization has been obtained through the simultaneous measurement of the total cross section [6] and implies an overall error of 5%. Our measured values of the differential cross section in the region $0.21 < -t < 0.50$ GeV² of ref. [4] are also given in Table 1. They are normalized to the present data in the overlap region $0.21 < -t < 0.32$ GeV²

and an additional overall normalization error of 3% should therefore be attributed to them. The data of Table 1 are displayed in fig. 2. Since the widths of the collinearity distributions of fig. 1 imply a very good t -resolution ($\Delta t = 0.006 \sqrt{-t}$), no smearing correction has therefore been applied when fitting the observed t -distribution.

In the region $0.03 < -t < 0.15 \text{ GeV}^2$ the distribution can be well described by a simple exponential $\exp(bt)$ with a constant slope parameter b . The result of a least-squares fit is $b = 15.2 \pm 0.2 \text{ GeV}^{-2}$ and is represented as a solid line in fig. 2. The χ^2 of the fit was 26 for 22 degrees of freedom. At these small t -values, the contribution of the Coulomb scattering amplitude and especially of its interference with the nuclear amplitude becomes non-negligible. At $-t = 0.03 \text{ GeV}^2$ they amount to 1.3% of the pure nuclear cross section, assuming $\rho = 0.15$ (ρ is the ratio of the real to imaginary part of the scattering amplitude at $t = 0$). The two terms due to Coulomb scattering have been added in the above fit and in all subsequent fits in the intervals below $-t = 0.15 \text{ GeV}^2$. Their effect is to lower the value of the slope parameter by 0.1 GeV^{-2} .

Our new value for the slope parameter b at small t is lower than our earlier measurements [1,4] with low statistics (about a thousand events). Part of the difference is accounted for by the correction for the beam momentum which amounts to $\sim 0.4 \text{ GeV}^{-2}$. Furthermore the large statistics of the present measurement allowed a more precise understanding of the geometrical acceptance in the region where the vacuum chamber is limiting the θ_H interval; a more restrictive θ_H cut was eventually applied to the data. Successive θ_H cuts on our previous data did not reveal - within the much poorer statistics - any variation of the resulting slope parameter.

For comparison with lower energy data we have also done a fit in the interval $0.03 < -t < 0.10 \text{ GeV}^2$. The result $b = 15.3 \pm 0.3 \text{ GeV}^{-2}$ confirms that within the given statistics there is no significant quadratic dependence on t in the exponential when fitted in the region $0.03 < -t < 0.15 \text{ GeV}^2$. The value of the slope in this interval differs very significantly by about two units from the slope value in the interval

$0.21 < -t < 0.50 \text{ GeV}^2$ [4,5], which when corrected for the true beam momentum amounts to $13.4 \pm 0.3 \text{ GeV}^{-2}$. The result of the constant slope fit in that region is represented by a dashed line in fig. 2. This change in slope in the vicinity of $-t = 0.15 \text{ GeV}^2$, was already observed at the ISR in pp [7] and $\bar{p}p$ [8] scattering and can also be seen in fig. 2. The experimental data points beyond $-t = 0.20 \text{ GeV}^2$ stay clearly above the (dotted) extrapolation of the line fitted in the region $0.03 < -t < 0.15 \text{ GeV}^2$. An exponential $\exp(bt)$ fit to the present data in the region $0.15 < -t < 0.32 \text{ GeV}^2$ gives $b = 14.2 \pm 0.4 \text{ GeV}^{-2}$; when the fit is limited to the overlap region with the data of ref. [4] ($0.21 < -t < 0.32 \text{ GeV}^2$) we find $b = 13.6 \pm 0.8 \text{ GeV}^{-2}$, compatible with the above mentioned value for the region $0.21 < -t < 0.50 \text{ GeV}^2$. A summary of our slope values in the different t -intervals is given in table 2.

In summary, the covered t -range of $0.03 < -t < 0.50 \text{ GeV}^2$ comprises two clearly distinct intervals where the differential cross section is well described by an exponential $\exp(bt)$ with constant slope parameter b . When a fit over the full t -range is done where one leaves as free parameters the two constant slopes and the t -value t_0 at which the change of slope occurs, one finds $-t_0 = 0.14 \pm 0.02 \text{ GeV}^2$. The χ^2 of this fit is 1.0 per degree-of-freedom. This fit was performed to extract the t -value where the local slope varies most rapidly with t ; it does not necessarily imply the actual existence of a sharp break. Other parametrizations of the t -distribution have also been tried, for instance an exponential with a quadratic dependence on t . A fit of the form $\exp(Bt + Ct^2)$ over the t -region $0.03 < -t < 0.15 \text{ GeV}^2$ does not give a value for C that differs significantly from zero. A significant quadratic dependence can only be obtained from fits over a t -region that extends up to at least 0.3 GeV^2 . These fits have a χ^2 which are reasonably good but imply a linear decrease of the local slope ($B - 2|C|t$) over the full t -range which is not supported by the data.

Our result on the slope parameter at low momentum transfer ($0.03 < -t < 0.10 \text{ GeV}^2$) is presented in fig. 3 together with lower energy data on pp and $\bar{p}p$ scattering in the same range of momentum transfer [7-9]. Assuming a $\log s$ dependence of the slope from the ISR to the Collider, the coefficient of the $\log s$ term would be $\sim 0.5 \text{ GeV}^{-2}$.

We wish to express our gratitude to the staff of the CERN Accelerator Divisions for the help in the installation and survey of this experiment and for the successful operation of the SPS Collider during the dedicated high- β run when this measurement was performed. Special thanks go to P. Faugeras of the SPS Division who carefully designed the high- β optics. We are very grateful to R. Maleyran of the SPS Division who took care of the design and installation of the pots. The continued effort of the technicians at CERN, INFN and NIKHEF who contributed to this experiment is warmly acknowledged.

REFERENCES

- [1] R. Battiston et al., UA4 Collaboration, Phys. Lett. 115B (1982) 333.
- [2] J. Buskens et al., Nucl. Instr. and Meth. 207 (1983) 365.
- [3] B. de Raad and W.C. Middelkoop (CERN/SPS Division), private communication.
- [4] R. Battiston et al., UA4 Collaboration, Phys. Lett. 127B (1983) 472.
- [5] G. Arnison et al., UA1 Collaboration, Phys. Lett. 121B (1983) 77 and 128B (1983) 336.
- [6] M. Bozzo et al., UA4 Collaboration, CERN/EP 84-91, submitted to Phys. Lett. B.
- [7] G. Barbiellini et al., Phys. Lett. 39B (1972) 663.
- [8] M. Ambrosio et al., Phys. Lett. 115B (1982) 495.
- [9] U. Amaldi et al., Phys. Lett. 66B (1977) 390;
N. Amos et al., Phys. Lett. 120B (1983) 460 and 128B (1983) 343;
D.S. Ayres et al., Phys. Rev. D15 (1977) 3105.
L. Baksay et al., Nucl. Phys. B141 (1978) 1.
V. Bartenev et al., Phys. Rev. Lett. 31 (1973) 1088.

TABLE CAPTIONS

Table 1 Differential elastic cross section.

Table 2 Average slope values and χ^2 for fits in different regions of
four-momentum transfer.

TABLE 1

-t GeV ²	dσ/dt mb/GeV ²	Error mb/GeV ²	-t GeV ²	dσ/dt mb/GeV ²	Error mb/GeV ²
.0325	124.71	1.57	.1775	14.30	.94
.0375	113.40	1.45	.1825	13.94	.93
.0425	105.64	1.44	.1875	12.87	.90
.0475	99.20	1.44	.1925	13.36	.92
.0525	89.72	1.41	.1975	11.72	.87
.0575	81.86	1.38	.2025	9.71	.80
.0625	78.59	1.38	.2075	9.16	.78
.0675	72.13	1.35	.2125	9.14	.78
.0725	66.94	1.33	.2175	8.43	.76
.0775	63.53	1.32	.2225	7.91	.74
.0825	58.32	1.28	.2275	7.51	.72
.0875	54.00	1.25	.2325	6.75	.69
.0925	47.53	1.19	.2375	5.47	.62
.0975	44.57	1.17	.2425	5.82	.65
.1025	39.99	1.13	.2475	6.53	.69
.1075	40.57	1.15	.2525	4.84	.60
.1125	36.49	1.10	.2575	3.85	.53
.1175	36.17	1.11	.2625	3.96	.54
.1225	30.08	1.02	.2675	4.38	.57
.1275	29.29	1.02	.2725	3.96	.55
.1325	26.95	.99	.2775	3.23	.50
.1375	25.96	.98	.2825	3.02	.48
.1425	22.05	.91	.2875	3.52	.52
.1475	20.49	.89	.2925	2.84	.47
.1525	21.36	1.10	.2975	2.63	.46
.1575	20.04	1.08	.3025	2.89	.48
.1625	17.95	1.03	.3075	2.51	.45
.1675	16.80	1.00	.3125	2.61	.46
.1725	15.42	.97	.3175	2.63	.47
.215	8.19	.30	.365	1.182	.091
.225	7.28	.28	.375	.842	.077
.235	6.76	.27	.385	.787	.075
.245	5.95	.26	.395	.853	.079
.255	4.92	.24	.405	.643	.069
.265	4.57	.23	.415	.612	.067
.275	3.58	.21	.425	.499	.061
.285	3.13	.14	.435	.495	.061
.295	3.09	.14	.445	.439	.058
.305	2.48	.13	.455	.367	.053
.315	2.25	.12	.465	.308	.049
.325	1.97	.11	.475	.265	.045
.335	1.81	.11	.485	.236	.043
.345	1.56	.10	.495	.151	.035
.355	1.42	.10			

TABLE 2

t-range (GeV^2)	Slope b (GeV^{-2})	$\chi^2/\text{d.o.f.}$
0.03 - 0.10	15.3 ± 0.3	10/12
0.03 - 0.15	15.2 ± 0.2	26/22
0.15 - 0.32	14.2 ± 0.4	18/32
0.21 - 0.32	13.6 ± 0.8	13/20
0.21 - 0.50	13.4 ± 0.3	23/27

FIGURE CAPTIONS

- Fig. 1 Vertical displacements of the antiproton (a) and proton (b) trajectories at the centre of the crossing region. Collinearity plots for the vertical (c) and horizontal (d) plane.
- Fig. 2 Differential cross section of proton-antiproton elastic scattering at $\sqrt{s} = 546$ GeV. The lines drawn are the results of simple exponential fits $\exp(bt)$ and are described in the text. The present data and the data of ref. [4] are shown as open and solid circles respectively.
- Fig. 3. The forward slope ($-t < 0.10$ GeV²) at the Collider compared to proton-proton and proton-antiproton results at lower energy in the same t -range.

Fig. 1

Fig. 2

Fig. 3