RESTRICTION ON AMOUNT OF ANTIMATTER IN EARLY UNIVERSE FROM
 $\bar{p}$ - ^4He REACTION DATA

F. Balestra⁵, Yu. A. Batusov¹, G. Bendiscioli⁴, M. P. Busa⁵,
L. Busso⁵, I. V. Falomkin¹, L. Ferrero⁵, V. Filippini⁴,
G. Gervino⁵, C. Guaraldo², E. Lodi Rizzini⁴, A. Maggiora²,
D. Panzieri⁵, G. Piragino⁵, G. B. Pontecorvo¹, A. Rotondi⁴,
M. G. Sapozhnikov¹, F. Tosello⁵, M. Vascon³, G. Zanella³, and
A. Zenoni⁴.

SUMMARY

A limit on the possible amount of antimatter in the early Universe has been deduced from $\bar{p}$ - ^4He reaction data obtained at 180 MeV.

(Submitted to Lettere al Nuovo Cimento)

-
- 1 - Joint Institute for Nuclear Research, Dubna, USSR.
 - 2 - Laboratori Nazionali di Frascati dell'INFN, Frascati, Italy.
 - 3 - Dipartimento di Fisica dell'Universita' di Padova, and INFN Sezione di Padova, Padova, Italy.
 - 4 - Dipartimento di Fisica Nucleare e Teorica dell'Universita' di Pavia, and INFN Sezione di Pavia, Pavia, Italy.
 - 5 - Istituto di Fisica Generale dell'Universita' di Torino, and INFN Sezione di Torino, Torino, Italy.

The standard baryon-symmetrical cosmology excludes the existence of a considerable quantity of antimatter in the Universe at $t > 10^3$ s (see, for example refs.(1,2)). On the other hand, the modern astrophysics as well as the theories of grand unification predict the existence of sources of antimatter (domains of antimatter, evaporation of primordial black holes, the decays of superheavy metastable particles, etc.) after that time (for review, see ref.(3)). Most of antimatter from these sources annihilated and today we can observe only the products of this reaction. As it was shown in refs.(3-5), if antiprotons annihilated on ^4He , this process could lead to a change in the concentration of the light elements, such as deuterium and ^3He .

Hence, to go back to the possible amount of antimatter in the early Universe, it is essential to measure the cross sections for the different reaction channels in the $\bar{p}$ - ^4He interaction, and to compare these data with the abundance of ^2H , ^3He and ^4He observed to day in the Universe ⁽³⁻⁵⁾. As the energy of the primordial antiprotons is uncertain, the $\bar{p}$ - ^4He cross sections should be measured at different energies down to at rest.

Recently we have obtained the first result on the interaction of $\bar{p}$ in flight (180 MeV) on ^4He ⁽⁶⁾, which allows us to evaluate the amount of antimatter in the early Universe on a more experimental basis than in the past. ⁽⁵⁾

The experimental apparatus, based on the use of a streamer chamber filled with helium at 1 atm and placed in a

magnetic field of 0.8 T, has been described in detail in ref.(7). The sensible volume of the chamber was $90 \times 70 \times 18^3$ cm³. Only the central part, 55 cm long, of the chamber has been considered for these measurements and in this fiducial volume 1097 $\bar{p}$ -⁴He reaction events at 180 MeV of kinetic energy in the laboratory have been photographed and analysed with a scanning efficiency of 99.7%. Besides the statistical errors we estimated a further systematic error of 2% due to scanning efficiency, target transparency and beam dose counting. With this technique we measured a reaction cross section $\sigma_r = \sigma_{tot} - \sigma_{el} = 233.3 \pm 7.1$ mb.

(The errors quoted here and in the following are statistical only).

Reactions involving the production of ³He are readily identified in the streamer chamber, since of all the possible annihilation reactions only these involve an even number of charged particle tracks in the final state. Using this very clear signature, we measured a cross section for ³He production $\sigma_{^3\text{He}} = 35.7 \pm 2.7$ mb.

The amount of ³He formed as a result of $\bar{p}$ -⁴He interactions (4) in the early Universe should be at a first approximation:

$$n_{^3\text{He}} = n_{^4\text{He}} R f_{^3\text{He}}^{\text{eff}} \quad (1)$$

where $n_{^4\text{He}}$ represents the concentration of ⁴He, $R = n_{\bar{p}} / n_p$ is the fraction of antiprotons in the early Universe,

$$f_{^3\text{He}}^{\text{eff}} = f_{^3\text{He}} + f_{^3\text{H}}$$

represents the effective output of ${}^3\text{He}$ in $\bar{p}-{}^4\text{He}$ annihilation. This relation takes also into account the output of tritium $f_{3\text{H}}$, because the subsequent decay of this nuclide in the Universe contributes to the ${}^3\text{He}$ concentration.

The output of ${}^3\text{He}({}^3\text{H})$ is defined as follows:

$$f_{3\text{He}({}^3\text{H})} = \sigma(\bar{p}-{}^4\text{He} \longrightarrow {}^3\text{He}({}^3\text{H}) + \text{anything}) / \sigma_{\text{tot}} \quad (2)$$

From our data one can estimate the total cross section of $\bar{p}-{}^4\text{He}$ interaction as $\sigma_{\text{tot}} = 1.5 \sigma_r = 350 \pm 10$ mb. Such a ratio holds in $\bar{p}-{}^2\text{H}$ interaction as well as in theoretical predictions for $\bar{p}-{}^3\text{He}$ interaction at our energy. (8) (9) Therefore we have

$$f_{3\text{He}} = \frac{\sigma_{3\text{He}}}{\sigma_{\text{tot}}} = 0.102 \pm 0.008$$

From this value it is possible also to evaluate the yield of tritium in $\bar{p}-{}^4\text{He}$ annihilation, assuming that

$$f_{3\text{H}} = f_{3\text{He}} r$$

where $r = \sigma^{\text{ann}}(\bar{p}p) / \sigma^{\text{ann}}(\bar{p}n) \simeq 1.32 \pm 0.05$. (8)

Thus, the estimate of the effective yield of ${}^3\text{He}$ is the following:

$$f_{3\text{He}}^{\text{eff}} = f_{3\text{He}} + f_{3\text{H}} = 0.237 \pm 0.014 \quad (3)$$

Assuming that the amount of ${}^3\text{He}$ created as a result of

$\bar{p}$ - ^4He annihilation could not exceed the abundance of ^3He observed at present, from eq.(1) we obtain the following restriction on the fraction of antimatter in the early Universe:

$$R < \frac{4}{3} \frac{X_{^3\text{He}}}{X_{^4\text{He}} f_{^3\text{He}}^{\text{eff}}} \quad (4)$$

where $X_{^3\text{He}} = (4.2 \pm 2.8) 10^{-5}$, $X_{^4\text{He}} = (0.23 \pm 0.02)$ are the observed mass abundances of ^3He and ^4He ⁽¹⁰⁾

Utilizing eqs(3-4) we finally obtain a restriction on the fraction of antimatter in the early Universe:

$$R < (1.03 \pm 0.70) 10^{-3}$$

(the error includes also the uncertainties in light element abundance).

This restriction on R exceeds by more than three orders of magnitude the earlier restriction ($R < 1$) that was based on the investigation of the permissible distortion of the Planck spectrum of the relic radiation. ^(3,11) Knowledge of this ratio put severe restrictions on the formation of primordial black holes, on the concentration of heavy metastable particles (on graviton concentration, for example) and on other parameters of sources of antimatter present in the Universe after the nucleosynthesis ^(3,4).

Although this upper limit of R refers to 180 MeV antiprotons, we are confident in this estimate, because a

preliminary analysis of our data taken at 50 Mev shows an increase of at least 60% in the ^3He production, and consequently a further decrease in R.

In effect, ^3H and ^3He are the only nuclides that can be produced in the $\bar{p}$ - ^4He annihilation in absence of initial or final state interactions. Hence, going toward the threshold for inelastic processes (25 MeV of incident energy in the laboratory) the production of these nuclides should be more and more favoured, the contribution from other channels being due mainly to the pionic final state interaction which is nearly independent of the incident energy.

We are grateful to V.P.Dzhelepov, S.A.Bunyatov, B.Pontecorvo, Ya.B.Zeldovich, V.M.Chechetkin and M.Yu.Khlopov for the interesting discussions and suggestions for this experiment.

REFERENCES

1. G.Steigman Ann.Rev.Astron.Astrophys. 14, 339 (1976).
2. Ya.B.Zeldovich and I.D.Novikov: Structure and Evolution of the Universe (Moscow, 1975).
3. V.M.Chechetkin, M.Yu.Khlopov, M.G.Sapozhnikov: Riv.Nuovo Cimento 5, 1 (1982).
4. V.M.Chechetkin, M.Yu.Khlopov, M.G.Sapozhnikov, Ya.B.Zeldovich: Phys.Lett.B118, 329 (1982).
5. I.V.Falomkin, G.B.Pontecorvo, M.G.Sapozhnikov, M.Yu.Khlopov, F.Balestra and G.Piragino: Nuovo Cimento A79, 193 (1984).
6. Dubna-Frascati-Padova-Pavia-Torino Collaboration: Submitted to Phys.Lett.B
7. Dubna-Frascati-Padova-Pavia-Torino Collaboration: Nucl.Instr. and Meth., in press.
8. R.Bizzarri, P.Guidoni, F.Marcelja, F.Marzano, E.Castelli and M.Sessa: Nuovo Cimento 22A, 225 (1974).
9. L.A.Kondratyuk and M.Zh.Shmatikov: Yad. Fiz. 38, 361 (1983); Sov. J. Nucl. Phys.38 (2), 216 (1983).
10. S.M.Austin: Progr. Part. Nucl.Phys.7, 1 (1981).
11. R.A.Sanyaev and Ya.B.Zeldovich: Astr.Space Sci. 7, 20 (1970).