CERN-TH.4023/84

HARMONIC EXPANSION AND DIMENSIONAL REDUCTION

IN G/H KALUZA-KLEIN THEORIES

R. Coquereaux

Centre de Physique Théorique, Section 2
CNRS, Luminy, Marseille

and

A. Jadczyk^{*)}

CERN - Geneva

A B S T R A C T

We propose a geometrical framework for harmonic expansion and dimensional reduction of matter fields in Kaluza-Klein theories with the most general G-invariant ansatz. Generalized Peter-Weyl and Frobenius theorems provide a basis for harmonic expansion, and a mechanism is shown by which the dimensional reduction of matter fields is then automatically accomplished. In particular, we discuss the dimensional reduction of tensor and spinor fields, and of the Laplace and Dirac operators.

^{*)} On leave from University of Wroclaw, Poland.

CERN-TH.4023/84

September 1984

I. Introduction.

Harmonic expansion and dimensional reduction are two mechanisms used in discussing the effective four-dimensional content of higher-dimensional field theories admitting a spontaneous compactification. A general discussion of these problems was given by Salam and Strathdee [1] while Witten [2] gave a broad discussion of the harmonic expansion and dimensional reduction of spinor fields and, in particular, of the difficulties in obtaining the effective four-dimensional chiral asymmetry. A space-time manifold was treated in those papers locally and no intrinsic geometrical meaning was given to the constructions. On the other hand, certain global aspects of the problem were studied by Römer [3] and Bleecker [4,5]; however, both authors restricted their discussion to the case of internal space being a group manifold, and spinors were not discussed at all in [4,5].

In this paper we consider a G/H Kaluza-Klein theory as formulated in [6]. As a geometrical model for the extended space-time we take a manifold E on which a global symmetry group G acts (G can be thought of as the internal symmetry group of a ground state, but the scheme is broad enough to apply to any Riemannian manifold with a compact group of isometries). Then E locally looks like a product $M \times S$ of space-time M and internal homogeneous space $S \cong G/H$. The space-time manifold is defined globally and unambiguously as the manifold of G -orbits (internal spaces). What is not unique is a local product representation of E as $M \times G/H$ - any two such representations differ by a gauge transformation with gauge group $N(H)/H$ (see [6,7]). All the analysis given in this paper is directed by the aim not to distinguish any particular product

representation of E (a global one may even not exist), and in doing so to exhibit an intrinsic geometrical meaning of the constructions.

The plan of the paper is as follows: we start with matter fields on E considered as sections of some equivariant vector bundle over E , that is of a vector bundle $F=F(E;F)$ with base E and typical fibre F , with G acting on F by bundle automorphisms. The set ΓF of all cross-sections of F is the space of matter-field configurations in the extended space-time E . We decompose ΓF according to the irreducible representations α of G and, for each α , define harmonics of type α . This constitutes the first step - harmonic expansion of fields (Sect.2). Then, in Sect.3. we show that the space of harmonics of a given type α (which are still fields on E) can be interpreted as the space of sections of an appropriate "effective" vector bundle $\tilde{F}_\alpha$ over M . This is the second step - dimensional reduction. After that we consider the particularly interesting situation when F has a structure group, say, R ; or, in other words, when F is a bundle associated to some principal bundle $U=U(E;R)$ via a certain representation ρ of R on F . The global symmetry group G is assumed to act on U by automorphisms. We use the results of [8,9] to find out the effective ("dimensionally reduced") gauge group $\tilde{R}$ and the effective principal bundle $\tilde{U}(M;\tilde{R})$ over M . We also find the effective representation ρ_α of $\tilde{R}$ on an appropriate F_α and show that the $\tilde{F}_\alpha$ can also be constructed as the bundle associated to the effective principal bundle $\tilde{U}$ via ρ_α . In Sect.4. we apply these results to discuss the case of U being the bundle of orthonormal frames of E endowed with a G -invariant metric, and in Sect.5 we discuss tensor and spinor fields and dimensionally reduced Laplace and Dirac operators.

2. Harmonic expansion

Let E be a manifold with a given action of a compact Lie group G . For instance E can be a multidimensional universe with G being its global symmetry group [6], but the analysis given here can be applied to many other situations as well. On E consider linear matter fields of some fixed type. If E is topologically trivial (e.g. if E is contractible, (see [10, Ch.4, Cor.10.3]) then these matter fields can be described by vector-valued functions i.e. by functions on E with values in some vector space F . In general, however, matter fields have to be described by cross-sections of a certain (not necessarily trivial) vector bundle $F=F(E;F)$ with base E and typical fibre F (example: vector fields on S^2). We suppose F is a finite dimensional vector space (real or complex). We also assume that the symmetry group G "knows" not only how to act on E (by diffeomorphisms), but also, by bundle automorphisms, on F^*). Thus each $a \in G$ maps a fibre F_y of F at $y \in E$ onto the fiber F_{ya} at $ya \in E$ by a one-to-one linear map: $F_y \ni u \rightarrow ua \in F_{ya}$ (we shall keep to the convention of using right actions of G on E and on F).

Let ΓF denote the space of all cross-sections ϕ of F . Then ΓF is an infinite dimensional vector space and its elements $\phi \in \Gamma F$ are called fields, or field configurations. Having given the action of G on E and on F there is a natural representation of G on ΓF . This induced representation T is defined by the formula

*) F is then called an equivariant vector bundle or, more specifically, G -vector bundle (see e.g. [11,12,13])

$$(T(a)\phi)(y) \doteq \phi(ya)a^{-1}. \quad (2.1)$$

We now want to decompose ΓF - the space of field configurations - into invariant subspaces $\Gamma_\alpha F$ corresponding to the irreducible representations α of the symmetry group G , with $\Gamma_\alpha F$ consisting of field configurations transforming according to the irreducible representation α . But first let us introduce the notation which will be used throughout the rest of the paper. For each irreducible representation α let $W_\alpha^{(*)}$ be the representation space of α (W_α is finite-dimensional since G is assumed to be compact), and let χ_α be the corresponding character. It is convenient to introduce $\hat{G}$ - the dual of G - which is the (discrete) space of (equivalence classes of) irreducible representations of G . Analogously as in [13, p.119] we can define, for every $\alpha \in \hat{G}$, a projection operator $\pi_\alpha : \Gamma F \rightarrow \Gamma F$ by the formula

$$(\pi_\alpha \phi)(y) \doteq d(\alpha) \int_G (T(a)\phi)(y) \chi_\alpha(a^{-1}) da, \quad (2.2)$$

where $d(\alpha) = \dim W_\alpha$, and da is the normalized Haar measure on G . The range of π_α will be denoted by $\Gamma_\alpha F$ and its elements will be called harmonics of type α . It can be shown, and the proof can be adapted from [11,13], that the algebraic direct sum $\bigoplus_{\alpha \in \hat{G}} \Gamma_\alpha F$ is dense in ΓF ; more precisely, every field configuration $\phi \in \Gamma F$ can be approximated with an arbitrary

*) Unless stated otherwise F and W_α are assumed to be complex vector spaces.

accuracy by a finite superposition of harmonics in the following sense: for every $\varepsilon > 0$ and for every compact $K \subset E$ there is a finite subset $A \subset \hat{G}$ and $\phi_A \in \bigoplus_{\alpha \in A} \Gamma_{\alpha} F$ such that $\sup_{y \in K} \|\phi(y) - \phi_A(y)\| < \varepsilon$. The norm here is any continuous norm on (the fibres of) F . This approximation theorem (which is akin to the classical Peter-Weyl theorem (see e.g. [14])) is sufficient for our purposes; we do not need to investigate in what sense the "inverse Fourier transform" formula

$$\phi(y) \sim \sum_{\alpha \in \hat{G}} \pi_{\alpha} \phi(y)$$

holds. In particular we are not interested in making ΓF into a Hilbert space - this because E is thought of as a model for (an extended) space-time of a hyperbolic signature, and in field theory it is the space of solutions of a hyperbolic differential equation rather than the space of all field configurations that carries a useful Hilbert space structure.

It is a general philosophy of any application of the Fourier analysis that, when the group G is correctly identified, the physically most important are the lowest modes, that is harmonics corresponding to a few of the lowest dimensional representations of G . It is therefore reasonable, and the Peter-Weyl theorem makes such a restriction also mathematically founded, to concentrate on the analysis of the elements of Γ_{α} , with $\alpha \in \hat{G}$ arbitrary but fixed^{*)}. This is what we will do in the

^{*)} It may happen that such a restriction is incompatible with field equations in E which may imply the necessity of having infinite towers of representations (see e.g. [15]). In such a case a reasonable approximation is needed to cut the tower.

following.

Let $w_i (i=1,2,\dots,d_\alpha)$ be a basis in W_α and let w^i be the dual basis in W_α^* - the dual of W_α . We can now represent the χ_α factor in (2.1) as

$$\chi_\alpha(a^{-1}) = \text{Tr}(\alpha(a^{-1})) = \sum_i \langle w^i, \alpha(a^{-1})w_i \rangle = \sum_i \langle \alpha^*(a)w^i, w^i \rangle, \quad (2.3)$$

where $\alpha^*(a) = \alpha(a^{-1})^T$ is the contragradient representation of G on W_α^* . It is therefore natural to introduce, for each $\phi \in \Gamma F$ and each $w \in W_\alpha^*$, the Fourier component of ϕ along w by

$$\phi_w(y) = d(\alpha) \int [T(a)\phi](y) \alpha^*(a)w \, da. \quad (2.4)$$

It follows then from (2.2 - 2.4) that

$$\pi_\alpha \phi = \sum_i \langle \phi_w, w^i \rangle, \quad (2.5)$$

w_i being any basis in W_α . In other words ϕ can be restored by knowing its Fourier components ϕ_w . We will now express this fact more precisely, introducing at the same time the concepts we shall need later on.

We denote by F_α the fibre bundle fibres of which the tensor products are $F_y \otimes W_\alpha^*$ of fibers F_y of F and W_α^* . F_α can be also thought of as the tensor product $F \otimes W_\alpha^*$ of two vector bundles: F and the trivial bundle $W_\alpha^* = E \times W_\alpha^*$. The field harmonics $\phi_w(y)$ defined in (2.4) are now cross-sections of the bundle F_α . We denote by $T \otimes \alpha^*$ the representation of G on ΓF_α^* defined by

) Observe that $\Gamma(F \otimes W_\alpha^)$ is naturally isomorphic to $\Gamma F \otimes W_\alpha^*$ - thus one can omit brackets in $\Gamma F \otimes W_\alpha^*$; also, we will often write W_α^* instead of W_α^* .

$$[(T \otimes \alpha^*)(a)\phi \otimes w](y) \doteq \phi(ya)a^{-1} \otimes \alpha^*(a)w(y). \quad (2.6)$$

This is a particular case of the general formula (2.1) provided that the action of G on the trivial bundle W_α^* is defined by

$$(y, w)a \doteq (ya, \alpha^*(a^{-1})w). \quad (2.7)$$

Comparing now the formulae (2.2) and (2.4), with T replaced by $T \otimes \alpha^*$ and $\chi_\alpha(a)$ put identically 1, we find that ϕ_w are G -invariant sections of F_α . We call $\Gamma_{\text{inv}} F_\alpha$ the space of these sections. One proves then (see [11, 13, 16]) that (2.5) gives an isomorphism between $\Gamma_\alpha F$ and $W_\alpha \otimes \Gamma_{\text{inv}} F_\alpha^*$). Because of this fact the term "harmonics" will also be used for the elements of $\Gamma_{\text{inv}} F_\alpha$.

We summarize the discussion: an arbitrary field configuration ϕ can be expanded into harmonics according to the irreducible representations of G ; we have

$$\Gamma F \sim \bigoplus_{\alpha \in \hat{G}} W_\alpha \otimes \Gamma_{\text{inv}} F_\alpha, \quad (2.8)$$

where $F_\alpha = F \otimes W_\alpha^*$ and $\Gamma_{\text{inv}} F_\alpha$ is the space of G -invariant sections of the bundle F_α .

) When F is a real vector space and α is a real irreducible representation, then $\Gamma_\alpha F$ is isomorphic to $W_\alpha \otimes \Gamma_{\text{inv}} F \otimes W_\alpha^$, where the tensor product $\otimes$ is taken over the commuting ring of α which is $\mathbb{R}, \mathbb{C}$ or $\mathbb{H}$ (see [11, also [17] for the concept of a commuting ring)

3. Dimensional reduction of a G-vector bundle

We have seen that the natural building blocks out of which every field configuration can be reconstructed are harmonics i.e. G-invariant (or "equivariant") sections of vector bundles $F_\alpha = F \otimes W_\alpha^*$, $\alpha \in \hat{G}$. The discussion which will now follow can be applied to the space of invariant sections of any equivariant vector bundle. Therefore we consider here $\Gamma_{\text{inv}} F$, F being an arbitrary equivariant vector bundle over E , and at the end we shall apply the results replacing F by F_α .

Let us recall the situation we deal with: G - a compact Lie group - acts on a manifold E and on a vector bundle $F = F(E; F)$ over E . In the following we shall always assume that E is a simple G -space i.e. that the stability groups of the points of E are all conjugated to a standard one, denoted by H . Every orbit is then isomorphic to G/H , and the space M of all orbits is a manifold. With this preparation the very essence of the dimensional reduction mechanism of matter fields can be expressed by the following result:

Theorem 3.1. There is a natural isomorphism between the space $\Gamma_{\text{inv}} F$ of invariant sections of an equivariant vector bundle F over E and the space of all sections of a certain vector bundle $\tilde{F}$ over M .

A proof of this result can be found in [18]. Here we indicate only how is the bundle $\tilde{F}$ built out. The construction goes in two steps: first an invariant subbundle F_0 of F is constructed, and next $\tilde{F}$ is defined

as the quotient F_o/G . To define F_o denote by G_y the stability group at $y \in E$. Then G_y acts by endomorphisms on F_y , and $(F_y)_o$ is defined as the subspace of F_y consisting of G_y -invariant vectors. The collection $F_o = \bigcup_y (F_y)_o$ is a G -invariant subbundle of F . One then shows that the quotient $\tilde{F} = F_o/G$ can be given the structure of a vector bundle over $M = E/G$.

As it was said in the beginning of this section this construction is now to be applied to the bundles $F_\alpha = F \otimes W_\alpha^*$ rather than to F itself, although the F of Sect.2 is a particular case of the F_α -family, namely $F = F_\alpha$ for α -the trivial representation of G .

Although the above recipe is rather straightforward, it has one serious disadvantage: it does not tell us how the effective structure group of $\tilde{F}_\alpha$ is to be found when the initial structure group of F is known. The rest of this section will be devoted to an alternative construction of the $\tilde{F}_\alpha$ which will make the problem of reduction of the structure (or "gauge") group explicit. In the construction below, as well as in the rest of this paper, we shall exploit the results of our earlier work [7-9]. For the reader's convenience let us first briefly recall the concepts and results which will be needed in the following.

We start, as before, with a simple G -space E , G being a compact Lie group. Let P be the submanifold of E consisting of all points with the stability group H (assumed connected):

$$P = \{y \in E : G_y = H\} .$$

One proves then that P is a principal bundle over M with structure group $N(H)|H$, $N(H)$ being the normalizer of H in G : $N(H)=\{a \in G: aHa^{-1}=H\}$. Suppose now $U=U(E;R)$ is a principal bundle over E with structure group R , and let G act also on U by principal bundle automorphisms. (Such a situation is, for instance, studied when considering symmetries of Yang-Mills fields). The action of G on U is characterized by a certain homomorphism $\lambda: H \rightarrow R$ [8]. By applying the same method which was used to define P , but now replacing E by U and G by $G \times R$ one constructs a submanifold $\tilde{U} \subset U$ which is a principal bundle over M with the effective structure group $\tilde{R} = N(H)|H$, where $H = \text{diag}(H, \lambda(H)) \subset G \times R$ is the standard stability group for the $G \times R$ action on U . One also proves [8] that $\tilde{R}$ is locally a product of $N(H)|H$ and the centralizer Z of $\lambda(H)$ in R^* .

With the above information at hand let us consider again our vector bundle F , but now assuming it to have a structure group R . Or, in other words, we assume there is given a representation ρ of R on a vector space F , and $\tilde{F} = \tilde{F}(E;F)$ is the bundle associated to U via the representation ρ . The standard notation is $\tilde{F} = U \times_R F$ or, better $U \times_{\rho} F$. We remind the reader (for more information see [22,23]) that $\tilde{F}$ is constructed out of the Cartesian product $U \times F$ by identifying those pairs (u,v) and (u',v') for which $u' = ur$ and $v' = \rho(r^{-1})v$ for some $r \in R$. The equivalence class of (u,v) is conveniently denoted as $u \cdot v$, and we have $ur \cdot v = u \cdot \rho(r)v$, which gives a formal justification for the notation used.

*) The earlier papers dealing with the problem of symmetric gauge fields [19-21] overlooked the $N(H)|H$ term.

We interpret $u \cdot v$ as "the object having co-ordinates v in the frame u ". Observe that the action of G on U induces, in a natural way, the action of G on F :

$$(u \cdot v)a \doteq ua \cdot v.$$

Thus F is automatically an equivariant vector bundle and the construction given by the Theorem 3.1. can be applied to F_α to give the vector bundle $\tilde{F}_\alpha$ over M . But now we are prepared to describe an alternative construction of $\tilde{F}_\alpha$: we shall construct it as a bundle associated to $\tilde{U}$ via a certain representation ρ_α of $\tilde{R}$ on a vector space F_α which we presently describe. First of all observe that $\rho \otimes \alpha^*$ is a representation of $R \times G$ on $F \otimes W_\alpha^*$. Denote by $F_\alpha = F \otimes_H W_\alpha^*$ the subspace of $F \otimes W_\alpha^*$ consisting of vectors invariant under the subgroup $\bar{H} = \text{diag}(H, \lambda(H)) \subset R \times G$. Then F_α is invariant under the action of $N(\bar{H})$ and, since $\bar{H}$ acts on $F \otimes_H W_\alpha^*$ trivially, we effectively obtain a representation of $N(\bar{H}) | \bar{H}$ on F_α . This representation will be denoted by ρ_α . We then have

Theorem 3.2. The bundle $\tilde{F}_\alpha$ constructed out of F_α according to the Theorem 3.1. is naturally isomorphic to the bundle $\tilde{U} \times_{\rho_\alpha} F_\alpha$ associated to $\tilde{U}$ via the representation ρ_α of $\tilde{R}$ on F_α .

The only interesting part of a proof of this theorem is to show how the isomorphism in question is constructed: given $q \cdot v \in \tilde{U} \times_{\rho_\alpha} F_\alpha$ we have $q \in U$ and $v \in F_\alpha \subset F \otimes W_\alpha^*$, so that we can consider $q \cdot v$ as an element of

$F\otimes W^*_\alpha$. It is easy to check that $q \cdot v$ is in fact in $(F\otimes W^*_\alpha)_0$ (see the discussion following the Theorem 3.1.). The isomorphism is then given by the map $\tilde{U} \times_{\rho_\alpha} F_\alpha \ni q \cdot v \rightarrow (q \cdot v)G \in (F\otimes W^*_\alpha)_0 / G$, where $(q \cdot v)G$ denotes the orbit (i.e. equivalence class under the quotient map) of $q \cdot v$ under the action of G on $(F\otimes W^*_\alpha)_0$. It is easy to see that the map above is well defined i.e. the result does not depend on the choice of representatives q and v in the class $q \cdot v \in \tilde{U} \times_{\rho_\alpha} F_\alpha$.

We end this section with a brief discussion of dimensional reduction of linear differential operators acting on the fields. Here we have the following result:

Theorem 3.3. Every G -invariant linear differential operator D on ΓF induces a linear differential operator $\tilde{D}_\alpha$ on $\tilde{F}_\alpha$.

The construction of $\tilde{D}_\alpha$ is straightforward: one first extends D to $D_\alpha = D \otimes I$ on $\Gamma F\otimes W^*_\alpha$ and observes that $\Gamma_{\text{inv}}(F\otimes W^*_\alpha)$ is invariant under $D \otimes I$. But $\Gamma_{\text{inv}}(F\otimes W^*_\alpha)$ is isomorphic to $\Gamma \tilde{F}_\alpha$ by the Theorem 3.1. Thus $\tilde{D}_\alpha$ is defined as the image of D_α , restricted to $\Gamma_{\text{inv}}(F\otimes W^*_\alpha)$, under this isomorphism. It remains to prove that $\tilde{D}_\alpha$ so defined is again a differential operator, and a proof of this fact can be found in [18].

4. Dimensional reduction of the bundle of orthonormal frames

So far we considered a quite general abstract vector bundle F over E . In particular we had to assume that some action of G on F is somehow given - it had to agree with the action of G on E but it did not follow from the latter. In the following we shall focus our discussion on the particular case of F being the bundle of tensors or spinors on E . For this we have to apply the reduction process $U \rightarrow \tilde{U}$ in a particular case of U being the bundle of orthonormal (or spin) frames. It is important to notice that the action of G on E essentially determines its action on tensors and spinors. But first we have to discuss in some details Riemannian geometry of simple G -spaces. We recall here the relevant concepts and facts; more information can be found in references [6-8].

Let us assume that E is equipped with a G -invariant metric g_E (we shall assume that the induced metric on orbits of G in E is either positive or negative definite). Then, because of G -invariance, g_E generates a metric g_M on $M=E/G$, gauge field A_μ on M with gauge group $N(H)|H$, and a certain number of scalar fields on M . The Lie algebra G of can be decomposed into $G=H+K+L$, where $N=H+K$ is the Lie algebra of N , $S=K+L$ can be identified with the tangent space to $S=G/H$ at the origin, and both decompositions $G=N+L$ and $G=H+S$ are reductive. We recall that H is assumed to be connected. Then K , which can be identified with the Lie algebra of $K=N(H)|H$, can be also characterized as the space of $\text{Ad}(H)$ -invariant elements of S .

We denote by OE the bundle of orthonormal frames of E . The structure

group of OE will be denoted by $O(\eta_E)$ or, shortly, $O(E)$. It consists of matrices Λ satisfying $\Lambda^T \eta_E \Lambda = \eta_E$ where η_E is the canonical diagonal form of g_E ; if E is a multidimensional universe then it is natural to take $\eta_E = \pm \text{diag}(-1, +1, +1, \dots, +1)$. Since G acts on E by isometries the action of G on E lifts automatically to OE ; indeed, point transformations induce transformations of tangent vectors and of frames and, being isometric, they map orthonormal frames into orthonormal ones. But OE and $O(E)$ are not yet our U and R of Sect.3 - the bundle OE is unnecessarily large and we shall first use the extra information we have to reduce its structure group. This can be done as follows: for every $p \in P$ define K_p and L_p as the subspaces of $T_p E$ spanned by the Killing vectors from K and L respectively. For an arbitrary $y \in E$ there always exist $p \in P$ and $a \in G$ such that $y = pa$; we define $K_y \doteq K_p a$ and $L_y \doteq L_p a$. Because $\text{Ad}(H)$ leaves the subspaces K and L of G invariant (with K being even pointwise invariant), our definitions are unambiguous.

Definition 4.1. An orthonormal frame $e_A(y)$ at $y \in E$ is called adapted if $e_A = (e_\mu, e_a)$ with $e_a \in K_y$, $e_\mu \in L_y$, and e_μ orthogonal to $S_y \doteq K_y \oplus L_y^*$.

The set of all adapted orthonormal frames will be denoted by AOE - it is a principal bundle with structure group $AO(E) = O(M) \times O(K) \times O(L)$, where $O(M)$ is the (pseudo-) orthogonal group of η_M - the standard flat metric of M , while $O(K)$ and $O(L)$ are orthogonal groups in $k = \dim(K)$ and $l = \dim(L)$ dimensions respectively. The group G acts on AOE by automorphisms. It is only now that we shall apply the dimensional reduction

*) For $p \in P$ the vectors $e_\mu(p)$ span the horizontal subspace of the induced $N(H)|_H$ connection A_μ .

mechanism discussed in Sect.3 by putting $U \equiv AOE$ and $R \equiv AO(E)$; but first we have to identify the homomorphism $\lambda: H \rightarrow R$. Let T_i be a basis in G with $T_a \in K$, $T_{\hat{a}} \in L$ and $T_{\alpha} = (T_{\hat{a}}, T_a)$. We denote by $\varepsilon_i(y)$ the fundamental vector fields on E (Killing vectors) generated by T_i . The vectors $\varepsilon_{\alpha}(y)$ then span the vertical subspace S_y , provided y is not too far from P . In a neighbourhood of P we can therefore always write

$$e_{\alpha}(y) = \varepsilon_{\beta}(y) \phi^{\beta}_{\alpha}(e; y), \quad (4.1)$$

and if e_{α} is replaced by $e'_{\alpha} = e_{\beta} \Lambda^{\beta}_{\alpha}$ then $\phi(e\Lambda) = \phi(e)\Lambda$. For $e_A \in AOE$, $\phi(e)$ has the block diagonal form $(\phi^{\alpha}_{\beta}) = (\phi^{\hat{a}}_{\hat{b}}, \phi^a_b)$. Now, let $e(p) \in AOE$. Then the action of any $h \in H$ leaves p invariant and thus rotates $e_A(p)$ by an orthogonal transformation; in fact it is only e_a 's which are rotated. Thus we have

$$e_a h^{-1} = e_b \lambda(e; h)^b_a, \quad h \in H,$$

$\lambda: H \rightarrow O(L)$ being a group homomorphism. Observe that

$$\lambda(e\Lambda; h) = \Lambda^{-1} \lambda(e; h) \Lambda.$$

Let A^i_j be the matrix of the adjoint representation of G on G . Then, since $\varepsilon h^{-1} = \varepsilon A(h)$, we find that

$$\lambda(e; h) = \phi^{-1}(e) A(h) \phi$$

Although it is by no means necessary, it is very convenient to choose a basis T_i in G in such a way that, for some $p_0 \in P$, the Killing vectors $\varepsilon_\alpha(p_0)$ are orthonormal. Then $\lambda(\varepsilon(p_0); h) = A(h)$ is the matrix of the adjoint representation of H in S . We now apply the results discussed in Sect.3 to $U = AOE$, $R = AO(E)$ and $\lambda = A$ to get the following result

Theorem 4.1. The effective "gauge" group $\tilde{AO}(E)$ is locally isomorphic to the product $N(H) | H \times O(M) \times O(K) \times Z$, where Z is the centralizer of $Ad(H)$ in $O(L)$.

Although we will not need this fact, it is worthwhile to notice that one can easily get rid of the $O(K)$ -factor of the effective gauge group. Indeed, we can always reduce the bundle by demanding that $e_{\hat{a}}(p)$ are fixed e.g. by a standard orthonormalization process of $\varepsilon_{\hat{a}}(p)$. That means that the effective internal gauge group is $N(H) | H \times_{loc} Z$. In the particular, well-known, case when E is a principal bundle, the internal spaces are group manifolds, H is trivial, L is trivial, Z is trivial, and $N(H) | H = G$ - as expected.

According to the algorithm of the dimensional reduction discussed in Sect.3 (for more details see [7-9]) the reduced bundle $\tilde{AOE}$ consists of those pairs $(p, e_A(p))$ for which $\lambda(e; h) = A(h)$. The following result gives an explicit description of the reduced frame bundle.

Theorem 4.2. The frames $e_A(p) \in \tilde{AOE}$ are characterized by Φ 's satisfying

$$i) \quad \Phi A(h) = A(h) \Phi$$

$$\text{ii) } \phi^T g \phi = \eta$$

where $g(p) = (g_{\alpha\beta})$ are the components of the metric in the ϵ -basis, and $\eta = \pm I$ depending on whether the metric in the orbit is positive or negative definite. Such frames exist for every $p \in P$ and any two such frames at the same $p \in P$ are connected by an orthogonal transformation $\Lambda_1 \times \Lambda_2$ with $\Lambda_1 \in O(K)$ and $\Lambda_2 \in Z$, Z being the centralizer of $\text{Ad}(H)$ in $O(L)$.

Proof: The non-evident part of the statement is the existence result.

To prove the existence of a ϕ satisfying i) and ii) one starts with an arbitrary adapted frame (i.e. the result of a standard orthonormalization of ϵ), with $e = \epsilon \phi$ (we omit the non-interesting e_μ -vectors, and observe that ϕ satisfies ii) and $\lambda(e) = \phi^{-1} A \phi$ is an orthogonal matrix. It follows that $\phi \phi^T$ commutes with $A(h)$. Let $\phi = \phi_o \Lambda$ be the polar decomposition of ϕ , with $\phi_o > 0$ and Λ orthogonal. Then $\phi_o = (\phi \phi^T)^{\frac{1}{2}}$ commutes with $A(h)$, and it also satisfies ii) since $\phi_o = \phi \Lambda^{-1}$ and Λ is an orthogonal transformation.

With the information given by the above theorem we can also explicitly describe the reduced structure group $\tilde{A} \tilde{O} E$. Indeed, let $e(p)$ and $e(p')$ be in $\tilde{A} \tilde{O} E$, p' and p being in the same fibre of P . Then we first choose $n \in N(H)$ such that $p' = pn$, and then define $\Lambda \in O(L)$ by the relation

$$e(p)n = e(p')\Lambda^{-1}$$

(we again omit e_μ 's and e_a 's, which are uninteresting). Since both e and e' are in $\tilde{A}\tilde{O}E$, Λ must satisfy the constraint that $A(n^{-1})\Lambda$ commutes with $A(H)$ on L . But n is unique only modulo $h \in H$. Thus (n, Λ) should be identified with $(hn, A(h)\Lambda)$. Consequently we have the following result:

Theorem 4.3. The effective structure group $\tilde{A}\tilde{O}(E)$ is the product $O(M) \times O(K) \times R_1$, where R_1 consists of pairs (n, Λ) with $n \in N(H)$ and $\Lambda \in O(L)$ such that $A(n^{-1})\Lambda$ commutes with $A(H)$; the pairs (n, Λ) and $(hn, A(h)\Lambda)$, $h \in H$, define the same element of R_1 .

We can now interpret the Φ as a cross-section of a bundle associated to $\tilde{A}\tilde{O}E$. Indeed, with $e \in \tilde{A}\tilde{O}E$ and (n, Λ) as in the Theorem 4.3 we find

$$\Phi(en\Lambda) = A(n)^{-1}\Phi(e)\Lambda,$$

and the condition i) of Theorem 4.2 ensures that the above transformation law effectively depends on the equivalence class of (n, Λ) only.

5. Dimensional reduction of tensor and spinor fields.

For calculating the dimensionally reduced Laplace and Dirac operators we shall need explicit expressions for the affine connection $\omega_{A C}^B(p)$, $p \in P$. These expressions will, in general, depend on a moving frame e_A which we will choose in $\tilde{A}\tilde{O}E$ at the points of P . More specifically we proceed as follows:

- 1^o) Choose a local cross-section $\sigma : x \rightarrow \sigma(x)$ of the bundle P .
- 2^o) At every point $\sigma(x) \in P$ choose an adapted frame $e_A(\sigma(x)) \in \tilde{A}\tilde{O}E$.
Then $(\sigma(x), e_A(\sigma(x)))$ is a local cross-section of the bundle $\tilde{A}\tilde{O}E$.
- 3^o) Extend $e_A(\sigma(x))$ to an open neighbourhood of σ in E defining $e_A(\sigma(x)e^\xi) = e_A(\sigma(x))e^\xi$ for ξ running through a neighbourhood of zero in S .

With the moving frame e_A introduced as above we write $e_\alpha = \epsilon_\beta^\alpha \Phi_\alpha^\beta$ as in (4.1), and from 3^o) we easily find

$$\epsilon_\alpha^\beta(\Phi_\delta^\gamma) = -C_{\alpha\gamma}^\beta \Phi_\delta^\gamma \quad (5.1)$$

Observe that now e_μ is a horizontal (i.e. orthogonal to the orbits of G in E) lift of a certain moving frame, denoted by ∂_μ , on M .

The Christoffel symbols in the (e_μ, ϵ_α) -basis have been given in [7].

Here we have to transform them to an orthonormal basis (e_μ, e_α) . The results of a straightforward calculation read as follows:

$\omega_{\mu, \nu\sigma}$ - the affine connection of M in the ∂_μ - basis,*)

$$\omega_{\mu, \alpha\nu} = -\omega_{\mu, \nu\alpha} = \omega_{\alpha, \mu\nu} = -\frac{1}{2}(\eta\Phi^{-1})_{\alpha\beta} F_{\mu\nu}^\beta,$$

$$\omega_{\alpha, \beta\mu} = -\omega_{\alpha, \mu\beta} = \frac{1}{2}(\Phi^T(D_\mu g)\Phi)_{\alpha\beta},$$

$$-\omega_{\mu, \alpha\beta} = \frac{1}{2}(\Phi^T(D_\mu g)\Phi)_{\alpha\beta} + (\eta\Phi^{-1}e_\mu(\Phi))_{\alpha\beta},$$

where $F_{\mu\nu}^\alpha \neq 0$ for $\alpha=\hat{a}$ only, and

$$F_{\mu\nu}^{\hat{a}} = \partial_\mu A_\nu^{\hat{a}} - \partial_\nu A_\mu^{\hat{a}} + \frac{1}{2}C_{\hat{b}\hat{c}}^{\hat{a}} A_\mu^{\hat{b}} A_\nu^{\hat{c}}$$

is the field strength of the $N(H)|H$ - gauge field,

$$D_\mu g_{\alpha\beta} = \partial_\mu (g_{\alpha\beta}) + C_{\alpha\hat{c}}^\delta A_\mu^{\hat{c}} g_{\delta\beta} + C_{\beta\hat{c}}^\delta A_\mu^{\hat{c}} g_{\alpha\delta},$$

is the covariant derivative of $g_{\alpha\beta}$, and $e_\mu(\Phi)$ is given by

$$e_\mu(\Phi) = \partial_\mu(\Phi) - A_\mu^{\hat{a}} \varepsilon_{\hat{a}}(\Phi).$$

We also used the flat metric η_{AB} of E to lower the index of ω_A^B :

$$\omega_{A,BC} = \eta_{BD} \omega_A^D C^D_C.$$

By the Theorem 4.3. the effective structure group $\tilde{AOE}$ is the pro-
duct $O(M) \times O(K) \times R_1$, where $R_1 \approx_{loc} N(H)|H \times Z$. We have already identified
the $O(M)$ - , and $N(H)|H$ - parts of the effective gauge field - they
are $\omega_{\mu\sigma}^\nu$ and $A_\mu^{\hat{a}}$ respectively. It still remains to find the $O(K) \times Z$ -part.

*) Notice that in spite of the notation ∂_μ is an anholonomic orthonormal moving frame of M.

A general construction of the effective, dimensionally reduced gauge field has been given in [8]. The construction given there in Proposition 3.5 is formal but the idea behind it is quite simple: the effective horizontal lift of a path from M to $\tilde{A}OE$ is obtained by a composition of a horizontal lift from M to P and a horizontal lift from P to $\tilde{A}OE$. With that geometrical picture it is easy to find the remaining $O(K) \times Z$ - part of the effective connection: with respect to the local moving frame chosen above we have $B_{\mu\beta}^{\alpha} = \omega_{\mu}^{\alpha}{}_{\beta}$, where we have identified the Lie algebra $\text{Lie } (O(K) \times Z)$ with a subalgebra of $S=K+L$. Explicitly, one has

$$B_{\mu} = - \phi^{-1} (e_{\mu} + \frac{1}{2} g^{-1} D_{\mu} g) \phi \quad (5.3)$$

With this preparatory knowledge at hand we now consider several examples. We start with the simplest one:

a) Laplace operator on scalar fields

A (real) scalar field on E is a section of the trivial bundle $F = E \times \mathbb{R}$ which is associated to the trivial representation of $O(E)$ on $\mathbb{R}$. According to the discussion in Sect.3 the effective fibre bundle $\tilde{F}_{\alpha}$ over M has the fibre $F_{\alpha} = F \otimes_{H_{\alpha}} W_{\alpha}^* = \mathbb{R} \otimes_{H_{\alpha}} W_{\alpha}^* = (W_{\alpha}^*)_0$, that is F_{α} consists of H -singlets in W_{α}^* . The effective field on M therefore gets the index $\phi \rightarrow \phi_L$ from the representation space W_{α}^* and is constrained by

$$\alpha(h)^L_M \phi_L = \phi_M, \quad h \in H, \quad (5.4)$$

where $\alpha(h)^L_M$ is the matrix of the representation α . Knowing the connection

coefficients ω_{AC}^B one easily finds the effective Laplace operator acting on $\underline{\phi}=(\phi_L)$:

$$\Delta_{\text{eff}} \underline{\phi} = g^{\mu\nu} \mathcal{D}_\mu \mathcal{D}_\nu \underline{\phi} + g^{\alpha\beta} T_\alpha T_\beta \underline{\phi} + v^\mu \mathcal{D}_\mu \underline{\phi}, \quad (5.5)$$

where $\mathcal{D}_\mu \underline{\phi} = \partial_\mu (\underline{\phi}) + A_\mu^{\hat{a}} T_{\hat{a}} \underline{\phi}$ is the $N(H)|H$ - covariant derivative, $\mathcal{D}_\mu$ contains also the gravitational part, and v^μ is given by

$$v_\mu = \frac{1}{2} \text{Tr}(g^{-1} \mathcal{D}_\mu g) = \frac{1}{2} \text{Tr}(g^{-1} \varepsilon_\mu(g)) \quad (5.6)$$

which measures a rate of change of the volume of the internal space. In the derivation of the above formulas we used the fact that G , being compact, is unimodular thus; in particular $C_{\alpha\beta}^\alpha = 0$, C_{jk}^i being the structure constants of G . In the particular case when $g_{\alpha\beta}$ comes from a fixed Killing metric g_{ij} on G the effective mass term in (5.5) is nothing but the Casimir operator of G acting on $(W_\alpha^*)_o$. Indeed owing to the constraint (5.4) we can extend the summation of the indices α, β in $g^{\alpha\beta} T_\alpha T_\beta$ to $g^{ij} T_i T_j$ which is the Casimir operator. In general however, $g_{\alpha\beta}$ is non-constant and the term $\mathcal{D}_\mu g_{\alpha\beta}$ contributes to the effective Laplacian.

b) One-forms

Here the vector bundle F is the cotangent bundle T^*E with the typical fibre $F = \mathbb{R}^{N^*} = \mathbb{R}^{m^*} \oplus \mathbb{R}^{k^*} \oplus \mathbb{R}^{1^*}$. The effective fibre bundle $\tilde{F}_\alpha$ over M has the fibre $F_\alpha = \mathbb{R}^{N^*} \otimes_H W_\alpha^* = [\mathbb{R}^{m^*} \otimes (W_\alpha^*)_o] \oplus [\mathbb{R}^{k^*} \otimes (W_\alpha^*)_o] \oplus [\mathbb{R}^{1^*} \otimes_H W_\alpha^*]$. In other words a 1-form ϕ_A on E gives rise to a multiplet $(\phi_{\mu L}, \phi_{aL}, \phi_{\hat{a}L})$ where each term has got an extra index from the represen-

tation space W_α^* constrained by $\alpha(h)_M^L \alpha_{\cdot L} = \phi_{\cdot M}$ for the first two members of the multiplet and

$$\alpha(h)_M^L \phi_{aL} = A(h)_a^b \phi_{bM}$$

for the last one. Each member of the multiplet will in general, split into submultiplets according to the colour charge of the gauge group $K=N(H)|H$.

c) Spinor fields

The group $O(E)$ is not simply connected and we denote by $\text{Pin}(E)$ its two-fold simply connected spin covering. A spin structure^{*)} on E consists of a principal bundle $\text{Pin}E$ over E with structure group $\text{Pin}(E)$ and of a covering bundle homomorphism $\text{Pin}E \rightarrow OE$ commuting with the group homomorphism $\sigma: \text{Pin}(E) \rightarrow O(E)$. The elements of $\text{Pin}E$ are called spin frames. Thus over each orthonormal frame $e \in OE$ there sit two spin frames which differ by the transformation " ϵ " $\text{Pin}(E)$. When a spin frame s is rotated by a transformation $\Lambda \in \text{Pin}(E)$ then the corresponding orthonormal frame e is rotated by an orthogonal transformation $\sigma(\Lambda) \in O(E)$. The group homomorphism σ is given explicitly by

$$\Lambda \Gamma_A \Lambda^{-1} = \Gamma_B \sigma(\Lambda)_A^B,$$

where Γ_A are generators of the Clifford algebra of η_E satisfying

*) For a discussion of spin structures see [24-26]. To avoid complications with orientation and time orientation we have replaced $\text{Spin}_+(E)$ by $\text{Pin}(E)$.

$\{\Gamma_A, \Gamma_B\} = 2\eta_{AB}$. As we have seen in Sect.4. the G-space structure of E allows us to reduce the bundle OE to the subbundle AOE of adapted frames, mes, the structure group being reduced from $O(E)$ to $AO(E)$. Here we will take $AO(E) = O(M) \times O(S)$ as there is no gain in a subtler reduction $O(S) \rightarrow O(K) \times O(L)$. We denote by APinE the counterimage of AOE under the covering homomorphism. The elements of APinE are called adapted spin frames. Thus over each adapted orthonormal frame there sit two adapted spin frames. The structure group of APinE is $APin(E) = (Pin(M) \times Pin(S)) / Z_2$, where the quotient identifies the two "minus identities" of $Pin(M)$ and $Pin(S)$ respectively^{*)}. Let $\partial_\mu(x)$ be a fixed orthonormal frame at $x \in M$, and consider the set of all spin frames at the points of the internal space S_x over x which project onto (e_μ, e_α) with e_μ being the horizontal lift of ∂_μ . This set of spin frames is a spin structure for S_x . Thus existence of a spin structure on E implies existence of a spin structures on the internal spaces S_x , $x \in M$, but not, in general, on M.

Let us assume that the action of G on AOE lifts to an action on AP in E^{**) ; then, since H is assumed to be connected, the homomorphism $\lambda: H \rightarrow O(S)$ induces a unique homomorphism $\tilde{\lambda}: H \rightarrow Pin(S)$. Repeating now the arguments used in Sect.4. we deduce that the resulting effective gauge group is now $(Pin(M) \times N(\tilde{H}) | \tilde{H}) / Z_2$ where $\tilde{H} = \text{diag}(H, \tilde{\lambda}(H)) \subset G \times Pin(S)$. Therefore M is endowed with a generalization of a $Spin^C$ - structure (for a discussion of $Spin^C$ -structures see e.g. [29,30]). To derive the}

^{*)} The action of the "minus identity" on a spin frame corresponds to a 2π -rotation of an orthonormal frame.

^{**) Action of Lie groups on spin manifolds in a connection with G-index is discussed in [27]. See also [28], and [29] for $U(1)$ actions.}

dimensionally reduced Dirac operator we use a local moving spin frame corresponding to the orthonormal moving frame introduced at the beginning of this section. Let F be a representation space (real or complex) of the Clifford algebra $C(E)$. The generators of the representation of $\text{Pin}(E)$ on F are then

$$\Sigma^{AB} = \frac{1}{4}[\Gamma^A, \Gamma^B]$$

and the Dirac operator is ^{*)}

$$D_E = \Gamma^A(e_A + \frac{1}{2}\omega_{A,BC}\Sigma^{BC}).$$

The typical fibre F_α of the dimensionally reduced spinor of type α is then $F \times_H W_\alpha^*$. Thus the spinor ψ gets an extra index $\psi = (\psi_L)$ of the representation space W_α^* and satisfies the constraint

$$(\alpha^*(T_\alpha^\wedge) - \frac{1}{2}C_{\hat{\alpha}, \beta\gamma} \Sigma^{\beta\gamma})\psi = 0,$$

where $T_\alpha^\wedge$ ($\hat{\alpha}=1,2,\dots,\dim H$) are the generators of H . The dimensionally reduced Dirac operator can be now written down as

$$D^{\text{eff}} = D_M + D_S + \Delta_1 + \Delta_2,$$

*) For more information on the Dirac operator on manifolds, in particular, in a connection with the index theorem see [31-35], also [36] for the Dirac operator on $G|H$.

with the four terms described as follows:

- D_M is the covariant Dirac operator on M. It is given by

$$D_M = \Gamma^\mu D_\mu$$

where D_μ is the covariant gravitational + gauge derivative:

$$D_\mu = \partial_\mu + \frac{1}{2} \omega_{\mu, \nu\sigma} \Sigma^{\nu\sigma} + A_\mu^{\hat{a}} \alpha^*(T_{\hat{a}}) + \frac{1}{2} B_{\mu, \alpha\beta} \Sigma^{\alpha\beta},$$

with B_μ defined in (5.3),

- D_S is the internal Dirac operator of type α

$$D_S = \Gamma^\alpha (-\Phi_\alpha^\beta \alpha^*(T_\beta) + \frac{1}{2} \omega_{\alpha, \beta\gamma} \Sigma^{\beta\gamma})$$

- Δ_1 is the anomalous non-minimal interaction term

$$\Delta_1 = \frac{1}{4} \gamma^\alpha \Phi^{-1\beta}_\alpha \eta_{\beta\gamma} F_{\mu\nu}^\gamma \Sigma^{\mu\nu},$$

- Δ_2 is analogous to the last term of (5.5)

$$\Delta_2 = \frac{1}{2} \gamma^\mu v_\mu,$$

with v_μ given by (5.6).

Let us consider now the case of M four-dimensional of signature $(-+++)$ and S - the typical internal space of signature $(++...+)$. The Clifford

algebra $C(E)$ of E is then isomorphic to a tensor product $C(E)=C(M)\otimes C(-S)$, the isomorphism being given by $\Gamma_\mu=\gamma_\mu\otimes I$, $\Gamma_\alpha=\gamma_5\otimes\gamma_\alpha$ with $\{\gamma_\mu,\gamma_\nu\}=2\eta_{\mu\nu}$ and $\{\gamma_\alpha,\gamma_\beta\}=-2\eta_{\alpha\beta}$, where $\gamma_5=\gamma_0\gamma_1\gamma_2\gamma_3$. Since $C(M)\simeq L(\mathbb{R}^4)$ - the 4×4 real matrix algebra it follows that the real spinor space F can be realized as the tensor product $F=\mathbb{R}^4\otimes_{\mathbb{R}} F'$ where F' is a representation space of $C(-S)$. The simplest example is the five-dimensional Kaluza-Klein theory. Here S is one-dimensional and the Clifford algebra $C(-1)$ is isomorphic to the algebra $\mathbb{C}$ of complex numbers. Thus $F=\mathbb{R}^4\otimes_{\mathbb{R}} \mathbb{C}\simeq \mathbb{C}^4$. The group G is in this example $U(1)$ and all its irreducible representations α are realized on $\mathbb{C}$. Since H is now trivial the effective fibre is therefore $F_\alpha=F\otimes\mathbb{C}=\mathbb{C}^4$. The effective structure group is then $\text{Pin}^c(3,1)\equiv(\text{Pin}(3,1)\times U(1))/\mathbb{Z}_2$.

REFERENCES

- [1] A. Salam and J. Strathdee, "On Kaluza-Klein theory", *Annals of Phys.*141, 316-352(1982)
- [2] E. Witten, "Fermion quantum numbers in Kaluza-Klein theory", Preprint, Princeton, October 1983
- [3] H. Römer, "Dimensional reduction, spinor fields and characteristic classes", Preprint THEP 82/6, Universität Freiburg, September 1982
- [4] D. Bleeker, "Harmonic analysis on the space-time gauge continuum", *Int.J.Theor.Phys.*22, 557-574(1983)
- [5] D. Bleeker, "Physics from the G-bundle viewpoint", *Int.J.Theor.Phys.*23, 735-750(1984)
- [6] R. Coquereaux and A. Jadczyk, "Geometry of multidimensional universes", *Commun.Math.Phys.*90, 79-100(1983)
- [7] A. Jadczyk, "Symmetry of Einstein-Yang-Mills systems and dimensional reduction", *J.Gem.and Phys.* (to appear)
- [8] A. Jadczyk and K. Pilch, "Geometry of gauge fields in a multidimensional universe", *Lett.Math.Phys.*8, 97-104(1984)
- [9] R. Coquereaux and A. Jadczyk, "Symmetries of Einstein-Yang-Mills fields and dimensional reduction", *Commun.Math.Phys.*96, (1984)
- [10] D. Husemoller, "Fibre bundles", Springer Verlag, New York-Heidelberg-Berlin, 1975
- [11] G. Segal, "Equivariant K-theory", *IHES Publ.Math.*N^o34, 129-151(1968)
- [12] R.K. Lashof, "Equivariant bundles", *Ill.J.Math.*26, 257-271(1982)
- [13] N.R. Wallach, "Harmonic analysis on homogeneous spaces", Marcel Dekker, New York, 1973
- [14] J. Dieudonné, "Éléments d'analyse", Vol.5, Gauthier-Villars, Bordeaux, 1975
- [15] L. Palla, "On dimensional reduction of gauge theories; Symmetric Fields and Harmonic expansions", *Z.Phys.*C24, 195-204(1984)
- [16] R. Bott, "The index theorem for homogeneous differential operators, in Differential and Combinatorial Topology", A symposium in Honor

- of Marston Morse, Steward S. Cairns Ed., Princeton University Press, 1965
- [17] A.A. Kirillov, "Elements of the Theory of Representations", Springer Verlag, Berlin-Heidelberg-New York, 1976
 - [18] J. Brüning and E. Heintze, "Representations of Compact Lie groups and elliptic operators", *Inv.Math.* 169-203 (1979)
 - [19] P. Forgacs and N.S. Manton, "Space-time symmetries in gauge theories", *Commun.Math.Phys.* 72, 15-35(1980)
 - [20] J. Harnad, S. Schnider and L. Vinet, "Group actions on principal bundles and invariance conditions for gauge fields", *J.Math.Phys.* 21, 2719-2724(1980)
 - [21] J. Harnad, S. Schnider and J. Tafel, "Group actions on principal bundles and dimensional reduction", *Lett.Math.Phys.* 4, 107-113(1980)
 - [22] S. Kobayashi and K. Nomizu, "Foundations of differential geometry", Vol.I, Interscience, New York, London, 1963
 - [23] J. Dieudonné, "Éléments d'Analyse", Tome IV, Gauthier-Villars, Paris 1971
 - [24] J. Milnor, "Spin structures on manifolds", *L'Enseignement Math.* 9, 198-203(1963)
 - [25] K. Bichteler, "Global existence of spin structures for gravitational fields", *J.Math.Phys.* 9, 813-815 (1968)
 - [26] R. Geroch, "Spinor structures of space-times in general relativity I". *J.Math.Phys.* 9, 1739-1744(1968)
 - [27] M. Atiyah and F. Hirzebruch, "Spin-manifolds" and group actions", in *Essays on Topology and Related Topics, Memoires dédiés à Georges de Thame*, ed. A. Haefliger and R. Narasimhan, Springer-Verlag, Berlin-Heidelberg-New York, 1970
 - [28] C. Chichlinsky, "Group actions on spin manifolds", *Trans.Am.Math.Soc.* 172, 307-315(1972)
 - [29] A. Hattori, "Spin^C-structures and S¹-actions", *Inv.Math.* 48, 7-31
 - [30] (1978)
 - [30] S.W. Hawking and C.H. Pope, "Generalized spin structures in quantum gravity", *Phys.Lett.* 73B, 42-44(1978)

- [31] R. Bott, "The index theorem for homogeneous differential operators" in Differential and Combinatorial Topology, A symposium in Honor of Marston Morse, ed. S. Cairns, Princeton University Press, Princeton 1963
- [32] R. Palais, "Differential operators on vector bundles", in Seminar on the Atiyah-Singer index theorem, ed. R. Palais, Princeton University Press, Princeton, 1965
- [33] N. Hitchin, "Harmonic spinors", Adv. in Math. 14, 1-55(1974)
- [34] H.B. Lawson, Jr., M.L. Michelson, "Clifford bundles, immersions of manifolds and the vector field problem, J.Diff.Geom. 15, 237-267 (1980)
- [35] M.F. Atiyah, I.M. Singer, "Dirac operators coupled to vector potentials", Proc.Natl.Acad.Sci.USA, 81, 2597-2600(1984)
- [36] K. Parthasaraty, "Dirac operator and the discrete series", Ann. Math. 96, 1-30(1972)