A 84 - 254

5

CERN
BIBLIOTHEQUE

ACADEMY OF SCIENCES OF THE USSR
P. N. LEBEDEV PHYSICAL INSTITUTE

I.E.Tamm Department of Theoretical Physics

High energy and
cosmic rays physics

Preprint N 254

QUANTUM ELECTRODYNAMICS IN EXTERNAL
CONSTANT FIELD

I.A.Batalin, E.S.Pradkin,
Sh.M.Shvartsman

CERN LIBRARIES, GENEVA

CM-P00067781

Moscow, 1984

QUANTUM ELECTRODYNAMICS IN EXTERNAL CONSTANT FIELD.

I.A.BATALIN, E.S.FRADKIN

P.N.LEBEDEV PHYSICAL INSTITUTE, MOSCOW, U S S R

SH.M.SHVARTSMAN

DEPARTMENT OF MATHEMATICAL ANALYSIS, TOMSK PEDAGOGICAL

INSTITUTE, TOMSK, U S S R

Generating functional of all quantum Green functions in an external field is found. The Green function of a scalar particle in a constant and uniform external field is calculated with the help of the modified perturbation theory. Its infra-red asymptotics is studied. It is shown that the enhancement of the pole takes place, the same as in the vacuum case.

INTRODUCTION

Electromagnetic processes in external fields have a great interest during last few years. This is accounted for, on the one hand, by the creation of the intensive electromagnetic fields in lasers and the appearance of electrons and photons of high energies, and, on the other hand, by possible astrophysical applications. In the above processes one can verify quantum electrodynamics in the domain of high energies and large fields. Studying this phenomena is of fundamental importance because a correct description of the phenomena in intensive external fields is connected with the necessity of going beyond the framework of the usual perturbation theory. For this reason the processes in e^2 -approximation in different external fields [1-14] have been studied in detail lately with the external field taken into account exactly (in this connection the constant uniform field and the plane wave field or their combination are of a special interest because in these external fields one can exactly solve the Dirac and Klein-Gordon equations and find the one-particle Green functions). However, the contribution of radiative corrections at high energies and intensive fields becomes essential. For this reason it becomes of a special interest to work out such methods of calculation of radiative corrections to the processes in an external field which, in their result, go essentially beyond the scope of the usual perturbation theory with the external field kept exactly.

This is due to the fact that the generalized (with the external field being taken into account exactly) Feynman Diagram technique encounters essential difficulties caused by the necessity of correct calculation of the vertex function in a selfconsistent approximation, since otherwise the obtained results are gauge noninvariant, and the corresponding generalized Ward identities in the Presence of external

field [15-17] are not correct.

The most adequate method for solving this problem is the functional method developed in papers [17, 19]. With the help of this method a closed expression for the generating functional of the Green function is obtained in [16, 17, 20] and the modified perturbation theory is constructed.

The method of modified perturbation theory proposed in [16, 17, 20] leads to the result which goes far beyond the scope of the usual perturbation theory. This method appeared to be especially fruitful in quantum electrodynamics, because in the first order of the modified perturbation theory the contribution of the soft photons, both virtual and real, is summed completely. In particular, the first order of the modified perturbation theory gave the possibility to obtain the correct asymptotic behaviour of the Green function and the cross section of the processes in QED [16, 20, 21] in the double logarithmic approximation. Not long ago [22] the method of the modified perturbation theory was also used in statistic quantum electrodynamics.

These general results (which are beyond the scope of the usual perturbation theory) will allow us to essentially advance in solving the problem of finding quantum Green function in the presence of a real external field.

In the present paper we consider the calculation of the one-particle Green function of a scalar field in a constant uniform external field by means of the modified perturbation theory. The Green function is found in the representation of eigenfunctions of the Klein-Gordon equation in a constant field. Such functions were first obtained in papers [12, 13]. It was shown there that Dirac's eigenfunctions (in the spinor case) in a constant field diagonalize not only Green's function without radiative correction, but also Green's function in the e^2 -approximation (see also [8]). The use of eigenfunc-

tions appears fruitful in the modified perturbation theory because they contain considerable information about the dynamics of the system in a constant field.

I. GENERATING FUNCTIONAL IN QED WITH EXTERNAL FIELD

The principle value of the functional method is the generating functional of all Green's functions

$$Z[\xi^*, \xi, J] = C_V^{-1} \cdot \langle 0_{out} | S[\xi^*, \xi, J] | 0_{in} \rangle \quad (1)$$

where

$$S[\xi^*, \xi, J] = T \exp \left\{ i \int \mathcal{L}_s(x) d^4x \right\},$$

$$\mathcal{L}_s(x) = J_\mu(x) A^\mu(x) + \xi^*(x) \psi(x) + \psi^+(x) \xi(x)$$

$C_V = \langle 0_{out} | 0_{in} \rangle$ is the probability amplitude for the vacuum to remain the vacuum, $|0\rangle_{in}$, $|0\rangle_{out}$ are initial and final vacuum states, $J_\mu(x)$ is the source of electromagnetic field $A^\mu(x)$, $\xi(x)$ and $\xi^*(x)$ are sources of scalar fields $\psi^+(x)$ and $\psi(x)$ respectively. Operators are taken in the usual Heisenberg representation (i.e. the interaction between the fields is taken into account in the absence of the external sources) and satisfy the equations

$$[\mathcal{D}_\mu(x) \mathcal{D}^\mu(x) - m^2] \psi(x) = 0, \quad \mathcal{D}_\mu(x) = i \partial_\mu - e A_\mu(x) - e A_\mu^{ext}(x),$$

$$[\mathcal{D}_\mu^*(x) \mathcal{D}^{*\mu}(x) - m^2] \psi^+(x) = 0, \quad \mathcal{D}_\mu^*(x) = i \partial_\mu + e A_\mu(x) + e A_\mu^{ext}(x),$$

$$\left[\square g_{\mu\nu} + (d_e - 1) d_e^{-1} \partial_\mu \partial_\nu \right] A^\nu(x) = I_\mu(x), \quad \square = -\partial_\mu \partial^\mu,$$

$$I_\mu(x) = e \psi^\dagger(x) \mathcal{J}_\mu(x) \psi(x) - e \left(\mathcal{J}_\mu^*(x) \psi^\dagger(x) \right) \psi(x) \quad (2)$$

A_μ^{ext} is an external electromagnetic field, d_e is a gauge parameter. Using the method proposed in [16, 17, 20] one may show that the solution of the functional equations for the generating functional (1) resulting from (2) has the form:

$$\begin{aligned} Z[\xi^*, \xi, J] = & C \exp \left\{ \Pi \left(A^{\text{ext}} + \frac{\delta}{i\delta J} \right) + \right. \\ & \left. + i \int d^4x d^4y \xi^*(x) G(x, y | A^{\text{ext}} + \frac{\delta}{i\delta J}) \xi(y) \right\}^{(3)} \\ & \exp \left\{ \frac{i}{2} \int d^4x d^4y J_\mu(x) D^{\mu\nu}(x-y) J_\nu(y) \right\} \end{aligned}$$

where $G(x, y | a)$ is the causal Green function of the scalar particle in the "external" field $a_\mu(x) = A_\mu^{\text{ext}}(x) - i\delta/\delta J^\mu(x)$ satisfying the equation

$$\left[(i\partial_\mu - e a_\mu(x)) (i\partial^\mu - e a^\mu(x)) - m^2 \right] G(x, y | a) = \delta^{(4)}(x-y) \quad (4)$$

$\Pi(A^{\text{ext}} + \frac{\delta}{i\delta J})$ is the polarization correction to the action of the classical electromagnetic field.

$$\Pi(A^{\text{ext}} + \frac{\delta}{i\delta J}) = \text{Sp} \ln \left[G(A^{\text{ext}} + \frac{\delta}{i\delta J}) / G(0) \right] \quad (5)$$

C is a constant to be determined by the condition: $Z[0,0,0] = 1$.

Further it is convenient to find a functional representation for the Green function in an arbitrary field $a_\mu(x)$. To do this let us write $G(x, y|a)$ in the form of the inverse operator

$$G(x, y|a) = -i \int_0^\infty dv \exp\{i(\pi^2 - m^2 + i\varepsilon)v\} \delta^{(4)}(x-y), \quad (6)$$

$$\pi_\mu = i\partial_\mu - e a_\mu.$$

Following the paper [20] let us write the Green function (6) in the form of functional integral

$$G(x, y|a) = -i \int \frac{d^4 p}{(2\pi)^4} e^{-ip(x-y)} \int_0^\infty dv e^{-i(m^2 - i\varepsilon)v} \int D^4 t \cdot \exp\left\{-i \int_0^v [t_\mu(v') t'^\mu(v') - 2p_\mu t'^\mu(v')] dv'\right\} \cdot \phi(x, p, v|t) \quad (7)$$

where the function $\phi(x, p, v|t)$ satisfies the equation:

$$-i \frac{\partial \phi}{\partial v} = 2 \pi_\mu t'^\mu(v) \phi, \quad \phi(v=0) = 1$$

$D^4 t$ is the normalized volume element in the functional space

$$D^4 t = \prod_{v'=0}^v d^4 t(v') \left[\int \prod_{v'=0}^v d^4 t(v') \exp\left\{-i \int_0^v t_\mu(v') t'^\mu(v') dv'\right\} \right]^{-1}$$

Let us consider the subsidiary function $\tilde{\phi}(x, p, v|A^{\text{ext}}, u)$ satisfying the equation

$$-i \frac{\partial \tilde{\phi}}{\partial v} = (2 \pi_\mu t'^\mu(v) + u_\mu(v) t'^\mu(v)) \tilde{\phi}, \quad \tilde{\phi}(v=0) = 1$$

Note that $\tilde{\Phi}(u=0) = \Phi$. It is not difficult to find the solution of this equation

$$\tilde{\Phi} = \exp \left\{ -2i \int_0^v \left[e a_\mu(x - 2 \int_0^v t(\lambda) d\lambda) + \frac{1}{2} u_\mu(v') \right] t^\mu(v') dv' \right\}$$

Separating the external field A_μ^{ext} and the "radiative field" $\frac{\delta}{\delta J^\mu}$ from a_μ and making some simple transformations we obtain [20]

$$G(x, y | A^{\text{ext}} + \frac{\delta}{\delta J}) = -i \int \frac{d^4 p}{(2\pi)^4} e^{-ip(x-y)} \int_0^\infty dv e^{-i(m^2 - i\varepsilon)v} \exp \left\{ -2e \int_0^v P^\mu(v') \frac{\delta}{\delta J^\mu(x(v'))} dv' \right\} Y(x, p, v | A^{\text{ext}}, u) \Big|_{u=0} \quad (8)$$

$$Y(x, p, v | A^{\text{ext}}, u) = \int D^4 t \exp \left\{ -i \int_0^v \left[\mathcal{L}(v' | t, A^{\text{ext}}) - u_\mu(v') t^\mu(v') \right] dv' \right\}$$

where

$$\mathcal{L}(v' | t, A^{\text{ext}}) = \left[t_\mu(v') - 2p_\mu + 2e A_\mu^{\text{ext}}(x - 2 \int_0^{v'} t(\lambda) d\lambda) \right] t^\mu(v')$$

is the classical action of the particle in the external field A_μ^{ext}

$$P^\mu(v') = -i \frac{\delta}{\delta u_\mu(v')} , \quad x(v') = x - 2 \int_0^{v'} P(\lambda) d\lambda.$$

Substituting (8) into (3) we obtain for the generating functional

$Z[\xi^*, \xi, J]$ the expression

$$Z[\xi^*, \xi, J] = C \exp \left\{ \int d^4 x \int_{s_0}^\infty \frac{ds}{s} e^{-i(m^2 - i\varepsilon)s} \left(\langle 1 \rangle_{x,x} - \left\langle \exp \left(-2e \int_0^s P^\mu(s') \frac{\delta}{\delta J^\mu(x(s'))} ds' \right) \right\rangle_{x,x} \right) \right\} \times \quad (9)$$

$$\exp \left\{ \int d^4 x d^4 y \xi^*(x) \int_0^\infty dv e^{-i(m^2 - i\varepsilon)v} \left\langle \exp \left(-2e \int_0^v P^\mu(v') \frac{\delta}{\delta J^\mu(x(v'))} dv' \right) \right\rangle_{xy} \xi(y) \right\}$$

$$\exp\left\{\frac{i}{2} \int d^4x d^4y J^\mu(x) D_{\mu\nu}(x-y) J^\nu(y)\right\},$$

where

$$\begin{aligned} \langle A(\frac{\delta}{\delta u}) \rangle_{x,y} &= A(\frac{\delta}{\delta u}) \tilde{Y}(x,y,v|A^{\text{ext}},u)|_{u=0}, \\ \tilde{Y}(x,y,v|A^{\text{ext}},u) &= \int \frac{d^4p}{(2\pi)^4} e^{-ip(x-y)} Y(x,p,v|A^{\text{ext}},u), \\ D_{\mu\nu}(x) &= \int \frac{d^4k}{(2\pi)^4} D_{\mu\nu}(k) e^{ikx}, \\ D_{\mu\nu}(k) &= (g_{\mu\nu} k^2 + (d_l - 1) k_\mu k_\nu) \cdot \int_{-it_0}^{\infty} t e^{ik^2 t} dt. \end{aligned} \quad (10)$$

Some remarks about expression (9) are in order. It is known that the expression for $Z[\xi^*, \xi, J]$ contains divergencies before the programme of renormalization is carried out. It is convenient to carry out this programme after all the calculations being done. To do so it is necessary to regularize $Z[\xi^*, \xi, J]$ (9), so that the divergencies be absent while calculating. As in papers [16, 20] this can be done by introducing the cut-off S_0 for the proptime of the particle in the polarization term in (9) and the cut-off t_0 for the photon proptime in (19). As long as S_0 and t_0 are not zero, the whole expression (9) is finite. After carrying out the programme of renormalization it is necessary to remove the regularization by turning S_0 and t_0 into zero. The resulting renormalized expressions should not depend upon S_0 and t_0 .

In (9) one may carry out the partial functional differentiation over the sources J^μ of the electromagnetic field A_μ by expanding (9) into a series in powers of the sources ξ^* and ξ :

$$\begin{aligned}
 Z[\xi^*, \xi, J] &= Z[0, 0, J] + \sum_{n=1}^{\infty} \frac{1}{n!} \prod_{k=1}^n \int d^4 x_k d^4 y_k \xi^*(x_k) \xi(y_k) \\
 &\int_0^{\infty} dv_k e^{-i(m^2 - i\varepsilon)v_k} \cdot \exp \left\{ \sum_{n_1, n_2=1}^n A_{n_1, n_2} + \sum_{n_1=1}^n B_{n_1}(J) - \right. \\
 &- \int d^4 z \int_0^{\infty} \frac{ds}{s} e^{-i(m^2 - i\varepsilon)s} \left[\exp \left[\sum_{n_1=1}^n R_{n_1} - 2e \int_0^s \mathcal{G}^{\mu}(s') \frac{\delta}{\delta J^{\mu}(z(s'))} \frac{ds'}{s'} \right] \cdot \right. \\
 &\left. \tilde{Y}(z, z; s | A^{\text{ext}}, v) \Big|_{v=0} - \tilde{Y}(z, z; s | 0, 0) \right] \prod_{\ell=1}^n \tilde{Y}(x_{\ell}, y_{\ell}; v_{\ell} | A^{\text{ext}}, u_{\ell}) \Big|_{u=0} \times \\
 &\times \exp \left\{ \frac{i}{2} \int d^4 x d^4 y J^{\mu}(x) D_{\mu\nu}(x-y) J^{\nu}(y) \right\}
 \end{aligned}$$

where

$$A_{n_1, n_2} = 2ie^2 \int_0^{v_{n_1}} dv' \int_0^{v_{n_2}} dv'' P_{n_1}^{\mu}(v') D_{\mu\nu}(x_{n_1}(v') - x_{n_2}(v'')) P_{n_2}^{\nu}(v''), \quad (12)$$

$$B_{n_1}(J) = -2ie \int_0^{v_{n_1}} dv' \int d^4 x J^{\mu}(x) D_{\mu\nu}(x - x_{n_1}(v')) P_{n_1}^{\nu}(v'), \quad (13)$$

$$R_{n_1} = 4ie^2 \int_0^s ds' \int_0^{v_{n_1}} dv' \mathcal{G}^{\mu}(s') D_{\mu\nu}(z(s') - x_{n_1}(v')) P_{n_1}^{\nu}(v'), \quad (14)$$

$$P_{n_1}^{\mu}(v') = -i \frac{\delta}{\delta u_{n_1, \mu}(v')} , \quad x_{n_1}(v') = x_{n_1} - 2 \int_{v'}^{v_{n_1}} P_{n_1}(\lambda) d\lambda, \quad (15)$$

$$\mathcal{G}^{\mu}(s') = -i \frac{\delta}{\delta v_{\mu}(s')} , \quad z(s') = z - 2 \int_{s'}^s \mathcal{G}(\lambda) d\lambda$$

and

$$Z[0,0,J] = C \cdot \exp\left(\Pi(A^{\text{ext}} + \frac{\delta}{i\delta J})\right) \exp\left\{\frac{i}{2} \int d^4x d^4y J^\mu(x) D_{\mu\nu}(x-y) J^\nu(y)\right\}$$

is the generating functional for the photon Green's functions. To

carry out the complete functional differentiation over the sources

J_μ one should expand $\Pi(A^{\text{ext}} + \frac{\delta}{i\delta J})$ into a series with respect to the functions $G(A^{\text{ext}} + \delta/i\delta J)$. In this way we obtain the following expression for $Z[\bar{\epsilon}^*, \bar{\epsilon}, J]$

$$Z[\bar{\epsilon}^*, \bar{\epsilon}, J] = Z[0,0,J] + \sum_{n=1}^{\infty} \frac{1}{n!} \prod_{K=1}^n \int d^4x_K d^4y_K \bar{\epsilon}^*(x_K) \bar{\epsilon}(y_K)$$

$$\begin{aligned} & \int d\nu_K e^{-i(m^2 - i\epsilon)\nu_K} \exp\left\{ \sum_{n_1, n_2=1}^n A_{n_1, n_2} + \sum_{n_1=1}^n B_{n_1}(J) + \right. \\ & + \int d^4z \int_{s_0}^{\infty} \frac{ds_0}{s} e^{-i(m^2 - i\epsilon)s} \tilde{Y}(z, z, s | 0, 0) \left. \right\} \cdot \\ & \left\{ 1 + \sum_{\tau=1}^{\infty} \frac{(-1)^\tau}{\tau!} \prod_{P=1}^{\tau} \int d^4z_P \int_{s_0}^{\infty} \frac{ds_P}{s_P} e^{-i(m^2 - i\epsilon)s_P} \right. \quad (16) \\ & \exp\left\{ \sum_{n_1=1}^n \sum_{\alpha=1}^{\tau} C_{n_1, \alpha} + \sum_{\alpha=1}^{\tau} B_{1\alpha}(J) + \sum_{\alpha, \alpha_1=1}^{\tau} A_{1\alpha, \alpha_1} \right\} \cdot \\ & \left. \times \prod_{\alpha=1}^{\tau} \tilde{Y}(z_\alpha, z_\alpha, s_\alpha | A^{\text{ext}}, \nu_\alpha) \Big|_{\nu=0} \right\} \cdot \end{aligned}$$

$$\prod_{\ell=1}^n \tilde{Y}(x_\ell, y_\ell, \nu_\ell | A^{\text{ext}}, u_\ell) \Big|_{u=0} \cdot \exp\left\{\frac{i}{2} \int d^4x d^4y J^\mu(x) D_{\mu\nu}(x-y) J^\nu(y)\right\},$$

where

$$C_{n_1, \alpha} = 4ie^2 \int_0^{\nu_{n_1}} dv' \int_0^{s_\alpha} ds' P_{n_1}^\mu(v') D_{\mu\nu}(x_{n_1}(v') - z_\alpha(s')) \mathcal{G}_\alpha^\nu(s') \quad (17)$$

$$\begin{aligned}
 B_{i\alpha}(J) &= -2ie \int_0^{S_\alpha} ds' \int d^4x J^\mu(x) D_{\mu\nu}(x - z_\alpha(s')) \mathcal{P}_\alpha^\nu(s'), \\
 A_{i\alpha, \alpha_1} &= 2ie^2 \int_0^{S_\alpha} ds' \int_0^{S_{\alpha_1}} ds'' \mathcal{P}_\alpha^\mu(s') D_{\mu\nu}(z_\alpha(s') - z_{\alpha_1}(s'')) \mathcal{P}_{\alpha_1}^\nu(s''), \\
 \mathcal{P}_\alpha^\mu(s') &= -i \frac{\delta}{\delta \mathcal{U}_{\alpha\mu}(s')} , \quad z_\alpha(s') = z_\alpha - 2 \int_{s'}^{S_\alpha} \mathcal{P}_\alpha(\lambda) d\lambda.
 \end{aligned} \quad (18)$$

The expression for $Z[\xi^*, \xi, J]$ will have the simplest form if we neglect the polarization effects. In this case $Z[0, 0, J]$ becomes trivial, and we will not write out this term in what follows.

Then

$$\begin{aligned}
 Z[\xi^*, \xi, J] &= \sum_{n=1}^{\infty} \frac{1}{n!} \prod_{k=1}^n \int d^4x_k d^4y_k \xi^*(x_k) \xi(y_k) \times \\
 &\cdot \int_0^{\infty} d\nu_k e^{-i(m^2 - i\varepsilon)\nu_k} \exp\left\{ \sum_{n_1, n_2=1}^n A_{n_1, n_2} + \sum_{n_1=1}^n B_{n_1}(J) \right\}^* \quad (19)
 \end{aligned}$$

$$\prod_{\ell=1}^n \tilde{Y}(x_\ell, y_\ell, \nu_\ell | A^{\text{ext}}, u_\ell) \Big|_{u=0} \cdot \exp\left\{ \frac{i}{2} \int d^4x d^4y J^\mu(x) D_{\mu\nu}(x-y) J^\nu(y) \right\}.$$

If we are interested only in the Green functions of the scalar particle, the expression for the generating functional becomes still more simplified

$$\begin{aligned}
 Z[\xi^*, \xi, J] &= \sum_{n=1}^{\infty} \frac{1}{n!} \prod_{k=1}^n \int d^4x_k d^4y_k \xi^*(x_k) \xi(y_k) \cdot \\
 &\exp\left\{ \sum_{n_1, n_2=1}^n A_{n_1, n_2} \right\} \prod_{\ell=1}^n \tilde{Y}(x_\ell, y_\ell, \nu_\ell | A^{\text{ext}}, u_\ell) \Big|_{u=0}.
 \end{aligned} \quad (20)$$

Using (16) one can write the one-particle quantum Bose Green's function in the form:

$$G(x, y) = - \int_0^\infty dv \exp \{ -i(m^2 - i\varepsilon)v + A_{1,1} \} \cdot \left(1 + \sum_{\tau=1}^\infty \frac{(-1)^\tau}{\tau!} \prod_{p=1}^\tau \int d^4 z_p \int_{s_0}^{s_p} \frac{ds_p}{s_p} e^{-i(m^2 - i\varepsilon)s_p} \cdot \exp \left\{ \sum_{\alpha=1}^\tau C_{1\alpha} + \sum_{\alpha, \alpha_1=1}^\tau A_{1\alpha, \alpha_1} - \int d^4 z \int_{s_0}^{s_p} e^{-i(m^2 - i\varepsilon)s} \langle 1 \rangle_{z, z} \right\} \prod_{\alpha=1}^\tau \tilde{Y}(z_\alpha, z_\alpha, s_\alpha | A^{\text{ext}}, v_\alpha) \right) \tilde{Y}(x, y, v | A^{\text{ext}}, u) \Big|_{u=0} \quad (21)$$

When neglecting polarization effects this expression takes an especially simple form:

$$G(x, y) = - \int_0^\infty dv \exp \{ -i(m^2 - i\varepsilon)v \} \cdot \langle \exp A_{1,1} \rangle_{x, y} \quad (22)$$

Consider now expression (20) in detail. Suppose we know a complete set of solutions of the eigenvalue problem.

$$\pi_\mu^{\text{ext}}(x) \pi_\mu^{\text{ext}}(x) \psi_{\{K\}}(x) = p^2 \psi_{\{K\}}(x), \quad \pi_\mu^{\text{ext}}(x) = i \partial_\mu \varphi A_\mu^{\text{ext}}(x) \quad (23)$$

where $\{K\}$ is a set of quantum numbers ($\{K\}$ contains the quantum number p^2). Let us call such functions eigenfunctions. In the constant uniform field such functions were first obtained in [12, 13].

Suppose, further, that the eigenfunctions form a complete orthonormalized system of functions with respect to the scalar product

$$(\varphi, \psi) \equiv \int d^4 x \varphi^*(x) \psi(x) \quad (24)$$

$$\begin{aligned} (\psi_{\{K\}}, \psi_{\{K'\}}) &= \delta_{\{K\}, \{K'\}} \\ \sum_{\{K\}} \psi_{\{K\}}(x) \psi_{\{K\}}^*(y) &= \delta^{(4)}(x-y) \end{aligned} \quad (25)$$

(If the eigenvalues are discrete, then $\delta_{\{k\}, \{k'\}}$ in (24) is Kronecker's symbol and if they are continuous, then $\delta_{\{k\}, \{k'\}}$ is the δ - function). With the help of the eigenfunctions one can construct the causal Green function of the Bose particle in the external field

A_μ^{ext} without radiative corrections:

$$G^{(0)}(x, y | A^{\text{ext}}) = \sum_{\{k\}} \frac{\psi_{\{k\}}(x) \psi_{\{k\}}^*(y)}{p^2 - m^2 + i\varepsilon} \quad (26)$$

which satisfies the equation

$$\{\pi_\mu^{\text{ext}}(x) \pi^{\text{ext}\mu}(x) - m^2\} G^{(0)}(x, y | A^{\text{ext}}) = \delta^{(4)}(x - y) \quad (27)$$

If we represent $(p^2 - m^2 + i\varepsilon)^{-1}$ in an exponential form, Green's function (26) can be written as

$$G^{(0)}(x, y | A^{\text{ext}}) = -i \int_0^\infty d\nu e^{-i(m^2 - i\varepsilon)\nu} \sum_{\{k\}} e^{ip^2\nu} \psi_{\{k\}}(x) \psi_{\{k\}}^*(y) \quad (28)$$

On the other hand, the Green function $G^{(0)}(x, y | A^{\text{ext}})$ can be obtained using expression (8)

$$G^{(0)}(x, y | A^{\text{ext}}) = -i \int_0^\infty d\nu e^{-i(m^2 - i\varepsilon)\nu} \tilde{Y}(x, y, \nu | A^{\text{ext}}, 0) \quad (29)$$

Comparing (28) and (29) we have

$$\tilde{Y}(x, y, \nu | A^{\text{ext}}, 0) = \sum_{\{k\}} e^{ip^2\nu} \psi_{\{k\}}(x) \psi_{\{k\}}^*(y) \quad (30)$$

Using (24) and (30) we obtain an important property of eigenfunctions

$$\int d^4x d^4y \psi_{\{k\}}^*(x) \tilde{Y}(x, y, \nu | A^{\text{ext}}, 0) \psi_{\{k'\}}(y) = e^{ip^2\nu} \delta_{\{k\}, \{k'\}} \quad (31)$$

Let us represent the sources $\xi^*(x)$ and $\xi(y)$ as a superposition of the eigenfunctions

$$\Xi^*(x) = \sum_{\{K\}} \alpha_{\{K\}}^* \psi_{\{K\}}^*(x), \quad \Xi(x) = \sum_{\{K\}} \alpha_{\{K\}} \psi_{\{K\}}(x)$$

Then the expression for the generating functional $Z[\alpha^*, \alpha]$ (20) has the form

$$Z[\alpha^*, \alpha] = \sum_{n=1}^{\infty} \frac{1}{n!} \sum_{\{K_1\}, \dots, \{K_n\}, \{K'_1\}, \dots, \{K'_n\}} \prod_{p=1}^n \alpha_{\{K_p\}}^* \alpha_{\{K'_p\}} \quad (32)$$

$$\int_0^{\infty} d\nu_p e^{-i(m^2 - i\varepsilon)\nu_p} \ll \exp \left\{ \sum_{n_1, n_2=1}^n A_{n_1, n_2} \right\} \gg_{K_1, \dots, K_n; K'_1, \dots, K'_n}$$

where

$$\begin{aligned} \ll A\left(\frac{\delta}{\delta u}\right) \gg_{K_1, \dots, K_n; K'_1, \dots, K'_n} &= \int d^4x_1 \dots d^4x_n d^4y_1 \dots d^4y_n \times \\ &\times \prod_{\ell=1}^n \psi_{\{K_\ell\}}^*(x_\ell) A\left(\frac{\delta}{\delta u}\right) \prod_{p=1}^n \tilde{Y}(x_p, y_p, \nu_p | A^{\text{ext}}, u_p) \Big|_{u=0^+} \cdot \psi_{\{K'_p\}}(y_p)^{(33)} \end{aligned}$$

Expression (32) allows one to find the representation of Green's functions in the basis of eigenfunctions.

II. GREEN'S FUNCTIONS IN CONSTANT FIELD

Consider the constant and uniform external field

$$A_m^{\text{ext}}(x) = -\frac{1}{2} F_m{}^\nu x^\nu \quad (1)$$

As seen from the above said the calculation of the generating functional and the Green functions reduces to the calculation of the functional integral (1.8), i.e. the function $\tilde{Y}(x, y, \nu | A^{\text{ext}}, u)$.

in the constant field (1) the functional integral (1.8) reduces to the Gaussian one. The function $Y(x, p, v|A^{\text{ext}}, u)$ was found in ref. [23].

$$Y(x, p, v|A^{\text{ext}}, u) = [\det \text{ch} eFv]^{-\frac{1}{2}} \exp \left\{ i \int_0^v dv' \pi^\mu(x, p, v') \cdot G_{\mu\nu}(v', v''; v) \pi^\nu(x, p, v'') \right\}, \quad (2)$$

where

$$\pi_\mu(x, p, v') = p_\mu + \frac{e}{2} F_{\mu\nu} x^\nu + \frac{1}{2} u_\mu(v')$$

$$G(v', v''; v) = \delta(v' - v'') + eF \exp\{2eF(v' - v'')\} \cdot [\varepsilon(v' - v'') + \text{th} eFv] \quad (3)$$

$$\varepsilon(v' - v'') = \begin{cases} 1, & v' > v'' \\ 0, & v' = v'' \\ -1, & v' < v'' \end{cases}$$

Thus, the integral over p_μ in (1.10) also reduces to the Gaussian form and we find:

$$\tilde{Y}(x, y, v|A^{\text{ext}}, u) = \frac{i}{(4\pi)^2} \exp \left\{ \frac{e}{2} x F y - i(x-y)L(v)(x-y) - \frac{1}{2} \text{Sp} \ln \frac{\text{sh} eFv}{eF} \right\} K(x-y, v|u), \quad L(v) = \frac{eF}{4} \text{ct} h eFv,$$

$$K(x-y, v|u) = \exp \left\{ \frac{i}{4} \int_0^v dv' dv'' u^\mu(v') \tilde{G}_{\mu\nu}(v', v''; v) u^\nu(v'') + \frac{i}{2} (x-y) \frac{eF \exp(eFv)}{\text{sh} eFv} \int_0^v \exp(-2eFv') u(v') dv' \right\}, \quad (4)$$

$$\tilde{G}(v', v''; v) = G(v', v''; v) - \frac{2eF}{\text{sh} eFv} \exp\{2eF(v' - v'')\}.$$

Substituting expression (4) into (1.21) we find quantum Green's function of the scalar particle.

$$\begin{aligned}
 G(x, y) = & \frac{i}{(4\pi)^2} \int_0^\infty dv \exp \left\{ -i(m^2 - i\varepsilon)v + A_{1,1} - \int d^4z \int_{S_0}^\infty \frac{ds}{s} \cdot \right. \\
 & \cdot e^{-i(m^2 - i\varepsilon)s} \langle 1 \rangle_{z,z} \left. \right\} \left(1 + \sum_{\tau=1}^\infty \frac{(-1)^\tau}{\tau!} \prod_{\rho=1}^\tau \int d^4z_\rho \int_{S_0}^\infty \frac{ds_\rho}{s_\rho} e^{-i(m^2 - i\varepsilon)s_\rho} \right. \\
 & \exp \left\{ \sum_{\alpha=1}^z C_{1\alpha} + \sum_{\alpha, \alpha_1=1}^z A_{1\alpha, \alpha_1} \right\} \frac{i^\tau}{(4\pi)^{2\tau}} \prod_{\alpha=1}^\tau \exp \left\{ -\frac{1}{2} S_\rho \ln \frac{shefs_\alpha}{eF} \right. \\
 & \left. + \frac{1}{4} \iint_{S_a} ds' ds'' \tilde{U}_a^\mu(s') \tilde{G}_{\mu\nu}(s', s''; S_a) \tilde{U}_a^\nu(s'') \right\} \Big|_{v=0} \cdot \\
 & \exp \left\{ i \frac{e}{2} x F y - i(x-y) L(v)(x-y) - \frac{1}{2} S_\rho \ln \frac{shefv}{eF} + \frac{i}{4} \iint_0^v dv' dv'' \cdot \right. \\
 & \left. U^\mu(v') \tilde{G}_{\mu\nu}(v', v''; v) U^\nu(v'') + \frac{i}{2} (x-y) \frac{eFv}{shefv} \int_0^v e^{-2eFv'} U(v') dv' \right\} \Big|_{u=0}
 \end{aligned} \tag{5}$$

If we perform the substitution $z_p + x = z'_p$ in the integral over z_p in (5) it is not difficult to see that the Green function $G(x, y)$ has the following structure:

$$G(x, y) = \exp \left\{ i \frac{e}{2} x F y \right\} G_1(x-y) \tag{6}$$

where $G_1(x-y)$ is the scalar function of the difference $x-y$ and of the constant tensor F . Hence it follows that $G_1(x-y)$ may depend only upon the arguments of the form $(x-y)K(F)(x-y)$, where $K(F)$ is an even function of the field.

The eigenfunctions of the problem (1.23) in the external field (1) can be found in a covariant (see Appendix):

$$\psi_{\{K\}}(x) = N_{\{K\}} \exp\{i\epsilon\phi(x) - i\rho_1(n-x) - i\rho_2(m-x) - \frac{1}{2}\rho^2\},$$

$$H_n(\rho) D_v[we^{i\frac{\pi}{4}\tau}] \quad (7)$$

where $\{K\} = \{\rho_1, \rho_2, n, \rho^2, \omega\}$, $\omega = \pm 1$, $n = 0, 1, 2, \dots$; the numbers ρ_1, ρ_2 are eigenvalues of the operators $i\partial/\partial(n-x)$ and $i\partial/\partial(m-x)$ respectively, $v = -\frac{1}{2}(1+i\lambda)$, $\lambda = [\rho^2 + |e|\mathcal{H}(2n+1)]/|e|\mathcal{E}$, $\mathcal{E}, \mathcal{H} = [(\mathcal{F}^2 + \mathcal{Y}^2)^{\frac{1}{2}} \mp \mathcal{F}]^{\frac{1}{2}}$; $\mathcal{Y}, \mathcal{F}$ are the invariants of the field (which we suppose are not zero), $\rho = (|e|\mathcal{H})^{\frac{1}{2}}((m+x) + \rho_2/|e|\mathcal{H})$, $\tau = (2|e|\mathcal{E})^{\frac{1}{2}}((n+x) + \rho_1/|e|\mathcal{E})$, $\phi(x) = -\frac{1}{2}\mathcal{E}(n+x)(n-x) - \frac{1}{2}\mathcal{H}(m+x)(m-x)$, $H_n(\rho)$ are Hermite polynomials, $D_v[we^{i\frac{\pi}{4}\tau}]$ are the parabolic cylinder functions, $n_{\pm}, m_{\pm}$ are the vectors which are solutions of the equations

$$Fn_{\pm} = \mathcal{E}n_{\mp}, \quad Fm_{\pm} = \mp\mathcal{H}m_{\mp}, \quad n_{\pm}^2 = \pm 1,$$

$$m_{\pm}^2 = -1, \quad (n_+n_-) = (n_+m_{\pm}) = (n_-m_{\pm}) = (m_-m_+) = 0;$$

$$N_{\{K\}} = \exp\{i\frac{\pi v}{4}\} \Gamma(-v) (\mathcal{H}/2\mathcal{E})^{\frac{1}{4}} (2^n n! \sqrt{\pi})^{-\frac{1}{2}}.$$

Using the integral representation for the parabolic cylinder functions, one may show that the eigenfunctions (7) form a complete orthonormal system of functions in the sense of (1.24) and (1.25). Note that the following important property of eigenfunctions (7) is valid.

$$\int dx dy \psi_{\{K\}}^*(x) \exp\{i\frac{\epsilon}{2}xFy - i(x-y)\Gamma(F)(x-y)\} \psi_{\{K'\}}(y) \sim \delta_{\{K\},\{K'\}} \quad (8)$$

Relation (8) permits us to prove that eigenfunctions (7) diagonalize not only the Green function $G^{(a)}(x, y | A^{ext})$, but also the quantum Green function $G(x, y)$ (5). Indeed, suppose that the function

$G_1(x-y)$ can be expanded into Fourier integral

$$G_1(x-y) = \int d\kappa_1 \dots d\kappa_\ell \cdot f(\kappa_1, \dots, \kappa_\ell) \exp\left\{-i(x-y) \sum_{j=1}^{\ell} \kappa_j K_j(F)(x-y)\right\} \quad (9)$$

Here ℓ is the number of the independent structure matrices $K(F)$ upon which the function $G_1(x-y)$ may depend. Then we have

$$\int d^4x d^4y \psi_{1\kappa}^*(x) G(x, y) \psi_{1\kappa'}(y) = \int d\kappa_1 \dots d\kappa_\ell \cdot f(\kappa_1, \dots, \kappa_\ell) \times \quad (10)$$

$$\int d^4x d^4y \psi_{1\kappa}^*(x) \exp\left\{i \frac{e}{2} x F y - i(x-y) \sum_{j=1}^{\ell} \kappa_j K_j(F)(x-y)\right\} \psi_{1\kappa'}(y).$$

Choosing in (8) $T(F) = \sum_{j=1}^{\ell} \kappa_j K_j(F)$ we obtain

$$\int d^4x d^4y \psi_{1\kappa}^*(x) G(x, y) \psi_{1\kappa'}(y) \sim \delta_{1\kappa, 1\kappa'} \quad (11)$$

Proceeding from this we can write

$$G(x, y) = \sum_{\omega=\pm 1} \sum_{n=0}^{\infty} \int \frac{dP_1 dP_2 dP^2}{(2\pi)^4} \psi_{1\kappa}(x) G_{1\kappa} \psi_{1\kappa}^*(y) \quad (12)$$

$$G_{1\kappa} = \int d^4x d^4y \psi_{1\kappa}^*(x) G(x, y) \psi_{1\kappa}(y) \quad (13)$$

Neglecting the polarization effects we obtain for $G(x, y)$

$$G_{1\kappa} = \frac{i}{(4\pi)^2} \int_0^{\infty} dv e^{-i(m^2 - i\varepsilon)v} \ll \exp A_{1,1} \gg_{\kappa, \kappa} \quad (14)$$

Let us rewrite the definition of the average $\ll \gg_{\kappa, \kappa}$ in a more convenient, for our subsequent work, form

$$\ll A\left(\frac{\delta}{\delta u}\right) \gg_{\kappa, \kappa} = \int d^4x d^4y \psi_{1\kappa}^*(x) \langle A\left(\frac{\delta}{\delta u}\right) \rangle_{x, y} \psi_{1\kappa}(y) \quad (15)$$

Relation (14) permits to develop the modified perturbation theory for $C_{\{K\}}$. To obtain the first order of the modified perturbation theory let us expand $G_{\{K\}}$ into the series with respect to radiative interaction. To within e^2 we have

$$G_{\{K\}} = \frac{i}{(4\pi)^2} \int_0^\infty dv e^{-i(m^2 - i\varepsilon)v} J_{\{K\}}(v) \times \quad (16)$$

$$(1 + \langle\langle A_{1,1} \rangle\rangle_{K,K} / J_{\{K\}}(v)) , \quad J_{\{K\}}(v) = \langle\langle 1 \rangle\rangle_{K,K}$$

According to [16, 17, 20] the first order of the modified perturbation theory consists in the replacement

$$1 + \langle\langle A_{1,1} \rangle\rangle_{K,K} / J_{\{K\}}(v) \rightarrow \exp\{ \langle\langle A_{1,1} \rangle\rangle_{K,K} / J_{\{K\}}(v) \}$$

Thus in the first order of the modified perturbation theory one has

$$\langle\langle \exp A_{1,1} \rangle\rangle_{K,K} = J_{\{K\}}(v) \exp\{ \langle\langle A_{1,1} \rangle\rangle_{K,K} / J_{\{K\}}(v) \} \quad (17)$$

and for the Green functions in this approximation we obtain

$$G_{\{K\}}^{(1)} = \frac{i}{(4\pi)^2} \int_0^\infty dv \cdot J_{\{K\}}(v) \exp\{ -i(m^2 - i\varepsilon)v + \frac{\langle\langle A_{1,1} \rangle\rangle_{K,K}}{J_{\{K\}}(v)} \} \quad (18)$$

Substituting the solutions (7) into (16) and using expression (A.25) we find

$$J_{\{K\}}(v) = i(4\pi)^2 \exp(ip^2 v) \quad (19)$$

Finally we have

$$G_{\{K\}}^{(1)} = - \int_0^\infty dv \exp\{ i(p^2 - m^2 + i\varepsilon)v + \frac{\langle\langle A_{1,1} \rangle\rangle_{K,K}}{J_{\{K\}}(v)} \} \quad (20)$$

III. CALCULATION OF THE $\ll A_{1,1} \gg_{K,K}$

Using the definition of $\ll A_{1,1} \gg_{K,K}$ (2.15) we can write

$$\ll A_{1,1} \gg_{K,K} = \int d^4x d^4y \psi_{iK}^*(x) \langle A_{1,1} \rangle_{x,y} \psi_{iK}(y), \quad (1)$$

$$\langle A_{1,1} \rangle_{x,y} = -\frac{2e^2}{(4\pi)^2} \exp\left\{i\frac{e}{2}xFy - i(x-y)L(v)(x-y) - \right. \quad (2)$$

$$\left. -\frac{1}{2}\zeta_p \ln \frac{sheFv}{eF}\right\} \cdot A_{1,1}(x-y, v) \quad (3)$$

$$A_{1,1}(x-y, v) = \int_0^y dv' dv'' a(x-y, v', v'', v) \quad (4)$$

$$a(x-y, v', v'', v) = P^M(v') D_{\mu\nu}(x(v') - x(v'')) P^{\nu}(v'') K(x-y, v|u)_{u=0} \quad (4)$$

where $K(x-y, v|u)$ is determined by (2.4). To find (4) let us use the integral representation of the photon Green function

$$D_{\mu\nu}(x(v') - x(v'')) = \int \frac{d^4k}{(2\pi)^4} D_{\mu\nu}(k) \exp\left\{2i \int_0^y t(\lambda; v', v'') k_\lambda P^\lambda(\lambda) d\lambda\right\} \quad (5)$$

$$t(\lambda; v', v'') = \theta(\lambda - v'') - \theta(\lambda - v')$$

As a result we obtain:

$$a(x-y, v', v'', v) = \int \frac{d^4k}{(2\pi)^4} D_{\mu\nu}(k) \exp\{iK\mathcal{L}k + 2i\varphi_K\} \times \quad (6)$$

$$\left\{-\frac{i}{2}\tilde{G}^{\mu\nu}(v', v'', v) + [K_\beta f_1^{\nu\beta}(v', v'', v) + f_3^\nu(v', v'')][K_\alpha f_2^{\mu\alpha}(v', v'', v) + f_3^\mu(v', v'')]\right\},$$

$$f_1(v', v'', v) = \int_0^y \tilde{G}(v'', \lambda; v) t(\lambda; v', v'') d\lambda,$$

$$f_2(v', v'', v) = \int_0^y \tilde{G}(v', \lambda; v) t(\lambda; v', v'') d\lambda,$$

$$f_3(v', v) = \frac{1}{2}(x-y) \frac{eF}{sheFv} \exp\{eF(v-2v')\}, \quad (7)$$

$$\mathcal{L} = \int_0^V \int d\lambda d\lambda' t(\lambda; v', v'') \tilde{G}(\lambda, \lambda'; v) t(\lambda'; v', v''),$$

$$q = \int_0^V f_3(\lambda, v) t(\lambda; v', v'') d\lambda$$

The integral over λ and λ' in (7) are not difficult to be calculated

$$\mathcal{L} = \frac{\text{sh} eFS \text{sh} eF(v-s)}{eF \text{sh} eFv}, \quad s = |v' - v''|$$

$$q = \frac{1}{4}(x-y) \frac{e^{eFv}}{\text{sh} eFv} \left[e^{-2eFv'} - e^{-2eFv''} \right] \quad (8)$$

$$f_1(v', v'', v) = \frac{1}{2} \varepsilon(v' - v'') + \frac{1}{2} \left(e^{-2eF(v' - v'')} - 1 \right) (\varepsilon(v' - v'') + c \text{th} eFv)$$

$$f_2(v', v'', v) = \frac{1}{2} \varepsilon(v' - v'') + \frac{1}{2} \left(e^{2eF(v' - v'')} - 1 \right) (\varepsilon(v' - v'') - c \text{th} eFv)$$

To integrate expression (6) over K_μ it is convenient to use representation (1.10) having rewritten it first in a more convenient form:

$$D_{\mu\nu}(K) = D_{\mu\nu}^d(K) + D_{\mu\nu}^x(K),$$

$$D_{\mu\nu}^d(K) = g_{\mu\nu} (K^2 + i\varepsilon)^{-1} = -ig_{\mu\nu} \int_{-i0}^{\infty} e^{iK^2 t} dt, \quad (9)$$

$$D_{\mu\nu}^x(K) = -K_\mu K_\nu (d-1) \int_{-i0}^{\infty} t e^{iK^2 t} dt.$$

In accordance with this division the expression (6) is convenient to be represented in the form of two terms, too

$$a(x-y, v', v'', v) = a^d(x-y, v', v'', v) + a^x(x-y, v', v'', v) \quad (10)$$

Then for $\alpha^d(x-y, v', v'', v)$ and $\alpha^x(x-y, v', v'', v)$ we obtain, respectively

$$\begin{aligned} \alpha^d(x-y, v', v'', v) = & -i \left[-\frac{i}{2} \tilde{G}_\mu^M(v', v'', v) + f_3^M(v', v) f_{3\mu}^M(v'', v) \right] \int_{-it_0}^{\infty} I dt \\ & - i f_1^{\mu\beta}(v', v'', v) f_{2\mu}^d(v', v'', v) \int_{-it_0}^{\infty} I_{\alpha\beta} dt - i \left(f_1^{\mu\beta}(v', v'', v) f_{3\mu}^M(v', v) + \right. \\ & \left. + f_2^{\mu\beta}(v', v'', v) f_{3\mu}^M(v'', v) \right) \int_{-it_0}^{\infty} I_\beta dt, \end{aligned} \quad (11)$$

$$\begin{aligned} \alpha^x(x-y, v', v'', v) = & -(de-1) \left\{ \left[-\frac{i}{2} \tilde{G}_\mu^{\mu\sigma}(v', v'', v) + f_3^M(v', v) f_3^\sigma(v'', v) \right] \times \right. \\ & \cdot \int_{-it_0}^{\infty} I_{\mu\sigma} dt + f_1^{\sigma\beta}(v', v'', v) f_2^{\mu d}(v', v'', v) \int_{-it_0}^{\infty} I_{\mu\sigma\alpha\beta} dt + \\ & \left. + \left(f_1^{\mu\sigma\beta}(v', v'', v) f_3^\mu(v', v) + f_2^{\mu\beta}(v', v'', v) f_3^\sigma(v'', v) \right) \int_{-it_0}^{\infty} I_{\mu\sigma\beta} dt \right\}, \end{aligned} \quad (12)$$

$$I_{\alpha_1 \dots \alpha_n} = \int \frac{d^4 k}{(2\pi)^4} K_{\alpha_1} \dots K_{\alpha_n} \exp\{ik \Lambda k + 2iqk\}, \quad (13)$$

$$\Lambda = \mathcal{L} + t$$

All integrals $I_{\alpha_1 \dots \alpha_n}$ involved in (11) and (12) are expressed by means of the integral I :

$$I = \int \frac{d^4 k}{(2\pi)^4} \exp(ik \Lambda k + 2iqk) = \frac{i}{(4\pi)^2} (\det \Lambda)^{-\frac{1}{2}} \exp(-iq \bar{\Lambda}^{-1} q) \quad (14)$$

$$I_\alpha = -(\bar{\Lambda}^{-1} q)_\alpha I, \quad I_{\alpha\beta} = \left[\frac{i}{2} \bar{\Lambda}_{\alpha\beta}^{-1} + (\bar{\Lambda}^{-1} q)_\alpha (\bar{\Lambda}^{-1} q)_\beta \right] \cdot I,$$

$$I_{\mu\alpha\beta} = - \left[\frac{i}{2} (\bar{\Lambda}_{\mu\alpha}^{-1} (\bar{\Lambda}^{-1} q)_\beta + \bar{\Lambda}_{\mu\beta}^{-1} (\bar{\Lambda}^{-1} q)_\alpha + \bar{\Lambda}_{\alpha\beta}^{-1} (\bar{\Lambda}^{-1} q)_\mu) \right] \cdot I$$

$$+ (\bar{\Lambda}^{-1} q)_\mu (\bar{\Lambda}^{-1} q)_\alpha (\bar{\Lambda}^{-1} q)_\beta \}. I,$$

$$\begin{aligned} I_{\mu\nu\alpha\beta} = & \left\{ -\frac{1}{4} \left(\bar{\Lambda}^{-1}_{\mu\nu} \bar{\Lambda}^{-1}_{\alpha\beta} + \bar{\Lambda}^{-1}_{\mu\alpha} \bar{\Lambda}^{-1}_{\nu\beta} + \bar{\Lambda}^{-1}_{\mu\beta} \bar{\Lambda}^{-1}_{\nu\alpha} \right) + \right. \\ & + \frac{i}{2} \left(\bar{\Lambda}^{-1}_{\mu\nu} (\bar{\Lambda}^{-1} q)_\alpha (\bar{\Lambda}^{-1} q)_\beta + \bar{\Lambda}^{-1}_{\mu\alpha} (\bar{\Lambda}^{-1} q)_\nu (\bar{\Lambda}^{-1} q)_\beta + \right. \\ & + \bar{\Lambda}^{-1}_{\mu\beta} (\bar{\Lambda}^{-1} q)_\nu (\bar{\Lambda}^{-1} q)_\alpha + \bar{\Lambda}^{-1}_{\nu\alpha} (\bar{\Lambda}^{-1} q)_\mu (\bar{\Lambda}^{-1} q)_\beta + \bar{\Lambda}^{-1}_{\nu\beta} (\bar{\Lambda}^{-1} q)_\mu (\bar{\Lambda}^{-1} q)_\alpha + \\ & \left. \left. + \bar{\Lambda}^{-1}_{\alpha\beta} (\bar{\Lambda}^{-1} q)_\mu (\bar{\Lambda}^{-1} q)_\nu \right) + (\bar{\Lambda}^{-1} q)_\mu (\bar{\Lambda}^{-1} q)_\nu (\bar{\Lambda}^{-1} q)_\alpha (\bar{\Lambda}^{-1} q)_\beta \right\} \cdot I. \end{aligned} \quad (15)$$

Substituting (15) into (11) and (12) we obtain after cumbersome calculations

$$\begin{aligned} Q^d(x-y, \nu', \nu'', \nu) = & (4\pi)^{-2} \int_{-it_0}^{\infty} (\det \Lambda)^{-\frac{1}{2}} \exp\{-i(x-y) \frac{\tau_1 \bar{\Lambda}^{-1}}{8Sh^2 eFV} (x-y)\} \times \\ & \left\{ -2i\delta(s) + 2^{-5}(x-y) \frac{2\bar{\Lambda}^{-2}}{Sh^2 eFV} (x-y) + \frac{i}{8} Sp \bar{\Lambda}^{-1} (1 + 4ef t \frac{cheF(\nu-2s)}{SheFV}) \right\}^{(16)} dt, \end{aligned}$$

$$Q^x(x-y, \nu', \nu'', \nu) = -i(4\pi)^{-2} (de-1) \int_{-it_0}^{\infty} t (\det \Lambda)^{-\frac{1}{2}} \exp\{-i(x-y) \frac{\tau_1 \bar{\Lambda}^{-1}}{8Sh^2 eFV} (x-y)\} \times$$

$$\left\{ (4t)^{-1} \delta(s) + i2^{-4}(x-y) \left[\bar{\Lambda}^{-2} \frac{eF\tau_1 cheF(\nu-2s)}{Sh^3 eFV} + \bar{\Lambda}^{-1} \frac{2e^2 F^2 ch2eFs}{Sh^2 eFV} \right. \right.$$

$$\left. - \bar{\Lambda}^{-1} \frac{\tau_4 sheF(\nu-2s)}{Sh^3 eFV} \right] (x-y) + 2^{-4} \left[\frac{1}{4}(x-y) \bar{\Lambda}^{-2} \frac{\tau_4}{Sh^2 eFV} (x-y) - \right.$$

$$\left. -i \text{Sp} \bar{\Lambda}^{-1} \frac{\text{sh} \epsilon F (v-2s)}{\text{sh} \epsilon F v} \right]^2 + \frac{1}{8} \text{Sp} \left[\bar{\Lambda}^{-1} \frac{2 \epsilon F \text{ch} \epsilon F (v-2s)}{\text{sh} \epsilon F v} - \right. \\ \left. - \bar{\Lambda}^{-2} \frac{\text{sh}^2 \epsilon F (v-2s)}{\text{sh}^2 \epsilon F v} \right] \} dt, \quad (17)$$

$$\tau_1 = \text{ch} 2 \epsilon F s - 1, \quad \tau_2 = 2 \epsilon F t + \text{sh} 2 \epsilon F s,$$

$$\tau_3 = \text{sh} 2 \epsilon F s + 2 \epsilon F t \text{ch} 2 \epsilon F s, \quad \tau_4 = \tau_1 + 2 \epsilon F t \cdot \text{sh} 2 \epsilon F s, \quad (18)$$

$$\eta = \tau_1 + 4 \epsilon F t \tau_3, \quad \Lambda = (2 \epsilon F)^{-1} (\tau_2 - \tau_1 \text{ch} \epsilon F v).$$

Note that α^d and α^x depend upon v' and v'' only in the combination $S = |v' - v''|$. Therefore, when integrating over v' and v'' in (3) one has

$$\int_0^v \int_0^v dv' dv'' \phi(s) = 2 \int_0^v (v-s) \phi(s) ds, \quad (19)$$

where $\phi(s)$ is an arbitrary function. Thus

$$A_{1,1}(x-y, v) = 2 \int_0^v (v-s) a(x-y, v, s) ds \quad (20)$$

If in accordance with the breaking of the photon function $D_{\mu\nu}(k)$ into the parts (9) one represents analogously $A_{1,1}(x-y, v)$ in the form of the sum of two terms, then $A_{1,1}^x(x-y, v)$ can be considerably simplified. For doing this note the identity

$$2i \frac{\partial I}{\partial s} = \left\{ \frac{1}{4} (x-y) \bar{\Lambda}^{-2} \frac{\tau_4}{\text{sh}^2 \epsilon F v} (x-y) - i \text{Sp} \bar{\Lambda}^{-1} \frac{\text{sh} \epsilon F (v-2s)}{\text{sh} \epsilon F v} \right\} \cdot I \quad (21)$$

Using this relation and integrating over S by parts, the expression for $A_{1,1}^x(x-y, v)$ may be represented in the form

$$\Lambda_{4,1}^X(x-y, v) = \frac{i}{2}(4\pi)^{-2} (de-1) \int_{-it_0}^{\infty} \frac{dt}{t} \left[\exp\left\{-i \frac{(x-y)^2}{4t}\right\} - 1 \right] \quad (22)$$

Finally we obtain

$$\begin{aligned} A_{4,1}(x-y, v) = & i v (8t_0)^{-1} + \frac{i}{2}(4\pi)^{-2} \int_0^v (v-s) ds \int_{-it_0}^{\infty} (\det \Lambda)^{-\frac{1}{2}} \times \\ & \exp\left\{-i(x-y) \frac{\tau_1 \Lambda^{-1}}{8sh^2 eFv} (x-y)\right\} \cdot \left\{ (x-y) \Lambda^{-2} \frac{2}{sh^2 eFv} (x-y) + \right. \\ & \left. + 4i \operatorname{Sp} \Lambda^{-1} (1+4eFt \frac{cheF(v-2s)}{sheFv}) \right\} + \frac{i}{2}(4\pi)^{-2} (de-1) \int_{-it_0}^{\infty} \frac{dt}{t} \left(e^{-i \frac{(x-y)^2}{4t}} - 1 \right) \end{aligned} \quad (23)$$

Substitute now (23) into (1). The result of this substitution is written in the form:

$$\begin{aligned} \langle\langle A_{4,1} \rangle\rangle_{K,K} = & \left\{ -i \frac{\alpha}{\pi} v t_0^{-1} \cdot J_{K,K}(L) + \frac{i\alpha}{2\pi} \int_0^v (v-s) ds \int_{-it_0}^{\infty} (\det \Lambda)^{-\frac{1}{2}} \times \right. \\ & \left[K^{\alpha\beta} J_{K,K;\alpha\beta} \left(L + \frac{\tau_1 \Lambda^{-1}}{8sh^2 eFv} \right) + 4i \operatorname{Sp} \Lambda^{-1} (1+4eFt \frac{cheF(v-2s)}{sheFv}) \right. \\ & \left. J_{K,K} \left(L + \frac{\tau_1 \Lambda^{-1}}{8sh^2 eFv} \right) \right] - \frac{\alpha}{4\pi} (de-1) \int_{-it_0}^{\infty} \frac{dt}{t} \left[J_{K,K} \left(L + \frac{1}{4t} \right) - \right. \\ & \left. \left. - J_{K,K}(L) \right] \right\} \left(\det \frac{sheFv}{eF} \right)^{-\frac{1}{2}}, \quad \alpha = \frac{e^2}{4\pi} \end{aligned} \quad (24)$$

where

$$J_{K,K'}(\Omega) = \int d^4x d^4y \psi_{1K}^*(x) \exp\left\{i \frac{e}{2} x F y - i(x-y) \Omega(F)(x-y)\right\} \psi_{1K'}(y) \quad (25)$$

$$J_{K,K';\alpha\beta}(\Omega) = \int d^4x d^4y (x-y)_{\alpha} (x-y)_{\beta} \psi_{1K}^*(x) \exp\left\{i \frac{e}{2} x F y - i(x-y) \Omega(F)(x-y)\right\} \psi_{1K'}(y) \quad (26)$$

Differentiating (25) over $\Omega^{\alpha\beta}$ one can find the connection between these two integrals:

$$J_{K,K';\alpha\beta}(\Omega) = i \frac{\partial J_{K,K'}(\Omega)}{\partial \Omega^{\alpha\beta}} \quad (27)$$

Making use of the integral representations for the parabolic cylinder functions and Hermit polynomials [24], the integrals (25) can be calculated to give

$$J_{K,K'}(\Omega) = i(4\pi)^2 [\det(16\Omega^2 - e^2 F^2)]^{-\frac{1}{4}} \exp\{-ip\chi(F)p\} \delta_{\{K\},\{K'\}} \quad (28)$$

$$\chi(F) = (2eF)^{-1} \ln \frac{4\Omega - eF}{4\Omega + eF},$$

and P_μ is a four-vector

$$P_\mu P^\mu = p^2, \quad P_\mu P^\mu = P_0^2 + P_1^2, \quad P_0^2 = P P P =$$

$$= p^2 + |e| \mathcal{H}(2n+1), \quad P_1^2 = P L P = -|e| \mathcal{H}(2n+1)$$

$$P_{\mu\nu} = (\mathcal{H}^2 + \mathcal{E}^2)^{-1} (F_{\mu\nu}^2 + g_{\mu\nu} \mathcal{H}^2), \quad L_{\mu\nu} = (\mathcal{H}^2 + \mathcal{E}^2)^{-1} (g_{\mu\nu} \mathcal{E}^2 - F_{\mu\nu}^2)$$

The integral $J_{K,K';\alpha\beta}(\Omega)$ enters into (24) together with the matrix $K = \Lambda^2 \mathcal{Z} / \text{sh}^2 eFv$. Differentiating (27) over $\Omega^{\alpha\beta}$ we obtain:

$$K^{\alpha\beta} J_{K,K';\alpha\beta}(\Omega) = 4 J_{K,K'}(\Omega) \left\{ P \frac{K}{16\Omega^2 - e^2 F^2} P - i s p \frac{2K\Omega}{16\Omega^2 - e^2 F^2} \right\} \quad (29)$$

From (27) we have

$$J_{K,K'}(L) = i(4\pi)^2 e^{ip^2 v} \delta_{\{K\},\{K'\}},$$

$$J_{K,K'}\left(L + \frac{\tilde{\tau}_1 \Lambda^{-1}}{8 \text{sh}^2 eFv}\right) = J_{K,K'}(L) (\det \Lambda)^{\frac{1}{2}} \left(\det \frac{\tilde{\tau}_2^2 - \tilde{\tau}_1^2}{4e^2 F^2}\right)^{-\frac{1}{4}} \quad (30)$$

$$\exp\{-ip\psi(s,t)p\}, \quad \psi(s,t) = (2eF)^{-1} \ln \frac{\tilde{\tau}_2 + \tilde{\tau}_1}{\tilde{\tau}_2 - \tilde{\tau}_1},$$

$$J_{K,K'}(L + \frac{1}{4t}) = J_{K,K}(L) \left(\det \frac{\tilde{\tau}_2^2 - \tilde{\tau}_1^2}{4e^2 F^2} \right)_{S=v}^{-\frac{1}{4}} \exp\{-ip\psi(v,t)p\}.$$

Note that $J_{K,K}(L) = J_{\{K\}}(v)$. Substituting (29), (30) into (24) we get

$$\begin{aligned} \frac{\langle\langle A_{1,1} \rangle\rangle_{K,K}}{J_{\{K\}}(v)} &= i \frac{\alpha}{\pi} v \tilde{t}_0^{-1} - i \frac{\alpha}{2\pi} \int_0^v (v-s) ds \int_{-it_0}^{\infty} M(s,t;P,F) dt + \\ &+ \frac{\alpha}{4\pi} (de-1) \int_{-it_0}^{\infty} t^{-1} [t^2 g(v,t;P,F) - 1] dt \equiv b_1(v;P,F) \end{aligned} \quad (31)$$

$$M(s,t;P,F) = g(s,t;P,F) \left\{ P \frac{2}{\tilde{\tau}_2^2 - \tilde{\tau}_1^2} P + \frac{i}{2} Sp \frac{R}{\tilde{\tau}_2^2 - \tilde{\tau}_1^2} \right\}, \quad (32)$$

$$R = eF [2eFt(3 - 2\operatorname{ch} 2eFs) + (1 - 8e^2 F^2 t^2) \operatorname{sh} 2eFs],$$

$$g(s,t;P,F) = \left[\det \left(\frac{\tilde{\tau}_2^2 - \tilde{\tau}_1^2}{4e^2 F^2} \right) \right]^{-\frac{1}{4}} \exp\{-ip\psi(s,t)p\}.$$

This expression diverges when $t_0 \rightarrow 0$. Before proceeding to the renormalization consider the case $F=0$. In this limit

$$M(s,t;P,0) = g(s,t;P,0) \left\{ P^2 \frac{(s+2t)^2}{2(s+t)^2} + \frac{i}{2(s+t)} \right\}$$

$$g(s,t;P,0) = \frac{\exp\{-ip^2 \frac{s^2}{s+t}\}}{(s+t)^2}$$

Thus

$$\frac{\langle\langle A_{1,1} \rangle\rangle_{K,K}}{J_{\{K\}}(v)} = -i \frac{3\alpha}{4\pi} v \tilde{t}_0^{-1} - i \frac{\alpha}{2\pi} \int_0^v (v-s) ds \int_{-it_0}^{\infty} e^{-ip^2 \frac{s^2}{s+t}} \times$$

$$p^2 \frac{s+2t}{(t+s)^3} dt - \frac{\alpha}{4\pi} \int_0^v ds \int_{-it_0}^{\infty} \frac{e^{-ip^2 \frac{s}{s+t}}}{(t+s)^2} dt + \quad (33)$$

$$+ \frac{\alpha}{4\pi} (de-1) \int_{-it_0}^{\infty} t^{-1} \left(\frac{t^2}{(t+v)^2} e^{-ip^2 \frac{v^2}{t+v}} - 1 \right) dt \equiv b_1(v; p)$$

which coincides with the results of [20]. Consider the Green function

$$G_{(1K)}^{(1)} \Big|_{F=0} = \int_0^{\infty} dv \exp \{ i(p^2 - m^2 + i\varepsilon)v + b_1(v; p) \} \quad (34)$$

Let us separate the terms linear in v from $b_1(v; p)$ and rewrite (33) in the following form

$$b_1(v, p) = -ivM_0(p) + N_0(v, p) \quad (35)$$

where

$$M_0(p) = \frac{3\alpha}{4\pi} t_0^{-1} + \frac{\alpha}{2\pi} \int_0^{\infty} ds \int_{-it_0}^{\infty} M(s, t; p, 0) dt, \quad (36)$$

$$N_0(v, p) = i \frac{\alpha}{2\pi} \int_0^v ds \int_{-it_0}^{\infty} M(s, t; p, 0) dt + iv \frac{\alpha}{2\pi} \int_v^{\infty} ds \int_0^{\infty} M(s, t; p, 0) dt$$

$$- \frac{\alpha}{4\pi} \int_0^v ds \int_{-it_0}^{\infty} g(s, t; p, 0) dt + \frac{\alpha}{4\pi} (de-1) \int_{-it_0}^{\infty} \frac{dt}{t} \left[t^2 g(v, t; p, 0) - 1 \right]. \quad (37)$$

Note that if we calculate the mass operator $\bar{M}_0(p^2, m^2)$ of a scalar particle in the ϵ^2 -approximation and make the substitution $m^2 \rightarrow p^2$ then we get just $M_0(p)$

$$M_0(p) = \bar{M}_0(p^2, m^2 \rightarrow p^2) \quad (38)$$

Therefore, the renormalization of $M_0(p)$ may be performed in a stan-

dard way by choosing the normalization point as $p^2 = m_e^2$ where m_e^2 is the experimental mass squared [20]. To do this let us represent $M_0(p)$ in the form

$$M_0(p) = M_0(p^2 = m_e^2) + (p^2 - m_e^2) \left. \frac{\partial M_0(p)}{\partial p^2} \right|_{p^2 = m_e^2} + M_{0R}(p) \quad (39)$$

where $M_{0R}(p)$ is the renormalized mass operator. Let us make in (34) the substitution

$$v = Z_2 v' \quad (40)$$

where Z_2 is the renormalization constant of the wave function, and δm^2 is the mass renormalization

$$\delta m^2 = m_e^2 - m^2 \quad (41)$$

Then

$$\begin{aligned} G_{1K}^{(4)}|_{F=0} &= Z_2 \int_0^\infty dv \exp\{i(p^2 - m_e^2)(Z_2 - 1)v + i\delta m^2 v - \\ &- i v M_0(p^2 = m_e^2) - i v \left. \frac{\partial M_0(p)}{\partial p^2} \right|_{p^2 = m_e^2} \cdot (p^2 - m_e^2) + \\ &+ i(p^2 - m_e^2)v - i v M_{0R}(p) + N_0(v; p) \end{aligned} \quad (42)$$

The constants Z_2 and δm^2 are chosen from the conditions

$$\delta m^2 = M_0(p^2 = m_e^2) \quad (43)$$

$$Z_2 - 1 = \left. \frac{\partial M_0(p)}{\partial p^2} \right|_{p^2 = m_e^2} \quad (44)$$

The integrals entering into (43) and (44) are easily calculated

$$\delta m^2 = \frac{3\alpha}{4\pi} m_e^2 \left[(m_e^2 t_0)^{-1} + \frac{3}{2} + \ln(\gamma m_e^2 t_0)^{-1} \right], \quad (45)$$

$$Z_2^{-1} = \frac{3\alpha}{4\pi} \left[\frac{1}{2} + \ln(\gamma m_e^2 t_0)^{-1} \right] \quad (46)$$

where γ is Euler's constant. The function $N_0(\nu, P)$ (37) is equal to

$$N_0(\nu, P) = -\frac{\alpha}{4\pi} d_e \ln(\gamma m_e^2 t_0)^{-1} + \frac{\alpha}{2\pi} (3-d_e) \ln(m_e^2 \nu) + Q(\nu, P) \quad (47)$$

$$Q(\nu, P) \rightarrow 0, \nu \rightarrow \infty$$

The diverging term $-\frac{\alpha}{4\pi} d_e \ln(\gamma m_e^2 t_0)^{-1}$ should also be related to the Z_2 -factor

$$\tilde{Z}_2 = Z_2 \exp\left(-\frac{\alpha}{4\pi} d_e \ln(\gamma m_e^2 t_0)^{-1}\right)$$

In e^2 -approximation we have

$$\tilde{Z}_2^{-1} = \frac{\alpha}{4\pi} \left[(3-d_e) \ln(\gamma m_e^2 t_0)^{-1} + \frac{1}{2} \right] \quad (48)$$

Let us return now to expression (31) and rewrite $b_i(\nu, P, F)$ in the following form

$$b_i(\nu, P, F) = -i \frac{\alpha}{\pi} \nu t_0^{-1} - i \frac{\alpha}{2\pi} \nu \int_0^\infty ds \int_{-it_0}^\infty M(s, t; P, F) dt +$$

$$+ N_0(\nu, P) + N(\nu, P, F) + i \frac{\alpha}{2\pi} \nu \int_\nu^\infty ds \int_0^\infty M(s, t; P, F) dt \quad (49)$$

$$N(\nu, P, F) = i \frac{\alpha}{2\pi} \int_0^\nu ds \int_0^\infty M_{IR}(s, t; P, F) dt +$$

$$+ \frac{\alpha}{4\pi} (d_e - 1) \int_0^\infty t [g(\nu, t; P, F) - g(\nu, t; P, 0)] dt, \quad (50)$$

$$M_{IR}(s, t; P, F) = M(s, t; P, F) - M(s, t; P, 0)$$

where $N_0(\nu, P)$ is determined by (37). In the last term in (49) and (50) we have everywhere put the lower proper-time-integration limit

equal to zero, on removing the regularization they are finite. One can verify this fact by expanding the integrals into power series with respect to the field and calculating the corresponding integrals. Consider the first two terms in (49) linear in V . Let us write

$M(s, t; P, F)$ in the form:

$$M(s, t; P, F) = M(s, t; P^2 - m_e^2, 0) + (P^2 - m_e^2) \frac{\partial M(s, t; P, 0)}{\partial P^2} \Big|_{P^2 = m_e^2} + M_R(s, t; P, F);$$

$$M_R(s, t; P, F) = M(s, t; P, F) - \exp(-im_e^2 \frac{s^2}{t+s})^{(51)}$$

$$(t+s)^{-3} \left\{ m_e^2 \frac{(s+2t)^2}{2(s+t)} + i \right\} - \exp\left\{ -im_e^2 \frac{s^2}{t+s} \right\} \left[\frac{(s+t)^2 + t^2}{t+s} - im_e^2 s^2 (s+2t)^2 (s+t)^{-2} \right].$$

Then the integrals over s and t in $M_R(s, t; P, F)$ are finite when $t_0 \rightarrow 0$ and we obtain for $b_1(v; P, F)$:

$$b_1(v; P, F) = -iv \delta m^2 - i(P^2 - m_e^2)(z_2 - 1)v - iv M_R(P, F)$$

$$+ i \frac{\alpha}{2\pi} v \int_v^\infty ds \int_0^\infty M(s, t; P, F) dt + i \frac{\alpha}{2\pi} \int_0^v s ds \int_0^\infty M_{1R}(s, t; P, F) dt \quad (52)$$

$$+ N_0(v, P) + \frac{\alpha}{4\pi} (de - 1) \int_0^\infty t g_R(v, t; P, F) dt,$$

$$M_R(P, F) = \frac{\alpha}{2\pi} \int_0^\infty ds \int_0^\infty M_R(s, t; P, F) dt,$$

(53)

$$g_R(s, t; P, F) = g(s, t; P, F) - g(s, t; P, 0).$$

where δm^2 and $Z_2 - 1$ are determined by (43) and (44). Thus we see that the procedure of renormalization in the presence of a constant field is the same as in the vacuum ($F = 0$) and the constants of renormalization do not depend upon the external field.

If we write the Green function $G_{1k}^{(1)}$ in the form

$$G_{1k}^{(1)} = \int_0^\infty dv \exp\{i(p^2 - m^2 + i\varepsilon)v + \theta_1(v, p, F)\} \quad (54)$$

and make the change of variables $v = Z_2 v'$ and the mass renormalization $m^2 = m_e^2 - \delta m^2$ we obtain

$$G_{1k}^{(1)} = \tilde{Z}_2 \cdot G_{1kR}^{(1)} \quad (55)$$

where $G_{1kR}^{(1)}$ is the renormalized Green function

$$G_{1kR}^{(1)} = \int_0^\infty dv \exp\{i(p^2 - m_e^2 + i\varepsilon)v + \theta_{1R}(v, p, F)\} \quad (56)$$

$$\begin{aligned} \theta_{1R}(v, p, F) = & -i v M_R(p, F) + N(v, p, F) + N_{0R}(v, p) + \\ & + i \frac{\alpha}{2\pi} v \int_0^\infty ds \int_0^\infty M(s, t; p, F) dt, \end{aligned} \quad (57)$$

$$N_{0R}(v, p) = N_0(v, p) + \frac{\alpha}{4\pi} d_e \ln(\gamma m_e^2 t_0)^{-1}$$

Note that if we calculate the mass operator in a constant field in the e^2 -approximation and write it in the form

$$M(p, F) = \int_0^\infty ds \int_{-it_0}^\infty M(s, t; p, F, m^2) dt \quad (58)$$

then the function $M(s, t; p, F)$ (32) is connected with $M(s, t; p, F, m^2)$ in the following way:

$$M(s, t; p, F) = M(s, t; p, F, m^2 \rightarrow p^2) \quad (59)$$

Then our result for $M(s, t; p, F)$ coincides with that of ref. [8], i.e. $M(s, t; p, F)$ is the kernel of the mass operator in the e^2 -approximation.

To find the infra-red asymptotics of the Green function one should calculate the asymptotic behaviour of $\mathcal{G}_R(V; P, F)$ when $V \rightarrow \infty$. The terms linear in V contribute to the position of the pole of the Green function. From (57) and (54) we see that the pole of the Green function is in the point.

$$P^2 - m_e^2 - M_R(P, F) = 0 \quad (60)$$

In the vacuum there occurs, besides, the enhancement of the pole near the point $P^2 = m_e^2$ [16, 17, 20] owing to the fact that the function $N_{OR}(V; P)$ behaves logarithmically when $V \rightarrow \infty$ i.e.

$$\lim_{V \rightarrow \infty} N_{OR}(V, P) \sim \frac{\alpha}{2\pi} (3 - d_e) \ln m_e^2 V$$

Let us find the asymptotic behaviour of $N_R(V; P, F)$ when $V \rightarrow \infty$

To do so we shall act in the following way: when $V \rightarrow \infty$ we write

$$\begin{aligned} N_R(V; P, F) &= \frac{N_R(V; P, F)}{\ln m_e^2 V} \cdot \ln m_e^2 V \stackrel{V \rightarrow \infty}{=} \\ &= f(P, F) \cdot \ln m_e^2 V \end{aligned} \quad (61)$$

where

$$f(P, F) = \lim_{V \rightarrow \infty} V \frac{\partial N_R(V; P, F)}{\partial V} \quad (62)$$

Substituting expressions (57) and (50) into (62) we obtain

$$\begin{aligned} f(P, F) &= \lim_{V \rightarrow \infty} \left\{ i \frac{\alpha}{2\pi} V^2 \int_0^\infty M(V, t; P, F) dt + \right. \\ &\quad \left. + \frac{\alpha}{4\pi} (d_e - 1) V \int_0^\infty t \frac{\partial g(V, t; P, F)}{\partial V} dt \right\} \end{aligned} \quad (63)$$

Thus, the Green function $G_{\{1, k\}}^{(1)}$ near the point $P^2 - m_e^2 - M_R(P, F) = 0$ has the form:

$$G_{\{K\}}^{(1)} \sim [p^2 - m_e^2 - M_R(p, F)]^{-(1+f(p, F))} \quad (64)$$

If $F=0$ then

$$f(p, 0) = \frac{\alpha}{2\pi} (3 - d_0) \quad (65)$$

and we obtain the well-known results [16, 20]. If $F \neq 0$, the function $f(p, F)$ depends upon the parameters: p^2 , $\chi = -(eFP)^2 m^{-6}$ and invariants $\mathcal{F}$ and $\mathcal{Y}$. As the function $f(p, F)$ is linear in the constant α it is necessary to substitute $p^2 \rightarrow m_e^2$ near the pole (60), as not to exceed the accuracy.

Consider the crossed field ($\mathcal{F} = \mathcal{Y} = 0$). In this case $f(p, F)$ depends upon one parameter χ and we have

$$f(p, F) = \begin{cases} \frac{\alpha}{2\pi} (3 - d_0), & \chi \ll 1 \\ \frac{\alpha}{12\pi} (2g + d_0), & \chi \gg 1 \end{cases} \quad (66)$$

For the constant magnetic field ($\mathcal{Y} = 0, \mathcal{F} < 0$) if $-\frac{\mathcal{F}}{m^6 e^2} \ll 1$, $\chi \gg -\frac{\mathcal{F}}{m^4 e^2}$ the results of the calculations for the function coincide with those for the crossed field.

APPENDIX

Here we find the eigenfunctions of Klein-Gordon's equation in a constant and uniform external field $A_\mu^{\text{ext}}(x) = -\frac{1}{2} F_{\mu\nu} x^\nu$. These functions satisfy the equations

$$\pi_\mu^{\text{ext}}(x) \pi^{\text{ext}, \mu}(x) \psi_{\{K\}}(x) = p^2 \psi_{\{K\}}(x). \quad (1)$$

Consider isotropic orthogonal eigenvectors of the tensor $F_{\mu\nu}$

$$F_{\mu\nu} n^\nu = \mathcal{E} n_\mu, F_{\mu\nu} \bar{n}^\nu = -\mathcal{E} \bar{n}_\mu, F_{\mu\nu} m^\nu = i\mathcal{H} m_\mu, \\ F_{\mu\nu} \bar{m}^\nu = -i\mathcal{H} \bar{m}_\mu, \mathcal{E}, \mathcal{H} = \left[(\mathcal{F}^2 + \mathcal{G}^2)^{\frac{1}{2}} \mp \mathcal{F} \right]^{\frac{1}{2}} \quad (2)$$

Vectors (2) are normalized by the conditions $(n\bar{n}) = -(m\bar{m}) = 2$.
If we introduce the vectors $n_\pm = \frac{1}{2}(n \pm \bar{n})$, $m_\pm = \frac{m \pm \bar{m}}{2}$,
 $= i \frac{\bar{m} - m}{2}$ satisfying the equations

$$F n_\pm = \mathcal{E} n_\mp, F m_\pm = \mp \mathcal{H} m_\mp, n_\pm^2 = \pm 1, \\ m_\pm^2 = -1, (n_+ n_-) = (n_\pm m_\pm) = (n_\pm m_\mp) = (m_- m_+) = 0$$

the tensor $F_{\mu\nu}$ and the field $A_\mu^{\text{ext}}(x)$ can be represented in the form

$$F_{\mu\nu} = \mathcal{E}(n_{-\mu} n_{+\nu} - n_{+\mu} n_{-\nu}) + \mathcal{H}(m_{-\mu} m_{+\nu} - m_{+\mu} m_{-\nu}), \quad (3)$$

$$A_\mu^{\text{ext}}(x) = -\frac{\mathcal{E}}{2} [n_{-\mu}(n_+ x) - n_{+\mu}(n_- x)] - \frac{\mathcal{H}}{2} [m_{-\mu}(m_+ x) - m_{+\mu}(m_- x)]. \quad (4)$$

Let us perform the gauge transformation of the potential

$$A_\mu^{\text{ext}}(x) = A_\mu^{\text{ext}}(x) + \partial_\mu \phi(x)$$

$$\phi(x) = -\frac{\mathcal{E}}{2} (n_+ x)(n_- x) - \frac{\mathcal{H}}{2} (m_+ x)(m_- x) \quad (5)$$

This transformation is due to the fact that in terms of the potential $A_\mu^{\text{ext}}(x)$ it is possible to separate the variables in the equation (1).
If we transform the functions in accord with the gauge transformation (5)

$$\psi_{\{k\}}(x) = \exp(ie\phi(x)) \psi'_{\{k\}}(x) \quad (6)$$

then the functions $\psi'_{1k_j}(x)$ satisfy the equation

$$\pi'^{\text{ext}}_m(x) \pi'^{\text{ext}}_{m'}(x) \psi'_{1k_j}(x) = \rho^2 \psi'_{1k_j}(x), \quad (7)$$

$$\pi'^{\text{ext}}_m(x) = i\partial_m - eA'_m(x)$$

Substitute (5) into (7). Then we have

$$\left[-\frac{\partial^2}{\partial(n+x)^2} - \left(i\frac{\partial}{\partial(n+x)} + e\mathcal{E}(n+x) \right)^2 + \frac{\partial^2}{\partial(m+x)^2} - \right. \\ \left. - \left(i\frac{\partial}{\partial(m+x)} + e\mathcal{H}(m+x) \right)^2 \right] \psi'_{1k_j}(x) = \rho^2 \psi'_{1k_j}(x) \quad (8)$$

Carrying out the separation of the variables in (8) we obtain

$$\psi'_{1k_j}(x) = N_{1k_j} \exp \left\{ -i p_1(n+x) - i p_2(m+x) - \frac{1}{2} \rho^2 \right\} H_n(\rho) D_\nu [\omega e^{i\frac{\pi}{4}} \tau],$$

$$\rho = (|e|\mathcal{H})^{\frac{1}{2}} [(m+x) + p_2/e\mathcal{H}], \quad \tau = (2|e|\mathcal{E})^{\frac{1}{2}} [(n+x) + p_1/e\mathcal{E}], \quad (9)$$

$$\nu = -\frac{1}{2}(1+i\lambda), \quad \lambda = [\rho^2 + |e|\mathcal{H}(2n+1)], \quad n=0,1,2,\dots; \quad \omega = \pm 1$$

$H_n(\rho)$ are the Hermite polynomials, $D_\nu[\omega e^{i\frac{\pi}{4}} \tau]$ are the parabolic cylinder functions, P_1 and P_2 are eigenvalues of the operational integrals of motion $i\partial/\partial(n+x)$ and $i\partial/\partial(m+x)$, respectively, N_{1k_j} is a normalizing factor. Let us verify that for the functions (9) the orthogonality condition (1.24) does take place and find in this way the normalizing factor. Using the orthogonality condition of the Hermite functions we have

$$(\psi_{1k_j}, \psi_{1k'_j}) = N_{1k_j}^* N_{1k'_j} (2\pi)^2 (2e^2 \mathcal{E} \mathcal{H})^{-\frac{1}{2}} \times \\ 2^n n! \sqrt{\pi} \delta_{n,n'} \delta(P_1 - P'_1) \delta(P_2 - P'_2) \cdot T, \quad (10)$$

$$T = \int_{-\infty}^{+\infty} d\tau D_{\nu*} [\omega e^{-i\frac{\pi}{4}\tau}] D_{\nu} [\omega' e^{i\frac{\pi}{4}\tau}] \quad (11)$$

to calculate the integral T let us make use of the integral representation for the parabolic cylinder functions [24]

$$D_{\nu} [\omega e^{i\frac{\pi}{4}\tau}] = \Gamma^{-1}(-\nu) \exp\{-i\frac{\pi\nu}{4} - i\frac{\tau^2}{4}\} \int_0^{\infty} t^{-\nu-1} \exp\{-i\frac{t^2}{2} - it\tau\} dt \quad (12)$$

As a result we obtain

$$T = 2|\epsilon| \mathcal{E} (2\pi)^2 [\Gamma(-\nu) \Gamma(-\nu^*)]^{-1} \exp\{i\frac{\pi}{4}(\nu^* - \nu)\} \delta_{\omega\omega'} \delta(\rho^2 - \rho'^2) \quad (13)$$

If the normalizing factor N_{1K} is equal to

$$N_{1K} = (2\pi)^{-2} \left(\frac{\mathcal{H}}{2\mathcal{E}}\right)^{1/4} (2^n n! \sqrt{\pi})^{-\frac{1}{2}} \Gamma(-\nu) e^{i\frac{\pi\nu}{4}} \quad (14)$$

the functions (9) are normalized by condition (1.24). Let us show that functions (9) form a complete set in the sense of (1.25)

$$\begin{aligned} \sum_{n=0}^{\infty} \sum_{\omega=\pm 1} \int \frac{d\rho_1 d\rho_2 d\rho^2}{(2\pi)^4} \psi_{1K}(x) \psi_{1K}^*(y) &= \left(\frac{\mathcal{H}}{2\mathcal{E}}\right)^{1/2} \cdot \\ \int d\rho_1 d\rho_2 \exp\{-i\rho_1(n-z) - i\rho_2(m-z)\} \sum_{n=0}^{\infty} \tilde{U}_n(\rho) \tilde{U}_n(\rho') \cdot \int_{-\infty}^{+\infty} J_{\rho^2} d\rho^2 & \quad (15) \\ J_{\rho^2} = e^{i\frac{\pi}{4}(\nu-\nu^*)} \Gamma(-\nu) \Gamma(-\nu^*) \sum_{\omega=\pm 1} D_{\nu} [\omega e^{i\frac{\pi}{4}\tau}] D_{\nu*} [\omega e^{-i\frac{\pi}{4}\tau'}] & \\ \nu^* = -\nu - 1, \quad \tau' = (2|\epsilon| \mathcal{E})^{\frac{1}{2}} [(n+y) + \rho_1/\epsilon \mathcal{E}] & \quad (16) \end{aligned}$$

$$\rho' = [|\epsilon| \mathcal{H}]^{\frac{1}{2}} [(m, y) + P_2 / \epsilon \mathcal{H}],$$

$U_n(\rho) = (2^n n! \sqrt{\pi})^{-\frac{1}{2}} \exp(-\frac{1}{2} \rho^2) H_n(\rho)$ are the Hermite functions. Let us find $\int_{-\infty}^{\infty} J_{P_2} d\rho^2$. To do this let us use the identity $\Gamma(-\nu) \Gamma(-\nu^*) = -\pi / \sin(\pi \nu)$ and introduce the new variable $\nu = -\frac{1}{2} - \frac{i}{2} [\rho^2 + |\epsilon| \mathcal{H} (2n+1)] / |\epsilon| \mathcal{E}$.

Then

$$\int_{-\infty}^{+\infty} J_{P_2} d\rho^2 = +i 2 |\epsilon| \mathcal{E} \int_{-\frac{1}{2} - i\infty}^{-\frac{1}{2} + i\infty} J_{\nu} d\nu \quad (17)$$

$$J_{\nu} = \pi \frac{\exp\{i\frac{\pi}{2}(\nu + \frac{1}{2})\}}{\sin \pi \nu} \sum_{\omega=\pm 1} D_{\nu}[\omega e^{i\frac{\pi}{4}} \tau] D_{-\nu-1}[\omega e^{-i\frac{\pi}{4}} \tau'] \quad (18)$$

The integral over ν is calculated with the help of the Cherry formula [24]

$$\begin{aligned} \int_{C-i\infty}^{C+i\infty} \frac{e^{i\frac{\pi}{2}(\nu + \frac{1}{2})}}{\sin \pi \nu} \sum_{\omega=\pm 1} D_{\nu}[\omega e^{i\frac{\pi}{4}} \tau] D_{-\nu-1}[\omega e^{-i\frac{\pi}{4}} \tau'] &= \\ &= -4\pi i \delta(\tau - \tau'), \quad -1 < \text{Re } C < 0 \end{aligned} \quad (19)$$

Thus, choosing $C = -\frac{1}{2}$ in (17) we obtain

$$\int_{-\infty}^{+\infty} J_{P_2} d\rho^2 = -(2|\epsilon| \mathcal{E})^{\frac{1}{2}} (2\pi)^2 \delta(n, z) \quad (20)$$

We see that this expression does not depend upon the quantum numbers. Therefore one may integrate over P_2 and sum over n in (15) using the condition of completeness of the Hermite functions

$$\sum_{n=0}^{\infty} U_n(\rho) U_n(\rho') = \delta(\rho - \rho') = (|\epsilon| \mathcal{H})^{-\frac{1}{2}} \delta(m, z) \quad (21)$$

After integrating over P_1 we obtain the condition of completeness (1.25)

$$\sum_{\{K\}} \psi_{\{K\}}(x) \psi_{\{K\}}^*(y) = \delta^{(4)}(x-y) \quad (22)$$

To calculate the integral

$$J_{K,K'}(\Omega) = \int d^4x d^4y \psi_{\{K\}}^*(x) \exp\{i \frac{e}{2} x f y - i z \Omega(F) z\} \psi_{\{K'\}}(y) \quad (23)$$

which occurs when calculating $\ll A_{1,1} \gg_{K,K}$ let us use the method of ref. [12, 13]. As a result we obtain

$$J_{K,K'}(\Omega) = i(4\pi)^2 [\det(16\Omega^2 - e^2 F^2)]^{-\frac{1}{4}} \times \exp\{ip\chi(F)p\} \delta_{\{K\},\{K'\}}, \quad \chi(F) = (2eF)^{-1} \ln \frac{4\Omega - eF}{4\Omega + eF}. \quad (24)$$

Proceeding from this expression we shall calculate the integral occurring in (2.24).

$$J_{K,K';\alpha\beta}(\Omega) = i \frac{\partial J_{K,K'}(\Omega)}{\partial \Omega^{\alpha\beta}} = i J_{K,K'}(\Omega) \times \frac{\partial}{\partial \Omega^{\alpha\beta}} \left\{ -ip\chi(F)p - \frac{1}{4} \text{Sp} \ln(16\Omega^2 - e^2 F^2) \right\} \quad (25)$$

Let $\tilde{F}(\Omega)$ be an arbitrary function of Ω which can be expanded in to a series in Ω

$$\tilde{F}_{\mu\nu}(\Omega) = \sum_{n=0}^{\infty} \frac{1}{n!} \tilde{F}^{(n)}(0) \Omega_{\mu\nu}^n$$

Let us find the derivative of $\tilde{F}_{\mu\nu}(\Omega)$ with respect to $\Omega^{\alpha\beta}$

$$\frac{\partial \tilde{F}_{\mu\nu}(\Omega)}{\partial \Omega^{\alpha\beta}} = \sum_{n=1}^{\infty} \frac{1}{n!} \tilde{F}^{(n)}(0) \sum_{K=0}^{n-1} \Omega_{\mu\alpha}^K \Omega_{\beta\nu}^{n-K-1} \quad (26)$$

Here we make use of the relation:

$$\frac{\partial \Omega_{\mu\nu}^n}{\partial \Omega^{\alpha\beta}} = \sum_{K=0}^{n-1} \Omega_{\mu\alpha}^K \Omega_{\beta\nu}^{n-K-1} \quad (27)$$

Expression (25) contains the derivatives of two types: $\frac{\partial}{\partial \Omega^{\alpha\beta}} P \tilde{F}(\Omega) P$ and $\frac{\partial}{\partial \Omega^{\alpha\beta}} Sp \tilde{F}(\Omega)$. Using (26) we have

$$\begin{aligned} \frac{\partial}{\partial \Omega^{\alpha\beta}} Sp \tilde{F}(\Omega) &= \tilde{F}'_{\beta\alpha}(\Omega) \\ \frac{\partial}{\partial \Omega^{\alpha\beta}} P \tilde{F}(\Omega) P &= \sum_{n=1}^{\infty} \frac{\tilde{F}^{(n)}(0)}{n!} \sum_{K=0}^{n-1} (F \Omega^K)_{\alpha} (\Omega^{n-K-1} P)_{\beta} \end{aligned} \quad (28)$$

But the expressions (28) are included in (2.24) together with the factor $K = \Lambda^2 \eta / sh^2 e F \nu$. Then we obtain finally from (28):

$$\begin{aligned} K^{\alpha\beta} \frac{\partial}{\partial \Omega^{\alpha\beta}} Sp \tilde{F}(\Omega) &= Sp [K \tilde{F}'(\Omega)] \\ K^{\alpha\beta} \frac{\partial}{\partial \Omega^{\alpha\beta}} P \tilde{F}(\Omega) P &= P \tilde{F}'(\Omega) K P. \end{aligned} \quad (29)$$

Using relations (29) we have

$$\begin{aligned} K^{\alpha\beta} J_{K,K';\alpha\beta}(\Omega) &= 4 J_{K,K'}(\Omega) \left[P \frac{K}{16\Omega^2 - e^2 F^2} P - \right. \\ &\quad \left. - i Sp 2K\Omega / (16\Omega^2 - e^2 F^2) \right] \end{aligned} \quad (30)$$

Substituting the expressions

$$\Omega(F) = L(v), \quad \Omega(F) = L(v) + \frac{\tilde{\epsilon}_1 \tilde{\Lambda}^{-1}}{8sh^2 eFv}, \quad \Omega(F) = L(v) + \frac{1}{4t}$$

instead of $\Omega(F)$ into (24) we obtain the relations (2.30).

REFERENCES

- 1 Ritus V.I. (1979), Proceedings of P.N.Lebedev Physical Institute, v.111.
- 2 Nikishov A.I. (1979), Proceedings of P.N.Lebedev Physical Institute, v.111.
- 3 Batalin I.A., Shabad A.E. (1971), Zh. Eksp. Teor. Fiz. 60, 894.
- 4 Wu-Yang Tsai, Yildiz (1973), Phys. Rev. D8, 3446.
- 5 Wu-Yang Tsai (1973), Phys. Rev. D8, 3460.
- 6 Sokolov A.A., Ternov I.M. (1974), "Relativistic electron", "Nauka", Moscow.
- 7 Baier V.N., Katkov V.M., Fadin V.S. (1973), "Radiation of relativistic electron", Atomizdat, Moscow.
- 8 Baier V.N., Katkov V.M., Strakhovenko V.M. (1974), Zh. Eksp. Teor. Fiz. 67, 453.
- 9 Baier V.N., Katkov V.M., Strakhovenko V.M. (1975), Zh. Eksp. Teor. Fiz. 68, 405.
- 10 Ritus V.I. (1972), Ann. Phys. 69, 555.
- 11 Ritus V.I. (1972), Problems of Theoretical Physics, "Nauka", Moscow.
- 12 Ritus V.I. (1970), Pis'ma Zh. Eksp. Teor. Fiz. 12, 416.
- 13 Ritus V.I. (1978), Zh. Eksp. Teor. Fiz. 75, 1560.
- 14 Volfengaut J.J., Gavrilov S.P., Gitman D.M., Shvartsman Sh.M. (1981), Yad. Fiz. 33, 743.
- 15 Fradkin E.S. (1955), Zh. Eksp. Teor. Fiz. 29, 258.
- 16 Fradkin E.S. (1965), Proceedings of P.N.Lebedev Physical Institute, v.29.

17. Fradkin E.S. (1966), Nucl. Phys. 76, 588.
18. Fradkin E.S. (1954), Dokl. Acad. Nauk SSSR, 98, 47.
19. Fradkin E.S. (1954), Dokl. Acad. Nauk SSSR, 100, 897.
20. Fradkin E.S. (1965), Acta Phys. Hung XIX, 1975.
21. Milekhin G.A., Fradkin E.S. (1963), Zh. Eksp. Teor. Fiz. 46, 1925.
22. Babichenco A.S., Shabad A.E. Preprint PhIAN, (1984), N 127.
23. Batalin I.A., Fradkin E.S. (1970), Teor. Mat Fiz. 5, 190.
24. Bateman H., Erdelyi A. (1953), "Higher transcendental Functions" vol. 2.

Т - 10966

Подписано в печать 21 мая 1984 года

Заказ № 473. Тираж 100 экз. п/л 2,1

Отпечатано в Отделе научно-технической
информации ФИАН СССР

Москва, В-312, Ленинский проспект, 53