10 JUN 1985

TRI-PP-85-4
Feb 1985

Lattice Results on the Meson Electric Form Factor

W. Milcox and R.M. Woloshyn

TRIUMF, 4004 Wesbrook Mall, Vancouver, B.C., Canada V6T 2A3

Abstract

A calculation is outlined and results presented for the electric form factor, measured at two values of the momentum, of the pseudo-Goldstone meson within the staggered formulation of lattice fermions.

(Submitted to Physical Review Letters)

CERN LIBRARIES, GENEVA

CERN-P00067388

TRI-PP 85-4
C2

- 2 -

As numerical simulations of quantum chromodynamics (QCD) on the lattice improve it is necessary to devise calculations which provide more detailed tests of the theory. One important area for study is hadron internal structure.¹⁻³ Electromagnetic properties can provide clean and experimentally accessible information for this purpose. In this letter we discuss the lattice calculation of the vector current-hadron vertex function and present our first results for the pseudoscalar meson electric form factor.

The staggered scheme for putting fermions on the lattice is used.^{4,5} The fermion action in terms of single component fermion fields $\bar{\chi}$, χ and gauge field matrices U_μ takes the form (suppressing color indices)

$$S_F = \frac{1}{2} \sum_{x,u} a_\mu(x) [\bar{\chi}(x) U_\mu(x) \chi(x+a_\mu) - \bar{\chi}(x+a_\mu) U_\mu^\dagger(x) \chi(x)] + m \sum_x \bar{\chi}(x) \chi(x), \quad (1)$$

where a is the lattice spacing and a_μ is the unit vector in the μ -direction ($\mu=1,\dots,4$). The quantity $a_\mu(x)=(-1)^{\sum_{\nu < \mu} x_\nu}$. Local phase transformation of the fermion field, $\chi(x) \rightarrow e^{i\omega(x)} \chi(x)$, $\bar{\chi}(x) \rightarrow \bar{\chi}(x) e^{-i\omega(x)}$ yields a vector current

$$j_\mu(x) = - \frac{\delta S_F(x)}{\delta \Delta\omega(x)}, \quad (2a)$$

$$= - \frac{i}{2} a_\mu(x) [\bar{\chi}(x) U_\mu(x) \chi(x+a_\mu) + \bar{\chi}(x+a_\mu) U_\mu^\dagger(x) \chi(x)], \quad (2b)$$

where $\Delta\omega(x)=\omega(x+a_\mu)-\omega(x)$. This is conserved in the sense that the ensemble average $\langle \sum_\mu (j_\mu(x+a_\mu) - j_\mu(x)) \rangle = 0$.

The interpretation of the fermion degrees of freedom in (1) is usually given in terms of flavored quark fields defined on hypercubes in

the lattice. The hadron correlation functions, to be identified with continuum matrix elements, are constructed from interpolating fields made up from these non-local flavored quark fields. It has been shown that the two-point function, relevant for mass calculations, can be cast in a form that can be constructed using only local χ -field bilinear operators.^{6,7} This is, of course, advantageous for numerical calculations. We have shown that our three-point function can also be calculated in a form that involves only local χ -field bilinear operators as interpolating fields. The derivation will be given elsewhere and here we only quote the final results used in our numerical study.

We want to construct current matrix elements for flavor nonsinglet meson states with nonzero electric charge. Specifically, we consider the pseudo-goldstone meson associated with the exactly conserved (in the zero mass limit) flavor non-singlet axial current.⁵ The usual flavor structure associated with the staggered fermions is not useful for constructing charged states since an "electric charge" defined within these flavors is not conserved.⁸ The conserved vector current, (2b), if interpreted as nondynamical electric charge, assigns identical charges to all four staggered fermion flavors. We therefore introduce two sets of χ fields (labelled by u and d , with charges q^u and q^d , $q^u - q^d = 1$) from which we construct charged meson interpolating fields and the conserved electromagnetic currents $j_{(x)}^u$ and $j_{(x)}^d$, based on (2b).

The three-point function which we calculate is (with color indices suppressed)

$$A(\vec{p}, \vec{q}; t_z, t_x) = \langle 0 | \sum_{\vec{z}} e^{-i\vec{p} \cdot \vec{z}} (-1)^{\vec{z}} \bar{\chi}^d(\vec{z}, t_z) \chi^u(\vec{z}, t_z) \times \sum_{\vec{x}} e^{i\vec{q} \cdot \vec{x}} \rho(\vec{x}, t_x) \bar{\chi}^u(0) \chi^d(0) | 0 \rangle, \quad (3)$$

where $\rho(\vec{x}, t_x) = i q^u j_{\mu}^u(x) + i q^d j_{\mu}^d(x)$ and $(-1)^{\vec{z}}$ means $(-1)^{z_1 + z_2 + z_3}$. We also need the two-point function⁹

$$G(\vec{p}; t_z) = \langle 0 | \sum_{\vec{z}} e^{-i\vec{p} \cdot \vec{z}} (-1)^{\vec{z}} \bar{\chi}^d(\vec{z}, t_z) \chi^u(\vec{z}, t_z) \bar{\chi}^u(0) \chi^d(0) | 0 \rangle. \quad (4)$$

For $t_x, (t_x - t_z) \gg 1$

$$A(\vec{p}, \vec{q}; t_z, t_x) \rightarrow \left\{ Z(p) Z(p') \frac{(1 + e^{E_p})}{(1 + e^{-E_{p^d}})} \frac{(1 + e^{-E_{p^a}})}{(1 + e^{E_{p^a}})} \right\}^{1/2} \times e^{-E_{p'} t_x} e^{-E_p (t_z - t_x)} \times \langle \pi^+(\vec{p}) | \rho(0) | \pi^+(p') \rangle, \quad (5)$$

where

$$\langle \pi^+(\vec{p}) | \rho(0) | \pi^+(p') \rangle = \frac{(E_p + E_{p'})}{2\sqrt{E_p E_{p'}}} F(q), \quad (6)$$

with $E_p = \left\{ \vec{p}^2 + M^2 \right\}^{1/2}$ and $\vec{p}' = \vec{p} - \vec{q}$. We use $|\pi^+(\vec{p})\rangle$ to label the pseudoscalar meson ground state with unit charge and degenerate (light

valence quarks. The quantity $F(q)$ is the electric form factor and the quantity $Z(p)$ is related to the two point function by

$$G(\vec{p}; t_z) \xrightarrow[t_z \gg 1]{} Z(p) e^{-E_p t_z}. \quad (7)$$

Numerical calculations were done in a model for QCD using only SU(2) color to save on computing time. The lattice size was $10^2 \cdot 20 \cdot 16$ with the current carrying momentum in the 3-direction. Thirty-two gauge field configurations were prepared using a heat bath Monte Carlo¹⁰ in quenched approximation and the Wilson gauge field action¹¹ at $\beta = 2.3$. The gauge fields were constructed on a $10^3 \cdot 16$ lattice and then doubled in the 3-direction.

Quark propagators were calculated using the conjugate-gradient algorithm.¹² Anti-periodic boundary conditions were used on fermion fields in the spatial directions. However the fermion coupling was put equal to zero across the time boundary of the lattice. This is similar, but not identical, to the boundary condition used by Bernard et al.¹³ The advantage of this choice is that we see simple exponential fall off of the correlation function over a large time interval. When calculating the form factor the problem of non-vacuum contamination can be corrected by taking a geometric mean of correlation functions calculated for appropriate momenta.

The three-point function was calculated as the derivative of a two-point function with the charge operator acting as a source.^{14,15} This means that in the two-point function one of the quark propagators is calculated not with the action S_F of Eq. 1 but with

$$S_F(\alpha) = S_F - \alpha \sum_{\vec{x}} e^{i\vec{q} \cdot \vec{x}} \rho(\vec{x}, t_x). \quad (8)$$

The derivative with respect to α (at $\alpha=0$) gives the three point function. This derivative is obtained numerically by calculating two-point functions at $\alpha=0$ and $\alpha=0.05$. A check of this procedure at $q=0$, where the three and two-point functions are related,⁹ indicates that our derivative is good to within a few percent.

Propagators, with and without the source, were calculated for three different spatial starting points in each of the thirty-two gauge field configurations. All calculations were done for the quark mass parameter $m_a=0.025$. The data summed over all configurations are shown in Fig. 1 for the two-point function $G(0;t_z)$ and the three-point function $A(0,\vec{q};t_z,t_x)$. The three-point function was calculated at two values of momentum, for $q=\pi/10$, the lowest nonzero value on our lattice and for

$q=\pi/5$. The charge operator is located at $t_x=4$. This actually means it involves lattice points at time steps four and five. From the two-point function we infer that the pseudoscalar meson mass $M_a=0.44 \pm 0.01$. As expected from (5) the three-point function falls with the same slope as $G(0;t_z)$.

The form factor is extracted from a combination of two- and three-point functions at large time separations:

$$\left\{ \frac{A(0,\vec{q};t_z,t_x) A(\vec{q},\vec{q};t_z,t_x)}{G(0;t_z) G(\vec{q};t_z)} \right\}^{1/2} \xrightarrow{(E_q+M)} \frac{F(q)}{2/E_q M}. \quad (9)$$

This not only simplifies the calculation but also corrects for nonvacuum contamination in the Z factors of (5) and (7). The results for F , plotted as a function of minkowskian four-momentum transfer squared $q^2 = \vec{q}^2 - (E_q - M)^2$, are shown in Fig. 2. These results were obtained by averaging (9) over time steps t_z equals five to nine. The errors are statistical only and were calculated by combining the uncertainties of the two- and three-point functions including the covariances between these quantities.¹⁶ The solid line in Fig. 2 is a monopole form factor¹⁷ $(1+Q^2 a^2/\lambda^2)^{-1}$ with $\lambda^2=1.05$. The conversion to physical units depends on the calculation of the lattice spacing. Using the value $a \approx 0.16$ fm obtained by Gutbrod and Montway,¹⁸ we infer an rms charge radius from the monopole form factor of about 0.38 fm. The older value $a \approx 0.23$ fm based on the evaluation of the Creutz ratio¹⁰ would give 0.55 fm.

The results presented here are, in a sense, preliminary. A number of important systematic effects remain to be studied. This includes the extrapolation in quark mass to physical values and the approach to the continuum. As far as finite lattice size effects are concerned, none were found in the detailed study of the meson spectrum in $SU(2)$ color by

Billoire et al.⁷ This is consistent with our finding that the meson is a compact object.

In conclusion, we have shown it is feasible to calculate directly an important physical observable, the electric form factor, for a meson on the lattice. Our results provide evidence that the quarks in a lattice meson are indeed localized in a compact object significantly smaller than the lattice volume.

We thank the Nuclear Research Centre, University of Alberta for use of their VAX 11/780. This work is supported in part by the Natural Sciences and Engineering Research Council of Canada.

Footnotes and References

- ¹R.M. Woloshyn and W. Wilcox, TRIUMF Report No. TRI-PP-84-91 (to be published);
- W. Wilcox and R.M. Woloshyn, *ibid.* TRI-PP-84-90 (to be published).
- ²K. Ishikawa, G. Schierholz, H. Schneider and M. Teper, Nucl. Phys. B227, 221 (1983).
- ³K. Barad, M. Ogilvie and C. Rebbi, Phys. Lett. 143B, 222 (1984).
- ⁴J. Kogut and L. Susskind, Phys. Rev. D11, 359 (1975);
- L. Susskind, Phys. Rev. D16, 3031 (1977);
- N. Kawamoto and J. Smit, Nucl. Phys. B192, 100 (1981).
- ⁵H. Kluberg-Stern, A. Morel and O. Napoly, Nucl. Phys. B220, 447 (1983).
- ⁶H. Hamber and G. Parisi, Phys. Rev. D27, 208 (1983);
- J.P. Gilchrist, G. Schierholz, H. Schneider and M. Teper, DESY Report No. DESY 84-021.
- ⁷A. Billoire, R. Lacaze, E. Marinari and A. Morel, CEN-Saclay Report No. SPHT/84-92.
- ⁸Here we mean the electric charge as defined, for example, in Ref. 5.
- ⁹The two- and three-point functions satisfy an integrated Ward identity $A(0,0;t_z,t_x) = G(0;t_z)$ for $t_z > t_x$ which has also been noticed by the authors of Ref. 3.
- ¹⁰M. Creutz, Phys. Rev. D21, 2308 (1980).
- ¹¹K.G. Wilson, Phys. Rev. D10, 2445 (1974).
- ¹²F.S. Beckman, in Mathematical Methods for Digital Computers, edited by A. Ralston and H.S. Wilf (Wiley, New York, 1960) p.62.
- ¹³C. Bernard, T. Draper, K. Olynyk and M. Rushton, Nucl. Phys. B220, 508 (1983).

¹⁴S. Gottlieb, P.B. Mackenzie, H.B. Thacker and D. Weingarten, Phys. Lett. 134B, 346 (1984).

¹⁵C. Bernard, in Gauge Theory on a Lattice: 1984, edited by C. Zachos, W. Celmaster, E. Kovacs and D. Sivers (National Technical Information Service, Springfield, VA, 1984) p.85.

¹⁶This is necessary since the fluctuations in the two- and three-point functions are correlated. See P.R. Bevington, Data Reduction and Error Analysis for the Physical Sciences (McGraw-Hill, New York, 1969)

Chapter 4 for a discussion of error propagation.

¹⁷This is a reasonable phenomenological form at low momentum transfer.

See S.R. Amendolia et al., Phys Lett. 146B, 116 (1984).

¹⁸F. Gutbrod and I. Montvay, Phys. Lett. 136B, 411 (1984).

Figure Captions

1. Plot of the two-point function $G(0;t_z)$ and the three-point function $A(0,\vec{t},t_x,t_z)$ (for $q = \pi/10$ and $q = \pi/5$) as a function of t_z . The solid lines are single exponential fits.
2. Plot of the electric form factor F versus the minkowskian four-momentum transfer squared (in lattice units). The solid line is a monopole form factor $(1+Q^2a^2/\lambda^2)^{-1}$ with $\lambda^2=1.05$.

Fig. 1

Fig. 2