CERN-TH.4184/85

POWER CORRECTIONS IN THE SEMILEPTONIC DETERMINATION OF $\sin^2\theta_W$ *)

J.L. Miramontes⁺⁾

CERN - Geneva

J. Sanchez Guillén and E. Zas

Facultad de Física
Universidad de Santiago de Compostela

A B S T R A C T

We analyze in general terms and estimate the twist-four corrections to the semileptonic measurement of θ_W relevant for testing the electroweak theory. We also extend to twist-four a recent proposal to remove the strange quark uncertainty by using y -distributions.

*) Work supported in part by CAICYT.

+) Permanent address: Facultad de Física, Universidad de Santiago de Compostela, Spain.

CERN-TH.4184/85

May 1985

1. INTRODUCTION

Testing the Weinberg-Salam theory beyond the leading order requires the measurement of its fundamental parameters with an uncertainty below 5%, the magnitude of the radiative corrections [1]. In the case of the electroweak mixing angle this precision has been reached recently by the collider [2], and it will soon be obtained in semi-leptonic measurements [3]. It is therefore urgent to control the hadronic corrections in these neutrino determinations of θ_w , which are the main theoretical uncertainties.

The hadronic contributions from the Kobayashi-Maskawa mixing matrix have been extensively treated in ref. [4] and a method has been recently proposed to reduce their largest uncertainties [5].

For higher twist, the first computation [6] claimed that its effect on θ_w could be as large as 10%, but the general treatment of ref. [4] showed that the estimate was incorrect for isoscalar targets. The same ref. [4] gives a rough estimate based on a naive description of twist-4 in combination with the quark mass effects, but a deeper discussion of higher twist corrections is certainly missing.

The purpose of this paper is to clarify the question, presenting what can be said right now in general about the twist-4 contributions and estimating their expected range of values. We also study the twist-4 modifications to the proposal in ref. [5]. We then improve that proposal to reduce simultaneously the uncertainties in both the twist-4 and the strange sea contributions.

In our analysis we shall use the expressions provided by the extended parton model developed to describe the higher twist contributions at tree level [7]. This allows the use of both the operator product expansion results and other ingredients as the standard counting rules. We remark that it is not possible to evaluate the relevant cross sections of the problem using only the OPE expressions for certain moments and hadronic models. The reason is that the Q^2 -differential cross section involves integrals of the structure functions at fixed Q^2 but not over the whole range in x . For a given neutrino energy E , the correct range is $Q^2/2ME < x < \min(Q^2/2M(E_H^m - M), 1)$ with E_H^m the minimum final hadronic energy, which are obviously not directly related to the moments given by the OPE, as assumed implicitly in recent estimates [8].

We consider u and d quarks only and we do not include logarithmic scaling violations, since they are not calculable at the twist-4 level at present. In the following we treat simultaneously both the twist-4 and the target mass contributions up to the order $1/Q^2$ and we shall refer to them as twist-4 effects in spite of their different origin.

In section 2 we present the analysis of the ratio between the neutral and charged current total neutrino cross sections. From the general isoscalar symmetry relations and considering power scaling violations only, we deduce a general logarithmic dependence of the twist-4 uncertainty on the lower experimental y -cut. To estimate the numerical value of the

corrections to $\sin^2 \theta_w$ we use some examples of standard hadronic models, the counting rules and some available experimental information. The idea is to obtain the possible range of the twist-4 corrections, which turn out to be of the same order of magnitude as the sea quark uncertainties.

In section 3 we deduce the twist-4 modifications to the y -differential cross sections. We conclude that this correction modifies substantially the parametrization proposed in ref. [5] and we suggest to parametrize together both the strange quark and the twist-4 contributions introducing only one more unknown parameter.

Section 4 is devoted to the conclusions.

2. TWIST-4 CONTRIBUTIONS TO $R^{\nu} = \sigma_{NC}^{\nu}/\sigma_{CC}^{\nu}$

The inclusive cross-section for neutrino scattering off an unpolarized target is given in terms of the standard structure functions, by

$$\frac{d\sigma^{\nu}}{dx dy} = \frac{ME}{2\pi} G_F^2 \left[\left(1 + (1-y)^2 - \frac{My}{E} x \right) F_2^{\nu}(x, Q^2) - y F_L(x, Q^2) + (1 - (1-y)^2) x F_3^{\nu}(x, Q^2) \right] \quad (1)$$

$$F_L(x, Q^2) = F_2(x, Q^2) - 2x F_1(x, Q^2)$$

where x is the usual Bjorken variable and y is the fraction of the neutrino energy carried away to the target; $y = (E_H - M)/E$ in the laboratory frame and is related to the squared 4-momentum

transfer by $Q^2 = 2MExy$. We denote by $F_i(x, Q^2)$ the full structure functions and not just their leading twist. Below we expand them up to $1/Q^2$ order and consider a non-vanishing longitudinal structure function. We assume an ideal experiment with fixed neutrino energy (actually one has a distribution of neutrino energies in each particular case).

The factor $1/ME$ plays the role of $1/Q^2$ in the usual inverse power expansion after integrating over x and y .

The experimental selection of neutral current events demands a lower cut in the y variable (actually in the final hadronic energy $E_H > E_H^m$) and so the total cross-section will also depend on the minimum y value, y_m .

There are two exact relations for the case of isoscalar targets that follow from the structure of the weak currents. First we have the Paschos-Wolfenstein relation [9] which relates the neutral and charged current F_3 structure functions

$$xF_3^{NC}(x, Q^2) = \left(\frac{1}{2} - \sin^2 \theta_w\right) xF_3^{CC}(x, Q^2) \quad (2)$$

and second a relation that holds in the limit of zero mixing angle [4]

$$d\tau^{CC} = 2 d\tau^{NC} \quad (3)$$

These two symmetry requirements were not satisfied by the early estimate of ref. [6].

The general fact that the structure functions can be ex-

panded in powers of $1/Q^2$ ($1/2MEy$ in our case) together with eq. (2) allows us to write the structure functions to order $1/Q^2$ ($1/2MEy$) for an isoscalar target as

$$\begin{aligned} \tilde{F}_2^{NC} &= \left(\frac{1}{2} - z + \frac{10}{9}z^2\right)(\tilde{F}_2^q + \tilde{F}_2^{\bar{q}}) + \frac{1}{2MEy} \frac{\tilde{F}_2^{NC}}{x} \\ \tilde{F}_2^{CC} &= \tilde{F}_2^q + \tilde{F}_2^{\bar{q}} + \frac{1}{2MEy} \frac{\tilde{F}_2^{CC}}{x} \quad (z = \sin^2 \theta_w) \end{aligned} \quad (4.a)$$

$$\tilde{F}_L^{NC} = \left(\frac{1}{2} - z + \frac{10}{9}z^2\right)\tilde{F}_L^{CC}, \quad \tilde{F}_L^{CC} = \frac{1}{2MEy} \frac{\tilde{F}_L^{CC}}{x} \quad (4.b)$$

$$\tilde{F}_3^{NC} = \left(\frac{1}{2} - z\right)\tilde{F}_3^{CC}, \quad \tilde{F}_3^{CC} = \tilde{F}_2^q - \tilde{F}_2^{\bar{q}} + \frac{1}{2MEy} \frac{\tilde{F}_3^{CC}}{x} \quad (4.c)$$

where we have explicitly distinguished the leading twist ($\tilde{F}_2^q$ and $\tilde{F}_2^{\bar{q}}$) and the twist-4 ($\tilde{F}_1$) contributions. In general all these functions depend both on x and, logarithmically, on Q^2 . Therefore in our approach (only considering power scaling violations) they only depend on x and the leading twist functions are $\tilde{F}_2^q = x(u+d)$ and $\tilde{F}_2^{\bar{q}} = x(\bar{u}+\bar{d})$. In this approach the twist-4 contribution to the neutral current longitudinal structure function has the same $z = \sin^2 \theta_w$ dependence as the leading twist terms in $\tilde{F}_2$ (as shown by the parton model and the OPE results). In fact, this last point is the only additional information contained in eqs. (4) which is not already present in eqs. (2,3). In the case of $\tilde{F}_2^{NC}$ there is an extra dependence in z . It has to be a second order polynomial in z and eq. (3) implies that $\tilde{F}_2^{NC} = \frac{1}{2}\tilde{F}_2^{CC}$ in the $z=0$ limit.

We calculate the ratio of the neutral to charged current total cross-sections substituting eqs. (4) into eq. (1), integrating over x and y and expanding the ratio keeping only terms up to order $1/ME$. We obtain

$$\begin{aligned} R^{\nu} = \frac{\sigma_{NC}^{\nu}}{\sigma_{CC}^{\nu}} = \frac{1}{2} - z \left\{ 1 + \frac{4 \ln y_m + (2-y_m)^2 - 1}{1-y_m} \frac{a_1}{2ME} \right\} + \\ + z^2 \left\{ \frac{5}{9} \left(1 + \frac{(1-y_m)^2}{3} \right) + \frac{1}{ME} \left(\frac{(1-y_m)(5-2y_m)}{3} a_L - \right. \right. \\ \left. \left. - \frac{4 \ln y_m + (2-y_m)^2 - 1}{2(1-y_m)} \left[a_2 + \frac{1}{3} (1-y_m)^2 a_3 \right] \right) \right\} \quad (5) \end{aligned}$$

where we have not included the antiquark contributions and we have used the minimal standard model $\rho = 1$ value. The parameters needed to describe the twist-4 contributions are

$$a_L = \frac{5}{36} \frac{\langle 2xM^2 F_2^q + \tilde{F}_L^{cc}/x \rangle}{\langle F_2^q \rangle} = -\frac{5}{36} \frac{\langle \tilde{F}_3^{cc} \rangle}{\langle F_2^q \rangle} \quad (6.a)$$

$$a_3 = -\frac{5}{36} \frac{\langle \tilde{F}_2^{cc}/x \rangle}{\langle F_2^q \rangle} \quad (6.b)$$

where $\langle \rangle$ denotes integration over the whole range of x . We have used eq. (3) to define the a_1 and a_2 coefficients as

$$\frac{\langle \tilde{F}_2^{NC}/x - \left[\frac{1}{2} - z + \frac{5}{9} z^2 \left(1 + \frac{(1-y)^2}{3} \right) \right] \tilde{F}_2^{cc}/x \rangle}{4 \langle F_2^q \rangle} = a_1 z + \left[a_2 - \frac{5}{36} \frac{\langle \tilde{F}_2^{cc}/x \rangle}{\langle F_2^q \rangle} (1-y)^2 \right] z^2 \quad (7)$$

Eq. (6.a) is obtained from the parton model and OPE results.

Taking into account the fact that the contribution from a_L is negligible as shown below, the twist-4 uncertainty on the value of $\sin^2\theta_w$ can be obtained from eq. (5). For the case of $y_m \ll 1$ the result is

$$\delta \sin^2\theta_w = \frac{z}{1 - \frac{40}{27}z^2} \frac{a_1}{2ME} \sqrt{1 + \left(\frac{a_3 + 3a_2}{3a_1}\right)^2 z^2} \left| 3 + 4 \ln y_m \right| + O(y_m) \quad (8)$$

Therefore, for a fixed neutrino energy E the uncertainty in the value of $\sin^2\theta_w$ increases when the lower y -cut decreases. This effect implies that the expected uncertainty for the case of, say, $E = 160$ GeV and $E_H > 2$ GeV is nearly double than that for the same neutrino energy and $E_H > 10$ GeV.

We remark that eq. (5) is the most general parametrization of the twist-4 contribution at this level. Thus the y_m dependence obtained is general in this sense.

For the task of evaluating the magnitude of $\delta \sin^2\theta_w$ we make use of the expressions provided by the extended parton model [7] which give the parameters in eq. (6) in terms of multipartonic correlation functions. At tree level there are two kinds of processes responsible for power corrections. These are the process with a correlation between a gluon and a quark, and that with two correlated quarks.

Quark-gluon processes are described by handbag-like diagrams (fig. 1.a) and modify the scaling limit expression for R only through the violation of the Callan-Gross relation ($F_L \neq 0$).

The main corrections are expected to come from the quark-quark processes [6] which are given by non-handbag diagrams ((fig. 1.b) and have thus a very different flavor structure. The expressions for the parameters are (see appendix).

$$a_L = \frac{60 \langle \Lambda^2 T_{\pm}^- \rangle - 5M^2 \langle x^2(u+d) \rangle}{54 \langle x(u+d) \rangle} \quad (9.a)$$

$$a_1 = \frac{-\langle x \nabla_2^+(\beta_i, x) F_{\pm_3, \pm_3}^-(\beta_i) \rangle}{\langle x(u+d) \rangle} \quad (9.b)$$

$$a_2 = \frac{1}{36 \langle x(u+d) \rangle} \left\langle \left[F_{\pm_1, \pm_1}^+(\beta_i) + 16 F_{\pm_3, \pm_3}^+(\beta_i) \right] x \nabla_1^+(\beta_i, x) + \right. \\ \left. + \left[F_{\pm_1, \pm_1}^-(\beta_i) + 36 F_{\pm_3, \pm_3}^-(\beta_i) \right] x \nabla_2^+(\beta_i, x) \right\rangle \quad (9.c)$$

$$a_3 = \frac{-5 \langle x \nabla_1^+(\beta_i, x) F_{\pm_3, \pm_3}^+(\beta_i) \rangle}{9 \langle x(u+d) \rangle} \quad (9.d)$$

where there is an implicit integration over β_i .

$\langle \Lambda^2 T_{\pm}^- \rangle$ is the normalization of the correlation function of a quark and a gluon with opposite helicities. (Actually it is the sum of the corresponding functions with an up and a down quark). It is the only unknown in the quark-gluon sector and it can be determined from the data on the observable $R = \sigma_L / \sigma_T$ [10] which measures the violation of the Callan-Gross relation. The analysis of the SLAC-MIT data requires a value for $\langle \Lambda^2 T_{\pm}^- \rangle$ between 0.01 and 0.03 GeV^2 .

$F_{M,N}^{\pm}(\beta_i)$ is the correlation function of two quarks with total momentum fraction $\beta_1 + \beta_2 = \beta_1' + \beta_2'$ opposite (-) or equal (+) helicity signs and flavor structure fixed by the

flavor matrices $M, N (F_{I,1} = F_{uu} + F_{dd} + 2F_{ud}$ and $4F_{I,3} = F_{uu} + F_{dd} - 2F_{ud})$. The evaluation of the integrals of these functions which contribute to eq. (10) demands the knowledge of both their normalization and dependence on the momentum fractions which are not available from the data.

By imposing the parton results to agree with the OPE ones [5] the normalization of the quark-quark correlation functions can be related to the matrix elements of certain four fermions local operators.

$$g^2 \langle P | \frac{1}{2} \bar{\Psi}_L M \gamma_\mu \lambda^a \Psi_L \bar{\Psi}_L N \gamma_\nu \lambda^a \Psi_L + \mu \leftrightarrow \nu + L \leftrightarrow R | P \rangle = \\ = \left[\not{p}_\mu \not{p}_\nu - \frac{M^2}{4} g_{\mu\nu} \right] \langle F_{M,N}^+ \rangle \quad (10.a)$$

$$g^2 \langle P | \frac{1}{2} \bar{\Psi}_L \dots \Psi_L \bar{\Psi}_R \dots \Psi_R + \mu \leftrightarrow \nu + R \leftrightarrow L | P \rangle = \left[\not{p}_\mu \not{p}_\nu - \frac{M^2}{4} g_{\mu\nu} \right] \langle F_{M,N}^- \rangle \quad (10.b)$$

where λ^a is a colour matrix. Consequently these normalizations can be estimated by using a specific model for the nucleon target denoted by $|P\rangle$. We use here the various hadronic models collected in ref. [11]. In these models the spinor of the quarks inside the nucleon is of the usual spherically symmetric form

$$\Psi(r) = \begin{bmatrix} f(r) \\ \vec{\sigma} \cdot \hat{r} g(r) \end{bmatrix} \chi \quad (11)$$

where χ is a two-component spinor. Assuming $SU(6)$ wave functions for the nucleons, one can express the normalization of the functions in eq. (10) in terms of two integrals over the

nucleon volume

$$I_1 = \int d^3r [|f(r)|^2 + |g(r)|^2]^2, \quad I_2 = \int d^3r [f(r)g(r)]^2 \quad (12)$$

Their values depend on the particular model. The resulting expressions for the normalizations of the functions in Eq. (9) are

$$\langle F_{I,I}^+ \rangle = -\frac{\pi\alpha_s}{M} \frac{32}{3} I_1, \quad \langle F_{I,I}^- \rangle = -\frac{\pi\alpha_s}{M} \frac{64}{3} \left[I_1 - \frac{4}{3} I_2 \right] \quad (13.a)$$

$$\langle F_{I_3,I_3}^+ \rangle = -\frac{\pi\alpha_s}{M} \frac{8}{9} I_1, \quad \langle F_{I_3,I_3}^- \rangle = \frac{\pi\alpha_s}{M} \frac{32}{9} \left[I_1 - \frac{10}{3} I_2 \right] \quad (13.b)$$

where $\alpha_s = 4\pi g^2$ is the strong coupling constant at hadronic scales. This is a phenomenological parameter and we use as typical values 0.5, as suggested by the MIT computations [12], and 0.8, as resulting from a fit to the spectrum with one-gluon corrections [13].

For the functional dependence of $F_{M,N}^\pm$, we use the one suggested by the standard counting rules for high s . In the case of a process with two active quarks, the minimum number of spectators in a nucleon is one and so a behaviour $F_{M,N}^\pm(\beta_i) \sim (1-\beta_1-\beta_2)$ for $\beta_1 + \beta_2 \rightarrow 1$ is expected. To account also for lower values of $\beta_1 + \beta_2$, we use the function $(\beta_1 + \beta_2)^2 (1-\beta_1-\beta_2)$ in our estimates independently of the helicity and flavor structure*.

* The order of magnitude of the results does not change if one uses a simpler function as $(1-\beta_1-\beta_2)$.

The results for $\delta \sin^2 \theta_w$ in the case of a neutrino energy of 100 GeV and a cut in the final hadronic energy of 2 GeV are collected in table I. We present in this table the extreme values of $\delta \sin^2 \theta_w$ in each model when α_s runs between 0.5 and 0.8. The results do not depend on the $\langle \Lambda^2 T_1^- \rangle$ value (actually it produces contributions of order 1% in the above given range of values). We have used a typical value of $\langle x(u+d) \rangle = 0.395$. The corrections are mainly produced by the quark-quark processes and this is the reason of the strong model dependence of the estimate, and not having a reliable estimate of the matrix elements, we present the results for a broad range of models.

The conclusion is that the twist-4 corrections to the $\sin^2 \theta_w$ value range between 0.001 and 0.015 for a typical $\sin^2 \theta_w = 0.22$ value. So they can be important even in case of large neutrino energy. Anyhow, they are of the same order as those arising from sea uncertainties (~ 0.008) and larger than the precision of the forthcoming semileptonic experiments [3] (~ 0.005) and therefore not negligible as suggested in ref. [4].

3. TWIST-4 CONTRIBUTIONS TO THE y -DIFFERENTIAL CROSS-SECTIONS

It was shown in ref. [2] that the neutral and charged current cross sections can be related in the case of an isoscalar target with only up and down quarks and zero Cabibbo angle. The resulting equation for the twist-2 contribution is

$$\frac{d\tau_{NC}^{(\bar{\nu})}}{dx dy} = \left(\frac{1}{2} - z + \frac{5}{9} z^2 \right) \frac{d\tau_{CC}^{(\bar{\nu})}}{dx dy} + \frac{5}{9} z^2 \frac{d\tau_{CC}^{(\nu)}}{dx dy} \quad (14)$$

The use of this equation together with a suitable parametrization of the strange quark contributions has been proposed [5] recently to avoid the large uncertainties arising from the strange quark component of the nucleons. The interest of this proposal makes it important to study the twist-4 modifications to eq. (14) improving the treatment of ref. [4]. We show that these contributions modify substantially the above mentioned proposal in an important range of the hadronic final energy value. At the same time our general treatment suggests an appealing modification to it reducing the uncertainties in the twist-4 contribution as well.

From the general hypothesis of sec.2, i.e. isoscalar symmetry and expanding the structure functions at tree level in $1/Q^2$ powers, the twist-4 contribution to the charged current cross section is described by the substitution

$$\frac{d\sigma_{cc}^{(5)}}{dy} \rightarrow \frac{d\sigma_{cc}^{(5)}}{dy} - \frac{G_F^2 ME}{2\pi} \left\{ -\frac{My}{E} \langle x F_2^q \rangle + \frac{1+(1-y)^2}{2MEy} \langle \frac{\tilde{F}_2^{cc}}{x} \rangle - \frac{y}{2ME} \langle \frac{\tilde{F}_L^{cc}}{x} \rangle + \frac{1-(1-y)^2}{2MEy} \langle \frac{\tilde{F}_3^{cc}}{x} \rangle \right\} \quad (15)$$

We emphasize the peculiar $1/y$ dependence of the twist-4 contributions which distinguishes them from the other corrections and which is going to play a crucial role when we compare them with the strange quark ones. Of course, the resulting expansion in $1/MEy$ presents convergence problems in the $y \rightarrow 0$ limit, but luckily the lower y -cut imposed by the selection of neutral current events assures the convergence when this cut is large

enough.

For the neutral current case, the substitution is

$$\frac{d\tau_{NC}^{(v)}}{dy} \rightarrow \frac{d\tau_{NC}^{(v)}}{dy} - \frac{G_F^2 M E}{2\pi} \left\{ -\left(\frac{1}{2} - z + \frac{10}{9} z^2\right) \left[\frac{M y}{E} \left\langle \frac{\tilde{F}_2^q}{x} \right\rangle + \frac{y}{2ME} \left\langle \frac{\tilde{F}_2^c}{x} \right\rangle \right] + \frac{1-(1-y)^2}{2ME y} \left\langle \frac{\tilde{F}_2^{NC}}{x} \right\rangle + \left(\frac{1}{2} - z\right) \frac{1-(1-y)^2}{2ME y} \left\langle \frac{\tilde{F}_3^{cc}}{x} \right\rangle \right\} \quad (16)$$

Thus, the resulting modification for the eq. (14) is described by adding the term

$$+ \frac{G_F^2}{4\pi} \left\{ \left\langle \frac{\tilde{F}_2^{NC}}{x} \right\rangle - \left[\frac{1}{2} - z + \frac{10}{9} z^2 \right] \left\langle \frac{\tilde{F}_2^{cc}}{x} \right\rangle \right\} \frac{1+(1-y)^2}{y} =$$

$$= (\alpha_1 z + \alpha_2 z^2) \frac{1+(1-y)^2}{y} \quad (17)$$

at the r.h.s. of the equation. We have used eq. (3) to define α_1 and α_2 .

The complete relation between y -distributions including the derived twist-4 effect and the strange quark terms of ref. [5] reads

$$\frac{d\tau_{cc}^{(v)}}{dy} = \left[\frac{1}{2} - z + \frac{5}{9} z^2 \right] \frac{d\tau_{cc}^{(v)}}{dy} + \frac{5}{9} z^2 \frac{d\tau_{cc}^{(v)}}{dy} + z \frac{1+(1-y)^2}{y} +$$

$$+ (g_s^2 s) f(y) - \frac{1}{B} \left[\frac{d\tau_{di-x}^{(v)}}{dy} + \frac{d\tau_{di-x}^{(s)}}{dy} \right] \quad (18)$$

where we have defined $\tau = \alpha_1 z + \alpha_2 z^2$ since it is going to be a free parameter as $(g_s^2 s)$ and B are.

We remind now, that in the scaling limit, $f(y) = 1 + (1-y)^2$. So the twist-4 contribution will dominate this strange quark term for low enough values of y and the proposal in ref. [5] becomes not valid in this range.

With the same ingredients used in section 2 we can estimate the critical $y(E_H)$ value below which the twist-4 term is dominant. In terms of the correlation functions the parameters α_1 and α_2 in eq. (17) are given by

$$\frac{\alpha_1}{G_F^2} = -\frac{1}{\pi} \langle x \nabla_2^+(\beta_i, x) \bar{\Psi}_{I_3, I_3}^-(\beta_i) \rangle \quad (19.a)$$

$$\begin{aligned} \frac{\alpha_2}{G_F^2} = \frac{1}{36\pi} \langle x \nabla_1^+(\beta_i, x) [\bar{\Psi}_{I, I}^+(\beta_i) - 4 \bar{\Psi}_{I_3, I_3}^+(\beta_i)] + \\ + x \nabla_2^+(\beta_i, x) [\bar{\Psi}_{I, I}^-(\beta_i) + 36 \bar{\Psi}_{I_3, I_3}^-(\beta_i)] \rangle \end{aligned} \quad (19.b)$$

Following sec. 2 the resulting values are between 0.02 and 0.1 GeV^2 for α_1/G^2 and between -0.01 and -0.1 GeV^2 for α_2/G^2 . Assuming a typical value of $\langle xs \rangle = 0.01$ for the strange sea quarks and $\sin^2 \theta_w = 0.22$, the twist-4 contribution to eq. (18) results larger than the strange quark ones for final hadronic energies below a value which lies between 4 GeV and 23 GeV. We remark that these values are above the usual experimental cuts. Therefore we conclude that the proposed parametrization of ref. [5] is only applicable to quite large values of the final hadronic energy.

Our general analysis gives the correct equation to be used in the spirit of ref. [5] in all the available range of data. We think that the use of the y -distributions as proposed in ref. [5] with the inclusion of the above derived twist-4 term can be a successful method to avoid the large strange sea and the polemical twist-4 uncertainties in the semileptonic determination of θ_W . We have shown that one only needs to introduce a new parameter τ to be fitted with (g_s^2) and B .

Another general result of our analysis is that the data with an extremely low final hadronic energy will be more affected by higher twist corrections. Therefore, too low final hadronic energies ought to be avoided to preserve eq. (18) from modifications coming from twist higher than four.

4. CONCLUSIONS

We have analysed the twist-4 correction in a general fashion, studying the contribution to the ratio of neutral to charged currents total cross sections and the modification to the y -differential cross sections. These expressions are expansions in powers of $1/ME$ and $1/MEy$ respectively which are good approximations in principle if a lower y -cut is imposed as it is actually performed in experiments.

We have obtained from general arguments that the twist-4 uncertainty in the case of the total cross sections ratio at the tree level is described in terms of only four parameters. We have also derived that the resulting uncertainty in the value of $\sin^2\theta_W$ obtained from this ratio depends logarithmically

on the lower y -cut imposed. Thus, at fixed neutrino energy, the lower the cut the larger the uncertainty.

We identify the quark-gluon contributions with an experimentally measurable violation of the Callan-Gross relation. We have also estimated the quark-quark contributions by using different hadronic models and the standard counting rules. The result is that the uncertainty on $\sin^2\theta_w$ from the total cross sections ranges between 0.001 for $E = 100$ GeV, $E_H > 2$ GeV and $\sin^2\theta_w = 0.22$, which is of the same order of magnitude as the sea quarks ones.

We have also studied the twist-4 modifications to the general relation between y -differential cross sections. It is described in terms of only two parameters and presents a peculiar dependence $\frac{1+(1-y)^2}{y}$. The estimate of this contribution shows that it modifies substantially the equation proposed in ref. [5] to avoid the strange sea uncertainties.

We propose the inclusion of the resulting twist-4 term in the above mentioned parametrization as a good method to avoid simultaneously both hadronic uncertainties in the semileptonic determination of θ_w . We also suggest that an extremely low cut in the final hadronic energy should be avoided since it would enhance contributions of twist higher than four.

ACKNOWLEDGEMENTS

We thank J.L. Cortés for discussions and CAICYT for financial support. One of us (J.L.M.) thanks the University of Santiago for a grant to visit CERN and M. Jacob for the hospitality.

TABLE 1

Caption: Extreme twist-4 contributions to $\sin^2\theta_w$ for the various hadronic models in the case of $\sin^2\theta_w = 0.22$, $E = 100$ GeV, $E_H > 2$ GeV and α_s and $\langle\Lambda^2 T^-\rangle$ between 0.5 and 0.8, and 0.01 GeV² and 0.03 GeV² respectively.

MODEL	$\delta\sin^2\theta_w$
MIT bag (A) [14]	0.001 - 0.002
MIT bag (B) [15]	0.004 - 0.006
MIT bag (C) [15]	0.006 - 0.010
Relativistic potential [16], [17]	0.003 - 0.006
Harmonic oscillator shell model [18]	0.009 - 0.015

APPENDIX

The resulting twist-4 contributions to the structure functions in the case of a general current $J_\mu = \bar{\psi} \gamma_\mu [M + \gamma_5 N] \psi$, with M and N flavor matrix, are

$$\begin{aligned} F_2 = & \frac{x^2}{Q^2} \int d[\beta_i] \left\{ \left[F_{M,M^+}^+(\beta_i) + F_{N,N^+}^+(\beta_i) \right] \nabla_1^+(\beta_i, x) + \left[F_{M,M^+}^-(\beta_i) - F_{N,N^+}^-(\beta_i) \right] \nabla_2^+(\beta_i, x) \right\} + \\ & + \frac{x^2}{Q^2} \left\{ 8\Lambda^2 T_{M^+M+N^+N}^-(x) + 2x \int dx_1 dx_2 \frac{\delta(x_1-x) - \delta(x_2-x)}{x^2-x_1} \Lambda^2 T_{M^+M+N^+N}^+(x_1, x_2) \right\} \end{aligned} \quad (A.1)$$

$$F_L = \frac{8x}{Q^2} \Lambda^2 T_{M^+M+N^+N}^-(x) \quad (A.2)$$

$$\begin{aligned} F_3 = & \frac{x^2}{Q^2} \int d[\beta_i] \left\{ \left[F_{M,N^+}^+(\beta_i) + F_{N,M^+}^+(\beta_i) \right] \nabla_1^-(\beta_i, x) - \left[F_{N,M^+}^-(\beta_i) - F_{M,N^+}^-(\beta_i) \right] \nabla_2^-(\beta_i, x) \right\} \\ & - \frac{2x}{Q^2} \int dx_1 dx_2 \frac{\delta(x_1-x) - \delta(x_2-x)}{x_2-x_1} \Lambda^2 T_{M^+N+N^+M}^+(x_1, x_2) \end{aligned} \quad (A.3)$$

with

$$\nabla_1^\pm(\beta_i, x) = \frac{\delta(x-\beta_2-\beta'_1) - \delta(x-\beta_2)}{\beta'_1(\beta_1-\beta'_1)} + (\beta_1, \beta_2 \leftrightarrow \beta'_1, \beta'_2) \pm \frac{\delta(x-\beta_2+\beta'_1)}{\beta'_2\beta'_1} \quad (A.4)$$

$$\nabla_2^\pm(\beta_i, x) = \frac{\delta(x-\beta_2-\beta'_1)}{\beta'_1\beta'_2} - \frac{\delta(x-\beta_2)}{\beta'_1(\beta_1+\beta_2)} \pm (\beta_1, \beta_2 \leftrightarrow \beta'_1, \beta'_2) \quad (A.5)$$

In these equations we have not included the target mass terms [19]. $F_{M,N}^\pm(\beta_i)$ are quark-quark correlation functions whose flavor subindex makes reference to a fermionic structure $\bar{\psi}_1 M \psi_2 \bar{\psi}_2 N \psi_1$, and the $+$ ($-$) sign corresponds to equal (opposite) helicity signs. $\Lambda^2 T_A^\pm(x)$ is the correlation function of a quark and a gluon with opposite helicity signs and total momentum

fraction x . $\Lambda^2 T_A^+(x_1, x_2)$ corresponds to the same correlation process with equal helicity signs, being $x_1(x_2)$ the incoming (outcoming) quark momentum fraction. In the quark-gluon cases the flavor subindex corresponds to the fermionic structure $\bar{\psi} A \psi$.

These correlation functions satisfy general relations derived from the equations of motion

$$\langle \Lambda^2 T_A^+ \rangle = 4 \langle \Lambda^2 T_A^- \rangle \quad (A.6)$$

$$\langle F_{A,I}^+ + F_{A,I}^- \rangle = 2 \left[3 \langle x_1 \Lambda^2 T_A^+ \rangle - 7 \langle x \Lambda^2 T_A^- \rangle \right] \quad (A.7)$$

and in the case of an isoscalar target these other relations must be satisfied

$$F_{I,I_3}^{\pm}(\beta_i) = F_{I_3,I}^{\pm}(\beta_i) = 0 \quad (A.8)$$

$$F_{I^+,I^-}^{\pm}(\beta_i) = 2 F_{I_3,I_3}^{\pm}(\beta_i) \quad (A.9)$$

REFERENCES

1. J. Panman; Talk given at the First Aspen Winter Conference Series (1985), CERN preprint EP/85-35.
2. A preliminar value of $\sin^2\theta_w = 0.227 \pm 0.009$ was recently presented by the UA2 group at the "V topical workshop on $p\bar{p}$ collider physics" (Aosta, 1985) and "XIII Winter meeting on fundamental physics" (Cuenca, 1985).
3. F. Dydak et al.; CERN preprint SPSC/83-49.
F. Bergsma et al.; CERN preprint SPSC/84-1.
4. C.H. Llewellyn Smith; Nucl. Phys. B228 (1983) 205.
5. J. Panman; Phys. Lett. B154 (1985) 221.
6. M. Gluck and E. Reya; Phys. Rev. Lett. 47 (1981) 1104.
7. J.L. Cortés and J. Sánchez Guillén; Phys. Lett. 120B (1983) 396.
R.K. Ellis, W. Furmanski and R. Petronzio; Nucl. Phys. B212 (1983) 29.
S. Wada; Phys. Lett. B119 (1982) 427.
J.L. Miramontes; Ph.D. Thesis (Santiago de Compostela, 1985)
J.L. Miramontes, J. Sánchez Guillén et al.; in preparation.
8. S. Fajfer and R. Oakes; Phys. Rev. D30 (1984) and Proceedings of XXII International Conference on high energy physics, Vol. I (1984) 236.
9. E.A. Paschos and L. Wolfenstein; Phys. Rev. D7 (1983) 91.
10. J.L. Cortés, J.L. Miramontes and J. Sánchez Guillén; Phys. Rev. D30 (1984) 46.
M.J. Herrero, J.L. Miramontes et al.; in preparation.

11. S. Fajfer and R. Oakes; Phys. Lett. 131B (1983) 192.
12. R.L. Jaffe and M. Soldate; Phys. Lett. 105B (1981) 467.
13. W. Celmaster; Ph.D. Thesis (Harvard 1977) and Phys. Rev. D15 (1977) 1984.
14. T. de Grand, R.L. Jaffe, K. Johnson and J. Kiskis; Phys. Rev. D12 (1975) 2060.
15. R.H. Hackman, N.G. Deshpande, D.A. Dicus and V.T. Teplitz; Phys. Rev. D18 (1978) 2537.
16. P. Leal Ferreira, J.A. Helayel and N. Zagmy; Il Nuovo Cim. 55 (1980) 215.
17. R. Ravndal; Phys. Lett. 113B (1982) 57.
18. P. Colic, J. Trampetic and D. Tadic; Phys. Rev. D26 (1982) 2286.
19. O. Nachtmann; Nucl. Phys. B63 (1973) 237.
H. Georgi and H.D. Politzer; Phys. Rev. D14 (1976) 1829.

FIGURE CAPTION

Fig. 1. Diagrams corresponding to a typical process with correlated quark-gluon (a) and correlated quark-quark (b).

(a)

(b)

fig. 1