The Compact Muon Solenoid Experiment
Conference Report

Mailing address: CMS CERN, CH-1211 GENEVA 23, Switzerland

09 October 2013 (v2, 18 October 2013)

Searches for Exotic Physics at CMS

Khristian Kotov for the ATLAS and CMS collaborations

Abstract

In this talk, the latest results from ATLAS and CMS collaborations on searches for exotic physics are reviewed. A variety of final state signatures are presented using up to 20/fb of data from the 8 TeV LHC run of 2012.

Presented at *Particle and Astro Protvino XXIX-th International Workshop on High Energy Physics NEW RESULTS and ACTUAL PROBLEMS in PARTICLE and ASTROPARTICLE PHYSICS and COSMOLOGY*

Searches for Exotic Physics at LHC

K. Kotov on behalf of ATLAS and CMS collaborations

*Department of Physics, Ohio State University,
Columbus, Ohio 43210, USA*

*E-mail: kkotov@cern.ch

The latest results from ATLAS and CMS collaborations on searches for exotic physics are reviewed. A variety of final state signatures are presented using up to 20 fb^{-1} of data from the 8 TeV LHC run of 2012.

Keywords: CMS; ATLAS; Exotic

1. Introduction

In this document we review the latest results from ATLAS and CMS collaborations on searches for exotic physics. The term exotic physics encompasses a variety of independent physics models addressing specific shortcomings of the Standard Model (SM). Most of these models predict new particles such as heavy gauge bosons, excitations of the SM particles, dark matter candidates, new SM partner particles. These new particles can show up as narrow resonances or broad excess in different event signatures. A variety of final state signatures are investigated using up to 20 fb^{-1} of data from the 8 TeV LHC run of 2012. The detailed description of ATLAS and CMS detectors can be found elsewhere.^{1,2}

2. Di-lepton signature

ATLAS and CMS collaborations conducted searches for high-mass resonances decaying to electron or muon pairs.^{3,4} Both collaborations interpreted their results in the context of the Sequential Standard Model (SSM) Z' with SM-like couplings⁵ and the Z' expected in grand unified theories.⁶ In addition to that ATLAS also probed model of spin-2 Randall-Sundrum (RS) graviton.⁷ In Fig. 1 one can see the simulation shows a good agreement with data.

No evidence for non-standard model physics is observed, and 95% confidence level (C.L.) upper limits are set on the mass of the Z'_{SSM} and superstring-inspired Z'_ψ . The Z'_{SSM} is excluded for masses below 2.86 TeV in the two channels combined. The superstring-inspired Z'_ψ is excluded for masses below 2.60 TeV.

In addition to the standard search for resonances, CMS collaboration examined^{8,9} the dilepton shape for a broad excess and interpreted results within the

Fig. 1. The invariant mass spectrum of $\ell^+\ell^-$ events. Left — ATLAS, right — CMS. The points with error bars represent the data. The histograms represent the expectations from SM processes: Z/γ^* , $t\bar{t}$ and other sources of prompt leptons (tW , diboson production, $Z \rightarrow \tau\tau$), and the multi-jet backgrounds.

Arkani-Hamed-Dimopoulos-Dvali (ADD) model¹⁰ of extra dimensions. Limits of up to 4.93 TeV at 95% C.L. are set on the fundamental scale M_S , depending on the number of extra dimensions and the validity range of the theory.

3. Signature with lepton and missing transverse energy

CMS collaboration published an update on the search for the W' boson. The results are interpreted in scope of the SSM, in the framework split universal extra dimensions (split-UED),¹¹ and in terms of four-fermion contact interaction (CI).¹²

The main observable in this search is the transverse mass (M_T) of the system of lepton and missing transverse energy (E_T^{miss}), calculated as $M_T = \sqrt{2p_T^\mu E_T^{miss}(1 - \cos(\phi^\mu - \phi^{E_T^{miss}}))}$. In Fig. 2 one can see the transverse mass distribution in MC and data along with two example W' signals.

No significant excess has been observed in the data. The results are interpreted within the SSM under two assumptions: new W' is right-handed and it does not interfere with the SM W boson or new W' is left-handed and have constructive (destructive) interference with the SM W boson. Depending on the assumption the exclusion limit on the mass of SSM W' varies from 3.35 TeV to 3.10 TeV (3.60 TeV). The split-UED model is also probed for two assumptions on the bulk mass parameter μ of the fermion field in five dimensions. The exclusion limits on mass of the Kaluza-Klein excitations of the W' boson are set at 1.7 TeV (3.7 TeV) for $\mu=0.05$ TeV (10.0 TeV). The four-fermion CI is ruled out up to the scale of $\Lambda=13.0$ TeV for muons and 10.9 TeV for electrons.

4. Searches for diboson resonances

The two new searches for the WZ and ZZ resonances were conducted by ATLAS collaboration.^{13,14} The extended gauge model (EGM) with a heavy W' boson⁵ and the Low Scale Technicolor (LSTC) model¹⁵ were used as benchmark models to interpret the WZ resonance search results. The search for ZZ resonances was interpreted in

Fig. 2. Observed transverse mass distributions for the electron (left) and muon (right) channels. The shaded area represents the total systematic uncertainty on the SM background. Simulated signal distributions for a W' with masses of 0.5 and 2.5 TeV are also shown, including detector resolution effects.

the context of the "bulk RS model"¹⁶ where the spin-2 RS Graviton (G^*) decays to a ZZ pair. The final state with three leptons and E_T^{miss} is considered in WZ search while in ZZ search events with two leptons and one or two jets are selected. Fig. 3 compares data with the SM predictions.

Fig. 3. Invariant mass of diboson system (left — ZZ candidates, right — WZ candidates).

No signal is observed and the data are consistent with a smoothly falling background. The lower limit on mass of the bulk RS graviton is set at 850 GeV. The EGM W' is excluded below 1.18 TeV.

5. Dijet signature

The final state with two jets is a common signature for a wide range of models of new physics. Both collaborations found the data to be consistent with the SM predictions.^{17,18} Fig. 4 shows good agreement of the dijet mass spectrum in data and MC. The resulting limits on the cross section of new physics are shown in Fig. 5.

Fig. 4. Dijet mass spectrum. Left — ATLAS, right — CMS.

Fig. 5. Observed 95% C.L. upper limits on $\sigma \times B \times A$ for dijet resonances of type gluon-gluon, quark-gluon, and quark-quark, compared to theoretical predictions for string resonances, excited quarks, axigluons, colorons, E6 diquarks, s8 resonances, W' and Z' bosons, and RS gravitons.

The SM QCD dijet production is the dominant irreducible background for the search. An independent analysis by the CMS collaboration¹⁹ explored b-tagging as a mean to mitigate QCD background. Similar limits were set on the cross section of the new physics models.

6. $t\bar{t}$ signature

Many models of new physics predict enhanced couplings to the third generation quarks.^{20–22} A new heavy resonance may decay to a pair of heavily boosted top quarks. In this case, hadronic top decay gives three jets which are often merged within one "fat" jet. The jet substructure algorithms²³ allow to tag such jets. ATLAS and CMS collaborations conducted searches for $t\bar{t}$ resonances in semileptonic^{24,25} and fully hadronic²⁶ final states where $t\bar{t}$ event can either be reconstructed from its decay products or tagged with the jet substructure algorithms.

Fig. 6 shows good agreement between data and the SM predictions in the invariant mass spectrum of the $t\bar{t}$ system. The topcolor Z' model²⁷ and RS model were used to interpret the results. Depending on the width of the topcolor Z' the lower limit on mass is set at 1.8 TeV ($\Gamma/m = 3\%$) and 2.1 TeV ($\Gamma/m = 1.2\%$). RS KK excitations of gluon with mass less than 2.54 TeV are ruled out.

Fig. 6. Reconstructed distributions for the invariant mass of the $t\bar{t}$ system. The shaded areas indicate the total systematic uncertainties.

7. $t+b$ signature

ATLAS and CMS collaborations performed a search for a W -like resonance decaying into final state of top and bottom quarks.^{28,29} Fig. 7 shows good agreement of the invariant mass of kinematically reconstructed top quark and a b quark in data and SM predictions. For the right-handed W' (no interference with the SM W) with SM-like coupling the best exclusion limit is set at 2.03 TeV.

8. $t+j$ signature

The CMS collaboration searched for pair produced excited top quarks (t^*) each decaying to a top quark and a jets.³⁰ The final state includes a lepton, E_T^{miss} , and at least six jets. The final state particles are first selected to form the two W

Fig. 7. Reconstructed W' invariant mass distributions.

candidates, then extra jets are added to form two top quark candidates, and finally two t^* candidates are constructed from the top candidates and the remaining jets. The kinematic fit constraints are based on the W mass, the top quark mass, and the requirement for both t^* candidates to have the same mass.

Fig. 8 show the mass distribution of the reconstructed t^* candidate. No bumps or excess is observed over the smoothly falling mass spectrum. The limit on t^* mass is set at 794 GeV.

Fig. 8. The reconstructed top+gluon mass spectrum for data (points) along with the background fit to the data (curve), and signal distribution (histogram) for the e +jets (left) and μ +jets (right) channels.

9. Monojet signature

Searches for a dark matter in events with one or two high p_T jets was performed by both collaborations.^{31,32} A new weakly interacting particle, a dark matter candidate, can be produced in pp collision and escape detection. Thus, it can only be inferred by the presence of large E_T^{miss} . Fig. 9 demonstrates a good agreement between data

and the SM predictions for the E_T^{miss} spectrum.

Fig. 9. Measured E_T^{miss} distribution compared to the predictions for SM backgrounds (stacked solid histogram) and the signal models (dashed lines).

The ADD framework of Large Extra Dimensions and effective WIMP model³³ were used to interpret the results. Limits of up to 5.67 TeV at 95% C.L. are set on the ADD fundamental scale depending on the number of extra dimensions. Exclusion contours on WIMP model are given in Fig. 10.

Fig. 10. Comparison of CMS MonoJet 90% C.L. upper limits on the nucleon cross section versus dark matter mass for the spin independent model with CDF, XENON100, CoGeNT and CDMSII results.

References

1. S. Chatrchyan *et al.*, *JINST* **3**, p. S08004 (2008).
2. G. Aad *et al.*, *JINST* **3**, p. S08003 (2008).
3. *Search for high-mass dilepton resonances in 20 fb⁻¹ of pp collisions at sqrt(s) = 8 TeV with the ATLAS experiment*, Tech. Rep. ATLAS-CONF-2013-017 (2013).

4. *Search for Resonances in the Dilepton Mass Distribution in pp Collisions at $\sqrt{s} = 8$ TeV*, Tech. Rep. CMS-PAS-EXO-12-061, CERN (Geneva, 2013).
5. G. Altarelli, B. Mele and M. Ruiz-Altaba, *Z.Phys.* **C45**, p. 109 (1989).
6. A. Leike, *Phys.Rept.* **317**, 143 (1999).
7. L. Randall and R. Sundrum, *Phys.Rev.Lett.* **83**, 3370 (1999).
8. *Search for Extra Dimensions in Dimuon Events in pp Collisions at $\sqrt{s} = 8$ TeV*, Tech. Rep. CMS-PAS-EXO-12-027.
9. *Search for Large Extra Spatial Dimensions in Dielectron Production with the CMS Detector*, Tech. Rep. CMS-PAS-EXO-12-031.
10. N. Arkani-Hamed, S. Dimopoulos and G. Dvali, *Phys.Lett.* **B429**, 263 (1998).
11. K. Kong, S. C. Park and T. G. Rizzo, *JHEP* **1004**, p. 081 (2010).
12. H. Terazawa, M. Yasue, K. Akama and M. Hayashi, *Phys.Lett.* **B112**, p. 387 (1982).
13. *Search for resonant $WZ \rightarrow \ell\nu\ell\ell$ production using $\sqrt{s} = 8$ TeV pp collisions with ATLAS*, Tech. Rep. ATLAS-CONF-2013-015, CERN (2013).
14. *Search for resonant ZZ production in the $ZZ \rightarrow \ell\ell q\bar{q}$ channel with the ATLAS detector using 7.2 fb^{-1} of $\sqrt{s} = 8$ TeV pp collision data*, Tech. Rep. ATLAS-CONF-2012-150.
15. E. Eichten and K. Lane, *Phys.Lett.* **B669**, 235 (2008).
16. K. Agashe, H. Davoudiasl, G. Perez and A. Soni, *Phys.Rev.* **D76**, p. 036006 (2007).
17. *Search for New Phenomena in the Dijet Mass Distribution updated using 13.0 fb^{-1} of pp Collisions at $\sqrt{s} = 8$ TeV collected by the ATLAS Detector*, Tech. Rep. ATLAS-CONF-2012-148 (2012).
18. *Search for Narrow Resonances using the Dijet Mass Spectrum with 19.6 fb^{-1} of pp Collisions at $\sqrt{s} = 8$ TeV*, Tech. Rep. CMS-PAS-EXO-12-059.
19. *Search for Heavy Resonances Decaying into $b\bar{b}$ and $b\bar{g}$ Final States in pp Collisions at $\sqrt{s} = 8$ TeV*, Tech. Rep. CMS-PAS-EXO-12-023.
20. J. L. Rosner, *Phys. Lett. B* **387**, p. 113 (1996).
21. C. T. Hill and S. J. Parke, *Phys. Rev. D* **49**, p. 4454 (1994).
22. K. Agashe *et al.*, *Phys. Rev. D* **77**, p. 015003 (2008).
23. S. D. Ellis, C. K. Vermilion and J. R. Walsh, *Phys.Rev.* **D81**, p. 094023 (2010).
24. *A search for $t\bar{t}$ resonances in the lepton plus jets final state with ATLAS using 14 fb^{-1} of pp collisions at $\sqrt{s} = 8$ TeV*, Tech. Rep. ATLAS-CONF-2013-052, CERN (2013).
25. *Search for $t\bar{t}$ resonances in semileptonic final state*, Tech. Rep. CMS-PAS-B2G-12-006.
26. *Search for Anomalous Top Quark Pair Production in the Boosted All-Hadronic Final State using pp Collisions at $\sqrt{s} = 8$ TeV*, Tech. Rep. CMS-PAS-B2G-12-005, CERN (Geneva, 2013).
27. R. M. Harris and S. Jain, *Eur.Phys.J.* **C72**, p. 2072 (2012).
28. *Search for $W' \rightarrow t\bar{b}$ in proton-proton collisions at a centre-of-mass energy of $\sqrt{s} = 8$ TeV with the ATLAS detector*, Tech. Rep. ATLAS-CONF-2013-050, CERN (2013).
29. *Search for narrow $t + b$ resonances in the leptonic final state at $\sqrt{s} = 8$ TeV*, Tech. Rep. CMS-PAS-B2G-12-010.
30. *Search for pair production of new physics resonances decaying to a top quark and jet in the lepton+jet decay channel*, Tech. Rep. CMS-PAS-B2G-12-014 (2013).
31. *Search for new physics in monojet events in pp collisions at $\sqrt{s} = 8$ TeV*, Tech. Rep. CMS-PAS-EXO-12-048.
32. *Search for New Phenomena in Monojet plus Missing Transverse Momentum Final States using 10 fb^{-1} of pp Collisions at $\sqrt{s} = 8$ TeV with the ATLAS detector at the LHC*, Tech. Rep. ATLAS-CONF-2012-147 (2012).
33. Y. Bai, P. J. Fox and R. Harnik, *JHEP* **1012**, p. 048 (2010).