DIRECT PHOTON PAIR PRODUCTION FROM PIONS AND PROTONS AT 200 GeV/c

NA3 COLLABORATION

CEN, Saclay¹-CERN, Geneva²-Collège de France, Paris³
Ecole Polytechnique, Palaiseau⁴ - Lab. de l'Accélérateur Linéaire, Orsay⁵
Università di Pisa and INFN, Pisa⁶.

J. Badier⁴, M. Bardadin-Otwinowska^{2, 5*}), C. Bemporad⁶, H. Bienvenu⁴,
J. Boucrot⁵, J. Bourotte⁴, O. Callot⁵, P. Charpentier¹, A.M. Cnops⁶,
M. Cohen⁵, F. Costantini⁶, M. Crozon³, D. Décamp⁵, P. Delpierre³,
J.F. Detoeuf¹, G.R. Giannini⁶, Y. Karyotakis⁴, P. Lariccia^{6**}),
J. Mas³, A. Michelini², M. Moniez⁵, E. Paré⁴,
G. Rahal-Callot² and A. Tilquin³.

ABSTRACT

We present results from a direct photon pair search performed with the NA3 spectrometer, using incident positive and negative beams at 200 GeV/c interacting with a Carbon target. The experiment is sensitive to photons with $p_T > 1.8$ GeV/c and $-0.4 < y^* < 1.0$, one in each arm of the apparatus. A 3 standard deviation signal is seen in π^- and p interactions. The cross-section is higher than the second order QCD calculations, but systematic errors are large.

Submitted to Physics Letters B

(*) Now at Université de Clermont-Ferrand, France
(**) Università di Perugia, Italy

1. INTRODUCTION

The search for direct photon pairs is closely related to the study of direct single photon production, which is the main topic of the NA3 experiment. Using π^- or π^+ beam on an isoscalar target, it is possible to isolate the annihilation diagram $q\bar{q} \rightarrow \gamma + g$, proportional to e_q^2 . If one replaces the outgoing gluon by a photon, the cross-section decreases, but the charge dependence is now proportional to e_q^4 , and the ratio between π^- and π^+ may increase up to 16 at large p_T . The first experimental result on direct photon pair production comes from the experiment R806 at the ISR in p-p collisions at $\sqrt{s} = 63$ GeV [1]: they find a 2 standard deviation signal for $M_{\gamma\gamma} > 8$ GeV/c². More recent results from experiment R808 [2] confirm the signal, with the same statistical significance, for $4 < M_{\gamma\gamma} < 6$ GeV/c².

2. EXPERIMENTAL SET-UP AND TRIGGER

The experiment was performed with the NA3 spectrometer. Detailed description of the set-up can be found in refs. [3] and [4]. The main features are a large acceptance ($-0.4 < y^* < 1.0$), a good charged track measurement, and a lead-scintillator calorimeter with a MWPC inside it for shower position and energy determination.

The photon pair trigger is defined as follows [4]:

- At least 1.5 GeV/c vertical transverse momentum (p_{TV}) in one strip in the upper and lower part of the calorimeter.
- No charged track in these two strips. This veto condition on charged particles reduces the trigger rate by more than a factor of 10, but does not cut to too great an extent the $\gamma\gamma$ events, due to their expected topology.

3. SELECTION OF EVENTS

The photon pair triggers are reconstructed in the following steps:

- Trigger confirmation using calorimeter information, to ensure a clean p_{TV} cut and to avoid threshold effects.

- Charged track reconstruction, requiring one vertex in the targets.
- Shower localisation and energy measurement, using the calorimeter and the shower chamber.

We perform the same analysis in the upper and lower part of the apparatus. We then define as "main photon" the neutral shower with the highest p_{TV} . We reject events with a charged track within ± 10 mrad vertical angle from this main photon.

As usual in a direct photon experiment, a large fraction of the events comes from π^0 (or η^0) decays. In order to understand this background, we must measure precisely the $\pi^0\pi^0$ cross-section as seen in the apparatus.

4. $\pi^0\pi^0$ EVENTS

In each part (up, down) of the apparatus, we combine the main photon with all other reconstructed neutral showers in the same part to select π^0 with the following criteria:

- Transverse momentum of the $\gamma\gamma$ pair greater than 1.8 GeV/c.
- Asymmetry $A < 0.85$. A is defined by the relation $A = |E_1 - E_2| / (E_1 + E_2)$, E_1 and E_2 being the measured energies of the two photons. This cut rejects most of the background due to spurious signals around the shower core in the MWPC inside the calorimeter [3, 4].
- Using the energy resolution ($0.22/\sqrt{E}$) of the calorimeter and the position of the two associated photons, we compute the most probable energies e_1 , e_2 of the photons, and the probability δ for this pair to come from a π^0 decay [4]. π^0 candidates are selected by:
 - a cut on the probability δ
 - a cut on the energy e_2 , which must be greater than 0.7 GeV since the MWPC inside the calorimeter is inefficient below this energy.

The background under the π^0 signal, due to random association, can be estimated using events with two π^0 candidates reconstructed with the same main photon: the ratio $2\pi^0/1\pi^0$ is 2%.

Due to the cut on charged particles, we cannot measure the full $\pi^0\pi^0$ cross-section. We can only obtain the "non-accompanied" $\pi^0\pi^0$ cross-section, and use it to compute the contamination to the $\gamma\gamma$ events. We parametrize the distribution of each π^0 in the pair by:

$$\frac{d^2\sigma}{dydp_T} = A (1-x_D)^F / p_T (p_T^2 + m^2)^N$$

$$\text{with } x_D = \sqrt{x_T^2 + (x_{\parallel} - x_0)^2}$$

m^2 is fixed to .96, x_0, F, N are fitted to reproduce our data when this cross-section is put in the Monte-Carlo. Table 1 gives the parameters for π^-, π^+, p ; the normalization constant A cannot be given since it is biased by the trigger conditions.

We also measure the transverse momentum correlation of the π^0 's, defined by the angle α between their $\vec{p}_T$. α has a gaussian-like distribution with a standard deviation σ which is p_T dependent. Fig. 1 gives the value of σ as a function of the transverse momentum p_T of one π^0 . The observed decreasing of σ is expected from pure kinematics, and has been already observed in ISR experiments [5]. Dynamical effects in the hard scattering can give an asymptotical behaviour of σ at high p_T , as shown by ISR data in a p_T range much larger than ours.

5. EXTRACTION OF THE $\gamma\gamma$ SIGNAL

We look for a direct photon independently in the upper and lower part of the apparatus. We request the p_T of the "main photon" to be greater than 1.8 GeV/c. In the following, p_T will always be that of the "main photon". We try to identify π^0 (same criteria as before with asymmetry cut at .8) and η^0 (second photon with transverse momentum greater than 500 MeV/c and mass of the pair between 500 and 600 MeV). We regroup π^0 and η^0 in the same class. The remaining events are classified as γ . Using Monte-Carlo simulation with our $\pi^0\pi^0$ cross-section and the LUND model with the $q\bar{q} \rightarrow \gamma\gamma$ process [6], we compute the detection efficiency ϵ and acceptance A defined as follows:

$A^\gamma(p_T)$ is the probability that a generated photon is seen in the apparatus; it includes the efficiency due to the veto on charged tracks, estimated to be 63% by the LUND simulation.

$A^\pi(p_T)$ is the probability that a generated π^0 is seen in the apparatus;

$\epsilon(\gamma\rightarrow\gamma)$ (resp $\epsilon(\gamma\rightarrow\pi)$) is the probability that a generated photon seen in the apparatus is classified as γ (resp π^0): $\epsilon(\gamma\rightarrow\gamma) + \epsilon(\gamma\rightarrow\pi) = 1$

$\epsilon(\pi\rightarrow\gamma)$ (resp $\epsilon(\pi\rightarrow\pi)$) is the probability that a generated π^0 seen in the apparatus is classified as γ (resp π^0): $\epsilon(\pi\rightarrow\gamma) + \epsilon(\pi\rightarrow\pi) = 1$

The ϵ are p_T dependent, typical values are (at $p_T = 2$ GeV/c):
 $\epsilon(\gamma\rightarrow\gamma) = .99$, $\epsilon(\pi\rightarrow\pi) = .35$. The last small value is due to the topology of the selected π^0 , p_T being that of one photon.

We are left with 4 kinds of events: $\gamma\gamma$, $\gamma\pi$, $\pi\gamma$, $\pi\pi$. Let O^{ij} be the number of observed events and N^{ij} the number of true events (i and j are γ or π). We can write:

$$O^{ij} = \sum_{k,l} \epsilon(k\rightarrow i) \epsilon(l\rightarrow j) N^{kl}.$$

In other words, the 4 observed numbers are related by a 4 x 4 matrix M to the 4 real numbers. The latter are obtained by applying the inverse matrix to the observed numbers. The ϵ being p_T dependent, this is carried out event by event: $n^{ij} = M^{-1} O^{kl}$. The n^{ij} are weights, and are furthermore corrected by the acceptance (which is also p_T dependent and is different for real γ and real π^0). The sum of these corrected weights gives the number R of real $\gamma\gamma$, $\gamma\pi$, $\pi\gamma$, $\pi\pi$ events produced in the experiment. Cross-sections are then obtained by dividing by the luminosity.

The total $\gamma\gamma$ cross-section is given in table 2 and displayed on fig. 2. The values are 3-standard deviation from zero in π^- and p ; as expected, π^- cross-section is higher than π^+ .

6. STABILITY OF THE RESULT

We must be sure that this result is stable when varying the analysis. We have performed the following checks:

- For Monte-Carlo events, the number R^{ij} of real events obtained is non zero only for $\gamma\gamma$ events in the $\gamma\gamma$ simulation and $\pi\pi$ events in the $\pi\pi$ simulation. This tests the correctness of the method and of the parametrization of the $\epsilon(i+j)$.
- We change the asymmetry cut for π^0 reconstruction between .7 and .85.
- We change the p_T cut to 1.9 GeV/c.
- We suppress the η^0 search.
- We vary by $\pm 5\%$ the values of $\epsilon(i+j)$.

In all cases, the statistical significance of the $\gamma\gamma$ signal remains the same, but the cross-section varies by as much as 30%. This is an estimation of the absolute normalisation error, due to the analysis - other sources are luminosity ($\pm 5\%$) energy scale ($\pm 10\%$) and acceptance, the later depending on the assumed distribution of charged particles around the photons, and on the graphs introduced in the event generator.

7. COMPARISON WITH QCD CALCULATIONS

As photon pairs coming from the decay of χ mesons are cut by our trigger conditions, only hard parton collisions may contribute to the observed cross-section.

A full second order QCD calculation of the $\gamma\gamma$ cross-section has been recently performed [7]. It includes different graphs in the following orders of α_s (fig. 3): order zero ($q\bar{q}+\gamma\gamma$), order one ($q\bar{q}+\gamma\gamma g$, bremsstrahlung in the $qg+\gamma q$ graph,...) and order two ($gg+\gamma\gamma$,...). As compared with the Born term, and within our p_T and y^* cuts, the cross-section is about 2 times higher for π^- and 3 times for π^+ and p . This is mainly due to the "Box diagram" $gg+\gamma\gamma$. Values of the predicted cross-section are given in table 3 in the same domain than our data; they are substantially smaller than the measured ones. Many effects can contribute to the observed discrepancy

between data and the QCD calculation described above:

- the intrinsic momentum of the interacting partons was not taken into account in the calculation and may substantially increase the predicted cross-section at low p_T (the decreasing of the ratio data/prediction observed in tables 2 and 3 when p_T increases, could be related to this point);
- the predicted cross-section may also vary substantially when changing the Q^2 value (energy scale), or using a different set of structure functions; this gives for instance a factor about 2 on the prediction of the single direct photon yield [8];
- for proton data, we used the acceptance given by the $\pi^- N (q\bar{q} \rightarrow \gamma\gamma)$ simulation, since the introduction of the dominant graph $gg \rightarrow \gamma\gamma$ in the LUND Monte-Carlo was not performed;
- finally, the normalization error on the data is $\sim 50\%$.

8. CONCLUSION

Using π^- and p beam at 200 GeV/c on Carbon target, we obtain a 3 standard deviation $\gamma\gamma$ signal for p_T of each γ greater than 1.8 GeV and $-.4 < y^* < 1$. Measured values are higher than recent calculations, but systematical errors on data and prediction are large. The signal observed in π^- interactions is the first experimental evidence for direct $\gamma\gamma$ production with incoming pions.

ACKNOWLEDGEMENTS

We want to acknowledge gratefully M. Fontannaz and G. Ingelman for fruitful discussions and access to their programs. We also thank the NA3 technical staff for the operation of the experiment.

TABLE 1
Parameters of the π^0 distribution

	π^-	π^+	p
F	3.8 ± 1	$1.9 \pm .5$	$6.1 \pm .9$
x_0	$-.01 \pm .02$	$-.05 \pm .04$	$-0.06 \pm .02$
N	$3.49 \pm .24$	$3.83 \pm .13$	$3.30 \pm .22$

TABLE 2
 $\gamma\gamma$ production cross-section (nb)

	π^-	π^+	p
$p_T > 1.8 \text{ GeV/c}$	$1.22 \pm .35$	$.35 \pm .64$	$1.48 \pm .38$
$1.8 < p_T < 2.0$	$.61 \pm .25$		$.74 \pm .25$
$2.0 < p_T < 2.5$	$.43 \pm .24$		$.57 \pm .23$
$p_T > 2.5$	$.18 \pm .14$		$.17 \pm .13$

TABLE 3
 $\gamma\gamma$ cross-section: predictions from [7] (nb)

	π^-	π^+	p
$p_T > 1.8 \text{ GeV/c}$	.20	.07	.05
$1.8 < p_T < 2.0$	.06	.03	.02
$2.0 < p_T < 2.5$	.09	.03	.02
$p_T > 2.5$	.06	.02	.007

REFERENCES

- [1] C. Kourkouvelis et al., Zeitschrift für Physik C, Particles and Fields, 16 (1982) 101-109.
- [2] J. T. Åkesson et al., R808 Collaboration, Contribution to the International Europhysics Conference on High Energy Physics, Bari (Italy), July 18-24, 1985.
- [3] J. Badier et al., Nucl. Instr. and Meth. 175 (1980) 319;
F. Costantini et al., Nucl. Instr. and Meth. 225 (1984) 463.
- [4] G. Rahal-Callot, Thesis, Université Paris-Sud, May 1985, unpublished.
- [5] J.J. Cobb et al., Phys. Rev. Lett. Vol. 40, No. 22 (1979) 1420;
C. Kourkouvelis et al., Phys. Lett. Vol. 85B, No. 1 (1979) 147.
- [6] H.U. Bengtsson, G. Ingelman, Computer Physics Communication 34 (1985) 251.
- [7] P. Aurenche et al., Preprint LPTHE Orsay 85/20, May 1985.
- [8] J. Badier et al., Direct Photon Production from Pions and Protons at 200 GeV/c. Contribution to the International Symposium on Lepton and Photon Interactions, Kyoto (Japan), August 1985.

FIGURE CAPTIONS

Fig. 1: Standard deviation σ of the angular distribution of π^0 's in a $\pi^0\pi^0$ pair observed in this experiment, as a function of the transverse momentum p_T of one π^0 .

Fig. 2: Direct two-photon cross-section as a function of the transverse momentum p_T of one photon:
Full circles: π^- interactions
Open circles: p interactions

Fig. 3: Some graphs used in the QCD calculations of ref. [7].

Fig. 1

Fig. 2

Fig. 3