EUROPEAN ORGANIZATION FOR NUCLEAR RESEARCH

CERN-EP/85-158

2 October 1985

LIGHT-ION INTERACTIONS AT ULTRARELATIVISTIC ENERGIES

E. Stenlund^{*)}

Division of Cosmic and Subatomic Physics,
University of Lund, Sweden

ABSTRACT

Recent results from the light-ion runs at the CERN Intersecting Storage Rings are presented. Some different aspects of alpha-alpha, alpha-proton, and deuteron-deuteron interactions are discussed: nuclear spectators in high transverse momentum interactions; nuclear stopping power; break-up reactions; and, finally, many-particle rapidity correlations.

Invited talk given at the
Second International Conference on Nucleus-Nucleus Collisions,
Visby, Sweden, 10-14 June 1985

^{*)} Present address: CERN, Geneva, Switzerland.

1. INTRODUCTION

In this paper I will try to give a survey of recent results from the light-ion runs at the CERN Intersecting Storage Rings (ISR), obtained by the R418 Collaborations [1] using the Split-Field Magnet (SFM) detector [2]. The data samples represented here are from three different runs: $\alpha\alpha$ at $\sqrt{s_{nn}} = 31.2$ GeV; αp at $\sqrt{s_{nn}} = 44.0$ GeV; and dd at $\sqrt{s_{nn}} = 31.2$ GeV. The incoming energies correspond to 500 A GeV ($\alpha\alpha$ and dd) and 1000 A GeV (αp) in a fixed-target mode.

Several different triggers were used to select the data, and three of these will be mentioned here. The minimum-bias trigger required at least one charged particle anywhere in the detector. The high- p_T trigger required a charged particle with $p_T > 3$ GeV/c in one of two independent trigger regions around $\theta = 45^\circ$. The fragmentation trigger required one or two charged particles in each arm of the detector, together with a veto on particles in the central region.

The paper will cover four different subjects: nuclear spectators in high- p_T $\alpha\alpha$ and αp interactions; nuclear stopping power in $\alpha\alpha$ and dd interactions; break-up reactions in αp collisions; and, finally, many-particle rapidity correlations in dd interactions. It should be stressed that some of the data shown here are preliminary and might be subject to future changes.

2. NUCLEAR SPECTATORS IN HIGH- p_T $\alpha\alpha$ AND αp INTERACTIONS

The subject of high- p_T hadron production from nuclei has attracted wide interest owing to the so-called ‘anomalous nuclear enhancement’ [3]. Results for the inclusive cross-section of the trigger hadron as a function of p_T have already been published [4]. The aim of the present study is to examine the role of the nuclear environment in hard interactions by correlating the properties of particles in the forward region—in particular spectator protons and leading protons emitted from the α particles—with the presence of a high- p_T trigger hadron.

The invariant distributions $E(dn_+/dp_L - dn_-/dp_L)$ for $\alpha\alpha$ interactions are shown in Figs. 1a and 1b, whilst those for αp interactions are displayed in Figs. 1c to 1f. The proton mass was assigned to all particles when calculating their energies. In the case of $\alpha\alpha$ the longitudinal momentum p_L is measured in the c.m. system; for αp interactions a system has been chosen where the α particle has the same momentum as in the $\alpha\alpha$ interactions. Furthermore, it is important to note that in the αp case the α particle moves along the positive p_L -axis. For comparison, the corresponding distributions for minimum bias events (the solid curves in the figures) are also shown. Here, only events with at least one charged particle with $|y| < 2$ were considered. The arrows in the figures indicate the p_L of the trigger hadron, which in Figs. 1a, 1c, and 1e is positively charged, and is negatively charged in Figs. 1b, 1d, and 1f.

The spectra look similar for high- p_T events and minimum-bias events, although differences are observed at both high and low p_L . Most prominent is the shrinkage of the spectator peak at +15 GeV/c in the case of a positively charged high- p_T trigger particle at positive p_L (Figs. 1a and 1c). In all other cases the spectator peak is either larger than or consistent with the peak observed for minimum-bias events.

If the observed long-range correlation between the charge of the high- p_T trigger hadron and the number of spectator protons is interpreted on the quark level, it can be concluded that there is a substantial contribution of valence quarks in the high- p_T hadrons. In the case of a positively charged trigger hadron a u-quark is more likely to be involved, thus decreasing the number of spectator protons observed.

3. NUCLEAR STOPPING POWER IN $\alpha\alpha$ AND dd COLLISIONS

The baryon densities achievable in central collisions of heavy nuclei at high energies constitute a topic of current interest [5].

We have now investigated the nuclear stopping power in $\alpha\alpha$ and dd collisions. It can be assumed that in collisions of isoscalar nuclei the neutron inclusive spectra are the same as those for protons. As for most detectors, protons cannot be extracted in a direct way. Fortunately, in the case of isoscalar beams and targets ($\alpha\alpha$ or dd scattering) it is possible to extract the proton rapidity spectra by means of a statistical method. The proton mass is assigned for both negative and positive particles, and spectra of negative secondaries are subtracted from those of positive secondaries. As a consequence, non-pair-produced proton spectra with a correct mass assignment are obtained. The influence on the result from the difference between positive and negative K-production and the production of heavier fragments was neglected. The outcome of such a procedure applied to $\alpha\alpha$ and dd minimum bias data is shown in Fig. 2 for three different multiplicity bins. In the smallest multiplicity bin it can be seen that the proton distribution is peaked at $|y| = 3.5$, coinciding with the beam rapidity (spectator protons), and that it falls rapidly when the central rapidity region is approached. Large errors at central rapidities result from the subtraction of two large numbers. When events with higher multiplicities are selected, a shift of some of the protons toward the central rapidity region is seen. In the $\alpha\alpha$ case the effect seems to be much stronger. Obviously, in $\alpha\alpha$ collisions the nucleons have a chance of making more rescatterings than in dd collisions, which causes most of the difference.

4. BREAK-UP REACTIONS IN αp COLLISIONS

Fragmentation (or break-up) of a nucleus is a soft process, involving small momentum transfers. At present it is not possible to tell anything definite about such a process from QCD. Goldhaber [6] has assumed that the momentum distributions of the fragments are governed by the intrinsic momenta of nucleons in the fragmenting nuclei prior to the break-up. Later, other models have been suggested [7].

In order to study the fragmentation process, events have been selected using the fragmentation trigger. Furthermore, events containing charged pions (or kaons) were rejected off-line, by demanding the absence of negative particles and of particles with rigidity < 10 GeV/c. Pulse-height information from scintillators placed downstream were used to discriminate between $Z = 1$ and $Z = 2$ fragments, and rigidity cuts were applied to identify them.

Measurements of longitudinal and transverse momentum distributions of p, d, t, and ^3He fragments in $\alpha\alpha$ collisions have been reported elsewhere [8]. It was shown that the p_T distributions reflect elastic scattering among the nuclear constituents as a main mechanism, whereas the p_L distributions are only weakly perturbed by this mechanism and reflect mainly the internal motion of the nuclear constituents.

In the present analysis, correlations in the momenta of the outgoing beam proton and a proton fragment from the α particle are studied in αp interactions. In Fig. 3, where the distribution of relative azimuthal angles ϕ_r is displayed, the correlation between the two protons is illustrated. It is seen that the two protons are likely to be scattered in opposite directions, i.e. if the beam is diverted to one side, the proton fragment is in most cases seen on the opposite side of the collision axis. If the proton fragments are divided into two groups according to whether

$90^\circ < \phi_r < 270^\circ$ or not, the two groups show a different behaviour concerning their p_T distributions. The protons with $\phi_r < 90^\circ$ or $\phi_r > 270^\circ$ show a rather narrow spectrum, close to what would be expected from the internal motion of the nucleons only, whereas the other group shows a broader spectrum. These results are in good agreement with the idea of quasi-elastic scattering being the main mechanism.

5. MANY-PARTICLE RAPIDITY CORRELATIONS IN dd INTERACTIONS

A study of two-particle rapidity correlations in $\alpha\alpha$, αp , and pp interactions has already been published [9], where, in the framework of a simple cluster model, the cluster widths as well as their multiplicities were found to be the same for all three cases. We are at present analysing dd data with a method that is partly new [10].

The data have been analysed using the normalized rapidity-gap correlation function given by

$$R_k(G) = F_k(G)/\tilde{F}_k(G) - 1 \quad ,$$

where $G \equiv \Delta y$ is the length of the rapidity gap, and the index k stands for gaps of the k^{th} order. A k^{th} -order gap is a gap between two particles, where exactly $k-1$ particles have rapidities in the interval between these two particles. The function $F_k(G)$ is the experimental distribution of such gaps, whereas the function $\tilde{F}_k(G)$ is the corresponding distribution calculated from the experimentally determined single-particle distributions. Thus the ratio $R_k(G)$ is defined in exact accordance with the frequently used normalized two-particle correlation function

$$R(y_1, y_2) = \varrho(y_1, y_2)/\varrho(y_1) \cdot \varrho(y_2) - 1 \quad .$$

The advantage of discriminating between gaps of different orders is that not only are the two particles defining the gap taken into account, but also the rest of the particles in the event are considered.

Figures 4 and 5 display the distributions for dd data, where only charged particles in the central rapidity region $|y| < 1.5$ are considered. In Fig. 4 the functions $F_k(G)$ and $\tilde{F}_k(G)$ for $k=1$ and $k=8$ are shown; Fig. 5 shows the corresponding $R_k(G)$. Preliminary results obtained from a simple cluster approach are also indicated in Fig. 5 [9, 10]. The solid curves show the results for a large cluster (multiplicity ≈ 4 , width ≈ 0.8) and the dashed curves correspond to a smaller one (≈ 1.4 , ≈ 0.2). It is seen that the gaps at the lowest order are well described by the larger clusters, whereas the gaps of the 8^{th} order are best described by a fairly small cluster. However, if all orders are analysed simultaneously, the data seem to contradict the approach.

The method used to analyse the dd data will also be used to study $\alpha\alpha$ and pp interactions.

6. SUMMARY AND CONCLUSIONS

Many different aspects of light-ion interactions produced at the CERN ISR can be, and have been, studied. The subjects discussed here are just a fraction of those studied so far. They have been chosen for two reasons: they are projects which are not yet finished; and they are, in my opinion, the topics which are most related to nuclear physics.

The continuation of these and similar studies at high energies is desirable, possibly with larger nuclei. The next opportunity for this will be at the CERN Super Proton Synchrotron in 1986.

Acknowledgements

I would like to thank all my colleagues in the R418 Collaborations, especially S. Garpman, M.A. Faessler and R. Szwed. The financial support from the Swedish Natural Science Research Council is gratefully acknowledged.

* * *

REFERENCES AND FOOTNOTES

- 1) The members of these Collaborations are: W. Bell, K. Braune, A. Breakstone, G. Claesson, H.B. Crawley, G.M. Dallavalle, K. Doroba, D. Drijard, F. Fabbri, M.A. Faessler, A. Firestone, H.G. Fischer, H. Frehse, R.W. Frey, S.Y. Fung, S. Garpman, W. Geist, G. Giacomelli, R. Gokieli, M. Gorbics, C. Gruhn, P. Hanke, M. Heiden, W. Herr, P.G. Innocenti, T.J. Ketel, E.E. Kluge, J.W. Lamsa, T. Lohse, I. Lund, R. Mankel, W.T. Meyer, G. Mornacchi, T. Nakada, I. Otterlund, M. Panter, B. Povh, H. Pugh, A. Putzer, K. Rauschnabel, B. Rensch, F. Rimondi, K. Schmelling, L. Schroeder, R. Sosnowski, E. Stenlund, T.J. Symons, M. Szczekowski, R. Szwed, O. Ullaland, R. Walczak, D. Wegener and M. Wunsch.
- 2) See, for instance, W. Bell et al., Phys. Lett. **128B** (1983) 349.
- 3) J.W. Cronin et al., Phys. Rev. **D11** (1975) 3105.
U. Becker et al., Phys. Rev. Lett. **37** (1976) 1731.
L. Kluberg et al., Phys. Rev. Lett. **38** (1977) 670.
D.A. Garbutt et al., Phys. Lett. **67B** (1977) 355;
R.L. MacCarthy et al., Phys. Rev. Lett. **40** (1978) 213.
D. Antreasyan et al., Phys. Rev. **D19** (1979) 764.
C. Bromberg et al., Phys. Rev. Lett. **42** (1979) 1202.
- 4) W. Bell et al., Phys. Lett. **112B** (1982) 271.
- 5) W. Busza and A.S. Goldhaber, Phys. Lett. **139B** (1984) 235.
- 6) A.S. Goldhaber, Phys. Lett. **53B** (1974) 306.
- 7) A. Schmidt and R. Blankenbecler, Phys. Rev. **D15** (1977) 3321.
S.E. Koonin, Phys. Rev. Lett. **39** (1977) 680.
R.L. Hatch and S.E. Koonin, Phys. Lett. **81B** (1979) 1.
J. Hüfner and M.C. Nemes, Phys. Rev. **C23** (1981) 2358.
H. Araseki and T. Fujita, Nucl. Phys. **A399** (1983) 434.
- 8) W. Bell et al., Nucl. Phys. **B245** (1985) 475.
- 9) W. Bell et al., Z. Phys. **C22** (1984) 109.
- 10) L.K. Mangotra, I. Otterlund and E. Stenlund, Studies of many-particle correlations in proton-nucleus interactions using distributions of rapidity gaps between particles, Cosmic and Subatomic Physics Report, LUIP 8503, Lund (1985).

Figure captions

- Fig. 1** : Energy flow of positive excess particles in $\alpha\alpha$ and αp interactions for high- p_T trigger events. The solid curves show the corresponding distributions for minimum-bias events.
- Fig. 2** : Proton rapidity distributions for $\alpha\alpha$ and dd interactions. The solid curves show the distributions averaged over all multiplicities.
- Fig. 3** : Distribution of relative azimuthal angle between the outgoing beam proton and a proton fragment from the α particle in αp break-up reactions. The distribution is corrected by a division of the same distribution obtained with two protons from different events.
- Fig. 4** : Gap-length distributions for 1st and 8th orders.
- Fig. 5** : Normalized rapidity-gap correlation function for 1st and 8th orders. The curves correspond to model calculations.

Fig. 1

Fig. 2

Fig. 3

Fig. 4

Fig. 5