**COMPOSITION OF PARTICLES EMITTED AT LARGE p_T AND
MEDIUM ANGLES IN pp, dd, AND $\alpha\alpha$ COLLISIONS AT
THE CERN INTERSECTING STORAGE RINGS**

(Ames¹⁾–Bologna²⁾–CERN³⁾–Dortmund⁴⁾–Heidelberg⁵⁾–LBL⁶⁾–Lund⁷⁾–Warsaw⁸⁾ Collaboration)

A. Breakstone^{1,a)}, G. Claesson^{7,b)}, H.B. Crawley¹⁾, G.M. Dallavalle⁵⁾, K. Doroba⁸⁾,
D. Drijard³⁾, F. Fabbri³⁾, A. Firestone¹⁾, M.A. Faessler³⁾, H.G. Fischer³⁾,
S.Y. Fung^{c)}, S. Garpman⁷⁾, W. Geist^{3,b)}, G. Giacomelli²⁾, R. Gokieli⁸⁾,
M. Gorbics¹⁾, C.R. Gruhn⁶⁾, P. Hanke⁵⁾, M. Heiden^{3,b)}, W. Herr⁵⁾, T.J. Ketel^{d)},
E.E. Kluge⁵⁾, J.W. Lamsa¹⁾, T. Lohse⁴⁾, I. Lund⁷⁾, R. Mankel⁴⁾, W.T. Meyer¹⁾, G. Mornacchi³⁾,
T. Nakada^{5,e)}, I. Otterlund⁷⁾, M. Panter³⁾, H. Pugh⁶⁾, A. Putzer⁵⁾, K. Rauschnabel⁴⁾,
B. Rensch⁵⁾, F. Rimondi²⁾, M. Schmelling⁴⁾, G.P. Siroli²⁾, R. Sosnowski⁸⁾, E. Stenlund⁷⁾,
M. Szczekowski³⁾, R. Szwed⁸⁾, O. Ullaland³⁾, R. Walczak⁸⁾, D. Wegener⁴⁾ and M. Wunsch⁵⁾

ABSTRACT

The relative yields of kaons and protons compared with the yield of pions at high p_T and polar angles θ away from 90° (in the range from 10° to 45°) are presented for pp, dd, and $\alpha\alpha$ interactions at a centre-of-mass energy $\sqrt{s_{NN}} = 31$ GeV per nucleon–nucleon collision. The measured particle ratios depend on the atomic mass number A of the beam particles and on θ . The A dependence of the ratios becomes stronger for larger p_T and is more pronounced at smaller polar angles.

(Submitted to Zeitschrift für Physik C)

-
- 1) Ames Laboratory and Physics Department, ISU, Ames, Ia., USA.
 - 2) Istituto di Fisica dell'Università and INFN, Bologna, Italy.
 - 3) CERN, Geneva, Switzerland.
 - 4) Institut für Physik der Universität Dortmund, Fed. Rep. Germany.
 - 5) Institut für Hochenergiephysik der Universität Heidelberg, Fed. Rep. Germany.
 - 6) NSD, LBL, University of California, Berkeley, Calif., USA.
 - 7) Division of Cosmic and Subatomic Physics, University of Lund, Sweden.
 - 8) Institute of Exp. Physics, Univ. of Warsaw and Institute for Nuclear Studies, Warsaw, Poland.
 - a) Present affiliation: University of Hawaii, Honolulu, Hi., USA.
 - b) Present affiliation: LBL, Berkeley, Calif., USA.
 - c) Physics department, Univ. of Calif., Riverside, Calif., USA
 - d) Natuurkundig Lab., Vrije Universiteit, Amsterdam, the Netherlands.
 - e) Present affiliation: SIN, Villigen, Switzerland.

1. INTRODUCTION

It was recently shown that the fraction of protons emitted at high transverse momentum ($p_T > 3$ GeV/c) in pp interactions at $\sqrt{s} = 62$ GeV depends strongly on the c.m.s. polar angle θ [1]. The measured θ dependence provides a good constraint for theoretical models [2] based on parton scattering in the framework of quantum chromodynamics (QCD).

In this paper we study the production of high- p_T pions, kaons, and protons at polar angles around 10° , 20° , and 45° in pp, dd, and $\alpha\alpha$ interactions, i.e. as a function of the atomic mass number. The data were taken at the CERN Intersecting Storage Rings (ISR) at a centre-of-mass energy $\sqrt{s_{NN}} = 31$ GeV per nucleon-nucleon collision. The results are compared with previous measurements at $\theta \approx 90^\circ$ for proton-nucleus interactions [3], which show that the high- p_T cross-section per nucleon increases as a function of the atomic mass number A [4], and that this effect is more pronounced for protons and antiprotons than for mesons.

2. APPARATUS AND DATA REDUCTION

The experiment was performed using the Split Field Magnet (SFM) detector. Charged-particle trajectories through the magnetic field are recorded by a system of multiwire proportional chambers (MWPCs), and some particle identification is provided by an array of scintillation counters for time-of-flight measurements. A trigger logic selects events with a charged high- p_T particle ($p_T \geq 1$ GeV/c) produced at $\theta \approx 10^\circ$, 20° , or 45° . Five threshold Cherenkov counters (four filled with freon-114 at atmospheric pressure, at 10° , 20° , and 45° ; and one, at 45° , with freon-13 at 7.5 atm) are used to identify high- p_T pions. No separation of kaons and protons is possible in the momentum range considered. The data were corrected for Cherenkov counter inefficiencies, decay in flight of pions and kaons and absorption in the Cherenkov counters and trigger telescopes [1]. More details concerning the set-up, the acceptance, and the trigger scheme have been described elsewhere [5].

The data were taken during 1983. The numbers of events in the different high- p_T data samples are summarized in Table 1.

3. EXPERIMENTAL RESULTS

The ratios of inclusive cross-sections for kaon-plus-proton to pion production,

$$R^+ = [\sigma(K^+) + \sigma(p)]/\sigma(\pi^+) , \quad R^- = [\sigma(K^-) + \sigma(\bar{p})]/\sigma(\pi^-) ,$$

are presented in Fig. 1 and in Table 2 for different regions of trigger angle and different incident particles as functions of p_T . The corrections applied to the ratios of the inclusive cross-sections for equally charged high- p_T particles are described in Ref. [1]. The systematic uncertainties due to these corrections are indicated in Fig. 1.

As can be seen from Table 2, the mean polar angle for pp interactions differs slightly from that for dd and $\alpha\alpha$ scattering. This is due to different magnetic field configurations of the SFM during pp and light-ion data-taking. We conclude from the results presented in Ref. [6] that the shift in θ has an important influence on R^+ and R^- at angles around 20° , whilst for the p_T range of the present analysis the ratios are nearly independent of θ near 45° . The angular dependence of

R_{pp}^+ is shown in Fig. 2a for different values of p_T . Near $\theta \approx 20^\circ$ the θ dependence can be parametrized by a simple universal function

$$f_+(\theta) \sim \exp(554 \times \theta^{-2.69})$$

(θ measured in degrees). The data points were multiplied by p_T -dependent factors to exhibit the universal θ dependence. In order to compare the ratios R^+ at a fixed angle in Fig. 1, the function $f_+(\theta)$ was used to interpolate the data given in Table 2 to $\theta = 23^\circ$. Similarly, the negative ratios R^- were interpolated to a fixed angle of 15° using a simple linear dependence,

$$f_-(\theta) = 0.013 \times \theta + \text{const.}$$

This is demonstrated in Fig. 2b. This time the data points are shifted by p_T -dependent values in order to exhibit a universal behaviour at small angles. The same functions f_+ and f_- were used for the interpolation of dd and $\alpha\alpha$ data.

3.1 The ratio R^+

For both $\theta = 50^\circ$ and 23° , $R_{\alpha\alpha}^+$ is significantly larger than R_{dd}^+ over the full p_T range. Furthermore, R_{pp}^+ is compatible with R_{dd}^+ for both angular regions. The FNAL data for $\theta \approx 90^\circ$ at $\sqrt{s} = 27$ GeV [3] are also shown in Fig. 1. They are in reasonable agreement with R_{pp}^+ as measured at $\theta = 50^\circ$, suggesting that R^+ has no strong θ dependence between 90° and 45° . However, from a comparison of data obtained at $\theta = 50^\circ$ and 23° (and $\approx 10^\circ$ for the pp case, Table 2), it can be concluded that for $\theta < 45^\circ$ the ratios R^+ are steeply increasing for decreasing θ . This is demonstrated in more detail in Fig. 2 for pp collisions.

3.2 The ratio R^-

The negative ratio R^- shows the same behaviour as that of R^+ for the 20° region, i.e. $R_{dd}^- < R_{\alpha\alpha}^-$. The systematic errors indicated in Fig. 1 are mainly due to the interpolation procedure described above. Since $\alpha\alpha$ and dd data were taken with the same magnetic field configuration, all systematic effects should therefore approximately cancel in the difference $R_{dd}^- - R_{\alpha\alpha}^-$. Hence the observation that R_{dd}^- is smaller than $R_{\alpha\alpha}^-$ is significant. For $\theta \approx 45^\circ$ no obvious difference between R_{pp}^- , R_{dd}^- and $R_{\alpha\alpha}^-$ is found. Again the FNAL measurement [3] of R_{pp}^- is in good agreement with our results, which excludes a strong θ dependence between 45° and 90° . Between 15° and 45° R^- is increasing as a function of θ , hence showing a θ dependence opposite to that of R^+ .

3.3 The A dependence

In order to illustrate the A dependence, we show in Fig. 3 the R^+ and R^- values (interpolated to fixed θ) for two fixed values of p_T as a function of $A_1 \times A_2$, where A_1 and A_2 are the atomic mass numbers of the colliding particles. The results are compared with those of Ref. [3]. One observes a significant increase of R^+ at all angles and of R^- at 15° . In addition, there is (apart from the pp data points) a very slight increase of R^- at 45° and 90° as a function of $A_1 \times A_2$; the ratio R_{pp}^- does not seem to follow this trend. An observation similar to the latter was already made in Ref. [3]. This effect is likely to be due to the fact that protons, in contrast to

deuterons and alpha nuclei, are not isospin singlets. The extrapolation of the measured ratios to $A = 1$ is therefore not expected to agree exactly with the pp result.

At fixed θ the slope of the $A_1 \times A_2$ dependence of R^+ is increasing with p_T . Furthermore, for R^+ the slope is larger for the 45° and 20° data than for the 90° data. However, we note that the data at 90° [3] were recorded in proton-nucleus collisions instead of in nucleus-nucleus collisions as in our case.

4. SUMMARY AND CONCLUSION

The A dependence of the particle ratios R^+ and R^- has been measured in the p_T range from 1 to 4 GeV/c at polar angles θ around 10° , 20° , and 45° , for pp, dd, and $\alpha\alpha$ collisions. For all A , the ratio R^+ increases with decreasing θ , whereas R^- decreases. Both ratios increase as a function of $A_1 \times A_2$, at all angles θ . (The only exception to this trend, $R_{\bar{p}p}^-$ at large angles, can most likely be explained as an isospin effect.) The A dependence becomes more pronounced at smaller θ ($\theta = 23^\circ$ and 15°), for both R^+ and R^- , compared with the A dependence at $\theta = 90^\circ$ and $\theta = 45^\circ$.

Until now, no completely satisfying theoretical explanation for the A dependence of the inclusive cross-sections as a function of p_T [3, 4, 7] ('Cronin effect') has been found, despite of a large number of theoretical models [8]. The degree to which multiple parton interactions or a possible nuclear dependence of the nucleon structure functions (including Fermi motion) contribute to the observed enhancement of high- p_T particle yields with increasing A has always been an open question. The recently discovered 'EMC effect' [9] has again drawn attention to this question. Two recent theoretical studies [10] seem to indicate that the Cronin effect cannot be explained by incorporating the EMC effect in a single hard scattering scheme. The authors of reference 10 come to the conclusion that multiple parton interactions or cumulative effects are needed to reproduce the Cronin effect.

The FNAL experiment which discovered the Cronin effect, observed significantly different A dependences for the production cross-sections at $\theta = 90^\circ$ of $\pi^\pm$, $K^\pm$, p , and $\bar{p}$ (compared at equal p_T). This flavour dependence of the Cronin effect has been an important test of theoretical models. The present experiment explored, for the first time, particle ratios away from $\theta = 90^\circ$ and for collisions of two nuclei. It has already been shown [2] that measurements of the θ dependence provide sensitive tests for production mechanisms of high- p_T hadrons in pp reactions. The θ and A dependences of the particle ratios presented in this paper should hence be important for the understanding of these mechanisms.

Acknowledgements

We are grateful to the British-Scandinavian Collaboration and the Science and Engineering Research Council (UK) for making the high-pressure Cherenkov counter available to us. This experiment was greatly helped by contributions from the SFM Detector Group. We are also indebted to the ISR Experimental Support Group. The Dortmund and Heidelberg Groups were supported by a grant from the Bundesministerium für Wissenschaft und Forschung of the Federal Republic of Germany; the Ames Group, from the US Department of Energy under contract W-7405-eng-82; and the Lund Group, from the Swedish Natural Science Research Council. One of us (MAF) thanks the Deutsche Forschungsgemeinschaft for a grant.

REFERENCES

- [1] A. Breakstone et al., Phys. Lett. **147B**, 237 (1984).
T. Lohse, Ph.D. thesis, Univ. of Dortmund, 1985.
- [2] A. Breakstone et al., Z. Phys. C **28**, 335 (1985).
- [3] D. Antreasyan et al., Phys. Rev. **D19**, 764 (1979).
- [4] J.W. Cronin et al., Phys. Rev. **D11**, 3105 (1975).
- [5] D. Drijard et al., Nucl. Phys. **B208**, 1 (1982).
- [6] M. Della Negra et al., Phys. Lett. **59B**, 481 (1975).
- [7] A.L.S. Angelis et al., Phys. Lett. **116B**, 379 (1982).
T. Åkesson et al., Nucl. Phys. **B209**, 309 (1982).
W. Bell et al., Phys. Lett. **112B**, 271 (1982).
- [8] G. Farrar, Phys. Lett. **56B**, 185 (1975).
J. Pumplin and E. Yen, Phys. Rev. **D11**, 1812 (1975).
P.V. Landshoff et al., Phys. Rev. **D12**, 3738 (1975).
A. Krzywicki et al., Phys. Rev. **D14**, 152 (1976).
S. Fredriksson, Nucl. Phys. **B111**, 167 (1976).
P.M. Fishbane and J.S. Trefil, Phys. Rev. **D12**, 2113 (1976).
J.H. Kühn, Phys. Rev. **D13**, 2948 (1976).
P.M. Fishbane et al., Phys. Rev. **D16**, 122 (1977).
A. Krzywicki et al., Phys. Lett. **85B**, 407 (1979).
D. Treleani and G. Wilk, Nuovo Cimento **60A**, 201 (1980).
V.V. Zmushko, Sov. J. Nucl. Phys. **32**, 127 (1980).
U.P. Sukhatme and G. Wilk, Phys. Rev. **D25**, 1978 (1982).
M. Lev and B. Peterson, Z. Phys. **C21**, 155 (1983).
M. Staszal and G. Wilk, Z. Phys. **C19**, 57 (1983).
- [9] J.J. Aubert et al., Phys. Lett. **123B**, 275 (1983).
A. Bodek et al., Phys. Rev. Lett. **50**, 1431 (1983), **51**, 534 (1983).
- [10] T. Ochiai, S. Daté and H. Sumiyoshi, Rikkyo University preprint, RUP-85-4 (1985).
S. Gupta and R.M. Godbole, Tata Institute of Fundamental Research, TIFR-TH 85-16 (1985).

Table 1

Numbers of events in different high- p_T data samples used for the analysis

θ Run	45°	20°	10°
$\alpha\alpha$	31 700	39 225	–
dd	21 979	16 571	–
pp	26 250	286 980	483 812

Table 2

Particle ratios R^+ and R^- as functions of p_T for different polar angles in pp, dd, and $\alpha\alpha$ interactions. The errors quoted are purely statistical.

$R^+ = \frac{\sigma(p) + \sigma(K^+)}{\sigma(\pi^+)} \text{ at } \theta \approx 50^\circ$				
p _T	⟨θ⟩ (deg.)	pp	dd	αα
1.59	50.6	0.73 ± 0.03	0.74 ± 0.08	1.03 ± 0.08
1.69	55.8			
1.95	49.5	0.83 ± 0.04	0.82 ± 0.05	1.05 ± 0.05
2.05	53.2			
2.03	53.2	0.72 ± 0.06	0.94 ± 0.05	1.12 ± 0.06
2.38	48.9			
2.49	51.5			
2.48	51.5			
2.76	48.5	0.86 ± 0.09	0.91 ± 0.07	1.21 ± 0.08
2.98	50.6			
3.02	50.6	0.72 ± 0.12	0.76 ± 0.17	1.04 ± 0.16
3.32	48.4			
3.82	49.9			
3.86	49.9			
$R^- = \frac{\sigma(\bar{p}) + \sigma(K^-)}{\sigma(\pi^-)} \text{ at } \theta \approx 45^\circ$				
p _T	⟨θ⟩ (deg.)	pp	dd	αα
1.59	43.9	0.35 ± 0.02	0.25 ± 0.02	0.35 ± 0.03
1.68	41.5			
1.96	44.0	0.35 ± 0.02	0.31 ± 0.02	0.30 ± 0.02
2.00	42.0			
2.35	44.4	0.34 ± 0.04	0.34 ± 0.02	0.39 ± 0.03
2.36	42.6			
2.37	42.6	0.35 ± 0.06	0.30 ± 0.03	0.38 ± 0.04
3.00	44.9			
2.81	43.2			
3.33	43.8			
3.41	43.8		0.27 ± 0.03	0.27 ± 0.03

Table 2 (Cont.)

$R^+ = \frac{\sigma(p) + \sigma(K^+)}{\sigma(\pi^+)} \text{ at } \theta \approx 20^\circ, 10^\circ$					
pT	$\langle\theta\rangle$ (deg.)	pp 10°	pp 20°	dd 20°	$\alpha\alpha$ 20°
0.9	11.8	1.86 ± 0.09			
1.1	11.2	2.15 ± 0.023			
1.3	12.4	2.00 ± 0.01			
1.3	22.9		1.10 ± 0.01		
1.5	12.4	2.19 ± 0.02			
1.5	21.9		1.29 ± 0.01		
1.7	12.4	2.08 ± 0.04			
1.7	21.2		1.49 ± 0.01		
1.9	12.4	2.08 ± 0.09			
1.9	20.9		1.61 ± 0.03		
1.9	27.9				2.16 ± 0.17
2.1	12.4	2.07 ± 0.21			
2.1	20.7		1.78 ± 0.04		
2.1	25.9			1.81 ± 0.08	1.76 ± 0.10
2.3	20.5		1.79 ± 0.06		
2.3	24.6			1.68 ± 0.06	2.21 ± 0.08
2.5	20.5		1.76 ± 0.10		
2.5	23.7			1.58 ± 0.05	2.19 ± 0.08
2.7	20.2		1.82 ± 0.17		
2.7	23.0			1.72 ± 0.08	2.19 ± 0.10
2.9	20.8		1.53 ± 0.21		
2.9	22.6			1.84 ± 0.11	2.56 ± 0.17
3.1	21.1		1.13 ± 0.26		
3.1	22.3			1.76 ± 0.17	1.98 ± 0.18
3.3	22.1			2.27 ± 0.39	2.21 ± 0.36

Table 2 (Cont.)

$R^- = \frac{\sigma(\bar{p}) + \sigma(K^-)}{\sigma(\pi^-)} \text{ at } \theta \approx 20^\circ, 10^\circ$					
pT	$\langle\theta\rangle$ (deg.)	pp 10°	pp 20°	dd 20°	$\alpha\alpha$ 20°
0.7	6.6	0.074 ± 0.005			
0.9	7.8	0.099 ± 0.002			
0.9	13.4		0.127 ± 0.003		
1.1	8.6	0.097 ± 0.002			
1.1	14.3		0.134 ± 0.002		
1.3	9.0	0.083 ± 0.004			
1.3	14.9		0.140 ± 0.003		
1.3	12.9			0.112 ± 0.020	
1.3	12.5				0.130 ± 0.008
1.5	9.4	0.081 ± 0.012			
1.5	15.5		0.148 ± 0.004		
1.5	13.5			0.107 ± 0.010	
1.5	13.1				0.160 ± 0.006
1.7	16.0		0.142 ± 0.005		
1.7	14.2			0.115 ± 0.009	
1.7	13.8				0.159 ± 0.006
1.9	16.3		0.120 ± 0.007		
1.9	14.6			0.127 ± 0.009	
1.9	14.3				0.151 ± 0.007
2.1	16.8		0.119 ± 0.012		
2.1	15.0			0.110 ± 0.012	
2.1	14.6				0.160 ± 0.011
2.3	17.1		0.088 ± 0.017		
2.3	15.5			0.075 ± 0.013	
2.3	15.0				0.145 ± 0.015
2.5	17.7		0.151 ± 0.038		
2.5	15.8			0.112 ± 0.026	
2.5	15.4				0.147 ± 0.027
2.7	17.3		0.142 ± 0.067		
2.7	16.4			0.073 ± 0.043	

Figure captions

- Fig. 1 Particle ratios R^+ and R^- for different c.m. polar angles θ as measured in pp, dd, and $\alpha\alpha$ interactions. The systematic errors are indicated in the figures. For comparison, the pp data from Ref. [3] are also shown.
- Fig. 2 Particle ratios R_{pp}^+ (a) and R_{pp}^- (b) as a function of θ for different values of the transverse momentum. Some of the symbols are shifted horizontally for better visibility. In the insert of (a) the (p_T -dependent) factors $B(p_T)$ are shown, by which the data were multiplied to normalize to the data at $p_T = 0.9$ GeV/c; in the insert of (b) the (p_T -dependent) numbers $C(p_T)$ are shown, by which the data were shifted to normalize to the data at $p_T = 0.7$ GeV/c.
- Fig. 3 Particle ratios R^+ and R^- at fixed transverse momenta p_T as functions of $A_1 \times A_2$ (see text) for different polar angles θ . The results from Ref. [3] are also shown.

Fig. 1

Fig. 2

Fig. 3