CERN-TH.4307/85

MONOJETTS IN THE STANDARD MODEL

S.D. Ellis ^{*)}

University of Washington, Seattle

R. Kleiss and W.J. Stirling

CERN - Geneva

A B S T R A C T

In a previous analysis we have shown that in the standard model the number of events with large missing transverse energy in $p\bar{p}$ collisions can be experimentally important. We now refine this analysis to confront the sample of monojet events recently reported by the UA1 collaboration. We show how an absolutely normalized calculation based on simple parton level cross-sections - which gives good agreement with the production cross-sections for observed W and Z events - appears to satisfactorily describe the monojet sample. We emphasize the various sources of uncertainty in the calculation.

^{*)} Research supported in part by the US Department of Energy, Contract N° DE-AC06-81ER-40048

1. INTRODUCTION

The importance of events with large missing transverse energy at the CERN collider - in particular the so-called "monojet events" - as a signature for physics beyond the Standard Model (SM) is by now well appreciated [1]. In this respect the fundamental question is to what extent there is an excess of observed events over SM expectations [2,3]. The UA1 collaboration have recently reported a total of 29 monojet events from the combined 1983 and 1984 data, and have demonstrated that at least in terms of overall rate, these can be accounted for by known SM processes ($Z \rightarrow \nu + \nu$, $W \rightarrow \tau + \nu$ etc) [4]. This background rate is obtained by calibrating to observed cross-sections ($Z \rightarrow e + e$, $W \rightarrow e + \nu$ etc) and necessarily involves a detailed Monte Carlo simulation of the events in the detector. Apart from the overall complexity, one drawback of this approach is that it becomes rather uncertain in kinematic regions where the "calibrating" events are sparse.

In this letter we show how a much simpler analysis - based on a perturbative calculation of W,Z production and decay - is also able to describe the monojet data. Although it is impossible for such an approach to reproduce the precise details of the experimental measurements, it does have several advantages. First, it allows an independent check on the UA1 SM estimate. Second, it gives an estimate of the missing energy cross-section in regions where at present it is impossible to calibrate from other processes and, third, it permits a direct extrapolation to the higher collision energies of future hadron colliders.

2. THE CALCULATION

The philosophy of our approach has already been described elsewhere ([3], hereafter referred to as I). To summarize, events with large missing transverse energy - balanced by jets - can arise from a wide variety of processes within the SM without being easily identifiable. While individual contributions may be small, the sum of the "many roads" can give a sizeable result. We shall restrict our attention to the following which, in the present context, are the most important:

$$\begin{aligned}
 p + \bar{p} &\rightarrow Z^0 + \text{jets} \\
 &\quad \downarrow \\
 &\quad \nu + \bar{\nu} \\
 \\
 p + \bar{p} &\rightarrow W + \text{jets} \\
 &\quad \downarrow \\
 &\quad \tau + \nu \\
 &\quad \quad \downarrow \\
 &\quad \quad \text{hadrons} + \nu \\
 \\
 p + \bar{p} &\rightarrow W + \text{jets} \\
 &\quad \downarrow \\
 &\quad \ell + \nu \\
 \\
 p + \bar{p} &\rightarrow W + \text{jets} \\
 &\quad \downarrow \\
 &\quad \tau + \nu \\
 &\quad \quad \downarrow \\
 &\quad \quad \ell + \nu
 \end{aligned}$$

(1)

Here ℓ represents a charged lepton (e or μ) which escapes detection for reasons described in more detail below. The associated production of 0,1 and 2 hadronic (i.e. quark and gluon) jets is included [5]. Since the energy of the τ lepton is always much larger than its mass, a collinear fragmentation approximation is valid. For hadronic τ decays we follow previous studies and sum over known scalar (π) and vector (ρ, A_1, ρ') meson resonances [6,7]. The fragmentation functions are calculated analytically using a meson- τ - ν effective interaction, incorporating masses and finite widths where appropriate. It has been pointed out that there remains approximately 10% of τ decays which cannot be attributed to known resonances [7,8]. We model this by a $q\bar{q}$ continuum, integrating over the $q\bar{q}$ invariant mass. The resulting fragmentation function is shown in Fig.1 with, for comparison, the corresponding function for $\tau \rightarrow e, \mu$.

Note that we do not include any contribution from the production and semi-leptonic decay of heavy quarks ($c, b, t, \dots$). Such events are in practice expected to be highly suppressed by the monojet cuts described below. Furthermore, it is difficult to give a reliable theoretical estimate for these processes given the uncertainties both in the production rate and in the fragmentation properties of heavy quarks. To this extent our predictions for missing energy events constitute a lower bound.

The above processes are treated perturbatively as described in 1. The missing transverse energy is associated with one or more neutrinos. In practice this missing transverse energy is inferred from the measured transverse energy in the event, which includes both the contribution from the hard scattering (the jets) and from the rest of the "underlying event". This measurement is performed with a finite resolution which for the UA1 detector is given approximately by a Gaussian distribution of width $\sigma = .7 \sqrt{E_T^{\text{tot}}}$ [9]. We model this by smearing the transverse energy using the theoretical definition

$$E_T^{\text{tot}} = E_T^{\text{hs}} + E_T^{\text{ue}} \quad (2)$$

where E_T^{hs} is the transverse hadronic energy generated in the hard scattering and E_T^{ue} is the contribution from the "underlying event". This latter contribution takes into account the transverse energy of the beam (spectator) fragments and also soft radiation from the incoming partons which participate in the hard scattering. In each event a value for E_T^{ue} is generated according to the empirical distribution $E^2 \exp(-bE)$ with $\langle E_T^{\text{ue}} \rangle = 3/b = 30$ GeV. This form leads to a total E_T , as defined above, which yields a good fit to the observed distribution in total transverse energy accompanying the UA1 1983 $W \rightarrow e + \nu$ events [10], as indicated in Fig.2. A measure of the uncertainty inherent in this procedure can be obtained by varying $\langle E_T^{\text{ue}} \rangle$ from 20 to 40 GeV. As we shall see, the cross-section for LARGE missing transverse energy (E_T^{miss}) does not depend significantly on the choice of $\langle E_T^{\text{ue}} \rangle$.

As with all such theoretical calculations, an important source of uncertainty arises from the different possible choices of structure functions and hard scattering scale Q . In the present calculation we gauge this uncertainty by using two "modes" - mode 1 uses Duke and Owens set 1 structure functions ($\Lambda = 200$ MeV) with $Q = M_V$ ($V = W$ or Z) and mode 2 uses Gluck Hoffman Reya structure functions ($\Lambda = 400$ MeV) also with $Q = M_V$. As we shall see these two choices give similar predictions.

This completes the list of ingredients for the calculation of missing energy events. To summarise, we calculate the cross-sections for W and Z production with up to 2 additional hadronic jets. We consider decay modes which give rise to missing transverse energy balanced by hadronic jets. The event selection

cuts, chosen to match as closely as possible those actually used by UA1 and described in detail below, are implemented on the leptons and partons (jets) in the final state. The missing energy is smeared according to the UA1 resolution as described above. The parameters (principally the structure function choice and the value of $\langle E_T^{ue} \rangle$) are varied to give a range of predictions. Finally, the cross-sections are combined with the appropriate integrated luminosity to give the total number of events.

3. W,Z PRODUCTION AND ELECTRONIC DECAY

Since we are attempting to perform an absolutely normalised calculation of the monojet rate, it is important to check that we are able to satisfactorily describe OBSERVED W and Z production. For definiteness we consider the electron decay modes of the weak bosons as measured by UA1 [11]. The relevant cuts on electrons and jets have been described in I. To summarise, we require electrons and jets to have a minimum transverse energy of 15 GeV and 6 GeV respectively, to have a pseudorapidity in the range $|\eta| < 2.5$ and to be separated from each other by at least one unit in the lego plot ($\Delta r = \sqrt{(\Delta\eta^2 + \Delta\phi^2)}$). The integrated luminosities are 136 nb^{-1} and 263 nb^{-1} at $\sqrt{s} = 546 \text{ GeV}$ and 630 GeV respectively [11]. The resulting expectation for the total number of $W \rightarrow e + \nu$ and $Z \rightarrow e + e$ events is shown in Table 1, together with the UA1 data. The agreement is clearly very good. Both the overall rate and the rate at large transverse momentum are consistent with observations. Table 1(b) also illustrates the point made in the Introduction. The number of missing energy events from $Z \rightarrow \nu + \nu$ can be crudely estimated by multiplying the observed $Z \rightarrow e + e$ rate by a factor of 6. (The actual enhancement will be larger because of the reduction in the observed electron rate due to the cuts.) Because of low statistics, calibration via observed events is rather uncertain at present for events with $E_T^{\text{miss}} > 20 \text{ GeV}$. Theoretically, about 6 events (monojets, dijets, ...) are expected.

For brevity, we have restricted our comparison to the total number of W and Z events. Other weak boson production properties - multiplicity of jets, jet angular and transverse energy distributions etc. - are also in good agreement with theoretical expectations [5,12].

4. MONOJETS

The 1984 UA1 monojet cuts can be summarised as follows: an event with a missing transverse energy vector (M), a large transverse energy jet (J) and possibly other jets (j) is accepted as a monojet event if

- (a) $E_T(M) > \max(4\sigma, 15 \text{ GeV})$ $\sigma = .7 \sqrt{E_T^{\text{tot}}}$
 $\phi(M, \text{vertical}) > 20^\circ$
- (b) $E_T(J) > 12 \text{ GeV}$
 $|\eta(J)| < 2.5$

and the additional jets satisfy

- (c) $E_T(j) < 12 \text{ GeV}$
 and if $8 \text{ GeV} < E_T(j) < 12 \text{ GeV}$
 then also $\phi(M, j) > 30^\circ$ and $\phi(J, j) < 150^\circ$

Here ϕ is the azimuthal angle between the three momenta. These are the cuts which are relevant to the present study. In practice there are of course other event validation criteria [4]. (Recall also the previous jet separation cut $\Delta r(J, j) \geq 1.0$.)

As described in I, the SM contribution to the monojet signal arises from events which cannot be identified as SM due to "lost" leptons. For neutrinos and hadronic decays of τ leptons it is easy to understand how they are lost. For the case of electrons and muons this situation arises either because the lepton is too soft ($E_T < E_T^{\min}$), has too large a rapidity ($\eta > \eta_{\max}$) or is too close to a jet ($\Delta r(\ell, J) < r_{\min}$). As discussed more fully in I, the values we use for the boundary values $E_T^{\min}, \eta_{\max}, r_{\min}$ are 10 GeV, 3.0, 0.4 for electrons and 3 GeV, 3.0, 0.0 for muons.

In the UA1 1984 data sample there are 23 events which survive the cuts described above [4]. The distribution of these events in E_T^{miss} is shown in Fig.3. When the 1983 data are analysed in the same way, there are an additional 6 events. For simplicity we calculate the SM monojet cross-section at the single energy $\sqrt{s} = 630 \text{ GeV}$ and so will restrict our comparison to the 1984 events. The results of the theoretical analysis - applying the cuts described above both

for "lost" leptons and event selection - are summarised in Tables 2 and 3. (Note that the E_T smearing procedure described above serves also to smear the p_T of the W in the case of $\tau \rightarrow 0$ jets events leading to a distribution with events beyond the naive missing E_T limit of $M_W/2$.) Table 2 gives the results for mode 1 and shows the contribution from the various roads described above. Table 3 gives the results for modes 1 and 2 and also shows the effect of varying $\langle E_T^{ue} \rangle$ by ± 10 GeV. As expected this last change only significantly effects events close to the 4σ cut. Increasing $\langle E_T^{ue} \rangle$ has the effect of increasing σ and so fewer events survive the cut. The E_T^{miss} distribution for mode 1 with $\langle E_T^{ue} \rangle = 30$ GeV is compared to the data in Fig.3.

Evidently these absolutely normalised calculations give slightly too many events below $E_T^{\text{miss}} = 30$ GeV. The most likely explanation for this is that we have not yet included any TRIGGER SELECTION cuts on the final state. Thus for a careful comparison with the data it is necessary that candidate missing E_T events pass not only the above final selection cuts but also some simulation of the trigger selection criteria which define the initial data sample. Because of inefficiencies and resolution effects in measuring the jet momenta, it is unlikely that all of the above events will pass these initial trigger selection criteria. To get some feeling for the size of this effect we can implement trigger cuts on the final state partons. For example, following the analysis of [13], we reduce all jet momenta by 20% and smear the resulting transverse momenta using a Gaussian of width $.2p_T^J$. The trigger criteria [9] are implemented on the resulting momenta and correspond to demanding that there is either one hard jet ($E_T > 25$ GeV) OR that the missing transverse energy calculated from the rescaled, smeared jet E_T 's shows a "right-left" imbalance of more than 17 GeV and there is a jet with $E_T > 15$ GeV. The effect of these trigger criteria is shown in Table 4 and by the dashed line in Fig.3. We see that the trigger is more than 95% efficient for $E_T^{\text{miss}} > 30$ GeV but considerably less efficient at lower E_T . We can therefore take as significant the agreement between the theoretical predictions and the data for $E_T^{\text{miss}} > 30$ GeV, as shown in Table 2.

An important component of our total rate is the contribution from events where the monojet is a τ -jet (i.e. the hadronic decay products of a τ). A further check on the standard model origin of the monojets is to confirm that these τ -jets are present in the data at the expected level. Not surprisingly, difficulties arise in attempting to quantify the " τ -ness" of a jet. The basic premise is that τ -jets are narrow, with a small invariant mass and a small

charged particle multiplicity. The UA1 collaboration have introduced a quantity which uses these properties to measure the τ -ness of a jet. Out of the total sample of 23 1984 monojets 9 are identified as τ -jet candidates [4]. Without a detailed fragmentation model and a full detector simulation we cannot quantify the SM monojets in the same way. However we can compute very easily the part of the theoretical cross-section in which the "bare" τ -jet is the observed monojet. This component is shown in Fig.4 together with the non- τ -jet part (i.e., where the monojet is an ordinary quark or gluon jet or a τ jet "hidden" in an ordinary jet). Evidently the ratio of τ -jets to non- τ -jets decreases as E_T^{miss} increases. This appears to be consistent with observations [4]. The number of expected τ -jet events BEFORE trigger cuts is shown in Table 2. The total number of such events (25.6) is reduced to 18.7 after the additional trigger cuts have been implemented. Note that this is not necessarily in conflict with the observed number (9) quoted above since, according to the UA1 analysis [4], a substantial fraction of τ monojets would fail the quite restrictive τ -likelihood criteria. In fact if we assume that the efficiency for selecting τ events in this way is about 50% - as suggested by the UA1 analysis - then we predict a total of 9.4 such τ events, of which 4.6 have $E_T^{\text{miss}} > 30$ GeV and 1.1 have $E_T^{\text{miss}} > 38$ GeV. The corresponding numbers of observed events are 9, 5, and 1 respectively [4].

5. DISCUSSION

The main result of the present analysis is that the UA1 1984 monojet events appear to be well described by SM processes from W and Z production and decay. Our calculation is based on a perturbative analysis of $W, Z + 0, 1, 2$ jet production. The predicted total rate of observed W and Z production agrees with the data. The addition of an empirical underlying event scalar transverse energy gives a good fit to the distribution in scalar transverse energy accompanying W bosons.

There are various uncertainties in the theoretical calculation of the monojet rate, especially the rate just above the threshold cuts. The two most important are - (a) the uncertainty in the underlying event average transverse energy, which has the effect of moving the cut on a steeply falling distribution, and (b) the uncertainty in imposing trigger selection cuts on the partonic final state.

Our implementation of the latter is surely only a crude approximation. Nevertheless, our "best guess" scenario gives a prediction for the total number of monojets in good agreement with the observed number.

More importantly, we have shown how the predictions for $E_T^{\text{miss}} > 30$ GeV are much more stable. We calculate 13.6 events - with an overall error of order ± 1 - and 12 events are observed. (Note that our result does not include the presumably small contribution from heavy quark production and decay. An analysis of these extra contributions is in progress.) We believe therefore that it is those monojets with $E_T^{\text{miss}} > 30$ GeV which, with more data, will provide the best test for the presence of new physics contributions.

Finally in Fig.5 we compare the cross-section for monojets with $E_T^{\text{miss}} > 30$ GeV at 630 GeV and 1.6 TeV. For simplicity we have used exactly the same cuts at the two energies although of course in practice the analysis will be different for different detectors. The much larger cross-section at the higher energy will allow a more a precise test of the standard model.

REFERENCES

- [1] See for example L.J. Hall, R.L.Jaffe and J.L.Rosner, Physics Reports 125 (1985) 103.
- [2] Previous estimates of standard model backgrounds include: P. Aurenche and R. Kinnunen, Annecy preprint LAPP-TH-108 (1984); A. Ballestrero and G. Passarino, Phys. Letters 148B (1984) 373, 378; J.R. Cudell, F. Halzen and K. Hikasa, Wisconsin-Madison preprint MAD/PH/237 (1985); E.W.N. Glover and A.D. Martin, Durham preprint DTP/85/10 (1985); K. Hikasa, Wisconsin-Madison preprint MAD/PH/261 (1985).
- [3] S.D. Ellis, R. Kleiss and W.J. Stirling, Phys. Letters 158B (1985) 341.
- [4] UA1 Collaboration, presented by J. Rohlf at the American Physical Society Division of Particles and Fields, Eugene, Oregon, August 1985; CERN preprint CERN-EP/85-160 (1985).
- [5] S.D. Ellis, R. Kleiss and W.J. Stirling, Phys. Letters 154B (1985) 435.
- [6] P. Aurenche and R. Kinnunen, reference 2.
- [7] E.W.N. Glover and A.D. Martin, reference 2.
- [8] F.J. Gilman and S.H. Rhie, Phys. Rev. D31 (1985) 1066.
- [9] UA1 Collaboration, presented by M. Mohammadi at the Conference on Collider Physics at Ultra-High Energies, Aspen, Colorado, January 1985; CERN preprint CERN-EP/85-52 (1985).
- [10] UA1 Collaboration, presented by J. Rohlf, Proceedings of the XXII International Conference on High Energy Physics, Leipzig, July 1984, Vol.II p.16.

- [11] UA1 Collaboration, presented by S. Geer at the Theoretical Advanced Study Institute in Elementary Particle Physics, Yale University, New Haven, June 1985; CERN preprint CERN-EP/85-163 (1985).
- [12] S. Geer and W.J. Stirling, Phys. Letters 152B (1985) 373.
- [13] R.M. Barnett, H.E. Haber and G.L.Kane, Lawrence Berkeley Laboratory preprint LBL-20102 (1985).

TABLE 1(a) : $W \rightarrow e + \nu$ events from the combined 1983 + 1984 data sample

	data [11]	mode 1	mode 2
total	172	177.6	171.0
$p_T(W) > 20 \text{ GeV}/c$	10	9.9	10.9

TABLE 1(b) : $Z \rightarrow e + e$ events from the combined 1983 + 1984 data sample

	data [11]	mode 1	mode 2
total	18	18.9	19.8
$p_T(Z) > 20 \text{ GeV}/c$	2	0.9	1.1

TABLE 2 : Number of monojet events at $\sqrt{s} = 630$ GeV from the various roads listed in Eqn.1. The results correspond to mode 1 with $\langle E_T^{ue} \rangle = 30$ GeV. The contributions labelled $\star$ are from "lost" electrons and muons from W (and $W \rightarrow \tau$) decay. The number of events with missing transverse energy greater than 30 GeV and 38 GeV are shown separately.

	4σ	> 30 GeV	> 38 GeV
$\tau + 0$	22.6	8.2	1.9
$\tau + 1$	5.1	2.4	0.9
$\tau + 2$	0.6	0.4	0.2
$\nu\nu + 1$	3.8	1.4	0.7
$\nu\nu + 2$	0.9	0.4	0.2
$\star + 1$	1.1	0.7	0.4
$\star + 2$	0.2	0.2	0.1
(total τ -jets	25.6	9.6	2.2)
total	34.3	13.6	4.4
DATA	23	12	5

TABLE 3 : Number of monojet events at $\sqrt{s} = 630$ GeV calculated using the different modes and different average underlying event transverse energies. The number of events with missing transverse energy greater than 30 GeV and 38 GeV are shown separately.

	4σ	> 30 GeV	> 38 GeV
mode 1	34.3	13.6	4.4
mode 2	34.2	13.7	4.6
mode 1 ($\langle E_T^{ue} \rangle = 20$ GeV)	39.3	13.4	4.2
mode 1 ($\langle E_T^{ue} \rangle = 40$ GeV)	30.1	13.6	4.5
DATA	23	12	5

TABLE 4 : Number of monojet events at $\sqrt{s} = 630$ GeV (calculated using mode 1 and $\langle E_T^{ue} \rangle = 30$ GeV) with and without the trigger cuts as described in the text. The number of events with missing transverse energy greater than 30 GeV and 38 GeV are shown separately.

	σ	> 30 GeV	> 38 GeV
without trigger cuts	34.3	13.6	4.4
with trigger cuts	24.9	13.0	4.3
DATA	23	12	5

FIGURE CAPTIONS

- Fig.1 The fragmentation function for τ decay, $\tau \rightarrow X + \text{neutrino(s)}$, showing the contribution of the various components X as described in the text. Here z is the fraction of the τ 's energy carried by the decay products.
- Fig.2 Scalar transverse energy accompanying W bosons. The data are the 1983 UA1 W sample [10]. The curves are the theoretical predictions using three different values for the average underlying event transverse energy.
- Fig.3 The distribution in missing transverse energy of the 1984 UA1 monojet sample [4]. The curves are the SM predictions with (dashed line) and without (solid line) trigger cuts as discussed in the text.
- Fig.4 As for Fig.3, but showing the contribution from the τ -jet and non τ -jet components, as defined in the text.
- Fig.5 Monojet cross-sections for $E_T^{\text{miss}} > 30 \text{ GeV}$ at $\sqrt{s} = 0.63 \text{ TeV}$ and 1.6 TeV .

Fig.1

Fig.2

Fig.3

Fig.4

Fig.5