06 JAN 1986

Numerical study of the lattice meson form factor

R.M. Woloshyn

TRIUMF

4004 Westbrook Mall, Vancouver, B.C., Canada V6T 2A3

and

A.M. Kobos

SuperComputing Services, The University of Calgary,
Calgary, Alberta, Canada T2N 1N4

Abstract

The electric form factor of the pseudo-Goldstone meson (the generic pion) is calculated in quenched lattice quantum chromodynamics with SU(2) color. Charge radii are calculated for different values of the bare quark mass. The results are in agreement with the physically reasonable expectation that heavier quarks have distributions of smaller radius.

(Submitted to Physical Review D)

CERN LIBRARIES, GENEVA

CM-P00067419

I. INTRODUCTION

Recently there has been considerable interest in determining lattice hadron sizes and internal structure.¹⁻⁴ The study of electromagnetic properties can be very useful for this purpose. In Ref. 1 the feasibility of a lattice calculation of the meson electric form factor was demonstrated. In Ref. 5 a detailed derivation of the lattice three-point function was given and it was shown how the form factor could be extracted from a combination of two and three-point functions. In this paper we use the results of Ref. 5 in a numerical survey of the pseudo-scalar meson electric form factor as a function of quark mass.

The staggered scheme^{6,7} for putting fermions on the lattice is used. The interpretation of the staggered fermion degrees of freedom is usually made in terms of flavored quark fields defined on hypercubes in the lattice.⁸⁻¹⁰ The two-point function, needed for mass calculations, can be cast in a form that is constructed using only local combinations of staggered fermion fields.⁸⁻¹¹ In Ref. 5 it was shown that the electric form factor can be extracted from a three-point function which involves only local staggered fermion field bilinear operators as interpolating fields. The relevant formulae are reviewed in Section II.

Section III contains details of the numerical work and the results are given in Section IV. The calculations are done in a model with SU(2) color (at $\beta=2.3$) for five values of the bare quark mass from $ma=0.025$ to $ma=0.4$. The meson masses we obtain agree very well with previous calculations. The electric form factor is calculated for a 0⁻ meson (pseudo-Goldstone boson) with an equal mass quark-antiquark pair. The results show clearly that the charge radius decreases as the quark mass increases.

Pseudoscalar mesons with unequal mass quarks and antiquarks are also considered. Allowing the charge density operator to act either on the light or on the heavy quark gives information on their relative distribution. We obtain the physically reasonable result that, when the quarks have very different masses, the heavy quark has a much smaller charge radius than the light quark. This provides good evidence that our measurement is indeed probing the meson structure.

Conclusions are given in Section V.

II. FORMALISM

In this paper we consider a charged pseudo-Goldstone meson (a generic pion) associated with the remnant continuous axial global symmetry present in the staggered scheme of lattice fermions.^{7,8} The usual flavor structure associated with the staggered fermions is not useful for constructing charged states since an "electric charge" defined within these flavors is not conserved.¹² We therefore introduce two sets of flavors, labelled by u and d , with charges q^u and q^d . The action takes the form (suppressing color indices)

$$S_F(u) = \bar{\chi} M \chi, \quad (1a)$$

$$= \frac{1}{2} \sum_{x,u,f=\{u,d\}} a_\mu(x) [\bar{\chi}^f(x) U_\mu(x) \chi^f(x+a_\mu) - \bar{\chi}^f(x+a_\mu) U_\mu^\dagger(x) \chi^f(x)] \\ + m_a \sum_{x,f} \bar{\chi}^f(x) \chi^f(x), \quad (1b)$$

$$\text{where } a_\mu(x) = (-1)^{\xi_\mu}, \quad \xi_\mu = \sum_{\nu < \mu} x_\nu$$

A conserved vector current can be derived using the Noether procedure.^{7,8} The result is (for each species, $f=u,d$)

$$j_\mu^f(x) = -\frac{1}{2} a_\mu(x) [\bar{\chi}^f(x) U_\mu(x) \chi^f(x+a_\mu) + \bar{\chi}^f(x+a_\mu) U_\mu^\dagger(x) \chi^f(x)]. \quad (2)$$

The usual way to expose the particle content of the staggered fermion theory is to construct non-local quark fields defined on hypercubes in the lattice. This is discussed in detail in many papers.⁸⁻¹¹ The end result is that the two-point function (which is essentially the correlation function of the local interpolating field

$$a_q(x)(-1)^t \bar{\chi}^d(\vec{x},t) \chi^u(\vec{x},t)$$

$$G_0(\vec{p};t) \equiv \frac{1}{N_c} \sum_{\vec{x},c} e^{-i\vec{p} \cdot \vec{x}} \text{tr} M^{-1}(\vec{x},t;0) [M^{-1}(\vec{x},t;0)]^\dagger, \quad (3)$$

where trace implies a sum over appropriate color and flavor indices, describes the propagator of a 0^- pseudo-Goldstone meson. The sum c is over Monte Carlo lattice gauge field configurations (N_c in number). We also use the correlation function in the $J=1$ channel

$$G_1(\vec{p};t) \equiv \frac{1}{N_c} \sum_{\vec{x},c} e^{-i\vec{p} \cdot \vec{x}} ((-1)^{x_1+(-1)^{x_2}+(-1)^{x_3}} \text{Tr} M^{-1}(\vec{x},t;0) \\ \cdot [M^{-1}(\vec{x},t;0)]^\dagger). \quad (4)$$

This describes the propagation of vector and axial-vector mesons.

In Ref. 5 it was shown that the matrix element of the charge density operator between pion states could be extracted from the lattice three-point function

$$A(\vec{p}, \vec{q}; t_2, t_1) = \sum_{\vec{x}} e^{-i\vec{p} \cdot \vec{x}} a_q(x_2) (-1)^{x_2} \langle \bar{\chi}^d(\vec{x}_2, t_2) \chi^u(\vec{x}_2, t_2) \\ \cdot \sum e^{i\vec{q} \cdot \vec{x}} 1_P(\vec{x}, t_1) \bar{\chi}^u(0) \chi^d(0) \rangle, \quad (5)$$

where $\rho(\vec{x}, t) = \sum_f q_f i_f(\vec{x}, t)$. Notice that this three-point function just involves local interpolating fields.

The most convenient way to determine the electric form factor from the Monte Carlo data is to take the combination

$$\mathcal{R} = \left\{ \frac{A(0, \vec{q}; t_2, t_1) A(\vec{q}, \vec{q}; t_2, t_1)}{G_0(0; t_2) G_0(\vec{q}, t_2)} \right\}^{1/2} \frac{(E_q + M_0)}{0 < t_1 < t_2} \frac{F(q)}{2\sqrt{E_q M_0}}, \quad (6)$$

where the electric form factor¹³ is defined by

$$\langle \pi^+(p) | \rho(0) | \pi^+(p') \rangle = (E_p + E_{p'}) F(q). \quad (7)$$

III. COMPUTATIONAL DETAILS

Numerical calculations were done in a model for QCD using only SU(2) color. The lattice size was $10^2 \cdot 20 \cdot 16$ with the current carrying momentum in the 3-direction. Twenty gauge field configurations, prepared with a heat bath Monte Carlo¹⁴ in quenched approximation, were used. The gauge fields were constructed using the Wilson gauge field action¹⁵ at $\beta = 2.3$ on a $10^3 \cdot 16$ lattice and then doubled in the 3-direction.

Fermion propagators were calculated using the conjugate gradient algorithm.¹⁶ Iterations were carried out until the maximum change in the correlation function $G_0(0; t)$ at any time point was less than 0.05% in four iterations. The absolute value squared of the residual vectors was less than 10^{-11} to 10^{-10} .

Anti-periodic boundary were used on fermion fields in the spatial directions. However, the fermion coupling was put equal to zero across the time boundary of the lattice. The advantage of this choice is that correlation functions have a simple exponential fall off at large time separation. Furthermore, charge conservation takes a very simple form.

The disadvantage of our boundary condition is that non-vacuum contributions can contaminate matrix elements when operators are placed near the time edges.¹⁷ Fortunately the combination $\mathcal{R}$ of Eq. (6) removes all the time dependence and the factors that refer to the time boundary. We expect the form factor extracted using Eq. (6) to be free of time boundary effects.

The three-point function was calculated as the derivative of a two-point function with the charge operator acting as a source.^{5,18} A check of this procedure was made at $q=0$ where the three and two-point functions are related.¹⁹ It indicates that, at the lightest quark mass, our derivative is accurate to within 2 to 3 percent and becomes better when the quark mass is increased.

The standard statistical error was calculated at each time value of the two-point functions using the whole sample of twenty configurations. Meson masses were determined by fitting $G(0; t)$ with ae^{-bt} for times three through ten steps removed from the hadron creation operator. The uncertainty in the mass was determined from the error matrix of the least squares fit. Statistical errors in the combination of three and two-point functions $\mathcal{R}$ were calculated including covariances between the different factors.²⁰ Again all twenty configurations were included in a single sum. The form factor was obtained from an average (weighted by statistical errors) of $\mathcal{R}$ for times five through nine steps from the initial hadron creation operator. The charge density operator is placed at time step four, i.e., it acts between time steps four and five.

IV. RESULTS

The pseudoscalar meson masses M_0 for equal mass quark and antiquark is shown in Fig. 1 as a function of bare quark mass m_a . The mass in t.e

$J=1$ channel is also shown. This was obtained by fitting the two-point function $G_1(0,t)$ with a single exponential, essentially ignoring a small admixture of 1^+ state. Although this is an approximation it should be adequate for our purposes. Where we can make a direct comparison (at $m_a=0.1, 0.2$) our pseudoscalar and vector masses agree very well with those of Billoire et al.¹¹

In Fig. 2 we plot $M_0^2 a^2$ versus the bare quark mass m_a . If PCAC holds this relation should be linear with $M_0^2 a^2$ extrapolating to zero at zero quark mass.²¹ Indeed, we find that these conditions are satisfied at the smaller values of quark mass. The line is not a fit but is a guide for the eye.

Figure 3 shows a typical result for the combination of three and two-point functions R_{of} Eq. (6) versus the time t_2 at which the hadron annihilation operator acts. The charge density operator acts at time $t_1=4$. When $t_2 < t_1$, R is zero to within a few percent. This reflects current conservation and the fact that with our time boundary conditions there is a net charge present only for times between the action of the meson creation ($t=0$) and annihilation operators ($t=t_2$).

The electric form factor of the pseudoscalar meson was calculated at two values of momentum transfer, for $q=\pi/10$, the lowest momentum value on our lattice, and for $q=\pi/5$. The results plotted as a function of Minkowskian four-momentum transfer squared (in lattice units) are shown in Fig. 4. We see clearly that large quark masses lead to a more slowly decreasing form factor. This can be seen again in Fig. 5a where the form factor values at $q=\pi/10$ are plotted as a function of quark mass. To extract a root mean square radius R_{rms} the form factor values at $q=\pi/10$

are first parametrized using $F(Q^2) = (1+Q^2 a^2/\lambda^2)^{-1}$ then the relation $R_{rms}^2 = -6 dF/dQ^2$ is used on the monopole fit. The resulting charge radii are shown in Fig. 5b. The parameters λ^2 themselves show an interesting behavior. They are plotted in Fig. 6 as a function of quark mass. Also shown are the values of $\lambda^2/M^2 a^2$, which within errors are constant, although not equal to one. The idea that the low momentum transfer fall off of form factors is intimately related to vector mesons is an old one²² and it is amusing that, to some extent, we observe this connection in our calculations.

The above calculations were done keeping quarks with different flavors degenerate in mass. It is also interesting to consider the situation where one quark has a mass very different from the other. We have done a calculation for pseudoscalar mesons in which one of the quark masses is fixed at $m_a=0.4$ and the other is allowed to decrease to $m_a=0.025$. Form factors are calculated with the charge density operator acting either on the light or on the heavy quark. The results at one unit of momentum transfer ($q=\pi/10$) are shown in Fig. 7. The form factor of the lighter quark essentially tracks the form factor of a meson with a light quark-antiquark pair. The form factor of the (fixed-mass) heavy quark approaches unity as the light quark mass decreases. Parametrizing these form factors with monopoles and extracting the root mean square radius yields the results of Fig. 8. We observe the physically reasonable result that in a meson with unequal mass quarks the distribution of the heavy quark has a smaller radius than the distribution of the light quark. This is good evidence that our form factor is really reflecting the meson structure and is not, for example, an artifact of the finite lattice.

V. CONCLUSION

In this paper we have done a survey of the 0^- lattice meson (pseudo-Goldstone boson) electric form factor as a function of quark mass. The staggered scheme for lattice fermions was used and that calculations were done in a model for quenched lattice QCD with $SU(2)$ color.

The results clearly show that the meson charge radius decreases as the quark mass increases. Parametrizing the form factor with a monopole $(1+Q^2a^2/\lambda^2)^{-1}$ we find that the parameters λ^2 are proportional to the vector meson mass squared as suggested by vector dominance ideas. The constant of proportionality is not however equal to one.

The form factor for pseudoscalar mesons with an unequal mass quark-antiquark pair has also been calculated. Allowing the charge density operator to act either on the light or on the heavy quark shows that the distribution of the heavy quark has a smaller radius than that of the light quark. This result gives us considerable confidence that the form factor we calculate is reflecting the meson structure correctly.

The extension of the form factor calculations to $SU(3)$ color and the baryon sector would be very interesting and, ultimately, could provide one of the best quantitative tests of QCD.

ACKNOWLEDGEMENT

We thank SuperComputing Services at the University of Calgary for providing time on the Cyber 205 computer. This work is supported in part by the Natural Sciences and Engineering Research Council of Canada.

REFERENCES

- ¹W. Wilcox and R.M. Woloshyn, Phys. Rev. Lett. 54, 2653 (1985).
- ²B. Vellkson and D. Weingarten, Nucl. Phys., B249, 433 (1985).
- ³O. Martin, K. Moriarty and S. Samuel, "The Quark Model on the Lattice," Columbia University preprint.
- ⁴K. Barad, M. Oglivie and C. Rebbi, Phys. Lett., 143B, 222 (1984).
- ⁵W. Wilcox and R.M. Woloshyn, TRIUMF report no. TRI-PP-85-32.
- ⁶J. Kogut and L. Susskind, Phys. Rev., D11, 395 (1975);
L. Susskind, Phys. Rev., D16, 3031 (1977).
- ⁷N. Kawamoto and J. Smit, Nucl. Phys., B192, 100 (1981).
- ⁸H. Kluberg-Stern, A. Morel, O. Napoly and B. Petersson, Nucl. Phys., B220, 447 (1983).
- ⁹M.F.L. Golterman, "Staggered Mesons," University of Amsterdam report ITFA-84-18.
- ¹⁰J. Kogut, M. Stone, H.W. Wylid, S.H. Shenker, J. Shigemitsu and D.K. Sinclair, Nucl. Phys. B225, 326 (1983).
- ¹¹A. Billoire, R. Lacaze, E. Marinari and A. Morel, Nucl. Phys., B251, 581 (1985).
- ¹²See, for example, the definition of electric charge in Ref. 8.
- ¹³See C. Itzykson and J. Zuber, Quantum Field Theory, (McGraw-Hill, New York, 1980), p.160.
- ¹⁴M. Creutz, Phys. Rev., D21, 2308 (1980).
- ¹⁵K.G. Wilson, Phys. Rev., D10, 2445 (1974).
- ¹⁶F.S. Beckman, in Mathematical Methods for Digital Computers, edited by A. Ralstan and H.S. Wilf (Wiley, New York, 1960), p.62.
- ¹⁷C. Bernard, T. Draper, K. Olynyk and M. Rushton, Nucl. Phys., B220, 508 (1983).

- 18S. Gottlieb, P.B. Mackenzie, H.B. Thacker and D. Weingarten, Phys. Lett., 134B, 346 (1984);
- C. Bernard, in Gauge Theory on a Lattice: 1984, edited by C.Zachos, W. Celmaster, E. Kovacs and D. Silvers (National Technical Information Service, Springfield, VA, 1984), p.85.
- 19The two and three-point functions satisfy a sum rule $A(0,0;t_2,t_1)=G(0,t_2)$ for $t_2 > t_1$. This simple form is a consequence of our non-periodic boundary condition in time.
- 20This is necessary since the fluctuations in the two and three-point functions are correlated. See P.R. Bevington, Data Reduction and Error Analysis For The Physical Sciences, (McGraw-Hill, New York, 1969), Chap. 4 for a discussion of error propagation.
- 21See Ref. 8 and F. Gliozzi, Nucl. Phys., B204, 419 (1982).
- 22W.R. Frazer and J.R. Fulco, Phys. Rev., 117, 1609 (1960). See also W.C. Holladay, Phys. Rev., 101, 1198 (1956).

FIGURE CAPTIONS

1. Pseudoscalar (●) and vector (■) meson masses versus the bare quark mass.
2. Square of the pseudoscalar meson mass versus the bare quark mass.
3. The combination R of three and two-point functions (Eq. (6)) versus the time coordinate of the meson annihilation operator. The charge density operator acts between time step 4 and 5.
4. The pseudoscalar meson electric form factor versus Minkowskian four-momentum transfer squared $q^2 a^2$ for different values of the quark mass from $m_a=0.025$ to $m_a=0.4$.
5. (a) The pseudoscalar meson electric form factor at $q=\pi/10$ versus the quark mass.
(b) The meson root mean square radius R_{rms} versus the bare quark mass.
6. The λ^2 parameter from a monopole parametrization of the electric form factor versus the quark. The quantity λ^2 is shown both in lattice units (o) and divided by the vector meson mass squared (●).
7. The electric form factor for the light (●) and heavy (■) quark at $q=\pi/10$ of a pseudoscalar meson with an unequal mass quark-antiquark pair versus the light quark mass. The heavy quark mass is fixed at $m_a=0.4$. For comparison the meson form factor in a meson with an equal mass quark-antiquark pair (o) is also shown.
8. The root mean square radius R_{rms} for the light (●) and heavy (■) quark in a pseudoscalar meson with an unequal mass quark-antiquark pair versus the light quark mass. The heavy quark mass is fixed at $m_a=0.4$. For comparison the root mean square radius of a meson with an equal mass quark-antiquark pair (o) is also shown.

Fig. 1

Fig. 2

Fig. 3

Fig. 4

Fig. 5

Fig. 6

Fig. 7

Fig. 8