DESY 85-132
CERN-TH-4212/85
December 1985

TESTING THE $W\gamma$ COUPLING OF THE STANDARD MODEL AT $p\bar{p}$ -COLLIDERS

by

J. Cortés

CERN, Theory Division, Geneva

K. Hagiwara

Deutsches Elektronen-Synchrotron DESY, Hamburg

F. Herzog

Institut f. Theoretische Physik, Bern

ISSN 0418-9833

NOTKESTRASSE 85 · 2 HAMBURG 52

DESY behält sich alle Rechte für den Fall der Schutzrechtserteilung und für die wirtschaftliche Verwertung der in diesem Bericht enthaltenen Informationen vor.

DESY reserves all rights for commercial use of information included in this report, especially in case of filing application for or grant of patents.

To be sure that your preprints are promptly included in the
HIGH ENERGY PHYSICS INDEX ,
send them to the following address (if possible by air mail) :

DESY
Bibliothek
Notkestrasse 85
2 Hamburg 52
Germany

The κ -sensitivity in the production process $d + \bar{u} \rightarrow W^- + \gamma$ and that in the decay process $W^- \rightarrow \gamma + \ell^- + \bar{\nu}_\ell$, is known to be spectacular⁷⁾ due to the occurrence of spin independent zeros⁸⁾ in the tree amplitudes: If the photon coupling is that of a renormalizable gauge theory, implying $\kappa = 1$ for the W -boson, the amplitudes $M(d\bar{u} \rightarrow W^- \gamma)$ and $M(W^- \rightarrow \ell \ell^- \bar{\nu}_\ell)$, respectively have a zero in the physical region. This κ -sensitivity leads to very distinctive features for various distributions. In $W\gamma$ production this amplitude zero occurs at the partonic center-of-mass (c.m.) scattering angle $\cos\theta_{d\gamma}^* = \frac{1}{3}$:

$$\mathcal{M}(d\bar{u} \rightarrow W^- \gamma) \Big|_{\kappa=1} = 0 \quad \text{at} \quad \cos\theta_{d\gamma}^* = \frac{1}{3} . \quad (1.4)$$

In the W radiative decay case the amplitude zero shows up in the W restframe when the photon is anticollinear to the charged lepton:

$$\mathcal{M}(W^- \rightarrow \ell \ell^- \bar{\nu}_\ell) \Big|_{\kappa=1} = 0 \quad \text{at} \quad \cos\theta_{\ell\ell}^* = -1 . \quad (1.5)$$

For any values of κ other than the gauge theory value, $\kappa = 1$, these amplitude zeros disappear.

In this paper we make an extensive study of the "production" process (1.2) as well as of the "decay" process (1.3) in search of κ -sensitive observables: Our first concern will be to show feasible methods to actually reconstruct the above mentioned angular distributions from the observable final state, consisting typically of a high transverse momentum (p_T) isolated photon and a high p_T charged lepton with large missing p_T ($\cancel{p}_T$) from a neutrino (ν). This is a nontrivial task since we do not know ν longitudinal momentum in general. Furthermore we give an estimate of backgrounds from $W + \text{jet}$ events where jet fakes a photon, a discussion of W -finite width effects, and that of higher order QCD corrections.

The paper is organized as follows: In section II) we present differential cross-sections for the process $p\bar{p} \rightarrow \gamma + \ell^- + \bar{\nu}_\ell + \text{anything}$ in a minimally extended Standard Model with an arbitrary κ -value⁴⁾ for the W -boson. By using the narrow width approximation for the W -propagators, the partonic subprocess can be split into the "production" process (1.2) and the "decay" process (1.3). Because of the different kinematics we shall discuss these two processes in two separate subsections:

In subsection IIa) we present the differential cross-section for $p\bar{p} \rightarrow W^- \gamma$ followed by the decay $W^- \rightarrow \ell^- \bar{\nu}_\ell$. We propose a variable $\cos\theta_\ell^*$ to be used in plotting the data: This variable is constructed from the measurable photon and charged lepton momenta in such a way that it reduces to one of the two possible $W\gamma$ c.m. scattering angles. In subsection IIb) we show the differential cross-section for $p\bar{p} \rightarrow W^- X$ followed by the radiative decay $W^- \rightarrow \ell \ell^- \bar{\nu}_\ell$. Here we suggest to use a variable $\cos\theta_{\ell\ell}^*$ to plot data: This quantity is again constructed from the measurable photon and charged lepton momenta in such a way that it reduces to one of the possible opening angles $\cos\theta_{\ell\ell}^*$ allowed by the ambiguity in the neutrino longitudinal momentum. Numerical estimates at $p\bar{p}$ colliders are made for the $\cos\theta_\ell^*$ -distribution as well as for the $\cos\theta_{\ell\ell}^*$ -distribution and we find the κ -sensitivity in both cases to be significant. We further show that a cut in the cluster transverse mass $m_T(\ell\ell; \bar{\nu}_\ell)$ ⁹⁾ can separate "production" and "decay" processes in a simple and very effective way.

In section III we discuss the influence of the finite W -width and address the difficult question of higher order QCD correction effects. The effects of the finite W -width are found to be small in both cases. The higher order QCD corrections are expected to be small in the W radiative decay signals. The corrections for the $W\gamma$ production process can, however, be quite large and we propose a modification of the variable $\cos\theta_\ell^*$ to include transverse motion of the $W\gamma$ system. Although we argue that this modification is necessary to control higher order corrections, only an actual evaluation of the corrections would tell us if it is sufficient or not.

In section IV we discuss possible background problems to the proposed angular distributions. The most severe background comes from associated production of W and a jet¹⁰⁾ where the jet fakes a single isolated photon. This background is found to affect the analysis for the $\cos\theta_\ell^*$ -distribution significantly while it causes little problem in the radiative decay analysis.

In section V we summarize our findings and draw some conclusions. In the Appendix we present the explicit tree-level cross-section for the subprocess $d\bar{u} \rightarrow \ell \ell^- \bar{\nu}_\ell$ without making a narrow-width approximation for intermediate W propagators.

2a) The process $\bar{p}p \rightarrow W^+ \gamma X; W^- \rightarrow \ell^- \bar{\nu}_\ell$

The parton cross section for the "production" process (2.3) is expressed

$$d\hat{\sigma}_{q\bar{q}} = \frac{1}{2\hat{s}} \bar{\Sigma} \left| M(q\bar{q}' \rightarrow W^+ \gamma; W^- \rightarrow \ell^- \bar{\nu}_\ell) \right|^2 d\Phi \quad (2.7)$$

as

Here $\hat{s} = (\hat{p}_q + \hat{p}_{\bar{q}})^2$ and $d\Phi$ denotes the three-body phase space

$$d\Phi = (2\pi)^5 \delta[(\hat{p}_q + \hat{p}_{\bar{q}} - \hat{p}_\ell - \hat{p}_{\bar{\nu}})^2] \frac{d^3\hat{p}_\ell}{2E_\ell} \frac{d^3\hat{p}_{\bar{\nu}}}{2E_{\bar{\nu}}} \quad (2.8)$$

The spin and color sum-averaged matrix element squared can be extracted from the complete formula given in the appendix; for massless quarks it reads (see eq. (A13)):

$$\begin{aligned} \bar{\Sigma} |M(q\bar{q}' \rightarrow W^+ \gamma; W^- \rightarrow \ell^- \bar{\nu}_\ell)|^2 \\ = \frac{128\pi^6 \alpha^2}{\sin^2 \theta_W} \mathcal{B}(W^- \rightarrow \ell^- \bar{\nu}_\ell) |U_{q1}|^2 \delta[(\hat{p}_q + \hat{p}_{\bar{q}} - \hat{p}_\ell)^2 - m_W^2] \\ \cdot \left\{ (Q_1 + \frac{\hat{s}}{\hat{s} + \hat{q}})^2 \frac{1}{\hat{s}} \frac{1}{\hat{q}} \left[(2\hat{p}_{\bar{q}} \cdot \hat{p}_\ell)^2 + (2\hat{p}_1 \cdot \hat{p}_\ell)^2 \right] + O(1-\kappa) \right\} \end{aligned} \quad (2.9)$$

θ_W denotes the Weinberg angle, $\mathcal{B}(W^- \rightarrow \ell^- \bar{\nu}_\ell)$ the branching ratio, α the fine structure constant, U_{q1} the weak mixing matrix, Q_q the quark electric charge, $\hat{t} = (\hat{p}_q - \hat{p}_\ell)^2$ and $\hat{q} = (\hat{p}_{\bar{q}} - \hat{p}_{\bar{\nu}})^2$. The factorization of the charge dependent factor is manifest in (2.9) for the $\kappa = 1$ case which leads to the amplitude zero at $\cos \theta_{q\ell}^* = 1 + 2Q_q$ in the partonic c.m. frame. In the following we neglect both the Cabibbo rotation and the contribution from strange and heavier quark sea. The only surviving subprocess is there $\bar{u}u \rightarrow W^+ \gamma; W^- \rightarrow \ell^- \bar{\nu}_\ell$ with $|U_{ud}| = 1$. We set $m_W = 83 \text{ GeV}$, $G_F = (\pi\alpha/\sqrt{2} \sin^2 \theta_W m_W^2) = 1.17 \times 10^{-5} \text{ GeV}^{-2}$ and $\mathcal{B}(W^- \rightarrow \ell^- \bar{\nu}_\ell) = 1/12$ for definiteness.

The cross section (2.9) depends quadratically on κ , as can be seen explicitly from (A8). Therefore we will always plot distributions for three representative κ -values as it is enough to reconstruct distributions for arbitrary κ . We choose the κ -values $\kappa = 0, \pm 1$ for the "production" process. The cross-section for an arbitrary κ -value then reads

$$d\sigma(\kappa) = \frac{1}{2} \kappa(1+\kappa) d\sigma(1) + (1-\kappa)(1+\kappa) d\sigma(0) - \frac{1}{2} \kappa(1-\kappa) d\sigma(-1). \quad (2.10)$$

After integrating out the azimuthal angle of the photon momentum about the beam axis we have five independent kinematical variables in terms of which the phase space factor in (2.8) can be written as

$$\frac{d^3\hat{p}_\ell}{2E_\ell} \frac{d^3\hat{p}_{\bar{\nu}}}{2E_{\bar{\nu}}} = \frac{\pi}{8} d\hat{p}_{qT}^2 d\hat{y}_\ell d\hat{\phi}_\ell^2 d\hat{p}_{\bar{\nu}T}^2 d\hat{y}_{\bar{\nu}} \quad (2.11)$$

Here $\hat{p}_{qT}$ and $\hat{p}_{\bar{\nu}T}$ denote the transverse momentum of the photon and the charged lepton with respect to the beam axis, respectively, $\hat{y}_\ell$ and $\hat{y}_{\bar{\nu}}$ are their rapidity in the proton momentum direction, and $\hat{\phi}_\ell$ the azimuthal angle of $\vec{\hat{p}}_{qT}$ measured from $\vec{\hat{p}}_{\bar{\nu}T}$.

Shown in Figs. 2a and 2b are the transverse momentum spectrum of the photon and the charged lepton, respectively, expected at the Tevatron collider energy $\sqrt{s} = 2 \text{ TeV}$. The solid line represents $\kappa = 1$, the dashed one $\kappa = 0$ and the dashed-dotted one $\kappa = -1$. The photon distribution falls strongly with rising $\hat{p}_{qT}$; this tells us the importance of the photon identification at relatively smaller $\hat{p}_T$ values. The shape of the spectrum is mildly sensitive to κ ; the steepest fall-off is expected for the gauge theory value $\kappa = 1$. On the contrary, the charged lepton $\hat{p}_T$ distribution (Fig. 2b) with the cut $\hat{p}_{qT} > 10 \text{ GeV}$ has a peak around $\hat{p}_{\bar{\nu}T} \approx m_W/2$ which is a remnant of the well-known Jacobian peak¹⁷⁾.

In Figs. 3a and 3b we show the rapidity distribution for the photon and the charged lepton, respectively; in both figures we have imposed the cuts $\hat{p}_{qT} > 10 \text{ GeV}$ and $\hat{p}_{\bar{\nu}T} > 10 \text{ GeV}$. The photon spectrum (Fig. 3a) is asymmetric and has its maximum in the region of negative $\hat{y}_\ell$. This reflects the fact that the photon couples more strongly to the $\bar{u}$ quark than to the d quark and hence tends to be radiated in the $\bar{u}$, or $\bar{p}$ direction. These figures (Figs. 2 and 3) show that the overall production rate grows as $|\kappa-1|$ grows. Hence even the total rate measures κ , but with poor sensitivity. In view of the present theoretical uncertainties¹⁸⁾ in the precise estimate of the QCD radiative correction to the production rate, it would be desirable to look for quantities much more sensitive to the value of κ .

For the computations of the rest of this subsection we will always apply the following cuts:

$$p_{\ell T}, p_{\ell T} > 10 \text{ GeV}; |y_{\ell}|, |y_{\bar{\ell}}| < 3 \quad (2.12)$$

These cuts are imposed to account for an actual experimental situation where one is looking for an isolated high p_T photon and a high p_T charged lepton with typically large p_T .

One of the most spectacular prediction of the electroweak gauge theory is the presence of the polarization and momentum-independent angular zero in the constituent scattering amplitude for $q\bar{q} \rightarrow W\gamma^{7,8}$. The angular zero in the spin-averaged cross section disappears for non-gauge theories because only in the gauge theory case do all polarization amplitudes have a zero at the same phase space point¹⁹. This striking signal is almost smeared away in the collider c.m. frame as exemplified in the previous figures. We need to find differential cross sections which are sensitive to k but relatively insensitive to the shape of the constituent distribution which is responsible for the smearing.

Since we shall discuss a dip in the cross section we should anticipate a brief discussion on the possible background problem: We are looking for events showing a high p_T charged lepton and a high p_T photon isolated from each other as well as from other hadronic activities, and large p_T . Apart from the photon/isolated- γ misidentification problem to be discussed in section IV, we expect many events from $W \rightarrow \ell\bar{\ell} \nu_{\ell}$ decay processes; this process is interesting in itself and will be discussed in subsection IIB). An immediate guess to get rid of this background is to reject all the events where the $\gamma\ell$ invariant mass $m_{\gamma\ell}$ is smaller than m_W . In Fig.4a we show the $m_{\gamma\ell}$ distribution of the production process (solid line) and of the W radiative decay (dashed line); both curves are for $k = 1$, the standard model. Separation of the two contributions is not good in this variable and thus we would lose a majority of the signal in the $m_{\gamma\ell} > m_W$ cut. Instead, we propose to use the cluster transverse mass $m_T(\gamma\ell; \text{missing})$, defined as follows⁹

$$m_T^2(\gamma\ell; \text{missing}) = [(m_{\gamma\ell}^2 + |\vec{p}_{\gamma T}|^2 + |\vec{p}_{\ell T}|^2)^{1/2} + |\vec{p}_{\gamma T}|^2 - |\vec{p}_{\gamma T} + \vec{p}_{\ell T} + \vec{p}_{\text{miss}}|^2]^2 \quad (2.13)$$

where $\vec{p}_T$ denotes the missing transverse momentum vector about the beam axis. In our case there is one undetectable particle ($\bar{\nu}$) and hence $\vec{p}_T = \vec{p}_{\bar{\nu}T}$. In Fig.4b we plot the $m_T(\gamma\ell; \bar{\nu})$ distribution expected from the production process (solid curve) and the decay process (dashed curve); for this latter process the distribution shows the expected Jacobian peak located at $m_T = m_W$. Since the cluster transverse mass is bounded by its parent mass, we only need to worry about the finite width effect and the error in the $\vec{p}_T$ measurement. By accepting only those events which satisfy

$$m_T(\gamma\ell; \text{missing}) > 90 \text{ GeV} \quad (2.14)$$

we lose only about 40% of the signal while getting rid of most contributions from the W radiative decay. In the following, we adopt the cut (2.14) as well as the cuts (2.12) in all our numerical computations of this subsection.

Let us now return to the problem of looking for the angular zero, predicted by the electroweak gauge theories^{7,8}. The angular zero appears in the W c.m. frame. The problem is simply that although we can in principle measure the W transverse momentum from its leptonic decay, we cannot measure its longitudinal momentum. A typical momentum configuration for a lepton-photon-missing p_T event is shown in Fig.5 where the frame has been boosted until the lepton momentum gets purely transverse. We fix the z and x axis as the proton beam and the $-\vec{p}_{\ell}$ direction, respectively. The dashed line shows the possible location of $\vec{p}_{\bar{\nu}T}$ which does not necessarily lie in the x - z plane for a given $\vec{p}_{\bar{\nu}T}$. Now, it is straightforward to see that if we neglect the width of the W -boson, i.e. $\Gamma_W/m_W = 0$, we have only two solutions for the $\bar{\nu}$ momentum since the invariant mass of the lepton system should form m_W . Explicitly we have for the $\bar{\nu}$ rapidity,

$$y_{\bar{\nu}}(\pm) = y_{\ell} \pm \ln \left\{ 1 + \delta + [\delta(2+\delta)]^{1/2} \right\} \quad (2.15a)$$

with

$$\delta = m_W \left\{ m_W^2 - m_T^2(\gamma\ell; \bar{\nu}), 0 \right\} / 2 |\vec{p}_{\ell T}| |\vec{p}_{\bar{\nu}T}|, \quad (2.15b)$$

$$m_T^2(\gamma\ell; \bar{\nu}) = (|\vec{p}_{\ell T}| + |\vec{p}_{\bar{\nu}T}|)^2 - |\vec{p}_{\ell T} + \vec{p}_{\bar{\nu}T}|^2. \quad (2.15c)$$

The maximum symbol in Eq. (2.15b) is required when we allow for the finite width of W (see discussions in section 3). We specify the two solutions by + and -, which correspond to the solutions with $y_{\bar{\nu}} > y_{\ell}$ and $y_{\bar{\nu}} < y_{\ell}$, respectively. These two solutions coincide at the maximum value of the $\ell\bar{\nu}$ transverse mass, $m_{T-}(\ell; \bar{\nu}) = m_W$. Knowing the $\bar{\nu}$ four-momentum from (2.15), we can easily obtain the W four-momentum

$$p_W(\pm) = p_{\ell} + p_{\bar{\nu}}(\pm) \quad (2.16)$$

as shown in Fig. 5. W bosons with these two different momenta give exactly the same lepton momentum and ϕ_T . The rapidity of the W system in the laboratory frame is then determined for each case by

$$\begin{aligned} y_{\pm} &= \frac{1}{2} \ln \frac{E_W(\pm) + E_T + p_{Wz}(\pm) + p_{Tz}}{E_W(\pm) + E_T - p_{Wz}(\pm) - p_{Tz}} \\ &= \frac{1}{2} \ln \frac{E_{\ell} + p_{\ell z} + E_T + p_{Tz} + \sqrt{p_{T\perp}^2 + m_W^2} \cosh(y_{\bar{\nu}}(\pm))}{E_{\ell} - p_{\ell z} + E_T - p_{Tz} + \sqrt{p_{T\perp}^2 + m_W^2} \cosh(-y_{\bar{\nu}}(\pm))} \end{aligned} \quad (2.17)$$

It is now straightforward to obtain the photon scattering angle in the W c.m. frame measured from the proton beam direction (5.n. 3),

$$\cos\theta_{\pm}^* = \tanh(y_{\gamma} - y_{\pm}) \quad (2.18)$$

On an event by event basis we cannot tell which solution (+ or -) is actually realized. However, we may still expect the $\cos\theta_+^*$ and/or $\cos\theta_-^*$ distributions to be sensitive to the anomalous moment K , since part of the time θ_+^* or θ_-^* actually coincides with the true scattering angle. In Fig. 6a we show our predictions for the $\cos\theta_+^*$ distribution in pp collisions at $\sqrt{s} = 2$ TeV. Although we do not have a zero, the spectrum has a minimum around $\cos\theta_+^* = 0.1$ and its shape is rather sensitive to K . Shown in Fig. 6b are the $\cos\theta_-^*$ distribution, also at $\sqrt{s} = 2$ TeV. For $K = 1$ a cleaner dip is seen at $\cos\theta_-^* \approx 1/3$. The rate difference between $K = 1$ (solid curve), the gauge theory value, and $K = -1$ (dashed-dotted curve) in this region is an order of magnitude; accordingly, the K -sensitivity of the shape of the distribution is even more pronounced in the $\cos\theta_-^*$ spectrum.

The above difference between the $\cos\theta_+^*$ and the $\cos\theta_-^*$ distribution is telling us that most of the time the W boson actually prefers the "-" configuration and thus the corresponding $\cos\theta_-^*$ distribution shows more similarity to the $\cos\theta_{d\gamma}^*$ distribution. Still, part of the time the W takes the "+" configuration and the zero disappears. Contributions from the subprocesses where an antiquark in p and a quark in $\bar{p}$ collide also destroy the zero but they are found to be negligible.

One may examine a combination of the two variables, $\cos\theta_+^*$ and $\cos\theta_-^*$, in search of optimal sensitivity to K . One possibility²⁰⁾ is to calculate event by event the probabilities for the W to take one of the two possible configurations and to choose the higher probability solution. The problem we find in this approach is that the probability depends not only on parton distributions nontrivially but also on the matrix element squared itself, e.g., the value of K which we want to measure. Our $\cos\theta_-^*$ distribution is, on the other hand a purely experimental distribution, straightforward to calculate, and shows sufficient sensitivity to K . Its sensitivity to parton distributions enter only through a simple quantity, the relative importance of the sea-quark contribution (more precisely, the relative contribution from the subprocess where an antiquark in p and a quark in $\bar{p}$ collide) which is not too difficult to estimate quantitatively and is small anyway in $p\bar{p}$ collisions at $\sqrt{s} = 2$ TeV. Hence we choose $\cos\theta_-^*$ as a good example of the variable whose distribution is sensitive to K while insensitive to parton distributions.

2b) The process $p\bar{p} \rightarrow W^-; W^- \rightarrow \ell\bar{\nu}_\ell$

The parton cross section for the "decay" process (2.4) is given by

$$d\hat{\sigma}_{q\bar{q}} = \frac{1}{2\hat{s}} \overline{\sum} |M(q\bar{q} \rightarrow W^-; W^- \rightarrow \ell\bar{\nu}_\ell)|^2 d\Xi \quad (2.19)$$

Here $\hat{s} = m_W^2$ and $d\Xi$ denotes the three body phase space eq.(2.8) and eq.(2.11). The spin and color sum-averaged matrix element squared for an arbitrary κ is given in the appendix (eq. (A14)). In its simplified form it reads

$$\overline{\sum} |M(q\bar{q} \rightarrow W^-; W^- \rightarrow \ell\bar{\nu}_\ell)|^2 = \frac{128\pi^4\alpha^2}{\sin^2\theta_W} B(W^- \rightarrow \ell\bar{\nu}_\ell) |U_{q1}|^2 \cdot \delta[(p_1 + p_2)^2 - m_W^2] \left\{ \frac{(2p_1 \cdot p_2)[(2p_1 \cdot p_\ell)^2 + (2p_2 \cdot p_\ell)^2]}{(2p_1 \cdot p_2)(2p_1 \cdot p_{\bar{\nu}_\ell}) + (2p_2 \cdot p_{\bar{\nu}_\ell})^2} + O(1-\kappa) \right\} \quad (2.20)$$

As in the production case we shall neglect both the Cabibbo rotation and the contribution from strange and heavier quark sea. The only surviving subprocess is then $u\bar{u} \rightarrow W^-; W^- \rightarrow \ell\bar{\nu}_\ell$ with $|U_{ud}| = 1$.

According to the generalized null radiation zone theorem⁸⁾, the matrix element squared (2.20) has a zero for $\kappa = 1$ when the photon and antineutrino are collinear ($p_1 \cdot p_2 = 0$). Although the amplitude has the same factor ($p_1 \cdot p_2$) (see appendix), the cross section is linear and not quadratic in this factor because of a cancellation. This cancellation can easily be seen by comparing our result with the polarization averaged W decay distribution into a photon and a quark pair (qq)

$$W^-(p_W) \rightarrow 1(p_1) + \bar{q}'(p_{\bar{q}'}), \quad (2.21)$$

calculated by Grose and Mikaelian²¹⁾. The differential decay width can be expressed as

$$d\Gamma \sim [Q_1(p_1, p_{\bar{q}'} - q_{\bar{q}'}) - Q_2(p_1, p_{\bar{q}'})]^2 \frac{(p_W \cdot p_{\bar{q}'})^2 + (p_W \cdot p_1)^2}{(p_1 \cdot p_{\bar{q}'})^2 (p_1 \cdot p_{\bar{q}'})} \quad (2.22)$$

where Q_q and $Q_{\bar{q}'}$ denote quark and antiquark charge, respectively. We find the square of a charge dependent factor. The polarization averaged differential decay width of the process $W^- \rightarrow \ell\bar{\nu}_\ell$ can be obtained from (2.22) simply by replacing q by ℓ and $\bar{q}'$ by $\bar{\nu}_\ell$. Since $Q_{\bar{\nu}_\ell} = 0$, the charge factor in eq. (2.22) becomes simply $(p_1 \cdot p_{\bar{\nu}_\ell})^2$ while one of the powers is cancelled by the same factor in the denominator.

Because of this cancellation, the κ -sensitivity in the decay process is less pronounced as compared to the production process. We thus choose the κ -values $\kappa = -3, 1, 5$, to show our results. The cross section for an arbitrary κ -value is then obtained from the three curves via the formula

$$d\sigma(\kappa) = \frac{1}{32} (\kappa-1)(\kappa+3) d\sigma(5) - \frac{1}{16} (\kappa-5)(\kappa+3) d\sigma(1) + \frac{4}{32} (\kappa-5)(\kappa-1) d\sigma(-3) \quad (2.23)$$

In the following distributions we always show separately the contribution from the production process as far as it is compatible with the cuts to be specified below. Even though this contribution may be regarded as a background in a strict sense, we add it to the signal since we find no practical way of distinguishing between these two contributions.

Shown in Figs. 7a and 7b are the transverse momentum spectrum of the photon and the charged lepton, respectively, expected at $\sqrt{s} = 2$ TeV. The solid line represents $\kappa = 1$, the dashed one $\kappa = -3$ and the dashed-dotted one $\kappa = 5$. The following cuts have been imposed and will be applied in further distributions of this subsection:

$$p_{T\gamma}, p_{T\ell} > 10 \text{ GeV} ;$$

$$|y_\gamma|, |y_\ell| < 3 ;$$

$$30 \text{ GeV} < m_T(\gamma\ell; \text{missing}) < 90 \text{ GeV} ;$$

$$\cos \theta_{\gamma\ell} < 0.95 ,$$

(2.24)

The cut on $m_T(\ell\bar{\ell}; \text{missing})$ is a consequence of the discussion in subsection 2a); the cut on the photon charged lepton opening angle in the collider c.m. frame is made to avoid the mass singularity. From these figures we find that the p_T distributions are governed mainly by the kinematics of the W decay and are insensitive to K . The photon p_T distribution falls off steeply with rising p_T and thus the photon identification at relatively small p_T values (~ 10 GeV) is essential to gain statistics. On the other hand, we expect a rather flat p_T distribution for the charged lepton up to $p_T \sim 30$ GeV.

We show in Figs. 8a and 8b the rapidity distributions of the photon and the charged lepton, respectively. Again, we see little sensitivity to K . A marked difference between these rapidity distributions and those in the production region (see Fig. 3) is that here the rapidity distributions are more or less symmetrical about the $y = 0$ axis. It is, however, obvious that we cannot determine the K -value from these distributions: We should look for those distributions whose shapes are sensitive to K .

To attack this problem we study first the final state configurations in the W rest frame. The polarization averaged W decay distribution can be expressed conveniently in terms of two variables: the photon momentum fraction

$$X = 2 p_W \cdot p_\gamma / p_W^2 = 2 E_\gamma^* / m_W \quad (2.25)$$

and $\cos\theta_{\ell\ell}^*$. Here the superscript * refers to the W rest frame. For arbitrary K the normalized decay distribution reads

$$\frac{1}{\Gamma(W \rightarrow \ell\bar{\ell})} \frac{d\Gamma}{dx d\cos\theta_{\ell\ell}^*} = \frac{x}{4\pi} \frac{1}{[1 - (1 - \cos\theta_{\ell\ell}^*)(x/2)]^2} \sum_{\ell} (x, \cos\theta_{\ell\ell}^*) \quad (2.26a)$$

with

$$\begin{aligned} \sum_{\ell} (x, c) = & \frac{1+c}{1-c} \left[\frac{1+(1-x)^2}{x^2(1-x)} - \frac{1-c^2}{2(1-\frac{1-c}{2}x)^2} \right] \\ & + (1-x) \frac{(1+c)(c+\frac{1-c}{2}x)}{2(1-\frac{1-c}{2}x)^2} + (1-x)^2 \frac{1-x}{8} \left[1 + \frac{(1-c^2)(1+x)}{2(1-\frac{1-c}{2}x)^2} \right] \end{aligned} \quad (2.26b)$$

The distribution (2.26b) has a single zero at $c = \cos\theta_{\ell\ell}^* = -1$, only when $K=1$. Since any deviation from the gauge theory value of K results in filling up this zero, the $\cos\theta_{\ell\ell}^*$ distribution is expected to be sensitive to K .

We show in Fig. 9 the $\cos\theta_{\ell\ell}^*$ distribution integrated over the x -range $0.3 < x < 1$, a typical momentum fraction region of the photon in our analysis ($p_{\gamma T} > 10$ GeV eliminates every event with $x < 0.24$). Indeed, we find that the distribution is rather sensitive to K in the smaller $\cos\theta_{\ell\ell}^*$ region. However, the K -sensitivity is weaker than that found in the angular distribution of the W production process. This is partly because the distribution (2.26) has only a single zero instead of a double zero at $K=1$; we further find that the Jacobian factor $(1-x(1-\cos\theta_{\ell\ell}^*)/2)^{-2}$ plays an important role in partially compensating for the dip at $\cos\theta_{\ell\ell}^* = -1$. This Jacobian factor is responsible for the nonobservability of the dip for $K=1$ and also for the enhancement observed in this region for nongauge theory cases. Our primary goal is thus to find a measurable quantity in the $p\bar{p}$ collision which is most similar to this $\cos\theta_{\ell\ell}^*$ distribution.

An immediate candidate is the distribution in terms of the azimuthal angle $\phi_{\ell\ell}$ between the photon and the charged lepton which is boost invariant. When $\cos\theta_{\ell\ell}^* = -1$, $\phi_{\ell\ell}$ should be π although the converse statement does not hold. Shown in Fig. 10 are the expected distributions for the three K -values. The distributions are symmetric about $\phi_{\ell\ell} = \pi$ because of the absence of both the CP violation phase and the final state interaction phase. Although there is no angular zero in this distribution, the K -dependence of the spectrum around $\phi_{\ell\ell} = \pi$ follows qualitatively that of the $\cos\theta_{\ell\ell}^*$ distribution shown in Fig. 9. The two peaks at $\phi_{\ell\ell} = 18^\circ$ and 162° are simply due to the collinear singularity cut, $\cos\theta_{\ell\ell} < 0.95$ (see eq. (2.24)). This distribution may, however, suffer from large QCD correction (see discussions in section 3) since the azimuthal angle is not invariant under the transverse motion of W . It is therefore desirable to look for a variable which refers to the W -boson rest frame.

Just as in the case of the W production process studied in the previous subsection, there is a two-fold ambiguity in the longitudinal momentum of $\bar{\nu}$. In Fig. 11 we show the possible three-momentum configuration of the $\ell\bar{\ell}\bar{\nu}$ final states in the boosted frame where $\vec{p}_\ell + \vec{p}_{\bar{\ell}} \rightarrow \vec{p}_T$ has no longitudinal component. The dashed line shows the possible location of $p_{\bar{\nu}}$ for a given $\vec{p}_T$.

In the limit $\Gamma_W/m_W \rightarrow 0$ the neutrino momentum is determined up to two symmetric configurations shown in the figure by $\vec{p}_\nu$. Explicitly we find for the $\vec{p}_\nu$ rapidity in the collider c.m. frame

$$y_{\vec{p}_\nu}(\pm) = y_{\vec{p}_\ell} \pm \ln \left\{ 1 + \delta + [\delta(2+\delta)]^{1/2} \right\} \quad (2.27a)$$

where $y_{\vec{p}_\ell}$ denotes the rapidity of the $\vec{p}_\ell$ cluster and

$$\delta = \max \left\{ m_W^2 - m_T^2(x_\ell; \vec{p}), 0 \right\} / 2 \left[\vec{p}_T^2 + |\vec{p}_T + \vec{p}_{x_\ell}|^2 \right]^{1/2} \quad (2.27b)$$

with the cluster transverse mass $m_T(\ell; \vec{p})$ as defined in eq. (2.13). The maximum symbol in eq. (2.27b) is needed because our cut (2.24) allows $m_T(\ell; \vec{p}) > m_W$, where a part of "production" contribution exists (see Fig. 4b). We specify the two solutions by + and -, corresponding to the solution with $y_{\vec{p}} > y_{\vec{p}_\ell}$ and $y_{\vec{p}} < y_{\vec{p}_\ell}$, respectively. These two solutions coincide at the maximum value of the cluster transverse mass $m_T(\ell; \vec{p}) = m_W$.

It is now straightforward to obtain the W four-momentum for each solution

$$P_W(\pm) = p_\ell + p_{x_\ell} + p_{\vec{p}}(\pm) \quad (2.28)$$

The rapidity of the W in the laboratory frame is

$$y_\pm = \ln \frac{E_W(\pm) + P_{Wz}(\pm)}{[m_W^2 + |\vec{p}_\ell + \vec{p}_{x_\ell} + \vec{p}_{\vec{p}}|^2]^{1/2}} \quad (2.29)$$

The opening angle between the photon and the charged lepton in the W rest frame can then be expressed as

$$\cos \theta_{\ell x_\ell}^* = \frac{\sinh(y_\ell - y_\pm) \sinh(y_{x_\ell} - y_\pm) + \cosh y_{\vec{p}_\ell}}{\cosh(y_\ell - y_\pm) \cosh(y_{x_\ell} - y_\pm)} \quad (2.30)$$

In Fig. 12 the $\cos \theta_{\ell x_\ell}^*$ distribution is plotted for the process $\bar{p}p \rightarrow W^+ X; W^+ \rightarrow \ell \ell^+ \vec{p}_\ell$ at $\sqrt{s} = 2$ TeV, with the final state cuts (2.24). The sensitivity of the distribution to K is clearly seen. For the gauge theory case ($K=1$) there is a dip at $\cos \theta_{\ell x_\ell}^* = -1$ which is a remnant of the zero. The shape of the distributions for non-gauge theory cases ($K=3,5$) shows a clear difference, especially in the negative $\cos \theta_{\ell x_\ell}^*$ region. In low statistics experiments, it would be nice to study an integrated quantity. We define the ratio of the events with negative $\cos \theta_{\ell x_\ell}^*$ as

$$R = \frac{N(-1 \leq \cos \theta_{\ell x_\ell}^* < 0)}{N(-1 \leq \cos \theta_{\ell x_\ell}^* < 0.9)} \quad (2.31)$$

where N denotes the number of events in a specified region. We obtain from Fig. 12, after summing decay and production contributions, a prediction for an arbitrary K -value by using the formula eq. (2.23):

$$R(\sqrt{s} = 2 \text{ TeV}) = 0.264 \frac{1 + 0.015(1-K) + 0.049(1-K)^2}{1 + 0.011(1-K) + 0.016(1-K)^2} \quad (2.32)$$

The numerical uncertainties in the coefficients of eq. (2.32) are less than 5%. The ratio R is 20% for $K=1$, 30% for $K=5$ and $K=-3$, at $\sqrt{s} = 2$ TeV. The ratio R is rather insensitive to $\sqrt{s}$ below Tevatron energies, since the $\sqrt{s}$ dependence enters mainly through the relative importance of the production process contribution which is small at lower energies. At the CERN $\bar{p}p$ collider energies, we find

$$R(\sqrt{s} = 540 \text{ GeV}) = 0.238 \frac{1 + 0.016(1-K) + 0.053(1-K)^2}{1 + 0.011(1-K) + 0.016(1-K)^2} \quad (2.33)$$

No significant deviation from the above values is expected at $\sqrt{s} = 630$ GeV.

Unlike in the case of $\cos \theta_{\ell^+}^*$ and $\cos \theta_{\ell^-}^*$ distributions studied in the previous subsection, we find no significant difference between $\cos \theta_{\ell x_\ell}^*$ and $\cos \theta_{\ell^+ \ell^-}^*$ distributions. This can be understood since the bulk of the events have large cluster transverse mass (see Fig. 4b) for which the difference between two solutions is small (see eq. (2.27b)).

3) Effects of finite W-width and higher order QCD corrections

Up to now our presentation did heavily rely on the zero width approximation for the W-boson on one side and on lowest order QCD predictions on the other side. In this section we discuss the effect of finite width in a quantitative way whereas only qualitative statements will be made for higher order QCD corrections.

3a) Finite W-width effects

Rigorously speaking, the finite width of the W-boson does not allow to divide the process

$$d + \bar{u} \rightarrow \ell^- + \gamma + \bar{\nu}_\ell \quad (3.1)$$

into the two processes which we called "production" (2.3) and "decay" (2.4) processes. Indeed, there appears an interference between the "production" and "decay" amplitudes (see the appendix for detail). Since the "production" amplitude has an angular zero at $\cos\theta_{\ell\gamma}^* = 1/3$ and the "decay" amplitude has a zero at $\cos\theta_{\ell\gamma}^* = -1$, it is clear that the whole contribution should vanish in gauge theory when both two conditions

$$\cos\theta_{d\ell}^* = 1/3 \quad (3.2a)$$

$$\cos\theta_{\ell\gamma}^* = -1 \quad (3.2b)$$

are satisfied. (Remark that $\theta_{d\ell}^*$ is defined in the $W\ell$ c.m. frame whereas $\theta_{\ell\gamma}^*$ is defined in the W rest frame). In fact, these two conditions follow from the generalized null-radiation zone theorem⁸⁾ applied directly to the process (3.1).

Fortunately, the interference term vanishes in gauge theory when either one of the two conditions in eq. (3.2) is satisfied. It should also be small in the dip regions and hence we do not expect large corrections from the interference. We confirmed this expectation by repeating all the numerical computations with the exact finite width formula given in the appendix. Apart from trivial kinematical effects, e.g., the smoothing of the Jacobian peak in the cluster transverse mass distribution (see Fig. 4b), we found no significant deviation from our zero-width results presented so far. Especially, both the $\cos\theta^*$ and $\cos\theta_{\ell\gamma}^*$ distributions agree with the zero-width results presented in Fig. 6b and Fig. 12, respectively, within the numerical errors of our Monte Carlo calculation.

3b) Higher order QCD corrections

The problem of higher order QCD corrections is far more difficult to study quantitatively; only an explicit next-to-leading order calculation will actually tell us the significance of the corrections (F.n.4). However, we shall in the following give a necessary modification of the "production" variable $\cos\theta^*$ in the presence of the transverse motion of the $W\ell$ system and also give a plausible argument for the smallness of the radiative corrections to the $\cos\theta_{\ell\gamma}^*$ distribution in the "decay" process.

Higher order QCD corrections can be especially critical in the $W\ell$ production studies since we are interested in the dip structure of the distribution. Single as well as multiple gluon radiation²³⁾ is expected to give the $W\ell$ system a transverse momentum of about 10 GeV, which is comparable in magnitude to our transverse momentum cuts (2.12). Unless we take into account the transverse motion of the $W\ell$ system properly, it seems clear that the dip structure would be smeared away. It is at least necessary to define the scattering angle in the $W\ell$ c.m. frame to avoid large kinematical smearing. Indeed, the successful confrontation²⁴⁾ of the dijet angular distribution at the CERN $p\bar{p}$ collider with the lowest-order QCD prediction²⁵⁾ was achieved in the dijet c.m. frame, in particular, the Collins-Soper (CS) frame²⁶⁾. The CS frame is particularly convenient, and probably gives optimal cancellation among corrections, due to its symmetric treatment of the colliding beams.

Encouraged by the success of the dijet studies²⁴⁾, we propose a possible modification of the $\cos\theta^*$ variable (2.18) in the presence of the nonzero transverse motion of the $W\ell$ system:

$$\cos\theta_{\pm}^* = (p_{T\ell} / \hat{p}_{\pm}) \sinh(y_{\ell} - y_{\pm}) \quad (3.3a)$$

where $p_{T\ell}$ and y_{ℓ} are the photon transverse momentum and rapidity, respectively, $y_{\pm}$ is defined in (2.17), and

$$\hat{p}_{\pm} = (\hat{s}_{\pm} - m_W^2) / 2 \sqrt{\hat{s}_{\pm}} \quad (3.3b)$$

with

$$\hat{S}_\perp = (p_{\perp 1} + p_{\perp 2})^2 = m_W^2 + m_{\tilde{W}}^2 + 2\vec{p}_{\perp 1} \cdot \vec{p}_{\perp 2} [\cosh(\theta_{\perp 1} - \theta_{\perp 2}) - \cos \phi_{\perp 1,2}]. \quad (3.3c)$$

Either $\hat{\theta}_\perp^*$ or $\hat{\theta}_\perp^*$ agrees in the zero W width limit with the χ polar angle in the CS frame of the $W\tilde{W}$ system. This new variable reduces to our old one (2.18) in the zero transverse momentum limit of the $W\tilde{W}$ system. Hence the $\cos \theta_\perp^*$ distributions shown in Fig. 6 are also the $\cos \hat{\theta}_\perp^*$ distributions in the lowest order. Higher order corrections can, however, be significantly different between the two distributions. We examined the effect of purely kinematical smearings by giving both colliding partons the Gaussian transverse momentum distributions with $\langle p_\perp^2 \rangle = 7 \text{ GeV}^2$; the $\cos \hat{\theta}_\perp^*$ distribution received small corrections while the dip in the $\cos \theta_\perp^*$ distribution almost disappeared.

It is therefore important to look for a "good" variable whose distribution receives small corrections in the future next-to-leading order studies. Guided by the successful dijet studies, we propose the variable $\cos \hat{\theta}_\perp^*$ as a candidate. Since our interest is in the dip structure of distributions, it is not clear at present whether such a "good" variable exists or not.

On the other hand, the situation is much better in the W radiative decay studies, where we expect small corrections in the $\cos \theta_{ff}^*$ distributions. First of all, the variable $\cos \theta_{ff}^*$ (2.30) refers to the W -rest frame variable even in the presence of a W transverse motion. Hence gluon radiation can affect the distribution only through the W -polarization. On one hand this cannot influence the distribution near the zero at $\cos \theta_{ff}^* = -1$ because every polarization amplitude has a common zero at that point¹⁹⁾. On the other hand, the whole distribution is rather insensitive to W polarizations as can be seen from comparing the distribution from an unpolarized W (Fig. 9) and that of a longitudinal polarized W (Fig. 12). The main uncertainty thus enters through the extra contribution from the $W\tilde{W}$ production process (see Fig. 12), whose magnitude is small anyway in most of the $\cos \theta_{ff}^*$ range. Only in the very vicinity of the dip at $\cos \theta_{ff}^* = -1$, one needs a rough estimate of the correction for the $W\tilde{W}$ production contribution.

4) Backgrounds

In this section we study background contributions to the production process (2.3) as well as to the decay process (2.4). A large $p_\perp$ isolated charged lepton with large missing $p_\perp$ can come either from W or heavy quark pair production. A pair production of a heavy quark (top^{27,28} or even heavier quarks²⁹), however, typically ends up with multi-jet events and is unlikely to fake our signal. A significant background is expected from the process

$$p + \bar{p} \rightarrow W^- + \text{jet} + \text{anything} \quad (4.1)$$

$$\quad \quad \quad \downarrow \quad \quad \quad \downarrow$$

$$\quad \quad \quad \ell \quad \quad \quad \bar{\nu}$$

where a high $p_\perp$ jet resembles a high $p_\perp$ photon through χ/π^0 misidentification. Since the W -jet production cross section is roughly a factor of α_s/α larger than the $W\tilde{W}$ production cross section, and since the W dependence of the distributions is strongest in the dip region, the background contribution severely restricts the sensitivity of our measurement of the K values. On the other hand, the W radiative decay experiment is found to be rather free from large backgrounds. The main restrictive factor here is then statistics since the K -dependence in this process is milder than in the case of W production.

4a) Backgrounds for $\bar{p}p \rightarrow W^- \tilde{W} X$; $W^- \rightarrow \ell \bar{\nu} \ell$

In this subsection we study backgrounds for the production process (2.3). A severe background is expected from the W -jet production process (4.1) where a high $p_\perp$ jet takes a single high $p_\perp$ photon. The lowest order subprocesses which contribute to W^- jet production are

$$d + \bar{u} \rightarrow W^- + \text{gluon} \quad (4.2a)$$

$$\bar{d} + \text{gluon} \rightarrow W^- + u \quad (4.2b)$$

$$\bar{u} + \text{gluon} \rightarrow W^- + \bar{d} \quad (4.2c)$$

The production cross section for these subprocesses including the subsequent decay of $W \rightarrow \ell \bar{\nu}$ can be obtained from the appendix; only color factors and coupling constants have to be changed. (See also the published literature¹⁰⁾). We use the same parton distribution^{15,16} as in section 2 and for the QCD running coupling constant we take

$$\alpha_s(\hat{s}) = 12\pi / (33 - 2n) \ln(\hat{s}/\Lambda_n^2) \quad (4.3)$$

where n is the number of the active flavors and $\hat{s}$ the squared mass of the colliding parton system. The flavor number is increased by one at each threshold, $\sqrt{\hat{s}} = 2m_n$; the continuity of α_s is maintained by choice of the Λ_n . We choose $\Lambda_4 = 0.3$ GeV, $m_b = 5$ GeV, and $m_t = 35$ GeV for definiteness.

After imposing the cuts (2.12) and (2.14) by replacing the photon by the jet, we find at $\sqrt{s} = 2$ TeV a cross section of about 0.16 nb for the $W^- \rightarrow e\bar{\nu}_e$ channel. The background contribution is then obtained by multiplying this cross section by the probability $P_{\gamma/j}$ that a jet is misidentified as a single isolated photon. A precise value for $P_{\gamma/j}$ depends on details of the jet structure and of the detector. As for the cut of $p_{\gamma T} > 10$ GeV, a BNL study group reported²⁰⁾ values between 1/230 and 1/500, depending on the resolution of the γ shower detector. In the following, we choose a value

$$P_{\gamma/j} = 1/200 \quad (4.4)$$

for the background estimate. We choose this rather pessimistic value because of the uncertainties in our computation of the signal-to-background ratio: The γ /jet misidentification probability should be different for gluon and quark jets and hence between pp reactions where the "annihilation" subprocess (4.2a) dominates (at $\sqrt{s} = 2$ TeV about 70% of the total W -jet events come from (4.2a)) and pp reactions where the "Compton" subprocess (4.2b) and (4.2c) are more important. Furthermore, the QCD higher order effects including multigluon emission can affect somewhat differently the signal and background contributions. In actual experiments, a reliable estimate for the probability $P_{\gamma/j}$ is essential to estimate the background and hence to determine the value of K . It is therefore very important to understand its property in prompt photon experiments at hadron colliders.

In Fig. 13 we plotted again our predictions for the $\cos\theta^*$ distributions of the signal events for several values of K in pp collisions at $\sqrt{s} = 2$ TeV. In the same figures we show by the dashed line the background W -jet contribution multiplied by the factor $P_{\gamma/j} = 1/200$ (eg. (4.4)). The integrated cross section for the $K=1$ signal and the background are roughly the same, about 3 pb after summing up e^+ and μ^+ modes. Without including the QCD motivated K -factor

we expect 300 events for both signal and backgrounds with a nominal integrated luminosity of 10^{38} cm^{-2} . The shape of the distributions are, however, very different for small $|K-1|$. At $|K-1|=5$, the shape of the signal and the background become more or less the same. In order to determine the K -value from the observed distribution, which is the sum of the signal and background contributions, we need to estimate both the magnitude and the shape of the backgrounds. The shape of the background distribution can be determined rather well by the jet trigger experiments, while its magnitude is essentially determined by the factor $P_{\gamma/j}$. It seems to be difficult to obtain a bound stronger than $|K-1| < 2$ because the background dominates over the signal for $|K-1| < 2$ in the dip region where the K -dependence is strongest.

4b) Backgrounds for $p\bar{p} \rightarrow W^- X; W^- \rightarrow \gamma \ell^- \bar{\nu}_\ell$

As studied in sections 2 and 3, there are two advantages and one disadvantage in measuring K in W production followed by its radiative decay, $W^- \rightarrow \gamma \ell^- \bar{\nu}_\ell$. The two advantages are the relatively larger cross section and the insensitivity of the analysis on the transverse motion of the W . The disadvantage is the fact that the K -dependence of the signatures is not as spectacular as that for the $W\gamma$ production case. In the following we will find another advantage in studying W radiative decay, the large signal-to-background ratio.

In terms of observable momenta, the W radiative decay process can be specified by the events satisfying the following cuts (see (2.24)):

$$\begin{aligned} P_{\gamma T}, p_{\ell T} &> 10 \text{ GeV}; |\gamma_\ell|, |\gamma_\ell| < 3; \\ 30 \text{ GeV} &< m_1(\gamma\ell; \text{missing}) < 90 \text{ GeV}; \\ \cos\theta_{\ell\ell} &< 0.95 \end{aligned} \quad (4.5)$$

A significant background can again come from W -jet production processes (4.2) where the jet is misidentified as a single photon. We find that roughly one-third of the jet-plus- $\ell\bar{\nu}$ events has the cluster transverse mass $m_T(\ell\text{jet};\bar{\nu})$ smaller than 90 GeV. Furthermore, the contribution from W -jet production followed by the decay $W^- \rightarrow \tau\bar{\nu}_\tau$ is nonnegligible since a substantial portion of the events with a subsequent $\tau \rightarrow e$ or μ decay survives the cuts (4.5) when we replace a photon by a jet. In this evaluation we use the branching ratio $\text{BR}(\tau \rightarrow e) = 0.17$ and the purely left-handed $\tau \rightarrow e$ decay distribution calculated in the $V-A$ theory. The $\vec{p}_T$ in this case is the vector sum of all the neutrino transverse momenta. We find that the cascade $W \rightarrow \tau \rightarrow e$ decay contribution to the background is about 20% of the direct $W^- \rightarrow e\bar{\nu}$ contribution. The background shown below is the sum of these two contributions.

In Fig. 14 we show our predictions for the $\cos\theta_{\ell\ell}^*$ distributions of the radiative W decay signal events for several values of κ in pp -collisions at $\sqrt{s} = 2$ TeV. The dashed line represents the background contribution, again multiplied with $P_{Y/j} = 1/200$ (see eq.(4.4)) for the Y/jet misidentification probability. The signal curves are just the sum of the pure $W \rightarrow e\bar{\nu}$ (decay) contribution and the W -jet production contribution, shown separately in Fig. 12, which satisfy the final state cuts (4.5). Contributions from the cascade processes $W \rightarrow \tau\bar{\nu}; \tau \rightarrow e$ and W -jet $W \rightarrow \tau\bar{\nu}; \tau \rightarrow e$ are both found to be negligible. From this figure we find that the estimated background is always smaller than the signal cross sections, even in the dip region around $\cos\theta_{\ell\ell}^* = -1$ for the $\kappa=1$ case. The shape of the distribution is also very different between the signal and the background. Hence, a factor of one error in the estimate of the magnitude of the background (that is of the Y/jet confusion probability $P_{Y/j}$) does not severely restrict the measurement of κ .

5) Summary and Discussion

In this paper we examined the feasibility to measure the WW triple gauge coupling at proton-antiproton colliders through the process $\bar{p}p \rightarrow \ell\bar{\nu}_\ell X$. To quantify this measurement, we allow arbitrary real value for the anomalous magnetic moment⁴⁾ κ of the W -boson. In any gauge theory of the electroweak interaction, the κ value should be unity up to $O(\kappa)$ corrections. Hence any measurement of κ different from unity signals the failure of gauge theories in the weak interaction.

In the zero-width approximation for the W propagator the $\ell\bar{\nu}_\ell$ final state can originate either from the "production" process $\bar{p}p \rightarrow W^+ X$ with the subsequent $W^+ \rightarrow \ell\bar{\nu}_\ell$ decay or from the "decay" process $\bar{p}p \rightarrow W^+ X$ with the W^+ decaying radiatively into $\ell\bar{\nu}_\ell$. We found that these two processes can be kept apart rather effectively by imposing a cut in the cluster transverse mass variable, $m_T(\ell\bar{\nu};\bar{\nu})$.

The cross section of the subprocesses $d\bar{u} \rightarrow W^+ X$ and $W^- \rightarrow \ell\bar{\nu}_\ell$ vanish at specific kinematical configurations^{7,8)}, $\cos\theta_{d\bar{u}}^* = 1/3$ in the W c.m. frame and $\cos\theta_{\ell\bar{\nu}}^* = -1$ in the W rest frame, at $\kappa=1$ the gauge theory value. A unique reconstruction of these angular variables from the observable momenta, $\vec{p}_\ell, \vec{p}_{\bar{\nu}_\ell}$, and the missing p_T vector is not possible, allowing for the two-fold ambiguity in the W -longitudinal momentum in 1st zero width limit. We find, however, that κ -sensitive distributions can still be obtained in terms of the variables $\cos\theta_{\ell\bar{\nu}}^*$ (2.18) and $\cos\theta_{Y\ell}^*$ (2.30), which are defined in terms only of observable momenta in such a way that in most of the cases they coincide with the quantity $\cos\theta_{d\bar{u}}^*$ and $\cos\theta_{\ell\bar{\nu}}^*$, respectively, in the zero W -width limit. The κ -sensitivity of the $\cos\theta_{\ell\bar{\nu}}^*$ distribution near the dip at $\cos\theta_{\ell\bar{\nu}}^* = 1/3$ is found to be spectacular, giving a cross section for $|\kappa-1| = 2$ which is an order of magnitude higher than the one corresponding to $\kappa=1$, the gauge theory value.

The κ -sensitivity of the $\cos\theta_{Y\ell}^*$ distribution is found to be moderate due to the fact that the W -radiative decay distribution vanishes at $\kappa=1$ only linearly (and not quadratically as in the W -production case) when the angular zero point is approached.

The effects of the finite W width on both distributions are found to be negligible. A more serious problem is to estimate higher order QCD corrections. Especially large corrections are expected for the $\cos\theta^*$ distribution from the transverse motion of the W system. We argued that at least a modification of the variable, e.g. $\cos\tilde{\theta}^*$ as defined in eq.(3.3), is necessary to make the correction smaller. Whether such a modification of the variable is sufficient or not, we will learn only after serious next-to-leading order studies. The situation in the case of $\cos\theta^*$ distribution is much better, for which we presented a plausible argument for the smallness of the QCD radiative corrections. This is mainly due to the fact that the quantity $\cos\theta^*$ refers to the variable in the W rest frame irrespective to the W transverse motion.

Backgrounds due to the W -jet production where the jet is misidentified as a single isolated photon were also studied. We estimated the effects of such contributions by assuming the W -jet misidentification probability $P_{\gamma/j}$ to be 1/200. The background is found to affect the measurement of W -seriously in the production signal ($\cos\theta^*$ distribution) while it is expected to be less dangerous in the radiative decay signal ($\cos\theta_{\gamma\ell}^*$ distribution).

On the experimental side the most serious problem is the photon identification at relatively small p_T (> 10 GeV). It is also important to reduce the error of p_T in order for the $m_T(\ell\gamma, \nu)$ cut (to distinguish production from decay) to be fully effective. Once those conditions as well as the high luminosity requirement are met, we find it feasible to measure the WW coupling at pp colliders, especially in the W radiative decay signals. Adding up e^+ and μ^+ contributions, the expected event rates for $W=1$ at the Fermilab collider ($\sqrt{s} = 2$ TeV) are, after applying all the final state cuts (2.12, 2.14) and (2.24), respectively,

30 events for "production" signal
135 events for "decay" signal

with a nominal integrated luminosity of 10^{37} cm^{-2} . One order of magnitude smaller event rates are expected at the CERN collider. High luminosity is clearly the basic requirement for the measurements.

We conclude by listing advantages and disadvantages of the two signal processes in measuring W . The $W\gamma$ production signal gives a spectacular W dependence but has a smaller cross section, a larger background contamination and is most likely to be rather sensitive to QCD higher-order corrections. The W radiative decay signal gives a moderate W -dependence while it has a relatively larger cross section, a small background contamination and is rather insensitive to higher-order QCD corrections.

Part of the results had been reported at 4th Topical Workshop on Proton Antiproton Collider Physics³⁰⁾.

Acknowledgements

The authors wish to thank V. Barger, M. Ebel, C. Goebel, F. Halzen, K. Hikasa, S. Jacobs and Y. Okamoto for useful discussions. The work of J.C. is partially supported by CAICYT and Comisión de Intercambio Cultural. The work of F.H. is supported by the Swiss National Science Foundation and partially by the Max Geldner Stiftung. This work was started when all the authors were at the University of Wisconsin-Madison, where it was supported in part by the University of Wisconsin Research Committee with funds granted by the Wisconsin Alumni Research Foundation, and in part by the Department of Energy under contract DE-AC02-76ER00881.

Appendix

In this appendix we present the complete partonic cross section for the process $d\bar{u} \rightarrow \gamma \ell \bar{\nu}$; all fermion masses will be neglected. The Feynman diagrams of this process are shown in Fig.1.

Using the short-hand notation

$$p = p_d, \bar{p} = p_{\bar{d}}, k = p_{\gamma}, \ell = p_{\ell}, \bar{\ell} = p_{\bar{\ell}}, P = p + \bar{p} \quad (\text{A.1})$$

for the four-momenta of the particles involved, the amplitude can be written as

$$M = \frac{(4\pi\alpha)}{2\sin^2\theta_W} \left\{ \bar{v}(p) \gamma_{\rho} u(p) \frac{\epsilon^{\alpha}(k) J_{\alpha}^{\rho}}{(P-k)^2 - m_W^2 + im_W \Gamma_W} + \bar{u}(\ell) D_{\rho\alpha} v(\bar{\ell}) \frac{\epsilon^{\alpha}(k) J_{\alpha}^{\rho}}{P^2 - m_W^2 + im_W \Gamma_W} \right\} \quad (\text{A.2})$$

α is the fine structure constant, θ_W the Weinberg angle and $m_W(\Gamma_W)$ the mass (width) of the W-boson. The tensors $P_{\rho\alpha}$ and $D_{\rho\alpha}$ are given by

$$P_{\rho\alpha} = \left\{ \bar{F}_c \left[\gamma_{\rho} \frac{k-k'}{k \cdot P} \gamma_{\alpha} + \gamma_{\alpha} \frac{\bar{k}-\bar{k}'}{\bar{P} \cdot k} \gamma_{\rho} \right] + (1-n) \frac{k \cdot \bar{k} - k' \cdot \bar{k}'}{P \cdot k} \right\} \frac{1-\delta_5}{4} \quad (\text{A.3a})$$

$$D_{\rho\alpha} = \left\{ \bar{F}_c \left[\gamma_{\rho} \frac{k+k'}{k \cdot \bar{I}} \gamma_{\alpha} - \gamma_{\alpha} \frac{\bar{k}+\bar{k}'}{\bar{k} \cdot \bar{I}} \gamma_{\rho} \right] + (1-n) \frac{k \cdot \bar{k} - k' \cdot \bar{k}'}{P \cdot k} \right\} \frac{1-\delta_5}{4} \quad (\text{A.3b})$$

with two charge-dependent factors

$$\bar{F}_c = \frac{Q_u P \cdot k + Q_d \bar{P} \cdot k}{P \cdot k}, \quad \bar{F}_c' = \frac{k \cdot \bar{k}}{P \cdot k} \quad (\text{A.4})$$

The tensors $P_{\rho\alpha}$ and $D_{\rho\alpha}$ satisfy the current conservation conditions

$$k^{\alpha} P_{\rho\alpha} = k^{\alpha} D_{\rho\alpha} = 0$$

In addition we defined the lepton and quark currents,

$$J_{\ell}^{\rho} = \bar{u}(\ell) \gamma^{\rho} \frac{1-\delta_5}{2} v(\bar{\ell}), \quad J_P^{\rho} = \bar{v}(\bar{P}) \gamma^{\rho} \frac{1-\delta_5}{2} u(P) \quad (\text{A.5})$$

In order to write the amplitude M in the form (A.2) we made use of the partial fraction decomposition of the propagator product

$$\frac{1}{(P-k)^2 - m_W^2 + im_W \Gamma_W} \frac{1}{P^2 - m_W^2 + im_W \Gamma_W} = \frac{1}{2 P \cdot k} \left[\frac{k}{(P-k)^2 - m_W^2 + im_W \Gamma_W} - \frac{k}{P^2 - m_W^2 + im_W \Gamma_W} \right] \quad (\text{A.6})$$

and of the factorization property, as discussed in sections I and II. In this decomposition of the amplitude, which is called the radiation decomposition by Brodsky and Brown⁸⁾, it is evident that the amplitude has a zero at

$$F_c = \bar{F}_c' = 0$$

in the gauge theory case, $W=1$.

The squared matrix element, summed (averaged) over final (initial) spin and colors, reads

$$\sum_{\text{spins}} |M|^2 = \frac{(4\pi\alpha)^3}{4\sin^4\theta_W} \left\{ \frac{1}{[(P-k)^2 - m_W^2]^2 + m_W^2 \Gamma_W^2} \left[P \cdot k \frac{A[(P-k)^2 - m_W^2] + B m_W \Gamma_W}{P \cdot k} \right] + \frac{1}{[P^2 - m_W^2]^2 + m_W^2 \Gamma_W^2} \left[\bar{P} \cdot k \frac{A[P^2 - m_W^2] + B m_W \Gamma_W}{P \cdot k} \right] \right\} \quad (\text{A.7})$$

Here $\bar{P}$ denotes the contribution which we labeled "production" and its explicit form is given by:

$$P = \sum_{i=1}^6 \alpha_i H_i \quad (\text{A.8a})$$

where

$$\begin{aligned} \alpha_1 &= 4 \ell \cdot \bar{\ell} \quad , \\ \alpha_2 &= 2 (P \cdot \ell) (P \cdot \bar{\ell}) / (\ell \cdot \bar{\ell}) \quad , \\ \alpha_3 &= 2 (\bar{P} \cdot \ell) (\bar{P} \cdot \bar{\ell}) / (\ell \cdot \bar{\ell}) \quad , \\ \alpha_4 &= 4 (P \cdot \bar{\ell}) (P \cdot \ell) / (\ell \cdot \bar{\ell}) - 2 P \cdot \bar{P} \quad , \\ \alpha_5 &= 2 P \cdot (\bar{\ell} - \ell) \quad , \\ \alpha_6 &= 2 \bar{P} \cdot (\ell - \ell) \quad , \end{aligned} \quad (\text{A.8b})$$

and

$$\begin{aligned}
 H_1 &= \frac{1}{2N} \left\{ F_c^2 \frac{\hat{q}^2 + \hat{u}^2 + 4\hat{q}\hat{u}}{\hat{q}\hat{u}} + F_c(1-\kappa) \frac{\hat{q}-\hat{u}}{\hat{q}+\hat{u}} + (1-\kappa)^2 \frac{\hat{q}\hat{u}}{2(\hat{q}+\hat{u})^2} \right\}, \\
 H_2 &= -\frac{2\hat{q}\hat{u}}{N} \left\{ F_c^2 \frac{4\hat{q}\hat{u}}{\hat{q}\hat{u}} + F_c(1-\kappa) \frac{2\hat{q}\hat{u} + \hat{q}}{\hat{q}(\hat{q}+\hat{u})} + (1-\kappa)^2 \frac{\hat{u}}{2(\hat{q}+\hat{u})^2} \right\}, \\
 H_3 &= -\frac{2\hat{q}\hat{u}}{N} \left\{ F_c^2 \frac{4\hat{q}\hat{u}}{\hat{q}\hat{u}} - F_c(1-\kappa) \frac{2\hat{q}\hat{u} + \hat{u}}{\hat{u}(\hat{q}+\hat{u})} + (1-\kappa)^2 \frac{\hat{q}}{2(\hat{q}+\hat{u})^2} \right\}, \\
 H_4 &= \frac{\hat{q}\hat{u}}{N} \left\{ F_c(1-\kappa) \frac{\hat{q}(\hat{q}-\hat{u})}{\hat{q}\hat{u}(\hat{q}+\hat{u})} - (1-\kappa)^2 \frac{1}{2(\hat{q}+\hat{u})} \right\}, \\
 H_5 &= \frac{\hat{q}\hat{u}}{N} \left\{ F_c^2 \frac{4\hat{q}\hat{u}-2\hat{u}}{\hat{q}\hat{u}} + F_c(1-\kappa) \frac{\hat{q}+2\hat{q}}{\hat{q}(\hat{q}+\hat{u})} + (1-\kappa)^2 \frac{\hat{u}}{2(\hat{q}+\hat{u})^2} \right\}, \\
 H_6 &= \frac{\hat{q}\hat{u}}{N} \left\{ F_c^2 \frac{-4\hat{q}\hat{u}+2\hat{q}}{\hat{q}\hat{u}} + F_c(1-\kappa) \frac{\hat{q}+2\hat{u}}{\hat{u}(\hat{q}+\hat{u})} - (1-\kappa)^2 \frac{\hat{q}}{2(\hat{q}+\hat{u})^2} \right\}.
 \end{aligned} \tag{A.8c}$$

In these expressions we introduced the usual partonic Mandelstam variables

$$\begin{aligned}
 \hat{s} &= (p+\bar{p})^2 \\
 \hat{t} &= (k-p)^2 \\
 \hat{u} &= (k-\bar{p})^2
 \end{aligned} \tag{A.9}$$

and $N = 3$ for QCD.

Similarly the term $\mathbb{D}$ in (A.7) corresponds to the "decay" contribution and its explicit form can be obtained from $\mathbb{I}$ by the following substitutions:

$$\mathbb{D} = \mathbb{I} (p \rightarrow -\bar{l}, \bar{p} \rightarrow -\bar{l}, k \rightarrow k, l \rightarrow -\bar{l}, \bar{l} \rightarrow -\bar{l}, Q_u \rightarrow 1, Q_d \rightarrow 0) \tag{A.10}$$

The quantities A and B in eq. (A.7) refer to the real and imaginary part of the interference term, respectively. Their evaluation is simplified by the use of identities given by Sirlin³¹⁾. One finds for the real part

$$A = 8 F_c \bar{F}_c A_1 + 4 F_c(1-\kappa) A_2 + 4 \bar{F}_c(1-\kappa) A_3 + 4(1-\kappa)^2 A_4 \tag{A.11a}$$

with

$$\begin{aligned}
 A_1 &= -\frac{\bar{l} \cdot g(l, k, \bar{l}) \bar{l} \cdot (l-k)}{l \cdot k \cdot \bar{l}} - \frac{l \cdot \bar{l} \cdot (l+k) l \cdot (\bar{l}-k)}{l \cdot k \cdot \bar{l}} \\
 &+ \left\{ \frac{2\bar{l} \cdot \bar{l} \cdot l \cdot \bar{l} \cdot \bar{l} \cdot l + \bar{l} \cdot l \cdot (l+k) \bar{l} \cdot k + \bar{l} \cdot k \cdot \bar{l} \cdot (l+k) \bar{l} \cdot l - l \cdot \bar{l} \cdot (l+k) \bar{l} \cdot k + \bar{l} \cdot k \cdot l \cdot \bar{l} \cdot (l+k)}{l \cdot k \cdot \bar{l}} \right\} \\
 &+ \langle l \leftrightarrow -\bar{l}, \bar{l} \leftrightarrow -l \rangle \tag{A.11b}
 \end{aligned}$$

$$A_2 = \frac{1}{\bar{l} \cdot k} \left\{ 2l \cdot \bar{l} \cdot l \cdot \bar{l} \cdot (-2\bar{l} \cdot l \cdot \bar{l} + \frac{l \cdot \bar{l}}{l \cdot k} [l \cdot \bar{l} \cdot l \cdot \bar{l} - l \cdot \bar{l} \cdot k \cdot l] - \frac{\bar{l} \cdot l}{\bar{l} \cdot k} [\bar{l} \cdot \bar{l} \cdot l \cdot k - \bar{l} \cdot \bar{l} \cdot k \cdot l]) \right\} \tag{A.11c}$$

$$A_3 = A_2 (l \leftrightarrow -\bar{l}, \bar{l} \leftrightarrow -l) \tag{A.11d}$$

$$A_4 = (p \cdot k)^{-2} [p \cdot k \bar{p} \cdot l \cdot k \cdot \bar{l} + \bar{l} \cdot k l \cdot \bar{l} \cdot k \cdot l] \tag{A.11e}$$

for the imaginary part we find

$$B = 4 \epsilon_{\mu\nu\lambda\delta} p^\mu \bar{l}^\nu l^\lambda k^\delta [2 \bar{p}_c \bar{k}_c \mathcal{B}_1 + \bar{l}_c(1-\kappa) \mathcal{B}_2 + \bar{k}_c(1-\kappa) \mathcal{B}_3] \tag{A.12a}$$

with

$$\mathcal{B}_1 = -\frac{\bar{l} \cdot l + l \cdot \bar{l}}{\bar{l} \cdot k \cdot k \cdot \bar{l}} + \frac{\bar{l} \cdot l + l \cdot \bar{l}}{p \cdot k \cdot l \cdot \bar{l}} \tag{A.12b}$$

$$\mathcal{B}_2 = \frac{1}{p \cdot l} \left\{ \frac{\bar{l} \cdot l}{\bar{l} \cdot k} + \frac{l \cdot \bar{l}}{l \cdot k} \right\} \tag{A.12c}$$

$$\mathcal{B}_3 = \mathcal{B}_2 (l \leftrightarrow -\bar{l}, \bar{l} \leftrightarrow -l) \tag{A.12d}$$

In the case of the zero width approximation there is no interference between the production and the decay process; the terms A and B can thus be dropped safely. The production process will be given by

$$\overline{\Sigma} |M(q\bar{q} \rightarrow W^+ l^-; W^+ \rightarrow l^+ \bar{\nu})|^2 = \frac{(g \pi \alpha)^3}{4 \sin^4 \theta_W} \frac{\pi c}{m_W f_W} \overline{\mathbb{I}} \{ (l \cdot k)^2 - m_W^2 \} \tag{A.13}$$

and similarly for the decay process:

$$\overline{\Sigma} |M(q\bar{q} \rightarrow W^-; W^- \rightarrow l^- \nu)|^2 = \frac{(g \pi \alpha)^3}{4 \sin^4 \theta_W} \frac{\pi c}{m_W f_W} \overline{\mathbb{D}} \{ (l \cdot k)^2 - m_W^2 \}. \tag{A.14}$$

Footnotes

$$\overline{\Sigma} \left| \mathcal{H}(\nu \overline{\nu} \rightarrow W^+ \gamma; \omega \rightarrow \overline{\nu} \nu) \right|^2 = \frac{128 \pi^2 \alpha^2}{\sin^2 \theta_W} \frac{4 F_c [(\overline{P} \cdot \hat{e})^2 + (P \cdot \hat{e})^2]}{\mathcal{B}(\omega \rightarrow \overline{\nu} \nu)} \frac{\mathcal{G}[(P \cdot \hat{e})^2 - \omega_0^2]}{4 \hat{e} \cdot \hat{e}} \quad [A.15]$$

For the decay process we apply on eq. (A.15) the substitution rule (A.10) to get

$$\bar{\Sigma} \left| \mathcal{H}(\bar{q} \bar{q}' \rightarrow W^+ W^- \gamma \bar{\nu}) \right|^2 = \frac{128\pi^4 \alpha^2}{\sin^2 \theta_W} \frac{\bar{H}_c [(\bar{P} \cdot q)^2 + (P \cdot \bar{q})^2]}{B(W \rightarrow \bar{\nu} \nu) (K \cdot q)(K \cdot \bar{q})} \delta(P^2 - m_W^2). \quad (\text{A.16})$$

F.F.n. 3 The following formula (2.18) is obtained when the W^0 system has no transverse motion about the beam axis. In an actual experiment where the W^0 system has non-zero transverse momentum, more refined definition of the variable $\cos \theta_{\pm}^*$ is required to minimize the transverse momentum smearing effects. See discussions in Sect. III and eqs. (3.3a) - (3.3c) for the possible improvement of the variable.

F.n. 4 For first attempts toward the calculation of the higher order QCD corrections, see Ref.22).

References

- 1) UA1 collaboration, Phys. Lett. 122B (1983), 103; ibid 126B (1983), 398.
UA2 collaboration, Phys. Lett. 122B (1983), 476; ibid 129B (1983), 130.
- 2) S.L. Glashow, Nucl. Phys. 22 (1961), 579; S. Weinberg, Phys. Rev. Lett. 19 (1967), 1264; A. Salam, in Elementary Particle Theory, edited by N. Svartholm (Almqvist and Wiksell, Stockholm, 1968), p. 367.
- 3) See, e.g., P.Q. Hung and J.J. Sakurai, Nucl. Phys. B143 (1978), 81; J.D. Bjorken, Phys. Rev. D19 (1979), 335.
- 4) T.D. Lee and C.N. Yang, Phys. Rev. 128 (1962), 885.
- 5) W.A. Bardeen, R. Gastmans and B. Lautrup, Nucl. Phys. B46 (1972), 319.
- 6) See, e.g. R.W. Brown, in Proton-Antiproton Collider Physics - 1981, edited by V. Barger, D. Cline and F. Halzen (AIP, New York, 1982), p. 251.
- 7) K.O. Mikaelian, Phys. Rev. D17 (1978), 750; R.W. Brown, D. Sahdev and K.O. Mikaelian, Phys. Rev. D20 (1979), 1164; K.O. Mikaelian, M.A. Samuel and D. Sahdev, Phys. Rev. Lett. 43 (1979), 746.
- 8) Zhu Dongpei, Phys. Rev. D22 (1980), 2266; C.J. Goebel, F. Halzen and J.P. Leveille, Phys. Rev. D23 (1981), 2682; S.J. Brodsky and R.W. Brown, Phys. Rev. Lett. 49 (1982), 966; R.W. Brown, K.L. Kowalski and S.J. Brodsky, Phys. Rev. D28 (1983), 624; M.A. Samuel, Phys. Rev. D27 (1983), 2724.
- 9) V. Barger, A.D. Martin and R.J.N. Phillips, Phys. Lett. 125B (1983), 339.
- 10) See, e.g., M. Chaichian, M. Hayashi and K. Yamagishi, Phys. Rev. D25 (1982), 130; E26 (1982), 2534.
- 11) S. Drell and T.M. Yan, Ann. of Phys. 66 (1971), 578.
- 12) H. Aronson, Phys. Rev. 186 (1969), 1434; K.J. Kim and Y.S. Tsai, Phys. Rev. D7 (1973), 3710.
- 13) R. Robinett, Phys. Rev. D28 (1983), 1185; K. Hikasa, Phys. Lett. 128B (1983), 253.
- 14) H.D. Politzer, Nucl. Phys. B129 (1977), 301; D. Amati, R. Petronzio and G. Veneziano, Nucl. Phys. B140 (1978), 54; R.K. Ellis, H. Georgi, M. Machacek, H.D. Politzer and G.G. Ross, Nucl. Phys. B152 (1979), 285; A.H. Mueller, Phys. Rev. D18 (1978), 3705.
- 15) A.J. Buras and K.J.F. Gaemers, Nucl. Phys. B132 (1978), 249.
- 16) J.F. Owens and E. Reya, Phys. Rev. D17 (1978), 3003.
- 17) See, e.g., C. Quigg, Rev. Mod. Phys. 49 (1977), 297.
- 18) See, e.g., E.L. Berger, in Proceedings of the Workshop on Drell-Yan-Processes, 1982.
- 19) M. Hellmund and G. Ranft, Z. Phys. C12 (1982), 333.
- 20) S.A. Kahn, I.J. Kifian, M.J. Murtagh and F.E. Paige, "Monte Carlo study of $pp \rightarrow W^+ X$ at CBA", BNL informal report 3/83 (1983).
- 21) T.R. Grose and K.O. Mikaelian, Phys. Rev. D23 (1981), 123.
- 22) N.M. Monyenko and J.H. Reid, in Proc. of Workshop on nonperturbative QCD, edited by K.A. Milton and M.A. Samuel, Oklahoma State Univ., March 1983; M.L. Laursen and M.A. Samuel, Oklahoma State Univ. preprint No. 137 (1982); A. Sen, Phys. Rev. D28 (1983), 650; M.L. Laursen, M.A. Samuel, A. Sen and G. Tupper, Nucl. Phys. B226 (1983), 429; G. Tupper, Phys. Lett. 156B (1985), 406.
- 23) See, e.g., G. Altarelli, M. Greco, and G. Martinelli, Nucl. Phys. B246 (1984), 12; G. Altarelli, R.K. Ellis and G. Martinelli, Phys. Lett. 151B (1985), 457; Z. Phys. C27 (1985), 617.
- 24) UA1 collaboration: G. Arnison et al., Phys. Lett. 136B (1984), 294.
- 25) B.L. Combridge, J. Kripfganz and J. Ranft, Phys. Lett. 70B (1977), 234; R. Cutler and D. Sivers, Phys. Rev. D17 (1978), 196; J.F. Owens, E. Reya and M. Glück, Phys. Rev. D18 (1978), 1501.
- 26) J.C. Collins and D.E. Soper, Phys. Rev. D16 (1977), 2219.
- 27) V. Barger, A.D. Martin and R.J.N. Phillips, Phys. Lett. 125B (1983), 339, 343; Phys. Rev. D28 (1983), 145; R.M. Godbole, S. Pakvasa and D.P. Roy, Phys. Rev. Lett. 50 (1983), 1539; K. Hagiwara and W.F. Long, Phys. Lett. 132B (1983), 202; V. Barger, H. Baer, A.D. Martin and R.J.N. Phillips, Phys. Rev. D29 (1984), 887; V. Barger, H. Baer, K. Hagiwara, A.D. Martin and R.J.N. Phillips, Phys. Rev. D29 (1984), 1923.
- 28) UA1 collaboration: G. Arnison et al., Phys. Lett. 147B (1984), 493.
- 29) V. Barger, H. Baer, K. Hagiwara and R.J.N. Phillips, Phys. Rev. D30 (1984), 947.
- 30) J. Cortés, K. Hagiwara and F. Herzog, Proceedings of 4th Topical Workshop on Proton-Antiproton Collider Physics, (Bern, 1984), 209.
- 31) A. Sirlin, Nucl. Phys. B192 (1981), 93.

Figure captions

Fig.1 Feynman diagrams which contribute to the process $q\bar{q} \rightarrow \gamma \ell \bar{\nu}_\ell$ at tree level.

Fig.2 The p_T distribution of the photon (a) and the charged lepton (b) in the production process $p\bar{p} \rightarrow W^+ \gamma X$; $W^+ \rightarrow \ell \bar{\nu}_\ell$, with cut $p_{\ell T}, p_{\gamma T} > 10$ GeV at $\sqrt{s} = 2$ TeV.

Fig.3 The hadronic center-of-mass rapidity distribution of the photon (a) and the charged lepton (b) in the production process $p\bar{p} \rightarrow W^+ \gamma X$; $W^+ \rightarrow \ell \bar{\nu}_\ell$ with $p_{\ell T}, p_{\gamma T} > 10$ GeV. Curves below 10^{-39} cm²/GeV are smooth interpolations of fluctuating Monte Carlo results.

Fig.4 The photon-lepton invariant mass distribution (a) and the cluster transverse mass distribution (b) for the production process $p\bar{p} \rightarrow W^+ \gamma X$; $W^+ \rightarrow \ell \bar{\nu}_\ell$ (solid lines) and the decay process $p\bar{p} \rightarrow W^+ X$; $W^+ \rightarrow \ell \bar{\nu}_\ell$ (dashed lines) at $\sqrt{s} = 2$ TeV. Both curves are for $\kappa = 1$ and the cuts $p_{\ell T}, p_{\gamma T} > 10$ GeV, $|y_\ell|, |y_\gamma| \leq 3$ and $\cos\theta_{\ell\ell}^* \leq 0.95$ (only for decay) are implemented.

Fig.5 Schematic view of the possible three momentum configuration of a charged lepton, a photon, a neutrino and a W in the frame where $\vec{p}_\ell$ has only a negative x component. None of the three momenta but $\vec{p}_\ell$ necessarily lie in the x-z plane.

Fig.6 The $\cos\theta_{\ell\ell}^*$ (a) and $\cos\theta_{\ell\ell}^*$ (b) distributions of the production process $p\bar{p} \rightarrow W^+ \gamma X$; $W^+ \rightarrow \ell \bar{\nu}_\ell$ at $\sqrt{s} = 2$ TeV. The cuts applied are specified by eqs. (2.12), (2.14). For the definition of $\cos\theta_{\ell\ell}^*$ see eq. (2.18).

Fig.7 The p_T distributions of the photon (a) and the charged lepton (b) in the kinematical region $m_T(\gamma\ell;\bar{\nu}) < 90$ GeV at $\sqrt{s} = 2$ TeV. Contributions from "production" and "decay" subprocesses are shown separately. The final state cuts are specified in eq. (2.24).

Fig.8 The hadronic c.m. rapidity distributions of the photon (a) and the charged lepton (b) in the kinematical region $m_T(\gamma\ell;\bar{\nu}) < 90$ GeV at $\sqrt{s} = 2$ TeV. Contributions from "production" and "decay" subprocesses are shown separately. The final state cuts are specified in eq. (2.24).

Fig.9 The polarization averaged differential W decay distribution integrated over the region $0.3 < x < 1.0$ ($x = 2E_\gamma^*/m_W$) plotted against $\cos\theta_{\ell\ell}^*$ in the W rest frame.

Fig.10 Azimuthal angle $\phi_{\gamma\ell}$ ($\phi_{\gamma\ell}(\vec{p}_{\ell T}, \vec{p}_{\gamma T})$) distribution at $\sqrt{s} = 2$ TeV in the kinematical region $m_T(\gamma\ell;\bar{\nu}) < 90$ GeV. Same specifications as in Fig.7. The distributions are symmetric about $\phi_{\gamma\ell} = \pi$ in our approximation.

Fig.11 Schematic view of the possible three-momentum configuration of the $\ell\bar{\nu}$ final state in the frame where $\vec{p}_\ell + \vec{p}_\gamma$ has no longitudinal component. In a typical radiative decay event, $\vec{p}_{\ell T}, \vec{p}_{\gamma T}$ roughly balances this momentum. The two possible solutions for the neutrino three-momentum $\vec{p}_{\bar{\nu}}$ are also shown.

Fig.12 $\cos\theta_{\ell\ell}^*$ distribution (see. eq.(2.30)) at $\sqrt{s} = 2$ TeV in the kinematical region $m_T(\gamma\ell;\bar{\nu}) < 90$ GeV. Same specifications as in Fig.7. The production process contribution for $\kappa=1$ is the only "background" contaminating the decay contribution substantially, especially in the region of the angular zero $\cos\theta_{\ell\ell}^* = -1$.

Fig.13 $\cos\theta_{\ell\ell}^*$ -distribution for the production process $p\bar{p} \rightarrow W^+ \gamma X$; $W^+ \rightarrow \ell \bar{\nu}_\ell$ at $\sqrt{s} = 2$ TeV for different values of κ . The cuts are specified by eqs. (2.12) and (2.14). The dashed lines denote the estimated background from $p\bar{p} \rightarrow W^+ \gamma X$ where " γ " represents a jet that may fake photons.

Fig.14 $\cos\theta_{\ell\ell}^*$ distribution at $\sqrt{s} = 2$ TeV for different values of κ in the kinematical region $m_T(\gamma\ell;\bar{\nu}) < 90$ GeV. The contributions from decay and production processes which were shown separately in Fig.12 are added together. The cuts are specified by eq. (2.24). The dashed lines denote the estimated background from $p\bar{p} \rightarrow W^+ \gamma X$, where " γ " represents a jet that may fake photons.

39021

Fig. 1

39026

Fig. 2

39027

Fig. 3

39013

Fig. 4

Fig. 6

39012

Fig. 5

39032

Fig. 7

39014

Fig. 8

39015

Fig. 9

Fig. 10

39024

Fig. 12

39033

Fig. 11

39018

Fig. 13

39017

Fig. 14