INVESTIGATION OF THE W AND Q^2 DEPENDENCE OF CHARGED PION DISTRIBUTIONS
IN μp SCATTERING

The European Muon Collaboration

Aachen¹, CERN², DESY (Hamburg)³, Freiburg⁴, Hamburg (University)⁵,
Kiel⁶, LAL (Orsay)⁷, Lancaster⁸, LAPP (Annecy)⁹, Liverpool¹⁰,
Marseille¹¹, Mons¹², MPI (München)¹³, Oxford¹⁴, RAL (Chilton)¹⁵,
Sheffield¹⁶, Torino¹⁷, Uppsala¹⁸ and Wuppertal¹⁹

M. Arneodo¹⁷, A. Arvidson¹⁸, J.J. Aubert¹¹, B. Badelek^{18a)}, J. Beaufays²,
C.P. Bee^{8b)}, C. Benchouk¹¹, G. Berghoff¹, I. Bird⁸, D. Blum⁸,
E. Böhm⁶, X. de Bouard⁹, F.W. Brasse³, H. Braun¹⁹, C. Broll^{9a)},
S. Brown¹⁰, H. Brück^{19,k)}, H. Calen¹⁸, J.S. Chima¹⁵, J. Ciborowski^{5a)},
R. Clifft¹⁵, G. Coignet⁹, F. Combley¹⁶, J. Coughlan^{8b)}, G. D'Agostini¹¹,
S. Dahlgren¹⁸, F. Dengler¹³, I. Derado¹³, T. Dreyer⁴, J. Drees¹⁹,
M. Düren¹, V. Eckardt¹³, A. Edwards^{19c)}, T. Ernst⁴, G. Eszes^{9d)},
J. Favier⁹, M.I. Ferrero¹⁷, J. Figiel^{5a)}, W. Flauger³, J. Foster¹⁶,
E. Gabathuler¹⁰, J. Gajewski^{5a)}, R. Gamet¹⁰, J. Gayler³, N. Geddes¹⁴,
P. Giubellino¹⁷, P. Grafström¹⁸, F. Grard¹², J. Haas⁴, E. Hagberg¹⁸,
F.J. Hasert¹, P. Hayman¹⁰, P. Heusse⁷, C. Hoppe⁵, M. Jaffré⁷,
A. Jacholkowska⁷, F. Janata⁵, G. Jancso^{9d)}, A.S. Johnson^{14f)}, E.M. Kabuss⁴,
G. Kellner², V. Korbel³, J. Krüger¹⁹, S. Kullander¹⁸, U. Landgraf⁴,
D. Lanske¹, J. Loken¹⁴, K. Long^{14g)}, M. Maire⁹, P. Malecki¹³,
A. Manz¹³, S. Maselli¹³, W. Mohr⁴, F. Montanet¹¹, H.E. Montgomery^{2h)},
E. Nagy^{9d)}, J. Nassalski¹³ⁱ⁾, P.R. Norton¹⁵, F.G. Oakham¹⁵, A.M. Osborne²,
C. Pascaud⁷, B. Pawlik^{13e)}, P. Payre¹¹, C. Peroni¹⁷, H. Pessard⁹,
J. Pettingale¹⁰, B. Pietrzyk¹¹, B. Pönsgen⁵, M. Pötsch¹⁹, P. Renton¹⁴,
P. Ribarics^{9d)}, K. Rith⁴, E. Rondio^{5a)}, M. Scheer¹, A. Schlagböhmer⁴,
H. Schiemann⁵, N. Schmitz¹³, M. Schneegans⁹, M. Scholz¹, T. Schröder⁴,
M. Schouten¹³, K. Schultze¹, T. Sloan⁸, H.E. Stier⁴, M. Studt⁵,
G.N. Taylor¹⁴, J.M. Thénard⁹, J.C. Thompson¹⁵, A. de la Torre⁵,
J. Toth^{9d)}, L. Urban¹, L. Urban^{9d)}, W. Wallucks⁴, M. Whalley^{16j)},
S. Wheeler¹⁶, W.S.C. Williams¹⁴, S.J. Wimpenny^{10g)}, R. Windmolders¹²,
G. Wolf¹³.

(submitted to Zeitschrift für Physik C)

Abstract:

The W and Q^2 dependence of the fragmentation functions and of the average multiplicity of charged pions is investigated, using data from the NA9 experiment at the CERN SPS on muon-proton scattering at 280 GeV. A significant increase of pion production with increasing W is observed at fixed Q^2 , leading to a rise of the average charged pion multiplicity, linear in $\ln W^2$, and of the pion fragmentation function in the central region, i.e. at small $|x_F|$. This increase can be understood from the kinematic widening of the cms rapidity range proportional to $\ln W^2$ and the observed W independent height of the rapidity distribution. At fixed W , a rise of the average charged pion multiplicity with Q^2 is observed. This rise appears to be weaker than that observed for all charged hadrons implying a stronger rise with Q^2 for kaons and protons.

- a) Now at University of Warsaw, Poland.
- b) Now at RAL, Chilton, Didcot, England.
- c) Now at Jet, Joint Undertaking, Abingdon, England.
- d) Now at Central Research Institute for Physics of the Hungarian Academy of Science, Budapest, Hungary.
- e) Now at Institute of Nuclear Physics, Krakow, Poland.
- f) Now at SLAC, Stanford, California, USA.
- g) Now at CERN, Geneva, Switzerland.
- h) Now at FNAL, Batavia, Illinois, USA.
- i) Now at Institute for Nuclear Studies, Warsaw, Poland.
- j) Now at University of Durham, England.
- k) Now at DESY, Hamburg, Germany.
- +) Deceased.

1.) Introduction

In this paper we investigate the W and Q^2 dependence of the fragmentation functions of charged pions produced in deep inelastic muon proton scattering,

$$\mu^+ p \rightarrow \mu^+ + \text{hadrons.} \quad (1)$$

In the quark parton model QPM (Fig. 1) this process is described by the scattering of the muon, via virtual photon (γ_v) exchange, off a constituent quark (or antiquark) in the proton. In the centre of mass system (cms) of all final state hadrons the scattered quark goes forward (i.e. in the virtual photon direction) and the proton remnant goes backward. In the simplest case of scattering off a valence quark (Fig. 1), the target remnant is a diquark. The scattered quark and the target remnant then fragment into the final state hadrons, which can be classified as current fragments or target fragments respectively. The fragmentation of quarks and diquarks into hadrons of type h (here π^+, π^-) is described by the quark or diquark fragmentation functions, $D_q^h(x_F)$ or $D_{qq}^h(x_F)$, respectively.

The kinematic variables used to describe a leptonproduction event are: W (the total hadronic energy in the hadronic cms), Q^2 (the four-momentum transfer squared) and $x = Q^2/(2M\nu)$ (Bjorken- x), where $\nu = E_\mu - E'_\mu$, E_μ and E'_μ being the laboratory energy of the incident and outgoing muons, respectively, and M is the proton mass. The variables W , Q^2 and x are kinematically related via

$$W^2 = M^2 + Q^2 \left(\frac{1}{x} - 1 \right). \quad (2)$$

Thus there are two independent event variables, e.g. W and Q^2 . This is in contrast to e^+e^- annihilation, where $W^2 = Q^2 = s$.

The variables used to describe a final state hadron are: $z = E/\nu$ (the energy fraction carried by the hadron, $z > 0$), $x_F = p_L^*/p_{L\max}^*$ (Feynman x , the cms longitudinal momentum fraction with $-1 \leq x_F \leq 1$), $y = 0.5 \cdot \ln[(E^* + p_L^*)/(E^* - p_L^*)]$ (the cms rapidity) and p_T (the transverse momentum with respect to the virtual photon direction). Here E , E^* and p_L^* are the laboratory energy, the cms energy and the cms longitudinal momentum of the hadron, respectively, and $p_{L\max}^*$ the maximum kinematic value of p_L^* . For high W , the kinematic correlation between z and x_F is such that backward going hadrons ($x_F < 0$) have small z values, whereas for large positive x_F the two variables x_F and z are roughly equal.

The measured fragmentation function $D^h(x_F, x, Q^2)$ of hadrons h (normalized x_F distribution) is defined as

$$D^h(x_F, x, Q^2) = \frac{1}{N_{ev}(x, Q^2)} \cdot \frac{dN^h}{dx_F}(x_F, x, Q^2) \quad (3)$$

where $N_{ev}(x, Q^2)$ is the number of events at (x, Q^2) and N^h the number of hadrons h in these events. In general, a fragmentation function measured in leptonproduction depends, apart from being a function of the particle variable (e.g. x_F), on any pair of event variables via eq. (2). The fragmentation functions can also depend on the scaling variable v/E_μ . However a recent study [1] has shown that such a dependence, if it exists, is small and is neglected here. The fragmentation function can be decomposed into quark or diquark fragmentation functions, depending on whether the hadron h is a current (quark) or target (diquark) fragment. For instance for μp -scattering, if h is a current fragment:

$$D^h(x_F, x, Q^2) = \frac{\sum_q e_q^2 [q(x, Q^2) D_q^h(x_F, x, Q^2) + \bar{q}(x, Q^2) D_{\bar{q}}^h(x_F, x, Q^2)]}{\sum_q e_q^2 [q(x, Q^2) + \bar{q}(x, Q^2)]} \quad (4)$$

where $q(x, Q^2)$ and $\bar{q}(x, Q^2)$ are the quark and antiquark momentum distribution functions in the proton and e_q the quark charges. In the simple QPM where scaling (no Q^2 dependence) and factorization (no x dependence) hold, the quark and antiquark fragmentation functions depend only on x_F (or z).

The simplest method of selecting the current and target fragments is by selecting as current (target) fragments those hadrons which go forward ($x_F > 0$) (backward, $x_F < 0$) in the hadronic cms. However this separation is incomplete and at currently available values of W there is a sizeable kinematic overlap of the two types of fragments. The fragmentation process can best be described as follows: The forward-going quark and backward-going remnant in the partonic final state fragment into a continuous chain of hadrons, e.g. by the creation of quark-antiquark pairs from the colour field, which then rearrange themselves into hadrons. This leads to a cms rapidity distribution which at sufficiently high W is subdivided into a current fragmentation region (large positive y), a target fragmentation region (large negative y) and a central region, which transforms into a region of small $|x_F|$. The total kinematic rapidity range Δy increases like $\ln W^2$:

$$\Delta y = \ln \frac{W^2}{m_n \cdot m_\pi} = \ln(W^2/\text{GeV}^2) + 2.03 \quad (5)$$

(calculated for the simplest reaction $\mu p \rightarrow \mu n \pi^+$, where m_n and m_π are the neutron and pion masses respectively). At small W , the central region has not yet developed and the two fragmentation regions overlap. In any case there is no clear separation between the current and target fragments at finite W but rather a continuous transition in the central region. Thus the separation between the two types of fragments at $x_F = 0$ (i.e. $y = 0$) is only an approximation which improves as W increases. Therefore, a W dependence may occur in the measured quark or diquark fragmentation functions due to the W dependent overlap and separation problem.

Apart from the kinematic overlap just discussed, several physical effects may lead to a W and/or Q^2 dependence of the fragmentation functions, e.g.

- i) A W dependence is expected from the increase of particle production in the central region due to the kinematic widening of the rapidity range according to eq. (5). In fact it is known [2,3] that in leptonproduction the average particle multiplicity $\langle n \rangle$, which is the integral of the x_F -distribution (fragmentation function),

$$\langle n \rangle(W) = \int_{-1}^{+1} D(x_F, W) dx_F, \quad (6)$$

increases linearly with $\ln W^2$. This increase could be due to the above mentioned widening of the rapidity range with W . If the height of the rapidity distribution is roughly W independent in the central region, one would then expect that the x_F -distribution $D(x_F, W)$ increases with W at small $|x_F|$.

A particular model for the increase of the particle number with increasing W is the LUND string model [4] which describes the non-perturbative fragmentation of partons into hadrons.

- ii) A Q^2 dependence (scaling-violation) is predicted by perturbative QCD [5-7] and higher twist contributions [8]. For the quark fragmentation functions $D_q^h(z, Q^2)$ this Q^2 dependence is described in leading-order QCD by the Altarelli-Parisi equations [9]. In next to leading order factorization [6] no longer holds and the fragmentation functions also depend on x , i.e. $D(z, x, Q^2)$.
- iii) An x dependence is expected since the fragmentation of the various quark flavours contributes to the measured x_F -distribution with weights given

by the x dependent distribution functions, according to eq. (4). This x dependence implies a W or Q^2 dependence via eq. (2). In certain combinations of x_F -distributions such an x dependence can cancel, e.g. in the x_F -distribution $D^\pi = D^{\pi^+} + D^{\pi^-}$ of all charged pions, neglecting strange and charmed sea quarks:

$$D^{\pi^+} + D^{\pi^-} = D_u^{\pi^+} + D_u^{\pi^-}, \quad (7)$$

using eq. (4), the isospin relation $D_u^{\pi^\pm} = D_d^{\pi^\mp}$ and charge conjugation invariance. Therefore, to eliminate (or at least reduce) a W and/or Q^2 dependence originating from the quark distribution functions and not from the quark fragmentation functions themselves, we investigate below in section 3 the x_F -distributions of all charged pions.

The dependence of fragmentation functions on W and/or Q^2 has been studied previously in various νN , $\bar{\nu} N$ [10] and μp [11,12] scattering experiments. The results may be summarized as follows: in νN and $\bar{\nu} N$ scattering, a Q^2 dependence is observed if W is not restricted to a narrow range of values. In some of these studies this Q^2 dependence could be approximately described by leading-order perturbative QCD. If W is restricted, there is no clear Q^2 dependence. These observations and direct studies of the W dependence seem to indicate that, in the energy ranges accessible at present ($W \lesssim 15$ GeV), the fragmentation functions depend mainly on W and not on Q^2 , the distributions becoming narrower with increasing W . The Q^2 dependence as observed, when no restriction on W is applied, would then be a kinematic reflection of this explicit W dependence due to eq. (2), implying on average an increase of Q^2 with increasing W and vice versa.

In μp scattering, on the other hand, where a wider energy range ($W \lesssim 20$ GeV) is accessible, this collaboration has recently observed [12] a weak, but statistically significant Q^2 dependence of the z distributions of charged hadrons and of K^0 mesons also at fixed W . This Q^2 dependence occurs at small values of z and leads to an increase of the average multiplicities with Q^2 at fixed W .

The main purpose of this paper is to investigate further the explicit dependence of the fragmentation functions $D^\pi(x_F, W, Q^2)$ and average multiplicities $\langle n_\pi \rangle(W, Q^2)$ of charged pions produced in μp scattering on W and/or Q^2 . In particular the question arises as to whether the Q^2 dependence, which was recently observed in ref. [12] for all charged hadrons and for

K^0 mesons, is also seen for charged pions only. To answer these questions we study below $\langle n_\pi \rangle(W, Q^2)$ and $D^\pi(x_F, W, Q^2)$ as a function of W at fixed Q^2 and as a function of Q^2 at fixed W , thus avoiding the kinematic correlation between W and Q^2 .

2.) Experimental procedure

a.) Experimental set-up and data analysis

The data used for this analysis were taken by the EMC(NA9) experiment in the M2 muon beam at the CERN SPS, at an incident muon energy of 280 GeV. The apparatus consists of a double magnetic spectrometer [13,14].

A vertex spectrometer magnet (VSM) contains a 1m long liquid hydrogen target inside a streamer chamber. Tracks leaving the streamer chamber are detected in a set of proportional chambers and drift tubes. The scattered muon and fast hadrons enter the forward spectrometer magnet (FSM) which has a bending sense opposite to the VSM. A hodoscope trigger system allows the selection of scattered muons with angles greater than 0.5° . Proportional and drift chambers placed inside and downstream of the FSM provide the track detection for fast forward particles. The complete system detects tracks down to laboratory momenta of 200 MeV/c and gives almost 4π coverage in the hadronic cms.

The particle identification system consists of a set of two gas Cerenkov counters (one filled with neopentane, the other with nitrogen), two aerogel Cerenkov counters, a time of flight hodoscope system for low momentum particles, and a large Cerenkov counter filled with neon to identify fast particles in the forward spectrometer. This set-up allows the identification of roughly 50% of all charged hadrons. Details of the off-line pattern recognition, track and vertex fitting algorithms are given in ref. [14].

In order to restrict the event sample to regions where the corrections for event acceptance, smearing effects from the resolution of the apparatus, and radiative corrections are relatively small ($\leq 10\%$), the following selection criteria were applied to the data:

$$\begin{aligned} Q^2 &> 4 \text{ GeV}^2 \\ 16 \text{ GeV}^2 &< W^2 < 400 \text{ GeV}^2 \\ 20 \text{ GeV} &< \nu < 260 \text{ GeV} \end{aligned}$$

$$\begin{aligned} E_{\mu}' &> 20 \text{ GeV} \\ v/E_{\mu} &< 0.9 \\ \Theta_{\mu}' &> 0.5^{\circ} \end{aligned}$$

where Θ_{μ}' is the angle between the incident and the scattered muon in the laboratory system. The number of events remaining after these cuts was about 28000.

In the following, distributions of charged pions will be shown. All tracks from the primary vertex were taken as pions unless they were identified as electrons, kaons or protons in the Cerenkov or time of flight counters. The errors resulting from misidentification were then corrected. Since at the energies of this experiment about 80% of the final state hadrons are pions, only relatively small corrections are needed (see below). The correction procedure is described in the subsequent section.

This method of producing corrected pion distributions is different from that applied in a previous publication [15], where only pions identified in the Cerenkov and/or time of flight counters have been used and the data were then corrected by relatively large correction factors for the limited efficiency and acceptance of these counters.

b.) Correction method

The data presented below were corrected for the effects of limited acceptance, identification inefficiency, smearing introduced by the resolution of the apparatus, and inefficiencies of the various off-line processes such as vertex fitting and track reconstruction. These corrections were determined by a complete Monte-Carlo (MC) simulation of the experiment. Deep inelastic scattering events were generated using the LUND String Model [4] for the fragmentation process. Radiative effects due to QED processes and secondary interactions of the produced hadrons in the target material were taken into account. The scattered muon and the secondary hadrons were tracked through the apparatus and the effects of multiple Coulomb scattering were simulated. The effects of chamber and hodoscope inefficiencies and smearing were also included and the resulting computed coordinates of the simulated measurements in each detector were passed through the reconstruction program chain. In this manner the inefficiencies of the apparatus and losses due to imperfections in the off-line analysis were fully included. Then, each distribution was produced

three times: from the data (D_{DA}), from the generated MC events (D_{MC}), and from the MC events after the experimental simulation (D_{mod}). The corrected distribution D was then determined for a given bin as

$$D = D_{DA} \cdot \frac{D_{MC}}{D_{mod}} \quad (8)$$

The values of the correction function $\frac{D_{MC}}{D_{mod}}$ depend on the kinematic variables and on x_F or z . As an example Fig. 2 shows the correction function for positive and negative pions as a function of x_F , for all events with $Q^2 > 4 \text{ GeV}^2$ and $16 < W^2 < 400 \text{ GeV}^2$. The shaded areas indicate the statistical uncertainty which is large only in x_F regions where few particles occur in the data. The weighted mean of the correction function in the forward hemisphere is ~ 0.9 for both π^+ and π^- , whereas it is smaller for π^+ (~ 0.69) than for π^- (~ 0.87) in the backward hemisphere. This is expected due to the presence of unidentified protons in the backward hemisphere.

The errors on the experimental data presented below have been calculated from the statistical errors of the data and Monte Carlo samples. Systematic effects have been studied using different track selection criteria, and identification and correction methods. They could result in an additional normalisation uncertainty of up to 20%.

3.) Results

a.) Average multiplicities of charged pions

Fig. 3 shows the average multiplicity $\langle n_\pi \rangle$ of charged pions vs. W^2 for fixed intervals of Q^2 . It is seen that $\langle n_\pi \rangle$ rises linearly with $\ln W^2$ and that, at fixed W , it depends weakly on Q^2 . The following function has been fitted to the data points of Fig. 3:

$$\langle n_\pi \rangle = a + b \cdot \ln(W^2/\text{GeV}^2) \quad (9)$$

The results of the fit are given in Table 1. The systematic uncertainty on the parameter b was estimated to be $\sim 10\%$. The values of the slope b for the various Q^2 intervals are compatible with each other, whereas the values of a seem to show an increase with increasing Q^2 .

A W dependence is also observed (not shown) for π^+ and π^- in the forward

($x_F > 0$) and backward ($x_F < 0$) hemispheres separately. The results of the various fits of form (9) are given for all Q^2 in Table 2.

In order to investigate the Q^2 dependence of $\langle n_\pi \rangle$ at fixed W , we show in Fig. 4a $\langle n_\pi \rangle$ vs. Q^2 for fixed intervals of W . To keep W fixed and thus to exclude any reflection of the W dependence via eq. (2), the total W range has been subdivided into a relatively large number of intervals and correspondingly, because of finite statistics, only four Q^2 intervals have been chosen. It is seen that at fixed W there is a weak, but statistically significant Q^2 dependence of the average pion multiplicity, analogous to that observed previously for all charged hadrons and for K^0 mesons in this experiment [12]. The results of fits of the form

$$\langle n_\pi \rangle = c + d \cdot \ln(Q^2/\text{GeV}^2) \quad (10)$$

for the various W intervals are given in Table 3a. The slope d , which has a systematic error of ~ 0.07 , is seen to be statistically significantly different from zero; the variation of the parameter c with W shows the above mentioned strong W dependence of $\langle n_\pi \rangle$.

Figs. 4b and 4c show the average multiplicity of π^+ and π^- separately, as a function of Q^2 for the fixed intervals of W . The results of the corresponding linear fits of the form (10) are listed in Tables 3b and 3c.

A simultaneous fit to the W^2 and Q^2 dependence of the form

$$\langle n_\pi \rangle = a_1 + a_2 \cdot \ln W^2 + a_3 \cdot \ln Q^2 \quad (11)$$

has been carried out to the data of Fig. 4a. The fitted values for the parameters a_1 , a_2 and a_3 and the value of χ^2/NDF are listed in Table 4. A correlation term $\ln W^2 \cdot \ln Q^2$ is not needed to obtain a good fit, although the data points in Fig. 4a and the d values in Table 3a indicate a weakening of the Q^2 dependence with increasing W .

Table 4 also lists the corresponding values obtained in ref. [12] for all charged hadrons. The coefficient a_3 , i.e. the Q^2 dependence of the average multiplicity is seen to be larger for all charged hadrons than for charged pions only. From this one may indirectly conclude that the Q^2 dependence of the average charged hadron multiplicity gets a substantial contribution from protons, antiprotons and charged kaons. This is expected, if the Q^2 dependence is a QCD effect, i.e. due to the emission of hard gluons, and if gluons fragment more readily to kaons and baryons.

The observed Q^2 dependence can have two possible sources. Firstly, varying Q^2 at fixed W implies varying Bjorken- x (eq. (2)). The Q^2 dependence could thus come from the x and Q^2 dependent quark and antiquark distribution functions, which in the charged pion distribution and average multiplicity occur as weights to the quark or diquark fragmentation functions according to eq. (4). In the forward direction however, for all charged pions these weights almost cancel as discussed in section 1. Secondly, the observed Q^2 dependence could be directly due to a Q^2 dependence (scaling violation) of the quark or diquark fragmentation functions themselves, as predicted by QCD.

b.) x_F distributions of charged pions

We next examine the question of where in the x_F range the observed (Fig. 3) increase of charged pion production with increasing W occurs. Fig. 5 shows, for fixed intervals of x_F , the x_F -distribution $D^\pi(x_F, W^2, Q^2)$ of charged pions vs. W^2 for fixed intervals of Q^2 , and vs. Q^2 for fixed intervals of W^2 . The distributions are shown separately for forward ($x_F > 0$, Fig. 5a) and backward ($x_F < 0$, Fig. 5b) going charged pions. Two features are recognized from the plots:

- Independently of Q^2 , the distribution $D^\pi(x_F, W^2, Q^2)$ rises with W^2 at small $|x_F| \lesssim 0.1$, both in the forward ($x_F > 0$) and backward ($x_F < 0$) hemispheres. For larger $|x_F|$, on the other hand, no W dependence is observed. (No clear statement can be made for $|x_F| \gtrsim 0.6$ because of the larger statistical errors). Thus the increase of pion production with increasing W occurs predominantly in the central region.
- At fixed W and at any x_F the distribution $D^\pi(x_F, W^2, Q^2)$ is seen to be practically independent of Q^2 within the errors. The explicit Q^2 dependence observed in the average pion multiplicity (Fig. 4) is too weak to be clearly recognized in this distribution.

c.) Rapidity distributions of charged pions

The observed increase of pion production, which is proportional to $\ln W^2$ (Fig. 3) and occurs at small $|x_F|$ (Fig. 5), i.e. in the central region, could be due to the kinematic widening of the rapidity range, in conjunction with a roughly W independent central height of the rapidity distribution (see section 1). We therefore study in this section the cms rapidity distribution

$$D^{\pi}(y, W^2) = \frac{1}{N_{ev}(W^2)} \frac{dN^{\pi}}{dy}(y, W^2) \quad (12)$$

of charged pions directly.

Fig. 6 shows the rapidity distributions of all charged pions and separately of π^+ and π^- for 4 intervals of W^2 and $Q^2 > 4 \text{ GeV}^2$. It is seen qualitatively that the widths of the distributions increase with W^2 and that for higher W^2 a central plateau region starts to develop. Furthermore their central height $D^{\pi}(y=0, W^2)$ is roughly independent of W^2 for $W^2 > 16 \text{ GeV}^2$. Finally the distributions are not quite symmetric around $y=0$, i.e. they are somewhat different in the current and target fragmentation region.

Comparing the π^+ and π^- distributions in Fig. 6 it can be seen that they are rather similar in height at larger W^2 , whereas in the low W^2 range the π^+ distribution is higher. This is easily understood from the fact that in the central region (neglecting rare particles like kaons or antiprotons) π^+ and π^- mesons are produced in pairs in reactions like $\mu p \rightarrow \mu p + m(\pi^+\pi^-)$ and $\mu p \rightarrow \mu n \pi^+ + m(\pi^+\pi^-)$. The extra π^+ in the second reaction leads to an excess of π^+ in the distribution; its relative significance decreases with increasing energy, i.e. with increasing number of $\pi^+\pi^-$ pairs. Furthermore the π^+ distribution is seen to be somewhat broader than the π^- distribution in both hemispheres. This is probably due to the effect of charge retention as expected from the QPM [16].

In order to determine the width and height of the rapidity distributions more quantitatively, fits have been carried out to the measured distributions in Fig. 6, using an empirical fitting function for smoothing. As examples, the curves in Fig. 6 show the fits to the distributions for all charged pions. Because of the asymmetry of the rapidity distributions the fits were carried out in the forward and backward hemisphere separately. The fitted forward height turned out to be greater than the backward height by $\sim 10\%$ for all W intervals. From the two fits the width and central height of each distribution were then determined; for the height the mean of the two values obtained from the forward and the backward fit was taken.

The resulting width (at 1%, 5% and 50% of the total height) and central height of the rapidity distribution of charged pions are shown in Fig. 7 as functions of W^2 . The points are plotted at W^2 values which are averages over the events in each of the four W^2 intervals of Fig. 6. As observed already

qualitatively, the central height of the distribution is practically independent of W , whereas the width increases linearly with $\ln W^2$ as expected kinematically. In fact the width at 1% of the total height agrees rather well with the maximum kinematically allowed value given by eq. (5) and shown in Fig. 7 by the dashed straight line. (The small excess of the experimental points is probably due to the smearing of the rapidity distributions near the kinematic boundaries because of measurement errors). The corresponding results for π^+ and π^- separately are plotted in Fig. 8. The solid lines in Figs. 7 and 8 are fits linear in $\ln W^2$, the coefficients of which are listed in Table 5.

Thus, this analysis shows that the cms rapidity distribution widens with increasing W as $\ln W^2$, occupying the whole kinematically allowed rapidity range, and has an approximately W independent height. Both features together lead directly to the observed increase of the average pion multiplicity linear in $\ln W^2$ and to the observed rise of the x_F distribution at small $|x_F|$.

4.) Summary and conclusions

In this paper we have investigated the W and Q^2 dependence of fragmentation functions (x_F or z distributions) $D^\pi(x_F, W, Q^2)$ of charged pions produced in μp interactions at 280 GeV. In order to avoid the correlation between W and Q^2 due to the kinematic relationship (2), the distribution $D^\pi(x_F, W, Q^2)$ and its integral (= average multiplicity) have been studied as functions of W at fixed Q^2 , and as functions of Q^2 at fixed W .

The following results were obtained for the W dependence:

- At fixed Q^2 the average charged pion multiplicity $\langle n_\pi \rangle$ rises linearly with $\ln W^2$. This observation holds for both π^+ and π^- mesons, and for the forward and backward hemisphere separately.
- This increase of pion production with increasing W occurs in the central region: The charged pion fragmentation function $D^\pi(x_F, W, Q^2)$ rises (at constant Q^2) with W for small $|x_F| \lesssim 0.1$, but stays constant at higher $|x_F|$, both in the forward and backward hemispheres.
- A simple consistent picture for the observed increase can be obtained from a direct study of the cms rapidity distribution $D^\pi(y, W)$ of charged pions: The width of the distribution grows like $\ln W^2$ as expected kinematically,

whereas its height is practically W independent. This then leads directly to the $\ln W^2$ increase of the average pion multiplicity.

As for the Q^2 dependence, the following results were obtained:

- At fixed W , a weak but statistically significant rise of the average pion multiplicity with increasing Q^2 is observed. This Q^2 dependence is too weak to be seen, within present statistics, in the differential charged pion fragmentation function $D^\pi(x_F, W, Q^2)$ when studied at fixed W and x_F as a function of Q^2 .
- The Q^2 dependence of the average multiplicity is weaker for charged pions than previously observed for all charged hadrons. This may indicate that a sizeable contribution to this Q^2 dependence comes from protons, anti-protons and charged kaons.
- There are two possible origins of the observed Q^2 dependence. Firstly, it could be a consequence of the x (and Q^2) dependence of the quark and antiquark distribution functions, which in $D^\pi(x_F, W, Q^2)$ occur as weights to the quark or diquark fragmentation functions. This effect however is expected to largely cancel in the distribution and average multiplicity of all charged pions in the forward hemisphere. Secondly, the observed Q^2 dependence could be a direct consequence of a Q^2 dependence (scaling violation) of the quark or diquark fragmentation functions themselves, as predicted by QCD.

Acknowledgements

We would like to thank all the people in the various laboratories who have contributed to the construction, operation and analysis of this experiment. The support of the CERN staff in operating the SPS, muon beam and computer facilities is gratefully acknowledged.

References

- 1.) EMC, J.J. Aubert et al.: Preprint CERN-EP/85-72
- 2.) N. Schmitz: Review in Proc. 1981 Intern. Symp. on Lepton and Photon Interactions at High Energies, Bonn, p. 527 (1981)
S. Barlag et al.: Z.Phys. C11, 283 (1982); C14, 281 (1982) (erratum)
H. Grässler et al.: Nucl. Phys. B223, 269 (1983)
- 3.) EMC, M. Arneodo et al.: Nucl. Phys. B258, 249 (1985)
- 4.) B. Andersson et al.: Phys. Rep. 97, 31 (1983)
- 5.) J.F. Owens: Phys. Lett. 76B, 85 (1978)
T. Uematsu: Phys. Lett. 79B, 97 (1978)
- 6.) N. Sakai: Phys. Lett. 85B, 67 (1979)
G. Altarelli et al.: Nucl. Phys. B160, 301 (1979)
R. Baier, K. Fey: Z.Phys. C2, 339 (1979)
- 7.) G. Altarelli: Phys. Rep. 81, 1 (1982)
G. Ingelman: Z. Phys. C26, 483 (1984)
- 8.) E.L. Berger: Phys. Lett. 89B, 241 (1980); Z.Phys. C4, 289 (1980)
- 9.) G. Altarelli, G. Parisi: Nucl. Phys. B126, 298 (1977)
- 10.) J. Blietschau et al.: Phys. Lett. 87B, 281 (1979)
P. Allen et al., Phys. Lett. 96B, 209 (1980)
M. Derrick et al.: Phys. Lett. 91B, 470 (1980)
H. Deden et al.: Nucl. Phys. B198, 365 (1982)
D. Allasia et al.: Phys. Lett. 124B, 543 (1983)
WA21 collaboration: Proc. 15 Intern. Symp. on Multiparticle Dynamics, Lund, p. 764 (1984)
- 11.) EMC, J.J. Aubert et al.: Phys. Lett. 114B, 373 (1982)
- 12.) EMC, M. Arneodo et al.: Phys. Lett. 165B, 222 (1985)
- 13.) EMC, O.C. Alkofer et al.: Nucl. Instr. and Meth. 179, 445 (1981)
- 14.) EMC, J.P. Albanese et al.: Nucl. Instr. and Meth. 212, 111 (1983)
- 15.) EMC, M. Arneodo et al.: Phys. Lett. 150B, 458 (1985)
- 16.) EMC, J.P. Albanese et al.: Phys. Lett. 144B, 302 (1984)

Table 1: Fitted values of the coefficients a and b in the expression $\langle n_{\pi} \rangle = a + b \cdot \ln(W^2/\text{GeV}^2)$ for the average charged pion multiplicity, for different intervals of Q^2 (in GeV^2).

	a	b	χ^2/NDF
$4 < Q^2 < 10$	-0.68 ± 0.35	1.09 ± 0.06	4.3/6
$10 < Q^2 < 20$	-0.45 ± 0.18	1.04 ± 0.04	4.1/6
$20 < Q^2 < 50$	-0.06 ± 0.23	0.99 ± 0.05	4.3/6
$Q^2 > 50$	0.44 ± 0.63	0.91 ± 0.13	1.9/6
$Q^2 > 4$	-0.33 ± 0.13	1.03 ± 0.03	2.4/6

Table 2: Fitted values of the coefficients a and b in the expression $\langle n_{\pi} \rangle = a + b \cdot \ln(W^2/\text{GeV}^2)$ for the average multiplicities of π^+ and π^- in the forward and backward hemispheres, for all $Q^2 > 4 \text{ GeV}^2$.

	a	b	χ^2/NDF
π^+ forward	-0.11 ± 0.05	0.288 ± 0.009	10.8/6
π^+ backward	0.18 ± 0.04	0.209 ± 0.008	14.4/6
π^- forward	0.03 ± 0.05	0.242 ± 0.009	3.5/6
π^- backward	-0.42 ± 0.04	0.284 ± 0.008	9.5/6
π forward	-0.07 ± 0.07	0.529 ± 0.015	4.0/6
π backward	-0.24 ± 0.07	0.493 ± 0.014	8.3/6
π^+	0.05 ± 0.07	0.500 ± 0.014	1.6/6
π^-	-0.38 ± 0.06	0.527 ± 0.011	6.2/6
π	-0.33 ± 0.13	1.027 ± 0.025	2.4/6

Table 3: Fitted values of the coefficients c and d in the expression $\langle n_{\pi} \rangle = c + d \cdot \ln(Q^2/\text{GeV}^2)$ for the average multiplicity of (a) all charged pions, (b) π^+ and (c) π^- , for different intervals of W (in GeV).

a) π	c	d	χ^2/NDF
4 < W < 6	2.33±0.52	0.28±0.18	1.5/2
6 < W < 8	3.05±0.35	0.22±0.12	1.3/2
8 < W < 10	3.51±0.33	0.22±0.11	2.2/2
10 < W < 12	3.98±0.28	0.21±0.09	1.4/2
12 < W < 14	4.42±0.26	0.19±0.09	1.9/2
14 < W < 16	5.23±0.24	-0.01±0.08	0.9/2
16 < W < 18	5.27±0.21	0.10±0.08	0.17/2
18 < W < 20	5.73±0.19	0.01±0.07	0.8/2

b) π^+	c	d	χ^2/NDF
4 < W < 6	1.29±0.26	0.16±0.09	1.6/2
6 < W < 8	1.60±0.19	0.14±0.06	3.0/2
8 < W < 10	1.86±0.20	0.13±0.07	2.8/2
10 < W < 12	2.08±0.16	0.13±0.05	2.1/2
12 < W < 14	2.28±0.14	0.11±0.05	1.8/2
14 < W < 16	2.69±0.13	0.02±0.04	1.5/2
16 < W < 18	2.71±0.11	0.07±0.04	0.07/2
18 < W < 20	3.02±0.10	0.00±0.04	0.26/2

c) π^-	c	d	χ^2/NDF
4 < W < 6	1.00±0.23	0.13±0.08	1.9/2
6 < W < 8	1.40±0.16	0.10±0.05	0.33/2
8 < W < 10	1.54±0.15	0.12±0.05	1.6/2
10 < W < 12	1.87±0.13	0.09±0.04	0.76/2
12 < W < 14	2.13±0.12	0.07±0.04	1.8/2
14 < W < 16	2.55±0.11	-0.04±0.04	0.82/2
16 < W < 18	2.56±0.10	0.02±0.04	1.0/2
18 < W < 20	2.72±0.09	0.01±0.04	2.3/2

Table 4: Fitted values of the coefficients a_1 , a_2 and a_3 in the expression $\langle n_\pi \rangle = a_1 + a_2 \cdot \ln W^2 + a_3 \cdot \ln Q^2$ for the average charged pion multiplicity, together with the corresponding values for all charged hadrons (from ref. [12]).

	a_1	a_2	a_3^*	χ^2/NDF
charged pions	-0.66 ± 0.17	1.04 ± 0.03	$0.10 \pm 0.03 \pm 0.04$	22/29
charged hadrons	-0.30 ± 0.16	1.22 ± 0.03	$0.22 \pm 0.04 \pm 0.06$	26/22

*) The errors are statistical and systematic respectively.

Table 5: Results of linear fits to the $\ln W^2$ dependence of the central height $D^\pi(y=0)$ and the width Δy (at various heights) of the rapidity distributions of charged pions and of π^+ and π^- separately.

	a	b	χ^2/NDF
$\pi \quad D^\pi(y=0)$	1.29 ± 0.20	-0.01 ± 0.04	0.027/2
$\pi \quad \Delta y(50\%)$	-0.36 ± 0.60	0.83 ± 0.11	0.16/2
$\pi \quad \Delta y(5\%)$	1.58 ± 0.60	0.92 ± 0.12	0.58/2
$\pi \quad \Delta y(1\%)$	2.90 ± 0.84	0.85 ± 0.17	0.79/2
$\pi^+ \quad D^\pi(y=0)$	0.77 ± 0.07	-0.02 ± 0.01	0.12/2
$\pi^+ \quad \Delta y(50\%)$	-0.83 ± 0.76	0.95 ± 0.14	0.034/2
$\pi^+ \quad \Delta y(5\%)$	1.22 ± 1.41	1.02 ± 0.26	0.17/2
$\pi^+ \quad \Delta y(1\%)$	2.32 ± 1.63	1.01 ± 0.32	0.38/2
$\pi^- \quad D^\pi(y=0)$	0.53 ± 0.01	0.02 ± 0.01	0.54/2
$\pi^- \quad \Delta y(50\%)$	-0.30 ± 0.70	0.78 ± 0.14	0.063/2
$\pi^- \quad \Delta y(5\%)$	1.53 ± 1.29	0.86 ± 0.25	0.12/2
$\pi^- \quad \Delta y(1\%)$	2.48 ± 1.80	0.90 ± 0.35	0.10/2

Figure captions

- Fig. 1: Muoproduction of hadrons in the quark-parton model.
- Fig. 2: Correction function of (a) positive and (b) negative pions vs. x_F , for events with $Q^2 > 4 \text{ GeV}^2$ and $16 < W^2 < 400 \text{ GeV}^2$. The shaded areas indicate the statistical uncertainty.
- Fig. 3: Average multiplicity $\langle n_\pi \rangle$ of charged pions vs. W^2 for fixed intervals of Q^2 . Error bars are only shown for some of the points in order to avoid confusion.
- Fig. 4: Average multiplicity $\langle n_\pi \rangle$ of (a) charged pions and of (b) π^+ and (c) π^- separately vs. Q^2 for fixed intervals of W .
- Fig. 5: Fragmentation functions (normalized x_F distributions) $D^\pi(x_F, W^2, Q^2)$ of charged pions for fixed intervals of x_F are shown vs. W^2 for fixed intervals of Q^2 (left hand side), and vs. Q^2 for fixed intervals of W (right hand side). The distributions are shown separately (a) for forward ($x_F > 0$) and (b) for backward ($x_F < 0$) going pions.

Symbols for x_F -intervals:

for $x_F > 0$ (Fig. 5a)	for $x_F < 0$ (Fig. 5b)
● $0.0 < x_F < 0.05$	● $-0.05 < x_F < 0.0$
■ $0.05 < x_F < 0.1$	■ $-0.1 < x_F < -0.05$
▲ $0.1 < x_F < 0.2$	▲ $-0.2 < x_F < -0.1$
× $0.2 < x_F < 0.3$	× $-0.3 < x_F < -0.2$
◆ $0.3 < x_F < 0.4$	◆ $-0.4 < x_F < -0.3$
▼ $0.4 < x_F < 0.6$	▼ $-0.6 < x_F < -0.4$
+ $0.6 < x_F < 1.0$	

- Fig. 6: Center of mass rapidity distributions $D^\pi(y, W^2)$ of charged pions (circles) and of π^+ (squares) and π^- (triangles) separately for 4 intervals of W^2 and $Q^2 > 4 \text{ GeV}^2$. The curves show fits to the data points for all charged pions.
- Fig. 7: Width Δy at 1%, 5% and 50% of the total height (crosses, triangles, squares respectively) and central height $D^\pi(y=0, W^2)$ (full points) of the cms rapidity distribution of charged pions, as function of W^2 . The solid straight lines are linear fits to the data points. The dashed straight line is $\Delta y = \ln(W^2/\text{GeV}^2) + 2.03$, eq. (5).
- Fig. 8: Same as Fig. 7, but separately for (a) π^+ and (b) π^- .

Fig. 1

Fig. 2

Fig. 3

Fig. 4

forward pions

Fig. 5a

backward pions

Fig. 5b

Fig. 6

Fig. 7

Fig. 8