$B \rightarrow Kl^+l^-$ and Other Rare B -Meson DecaysN. G. Deshpande,^(a) G. Eilam,^(b) and A. Soni*Physics Department, University of California, Los Angeles, California 90024*

and

G. L. Kane^(c)*Theory Division, CERN, Geneva, Switzerland*

(Received 31 March 1986)

We calculate the rate and the CP -nonconservation asymmetry for the rare decay $B \rightarrow Kl^+l^-$ ($l = e$ or μ). In the standard model with three generations we find the branching ratio to be of the order of 10^{-6} , and the CP -nonconservation asymmetry at most about 1%. The decay proceeds via the flavor-changing loop structure of the standard model, and unlike $K \rightarrow \pi e^+e^-$ suffers few uncertainties and can therefore be used as a probe of new physics. We also comment on other rare decay modes of the B meson.

PACS numbers: 13.20.Jf, 11.30.Er, 12.15.Ji

One interesting type of physics that the Superconducting Super Collider (SSC) as well as other upcoming machines will allow is the study of rare- B -decay branching ratios down to the level of 10^{-7} to 10^{-8} .^{1,2} Furthermore, new detection techniques,² such as vertex detectors, and the fact that the B meson has a relatively large lifetime³ will be helpful in these experiments.

The interest in rare B decays^{1,4-9} stems from the possibility of measuring hitherto unmeasured higher-order corrections to the standard model, including non-Abelian couplings. Kaon decays are similar in these and in other respects, but suffer from uncertainties due to long-distance effects which are very difficult to disentangle reliably from short-distance contributions.¹⁰ In this Letter we consider a rare B decay, namely $B \rightarrow Kl^+l^-$ ($l = e$ or μ) which seems to be the most promising candidate for a measurable rare decay process, and which proceeds via the flavor-

changing weak transition in the standard model. With three generations, we find that the branching ratio for the inclusive decay $B \rightarrow Kl^+l^- + X$ ranges from around 2×10^{-6} to 3×10^{-6} as m_t changes between 40 and 240 GeV, with very little sensitivity to unknown mixing angles. The CP -nonconserving asymmetry which depends of course on mixing angles is also considered, and is found to be limited to about 1.5%. The exclusive mode $B \rightarrow Kl^+l^-$ ($l = e$ or μ) is about a factor of $\frac{1}{2}$ as abundant as the inclusive one and exhibits a similar asymmetry. Any significant deviation of the experimental results from our predictions would indicate a departure from the standard model¹¹ indicating interesting new physics such as the existence of a fourth generation, supersymmetry, flavor-changing neutral currents, etc. Other rare decay modes of B mesons,¹² including purely nonleptonic ones, will also be briefly discussed.

The matrix element for the quark-level process (see Fig. 1) is¹³

$$\mathcal{M}^{\text{inclusive}} = -i(G_F/2\sqrt{2})(\alpha/\pi)G_1\bar{u}_2\gamma^\mu(1-\gamma_5)u_1\bar{u}_4\gamma_\mu v_3, \quad (1)$$

where

$$G_1 = A_c(F_1^c - F_1^t) + A_u(F_1^t - F_1^c), \quad (2)$$

with $A_j = U_{j\ell}U_{jb}^*$, the U 's being elements of the Kobayashi-Maskawa matrix¹⁴ ($j = u, c, t$), and $A_u + A_c + A_t = 0$. In general two classes of diagrams, denoted by R and Λ (see Fig. 1) contribute to the form factors F_1^j . We use the calculation reported elsewhere,^{15,16} and immediately observe that for the top-quark contribution, $k^2 \approx 0$ can be safely assumed. The resulting values¹⁷ of F_1^j for $m_t = 40, 60, 80, 160$, and 240 GeV are respectively 0.62, 0.28, 0.05, -0.45 , and -0.70 . Let us now discuss the calculation of $F_1^j(k^2)$ with $j = c$ or u . First, note that if $\hat{m}_j^2 \ll 1$ with $\hat{m}_j = m_j/M_w$, the contribution of the unphysical Higgs scalar is negligible.¹⁸ Furthermore, it is straightforward to show that unlike the t -quark case, only the Λ diagrams contribute dominantly for $j = u, c$, since $|\ln \hat{m}_j^2| \gg \hat{m}_j^2$, and the logarithm is present in Λ , but absent from R . We then obtain

$$F_1^j(k^2) = -2e_j \int_0^1 dy \int_0^{1-y} dx [y(x-1) + 2x(x-1)]/[y + \hat{m}_j^2(1-y) - \hat{k}^2 x(1-x)], \quad (3)$$

FIG. 1. Diagrams Λ and R (where each stands for a set of diagrams) for $b \rightarrow s l^+ l^-$. ϕ is the unphysical Higgs boson required in R_ξ gauges.

TABLE I. Values of the coefficients a_i and integrands I_i appearing in branching ratios and asymmetries. F_l^j given in Eqs. (4) and (5) are to be evaluated at $k^2 = z m_b^2$.

i	a_i	I_i
1	$(\text{Re} A_c)^2$	$(\text{Re} F_l^j - F_l^j)^2$
2	$2 A_u \text{Re} A_c$	$(\text{Re} F_l^j - F_l^j)(\text{Re} F_l^j - F_l^j)$
3	A_u^2	$(\text{Re} F_l^j - F_l^j)^2$
4	$(\text{Im} A_c)^2$	$(\text{Im} F_l^j)^2$
5	$-2 A_u \text{Im} A_c$	$(\text{Re} F_l^j - F_l^j) \text{Im} F_l^j$
6	$(\text{Im} A_c)^2$	$(\text{Re} F_l^j - F_l^j)^2$
7	$(\text{Re} A_c)^2$	$(\text{Im} F_l^j)^2$
8	$2 A_u \text{Re} A_c$	$\text{Im} F_l^j \text{Im} F_l^j$
9	A_u^2	$(\text{Im} F_l^j)^2$
10	$2 A_u \text{Im} A_c$	$(\text{Re} F_l^j - F_l^j) \text{Im} F_l^j$

where y, x are Feynman parameters, $\hat{k}^2 = k^2/M_W^2$, and $e_j = \frac{2}{3}$ is the charge of j . Equation (3) leads to

$$\text{Re} F_l^j(k^2) = -4e_j \int dx x(1-x) \ln |\hat{m}_j^2 - \hat{k}^2 x(1-x)|, \quad (4)$$

and

$$\text{Im} F_l^j(k^2) = [-\frac{1}{2} e_j \pi (b_+^2 - b_-^2 - \frac{1}{3} b_+^3 + \frac{1}{3} b_-^3)] \theta(k^2 - 4m_j^2), \quad (5)$$

with $b_\pm = 1 \pm [1 - 4(m_j^2/k^2)]^{1/2}$.

Let us first consider the inclusive rate $B \rightarrow K l^+ l^- + X$ which we take to be equal to the quark-level process $b \rightarrow s l^+ l^-$. After squaring V in Eq. (1) and integrating over the three-body phase space one finds

$$\Gamma(B \rightarrow K l^+ l^- + X) = (G_F^2 M_b^5 / 192 \pi^3) (\alpha / 2\pi)^2 \int_{z_{\min}}^1 dz (1-z)^2 (1+2z) |G_1(k^2 = z m_b^2)|^2, \quad (6)$$

where $z_{\min} = 4(m_l/m_b)^2$ and G_1 is given in Eq. (2). We can rewrite the above equation as an expression for the inclusive branching ratio

$$B(B \rightarrow K l^+ l^- + X) = \frac{T e^4}{\pi^4} \sum_{i=1}^{10} a_i \int_{z_{\min}}^1 dz (1-z)^2 (1+2z) I_i, \quad (7)$$

where a_i, I_i are given in Table I, and T is defined through the B lifetime,¹⁹

$$\tau_B = 10^{-12} T \text{ sec}, \quad (8)$$

with $1 < T < 2$.³ It is obvious that since from experiment²⁰

$$B(b \rightarrow u) = 0.04y, \quad y \leq 1, \quad (9)$$

then the c and t quarks contribute more than the u quark. Although T has the indicated uncertainty at present, the branching ratio in Eq. (7) is practically independent of T , since T^{-1} appears in Kobayashi-Maskawa angles in the numerator of Eq. (7) through a_i .²¹ In Fig. 2 the inclusive branching ratio is shown (solid line) as a function of m_t for $s_8 = 0.1$, $c_8 > 0$, $y = 0.5$, and $T = 1$. s_2 and s_3 are given by²³

$$s_3 = (10^{-3}y/T)^{1/2}, \quad (10)$$

$$s_2 = (10^{-3}/T)^{1/2} [-c_8 y^{1/2} + (3 - s_8^2 y)^{1/2}].$$

As s_8, y, T vary over their allowed region (including a reversal of the sign of C_8) the branching ratio changes

FIG. 2. Branching ratios for the inclusive $B \rightarrow K l^+ l^- + X$ (solid line) and exclusive $B \rightarrow K l^+ l^-$ (dashed line) processes.

by at most 10%.

Let us now consider the CP -nonconserving asymmetry which should exhibit itself through a rate difference between B^+ and B^- :

$$a = \frac{\Gamma_{\bar{b}} - \Gamma_b}{\Gamma_{\bar{b}} + \Gamma_b} = \frac{a_5 \int dz (1-z)^2 (1+2z) (I_5 - I_{10})}{a_5 \int dz (1-z)^2 (1+2z) (I_5 - I_{10}) - \Gamma_b}, \quad (11)$$

where Γ_b is the inclusive width for $B \rightarrow Kl^+ l^- + X$ [Eq. (7)], and a_i, I_i were given in Table I. Since a_5 is proportional to A_u , only when $A_u \neq 0$ is there a possibility of CP asymmetry ($a \neq 0$), and when we vary s_8 , y , and T over their allowed region the asymmetry changes accordingly. From Eqs. (10) and (11) we obtain

$$a = \frac{2}{3} s_8^2 y^{1/2} s_8 R [-c_8 y^{1/2} + (3 - s_8^2 y)^{1/2}], \quad (12)$$

where

$$R = \frac{-\int dz (1-z)^2 (1+2z) (I_5 - I_{10})}{\int dz (1-z)^2 (1+2z) I_1}. \quad (13)$$

The maximum value allowed for the asymmetry is obtained for $s_8 = 0.87$, $c_8 < 0$, and $y = 1$. It is given by

$$\max(a) = 0.06R. \quad (14)$$

In Fig. 3 we plot the value of $\max(a)$ (solid line) as a function of m_t ; it is of the order of 1%–1.5%.

We now discuss the exclusive decay $B \rightarrow Kl^+ l^-$ which is much easier to reconstruct. The estimate of the exclusive mode is subject to an uncertainty due to the transition from the quark to the meson level. We use

$$\langle K^+ | \bar{s} \gamma_\mu b | B^+ \rangle = f^+ (k^2) (2p_2 + k)_\mu + f^- (k^2) k_\mu \quad (15)$$

(for definition of momenta see Fig. 1). Therefore, the matrix element for the exclusive process is

$$\mathcal{M}^{\text{exclusive}} = -i \left(\frac{G_F}{2\sqrt{2}} \right) \frac{\alpha}{\pi} G_1 f^+ (2p_2 + k)^\mu \bar{u}_4 \gamma_\mu v_3. \quad (16)$$

After squaring, integrating over phase space, and using $f^+ (k^2) = m_B^2 / (m_B^2 - k_2^2)$, we find

$$B(B \rightarrow Kl^+ l^-) = (Te^4 / 2\pi^4) \sum a_i \int_{z_{\min}}^1 dz (1-z) I_i, \quad (17)$$

where a_i, I_i are in Table I. The exclusive branching ratio and the corresponding asymmetry²⁴ are plotted in Figs. 2 and 3 (dashed lines). The rate is smaller by

FIG. 3. The maximum value of the CP -nonconservation asymmetry a [defined in Eq. (11)] vs m_t for the inclusive (solid line) and exclusive (dashed line) processes.

about a factor of 2 as compared to the inclusive case. Clearly, the largest uncertainty is due to our ignorance regarding the form factor in Eq. (15), i.e., our choice of f^+ as having a form motivated after a similar form which successfully parametrizes kaon¹⁰ and D decays.²⁵

We therefore find that the branching ratios are of the order of 10^{-6} , and measurable, while the asymmetries are bounded by 1% or so. As emphasized, we have estimated the long-distance (LD) contributions²⁶ to $B \rightarrow Kl^+ l^-$ and find them to be less than 2% of the short-distance (SD) contribution.²⁷ The calculation involves virtual photon emission preceded or followed by $B \rightarrow K$ transition. This transition is estimated from the effective nonleptonic Hamiltonian which is dominated by the penguin graph. Using vacuum saturation we find that the ratio of LD to SD amplitude is less than

$$\frac{2\alpha_s(m_b^2) F_B F_K \ln(m_t/m_b)}{m_p^2 \ln(m_t/m_c)} < 0.02.$$

Details of this calculation will be presented elsewhere. A source of background for the processes we have considered arising from the decay of charmonia produced in $B \rightarrow K\psi$ or $B \rightarrow K\psi\chi$ can be easily removed by cutting of the dilepton mass around the mass of ψ .

Finally we note that there are other rare decay modes of B mesons which proceed via the flavor-changing loop.²⁸ First, there are the $B \rightarrow K + \gamma + X$ (no charm) and²⁹ $B \rightarrow K^* + \gamma$ decays³⁰ which proceed through diagrams similar to Fig. 1.¹³ The branching ratios are of the order of 10^{-4} . Then there are exclusive modes which receive no contribution except for higher-order loops in the standard model, for example^{5,8,9} $B_d \rightarrow \phi K_S$, $B_s \rightarrow \phi\phi$. The branching ratios are difficult to estimate reliably. They may be well above 10^{-5} .⁸ All these decays should also exhibit CP nonconservation in the standard model (unless $U_{bu} = 0$) which will be hard to detect, since it is again expected to be at most of the order of 1% in the standard

model as discussed above.

The research of one of us (A.S.) was supported by the U. S. Department of Energy Outstanding Junior Investigator Program under Grant No. DE-AM03-76SF0034/PA-A5-AT03-815R40024/Task J. Work also was supported by U. S. Department of Energy Grant No. DE-FG06-85ER40224.

(a)Permanent address: Institute of Theoretical Science, University of Oregon, Eugene, OR 97403.

(b)On leave from Physics Dept., Technion, Haifa, Israel.

(c)On leave from Physics Dept., University of Michigan, Ann Arbor, MI 48109.

¹N. G. Deshpande, in *Proceedings of the Topical Conference on Supercollider Physics, Eugene, Oregon, 9-10 August 1985*, edited by D. Soper (World Scientific, Singapore, 1986), and in *Proceedings of the Workshop on Standard Model Physics at the Superconducting Super Collider, Los Angeles, 1986* (to be published).

²*Proceedings of the 1984 Summer Study on the Design and Utilization of the Superconducting Super Collider*, edited by R. Donaldson and J. Morfin (Division of Particles and Fields of the American Physical Society, New York, 1985), p. 161.

³D. E. Klem *et al.*, Phys. Rev. Lett. **53**, (1984); E. Fernandez *et al.*, Phys. Rev. Lett. **51**, 1022 (1983); N. S. Locker *et al.*, Phys. Rev. Lett. **51**, 1316 (1983); M. Althoff *et al.*, Phys. Lett. **149B**, 524 (1984); W. Bartell *et al.*, DESY Report No. 86-001, 1986 (to be published).

⁴M. Bander, D. Silverman, and A. Soni, Phys. Rev. Lett. **43**, 142 (1979).

⁵G. Eilam and J. P. Leveille, Phys. Rev. Lett. **44**, 1648 (1980); B. Guberina, R. D. Peccei, and R. Ruckl, Phys. Lett. **90B**, 169 (1980).

⁶G. Eilam, Phys. Rev. Lett. **49**, 1478 (1982).

⁷J. W. Cronin *et al.*, in Ref. 2; D. B. Cline, to be published.

⁸M. B. Gavela, A. Le Yaouanc, L. Oliver, O. Pene, J.-C. Raynal, M. Jarfi, and O. Lazrak, Phys. Lett. **154B**, 425 (1985).

⁹K. Whisnant, B.-L. Young, and J.-C. Su, Iowa State and Jilin University Report No. IS-J 1704, 1985 (to be published).

¹⁰For a review see J. F. Donoghue, E. Golowich, and B. Holstein, Phys. Rep. **131**, 319 (1986), and references therein.

¹¹For recent reviews see L. Wolfenstein, Carnegie-Mellon University Report No. CMU-HEP 86-3, 1986 (to be published); L.-L. Chau, Invited talk at the Meson 50 Conference, Kyoto, Japan, 15-17 August 1985 (to be published), and references therein; J. D. Bjorken, Fermilab Report No.

FNAL-CONF-85/45, 1985 (unpublished).

¹²L.-L. Chau and H. Y. Cheng, Phys. Rev. Lett. **53**, 1037 (1984), and Phys. Lett. **165B**, 429 (1986).

¹³A second term proportional $G_2 \sigma_{\mu\nu} k^\nu$ can be neglected for a virtual photon with momentum k (but is the only contribution for a real photon) since $G_2 = A_c(F_2^x - F_2^y) + A_u(F_2^y + F_2^x) \ll G_1$.

¹⁴M. Kobayashi and T. Maskawa, Prog. Theor. Phys. **44**, 652 (1973).

¹⁵N. G. Deshpande and G. Eilam, Phys. Rev. D **26**, 2463 (1982).

¹⁶N. G. Deshpande and M. Nazerimonfared, Nucl. Phys. **B123**, 390 (1983).

¹⁷See Eq. (5.20) and Table 2 in Ref. 16. Note that the effect of the non-Abelian coupling (R diagrams) becomes more important as m_t increases.

¹⁸We performed the calculations both in linear (Ref. 15) and in a nonlinear (Ref. 16) R_f gauges.

¹⁹We then find $\Gamma_B = (G_F^2 m_b^5 / 192 \pi^3) (1/64 T)$ GeV, in terms of the measurable quantity T , if $m_b = 4.5$ GeV (substituting m_B instead of m_b will change the constant $\frac{1}{64}$ in the above equation for Γ_B , but will not affect any of our results).

²⁰A. Chen *et al.*, Phys. Rev. Lett. **52**, 1084 (1984).

²¹The dominant terms in Eq. (7) are obviously $i=1, 6$ which are proportional to $|U_{bc}|^2$ thus canceling $T \propto 1/|U_{bc}|^2$. We therefore avoid uncertainties in theoretical expressions for Γ_B [see J. P. Leveille, in *Proceedings of the CLEO Collaboration Workshop on B decays*, Cornell University Report No. CLNS 51/505, CLEO 81/05, 1981 (unpublished); A. Soni, Phys. Rev. Lett. **53**, 1407 (1984)].

²² $s_i = \sin \theta_i$, $c_i = \cos \theta_i$.

²³P. H. Ginsparg, S. L. Glashow, and M. B. Wise, Phys. Rev. Lett. **50**, 1415 (1983); F. Gilman and J. Hagelin, Phys. Lett. **133B**, 443 (1983).

²⁴The expression for the asymmetry in the exclusive case is similar to the inclusive formula [Eq. (11)] except for $(1-z)^2(1+2z) \rightarrow 1-z$ in all integrals.

²⁵F. Bletzacker, H. T. Nieh, and A. Soni, Phys. Rev. D **16**, 732 (1977); J. Barnabeu and C. Jarlskog, Z. Phys. C **8**, 233 (1981).

²⁶For the corresponding situation in K decays see G. Eilam and M. Scadron, Phys. Rev. D **31**, 2263 (1985); L. Bergstrom and P. Singer, Phys. Rev. Lett. **55**, 2633 (1985).

²⁷For the corresponding situation in K decays see F. J. Gilman and M. B. Wise, Phys. Rev. D **21**, 3150 (1980).

²⁸M. A. Shifman, A. I. Vainshtein, and V. I. Zakharov, Nucl. Phys. **B120**, 316 (1977); and Zh. Eksp. Teor. Fiz. **45**, 1275 (1977) [Sov. Phys. JETP **45**, 670 (1977)].

²⁹Note that $B \rightarrow K\gamma$; therefore, the calculation of $B \rightarrow K^*\gamma$ may be more reliable.

³⁰B. A. Campbell and P. J. O'Donnell, Phys. Rev. D **25**, 1989 (1982).