CERN - TH. - 4495/86

THE GHOST VERTEX IN E. WITTEN'S STRING FIELD THEORY

by

Stuart Samuel¹
CERN - Geneva

ABSTRACT

We construct in terms of creation operators the ghost part of the vertex associated with the covariant bosonic string field theory recently proposed by E. Witten. It contains the couplings of the auxiliary fields which were introduced to make the free string theory local and to recast it as a BRST action. Combined with previous results all couplings (physical - physical, physical - auxiliary, auxiliary - auxiliary) are now known. We explicitly display these couplings through fourth level and check to this order the BRST invariance of the full vertex.

¹ A Sloan Foundation Fellow; On leave of absence from City College of New York, Physics Department, 138th and Convent Ave, NY, USA 10031

I. INTRODUCTION

Field theory in particle physics has been a powerful computational tool. Obtaining a field theory for strings is thus a desirable goal. Recently, E. Witten has made an interesting proposal [1] for the open bosonic string which has good chances [2] of achieving this goal (the three point couplings have been computed [2,3] and found to agree with the Veneziano couplings for on-shell physical states). A light-cone formulation has existed for many years [4], but a manifestly Lorentz covariant version is important for understanding the string vacuum structure, compactification and non-perturbative effects.

Free string field theory is well understood [5–32]: Auxiliary fields must be introduced [5–7] to make the theory local and the action is of the form $S = \int \Psi^\dagger Q \Psi$ [1,5,14,15] where Q is the BRS operator [33,34]. The nilpotency of Q ($Q^2 = 0$) gives rise to the gauge invariance $\delta\Psi = Q \Lambda$. The first quantized theory also had a gauge (=reparameterization) invariance. Fixing the gauge required the introduction of Faddeev–Popov ghosts [14,33,34]. In second quantization, the string field became a function of both the bosonic and the anticommuting ghost string coordinates. When the string field was expanded, the coefficients of the ghost degrees of freedom were the auxiliary fields [5–7,14]. The gauge symmetries, $\delta\Psi = Q \Lambda$, can be used to gauge away unphysical fields [6,21], if so desired.

Interactions have been introduced into this framework in Ref. [1]. The midpoint of the string is singled out so that each string has a right half and a left half. Two strings interact to produce a third string by pairwise joining halves [1]. The combined quadratic and trilinear terms constitute a Chern–Simons form in the space of string forms [1].

The vibrations of the string generate an infinite number of particle states or Reggeons. The vertex operator provides the couplings between three arbitrary Reggeons and hence is a concrete well-defined representation of the interactions. References [2] and [3] computed this vertex operator for string vibrations not involving the ghost excitations using methods developed some 10 years ago [4,35–40]. It was shown that the correct couplings of three physical states are obtained from E. Witten’s way of joining strings. The interactions involving the ghost coordinates have been described [1,2] in bosonized form (i.e. when the Faddeev–Popov anticommuting ghosts $c(\sigma)$ and $\bar{c}(\sigma)$ are expressed as exponentials of a boson field, ϕ) in which the operator, $\exp(-3i\phi(\pi/2)/2)$, appears. It is a singular object (“a fermion to the power 3/2”) and does not presently have a Fock space interpretation. Hence, the couplings of the auxiliary fields introduced in refs.[5–7,19] are still not known. The purpose of this note is to provide these couplings. This is done by using the original anticommuting ghosts, $c(\sigma)$ and $\bar{c}(\sigma)$ which have a Fock space interpretation. The vertex associated with these degrees of freedom is obtained. All interactions in E. Witten’s theory are thus known.

The vacuum $\langle + |$ satisfies $\langle + | c_0 = 0$ and $\langle + | \bar{c}_0 \neq 0$ (see ref.[46]). Equations (2) and (3) constitute the main result of this letter. Explicit couplings to the fourth level are displayed in Eqs.(5) and (6) below. The full vertex is the product of Eqs.(1) and (2).

The rest of the letter will explain the reason why Eqs.(2) and (3) are the ghost vertex.

The ghost vertex is actually fixed by BRST invariance. Refs. [44] and [46] showed that order g invariance requires the full vertex, $\langle V |$, to satisfy $\langle V | (Q^{(1)} + Q^{(2)} + Q^{(3)})$ where Q^r is the BRST charge acting on the r^{th} set of Fock space coordinates.

When the Q 's act on the vertex, the creation operators can be pushed to the left through the vertex so that only destruction operators appear. Three types of terms are generated: (A) terms bilinear in bosonic α_n oscillators and linear in c_n 's (B) terms bilinear in c_n 's and linear in $\bar{c}_n$'s and (C) terms linear in c_n 's. It can be shown that a necessary criterion for BRST invariance is [46]:

$${}_{123}\langle 0 | V^{\text{boson}} L_{-m}^r = {}_{123}\langle 0 | V^{\text{boson}} \left(- \sum_{n=0}^{\infty} \chi_{mn}^{rs} L_n^s + \chi_m^r \right), m > 0, \quad (4)$$

where $L_n = 1/2 : \sum_m \alpha_{n-m} \cdot \alpha_m :$ are the bosonic Virasoro generators and the χ_n^r are constants. If the vertex operator is quadratic in bosonic and ghost oscillators then Eq.(4) fixes the ghost action. If Eq.(4) is true then type (A) terms cancel if X in Eq.(4) is used in Eq.(2). Eq.(4) thus provides a way of computing the ghost vertex.

We have checked that Eq.(4) is satisfied for E. Witten's theory to 4th level. Non-trivial constraints on the N 's in Eq.(1) must be satisfied for the L_{-n} 's "to get reflected back". For an arbitrary quadratic vertex, the terms generated by pushing creation operators through the vertex cannot be recast in terms of L_n^r 's (for example, off-diagonal terms $\alpha_n^r \cdot \alpha_m^s$ must cancel). From Eq.(4), we obtain the ghost vertex to fourth level:

$$\begin{aligned} {}_{123}\langle + + + | V^{\text{ghost}} = {}_{123}\langle + + + | \exp & \left[\frac{4}{3\sqrt{3}} c_1^{(1)} (\bar{c}_0^{(2)} - \bar{c}_0^{(3)}) \right. \\ & + c_1^{(1)} \left(\frac{11}{27} \bar{c}_1^{(1)} + \frac{8}{27} (\bar{c}_1^{(2)} + \bar{c}_1^{(3)}) \right) + c_2^{(1)} \left(\frac{16}{27} \bar{c}_0^{(1)} - \frac{8}{27} (\bar{c}_0^{(2)} + \bar{c}_0^{(3)}) \right) \\ & + \frac{8.5\sqrt{3}}{3^5} c_1^{(1)} (\bar{c}_2^{(2)} - \bar{c}_2^{(3)}) - \frac{16.5\sqrt{3}}{3^5} c_2^{(1)} (\bar{c}_1^{(2)} - \bar{c}_1^{(3)}) - \frac{4.17\sqrt{3}}{3^5} c_3^{(1)} (\bar{c}_0^{(2)} - \bar{c}_0^{(3)}) \\ & + c_3^{(1)} \left(-\frac{16.5}{3^5} \bar{c}_1^{(1)} + \frac{8.5}{3^5} (\bar{c}_1^{(2)} + \bar{c}_1^{(3)}) \right) + c_2^{(1)} \left(-\frac{19}{3^5} \bar{c}_2^{(1)} - \frac{16.7}{3^5} (\bar{c}_2^{(2)} + \bar{c}_2^{(3)}) \right) \\ & + c_1^{(1)} \left(-\frac{16.5}{3^6} \bar{c}_3^{(1)} + \frac{8.5}{3^6} (\bar{c}_3^{(2)} + \bar{c}_3^{(3)}) \right) + c_4^{(1)} \left(-\frac{32.11}{3^6} \bar{c}_0^{(1)} + \frac{16.11}{3^6} (\bar{c}_0^{(2)} + \bar{c}_0^{(3)}) \right) \\ & \left. + (1) \rightarrow (2) \rightarrow (3) \rightarrow (1) + (1) \rightarrow (3) \rightarrow (2) \rightarrow (1) + \dots \right] \end{aligned} \quad (5)$$

For completeness, we also display the bosonic vertex to fourth level:

$$\begin{aligned}
 {}_{123}\langle 0|V^{\text{boson}} = g \delta(\sum_r k^r) {}_{123}\langle 0| \exp \left[\left(\frac{1}{2} k^{(1)} \cdot k^{(1)} - 1 \right) \ln \left(\frac{4}{3\sqrt{3}} \right) \right. \\
 + \frac{2}{3\sqrt{3}} \alpha_1^{(1)} \cdot (k^{(2)} - k^{(3)}) + \alpha_1^{(1)} \cdot \left(-\frac{5}{27} \frac{1}{2} \alpha_1^{(1)} + \frac{16}{27} \alpha_1^{(2)} \right) - \frac{1}{9} \alpha_2^{(1)} \cdot k^{(1)} \\
 + \frac{32\sqrt{3}}{3^5} \alpha_1^{(1)} \cdot (\alpha_2^{(2)} - \alpha_2^{(3)}) - \frac{2 \cdot 11 \sqrt{3}}{3^6} \alpha_3^{(1)} \cdot (k^{(2)} - k^{(3)}) \\
 + \alpha_1^{(1)} \cdot \left(\frac{32}{3^6} \alpha_3^{(1)} - \frac{16}{3^6} \alpha_3^{(2)} \right) + \alpha_2^{(1)} \cdot \left(\frac{13}{2 \cdot 3^5} \frac{1}{2} \alpha_2^{(1)} - \frac{2^6}{3^5} \alpha_2^{(2)} \right) + \frac{19}{2 \cdot 3^5} \alpha_4^{(1)} \cdot k^{(1)} \\
 \left. + (1) \rightarrow (2) \rightarrow (3) \rightarrow (1) + (1) \rightarrow (3) \rightarrow (2) \rightarrow (1) + \dots \right] .
 \end{aligned} \tag{6}$$

We can now check whether the ansatz in Eq.(3) works. Table 1 displays the relevant Neumann functions. Inspection shows that Eq.(5) calculated from Eqs.(2) and (3) yield the same result as Eq.(5) computed from the BRST criterion in Eq.(4).

One can then check whether terms (B) and (C) cancel. After quite a bit of algebra, we have verified this to fourth level. There are 3·39 different type (B) terms (the 3 comes from cycling the string labels: $(1) \rightarrow (2) \rightarrow (3) \rightarrow (1)$ and $(1) \rightarrow (3) \rightarrow (2) \rightarrow (1)$). The three × four type (C) terms are $< V \sum_r [0 \cdot c_1^r + (5 \cdot (26 - D)) / (2 \cdot 27) \cdot c_2^r + 0 \cdot c_3^r + (5 \cdot (26 - D)) / (27) \cdot c_4^r]$ which vanish if we inhabit a 26 dimensional world. The 3·59 different types of up-to-fourth-level (A) terms vanish because of Eq.(4) and (2). We have thus shown the BRST invariance of the vertex of Eqs.(1) and (2) through fourth level.

Equations (5) and (6) display all the interactions of three string states to the fourth level. One thus knows exactly how the low-lying string vibrational modes (particles) interact which should be sufficient for experimental purposes in the near future.¹

III. THE GHOST OVERLAP CONDITIONS

It is interesting to determine the ghost overlap delta function conditions. From the work of ref.[1], one suspects they are quite similar to those of the bosonic degrees of freedom: $\Pi_r \delta(x^r(\sigma) - x^{r+1}(\pi - \sigma))$, $0 \leq \sigma \leq \pi/2$, which is equivalent to the Goto-Naka conditions [48]:

$$\begin{aligned}
 {}_{123}\langle 0|V^{\text{boson}}(X^r(\sigma) - X^{r+1}(\pi - \sigma)) = 0 \\
 {}_{123}\langle 0|V^{\text{boson}}(P^r(\sigma) + P^{r+1}(\pi - \sigma)) = 0 , \quad 0 \leq \sigma \leq \frac{\pi}{2} ,
 \end{aligned} \tag{7}$$

As explained in ref.[3], Eq.(7) can be proven using the fact that, by construction [35-38], the six string process satisfies its own set of Goto-Naka conditions. The six string Neumann functions satisfy:

¹ We assume that the inverse square root string tension is of the order of the Planck scale so that the couplings in Eqs.(5) and (6) represent all relevant physics up to several Planck scales.

$$\begin{aligned} \frac{1}{n} z^{-n} \delta^{rs} - \frac{1}{n} (-z)^n \delta^{r+1s} - \sum_{m=0}^{\infty} (z^m \bar{N}_{mn}^{rs} - (-z)^{-m} \bar{N}_{m \ n}^{r+1s}) &= 0, \\ \alpha_0^r + \alpha_0^{r+1} + \sum_{s=1}^6 \sum_{m=1}^{\infty} m (z^m \bar{N}_{m0}^{rs} + (-z)^{-m} \bar{N}_{m \ 0}^{r+1s}) \alpha_0^s &= 0, \\ \sum_{s=1}^6 \sum_{m=0}^{\infty} \cos(m\sigma) (\bar{N}_{m0}^{rs} - (-1)^{-m} \bar{N}_{m \ 0}^{r+1s}) \alpha_0^s &= 0. \end{aligned} \quad (8)$$

where we have introduced $z = \exp(i\sigma)$. In the above, $n \geq 1$, $-\pi/2 \leq \sigma \leq \pi/2$ and the α_0^r 's are to be treated as c -numbers satisfying the constraint $\alpha_0^1 + \dots + \alpha_0^6 = 0$. Letting $c(\sigma) = \sum c_{-n} z^n \equiv c(z)$ and $\bar{c}(\sigma) = \sum \bar{c}_{-n} z^n \equiv \bar{c}(z)$ we can show that Eq.(8) implies:

$$\begin{aligned} \langle +++ | V^{\text{ghost}} [c^r(z) + c^{r+1}(-\frac{1}{z})] &= 0, \\ \langle +++ | V^{\text{ghost}} [\bar{c}^r(z) - \bar{c}^{r+1}(-\frac{1}{z})] &= 0, \end{aligned} \quad (9)$$

where $-\pi/2 \leq \sigma \leq \pi/2$. Hence c satisfies a condition similar to the Fubini-Veneziano field $P(\sigma) + X'(\sigma)$ [49] and $\bar{c}$ is similar to X . If one treats the BRST charge classically (i.e. the oscillators are treated as c -numbers) then the conditions in Eqs.(7) and (9) imply that $\langle V | (Q(1) + Q(2) + Q(3)) = 0$ (formally). Similar motivations were used to construct the BRST invariant light cone vertex [44].

Summarizing we find that ghosts and antighosts satisfy similar overlap conditions as the boson degrees of freedom, that is, the right half of one ghost variable is related to the left half of the next one. When bosonized, this is the picture of ref. [1].

The ghost vertex should contain three c -delta functions. By inspection we find $\langle V | [c_0^{(1)} + c_0^{(2)} + c_0^{(3)}] = 0$. This is the ghost analog of $\delta(p^{(1)} + p^{(2)} + p^{(3)})$ and similar to what happens in the light cone approach [44,46]. Since $\langle V | [c^{(1)}(z_0) = -c^{(2)}(z_0) = +c^{(3)}(z_0) = -c^{(1)}(z_0)]$ where $z_0 = \exp(i\pi/2)$

(similarly for $z_0^* = \exp(-i\pi/2)$), $\langle V | c^{(1)}(z_0) = 0 = \langle V | c^{(1)}(z_0^*)$ (it does not matter which of the three c 's is used because at $\exp(\pm i\pi/2)$ they are all equal (up to a possible minus sign)). We thus find that, when the ghosts are written as functions of a complex parameter $z (= \exp(i\sigma))$ and σ is analytically continued from the region $0 \leq \sigma \leq \pi$ to $-\pi \leq \sigma \leq \pi$, there is a delta function associated with the interaction point $\pi/2$ ($z = i$) and a delta function associated with its complex image at $-\pi/2$ ($z = -i$). This is almost the same as in refs. [1,44].

IV. THE " $\int$ " AND " $*$ "

In terms of the fundamental building blocks of ref.[1], Eqs.(1) and (2) represent the interaction term, $\int \Psi * \Psi * \Psi$. It would be desirable to have $\int$ and $*$ separately. For bosonized ghosts, this has been done by treating the ghost coordinates as a 27th coordinate [1,2]. We believe that $*$ can be obtained from the vertex using the following procedure: If $\Psi_3 = \Psi_1 * \Psi_2$ then

$$\Psi_3(3)|+\rangle_3 = {}_{12}\langle ++| \vee \left(\begin{array}{l} \alpha_n^3 \rightarrow -(-1)^n \alpha_{-n}^3, \\ c_n^3 \rightarrow -(-1)^n c_{-n}^3, \\ \bar{c}_n^3 \rightarrow (-1)^n \bar{c}_{-n}^3 \end{array} \right) \Psi_1(1) \Psi_2(2) |---\rangle_{123} , \quad (10)$$

where (3) denote the third oscillators and the (3) oscillators must be transcribed as indicated. The $| - \rangle$ vacuum satisfies $c_0 | - \rangle \neq 0$ and $\bar{c}_0 | - \rangle = 0$. To verify Eq.(10), one would have to show that $\int \Psi_1 * \Psi_2$ is equivalent to a twisted delta function overlap (see Eq.(10) of ref.[3]). Such a result has been shown for the bosonic degrees of freedom [2].

$\int$ is quite easy to obtain: Let

$$\begin{aligned} E &= \exp \left[- \sum_{n=1}^{\infty} (-1)^n \left(\frac{\alpha_n \alpha_n}{2n} - \bar{c}_n c_n \right) \right] , \\ P_1 &= (\bar{c}_1 - \bar{c}_3 + \bar{c}_5 - \bar{c}_7 + \dots) , \\ P_2 &= (\bar{c}_0 - 2\bar{c}_2 + 2\bar{c}_4 - 2\bar{c}_6 + \dots) . \end{aligned} \quad (11)$$

E is the exponential factor obtained by the delta function constraint which equates the left half of the string with its right half, which is the geometrical picture behind the string integral [1]. The string integral, itself, is

$$\int_{\text{string}} = \frac{i}{4} \int d^{26}x \langle + | E \bar{c}(z) \Big|_{z=i} \bar{c}(z) \Big|_{z=-i} = \int d^{26}x \langle + | P_2 P_1 E . \quad (12)$$

The first line in Eq.(12) shows that there is an antighost insertion at $z = i$ and $z = -i$, the complexified midpoints of the string. Results identical to Eq.(12) have been independently obtained in ref. [50]. The ghost insertion factors were fixed by the requirement of BRST invariance: The equation $\int Q \Psi = 0$ can be proven (it works for $D=26$ only) after quite a bit of algebra by pushing the creation operators in Q through E , P_1 and P_2 .

V. CONCLUSION

In this letter we have computed the ghost vertex operator in E. Witten's string theory, thus obtaining the auxiliary field interactions. The theory has now been completely recast in mathematical form, so that from now on, explicit calculation can be carried out. The exact way in which physical fields and auxiliary fields interact is known and we have explicitly displayed these interactions through 4th level in Eqs.(5) and (6).

Although we have obtained the complete vertex, many mathematical features of the $\int$ and $*$ operators remain to be proved: One would like to show (a) that $\int$ plus $*$ is equivalent to a twisted delta function, (b) the associativity of $*$, (c) the $K_n (= L_n - (-1)^n L_{-n})$ where these Virasoro generators are the full ones (boson + ghost) symmetries for $\int$ and $*$, (d) the BRST invariance for all levels (we have only verified it to 4th level), which would prove closure of the string gauge symmetry to order g , (e) the closure of the algebra to order g^2 , (f) to compute the tree amplitudes directly from the string field theory, (g) to extend the results to other string theories, (h) to make contact with the light cone theory [4,21] by appropriately fixing the gauge. With bosonized ghosts, D. Gross and A. Jevicki have verified (a), (b) and (c) [2]. It would be desirable to repeat these calculations with fermionic ghosts. In regard to (d) some powerful computational techniques have been developed in refs. [44,46] and may be applicable. Techniques in ref. [44] may be useful for establishing (e). (f) can be approached using the ideas in refs. [41,51,52] if the string field theory can be directly connected to Polyakov's approach [42]. Intuitive ways of obtaining string field theories for cases other than the open bosonic string have appeared in [1,50,53]. In short, although we now know explicitly the theory, there is much to be done.

ACKNOWLEDGEMENTS

I would like to thank A. Neveu and D. Sahdev for discussions.

REFERENCES

- [1] E. Witten, Nucl. Phys. B268 (1986) 253.
- [2] D. Gross and A. Jevicki, Princeton preprint (May, 1986).
- [3] S. Samuel, CERN preprint TH. – 4365/86/rev. (February, 1986 (revised June, 1986)).
- [4] M. Kaku and K. Kikkawa, Phys. Rev. D10 (1974) 1110, 1823.
- [5] W. Siegel and B. Zwieback, Nucl. Phys. B263 (1986) 105.
- [6] A. Neveu and P. C. West, Nucl. Phys. B268 (1986) 126;
A. Neveu, H. Nicolai and P. C. West, Nucl. Phys. B264 (1986) 573;
A. Neveu, J. H. Schwarz and P. C. West, Phys. Lett. 164B (1985) 51;
A. Neveu and P. C. West, Phys. Lett. 165B (1985) 63;
A. Neveu, H. Nicolai and P. C. West, CERN preprint TH.4297/85, to be published in Phys. Lett. B.
- [7] T. Banks and M. Peskin, Nucl. Phys. B264 (1986) 513.
- [8] K. Itoh, T. Kugo, H. Kunitomo and H. Ooguri, Prog. Theor. Phys. 75 (1986) 162.
- [9] D. Friedan, Phys. Lett. 162B (1985) 102; University of Chicago preprint EFI 85/27.
- [10] D. Friedan, E. Martinec and S. Shenker, Phys. Lett. 160B (1985) 55.
- [11] K. Bardakci, Berkeley University preprint LBL – 20024 (1985).
- [12] S. R. Das and M. Rubin, Fermilab preprint (1985).
- [13] M. Kaku and J. Lykken, CCNY preprint 85 – 412 (1985);
M. Kaku, Gauge Theory of Covariant Strings, CCNY preprint (1985);
Phys. Lett. B162 (1985) 97.
- [14] W. Siegel, Phys. Lett. 142B (1984) 276; 151B (1985) 391.
- [15] H. Aratyn and A. H. Zimerman, Racah Institute of Physics preprint (October, 1985), to be published in Phys. Lett. B.
- [16] E. G. Floratos, Y. Kazama and K. Tamvakis, Phys. Lett. 166B (1986) 295.

- [17] H. Aratyn and A. H. Zimerman, Racah Institute of Physics preprint (December, 1985), to be published in Phys. Lett. B.
- [18] L. Baulieu and S. Ouvry, Phys. Lett. 171B (1986) 57.
- [19] T. Banks, M. E. Peskin, C. R. Preitschopf, D. Friedan and E. Martinec, SLAC – PUB – 3853, (December, 1985).
- [20] S. Raby, R. Slansky and G. West, Los Alamos preprint LA – UR – 85 – 3794 (1985).
- [21] M. E. Peskin and C. B. Thorn, SLAC preprint PUB – 3801 (1985).
- [22] J. – L. Gervais, Ecole Normale Supérieure preprint LPTENS 85/35 (1985).
- [23] H. Terao and S. Uehara, Phys. Lett. 168B (1986) 70.
- [24] D. Pfeffer, P. Ramond and V. G. J. Rodgers, U. of Florida preprint UFTP – 85 – 019 (1985).
- [25] Y. Meurice, CERN preprint TH.4356 (1986).
- [26] H. Ooguri, Kyoto preprint KUNS – 819 – rev. (1985).
- [27] Y. Kazama, A. Neveu, H. Nicolai and P. C. West, CERN preprint TH.4301 (1985) to appear in Nucl. Phys. B;
CERN preprint TH.4418 (1986).
- [28] N. Ohta, Phys. Rev. Lett. 56 (1986) 440.
- [29] A. LeClair, Phys. Lett. 168B (1986) 53.
- [30] A. LeClair and J. Distler, Harvard preprint HUTP – 86/A008 (1986).
- [31] J. P. Yamron, Berkeley preprint UCB – PTH – 86/3 (1986).
- [32] G. D. Date, M. Gunaydin, M. Pernici, K. Pilch and P. van Nieuwenhuizen, Phys. Lett. 171B (1986) 182.
- [33] M. Kato and K. Ogawa, Nucl. Phys. B212 (1983) 443.
- [34] S. Hwang, Phys. Rev. D28 (1983) 2614.
- [35] S. Mandelstam, Nucl. Phys. B64 (1973) 205.
- [36] S. Mandelstam, Nucl. Phys. B69 (1974) 77.

- [37] E. Cremmer and J. – L. Gervais, Nucl. Phys. B76 (1974) 209.
- [38] E. Cremmer and J. – L. Gervais, Nucl. Phys. B90 (1975) 410.
- [39] J. – L. Gervais and B. Sakita, Phys. Rev. Lett. 30 (1973) 716.
- [40] M. B. Green and J. H. Schwarz, Nucl. Phys. B218 (1983) 43;
L. Brink, M. B. Green and J. H. Schwarz, Nucl. Phys. B219 (1983) 437;
M. B. Green and J. H. Schwarz, Nucl. Phys. B243 (1983) 475.
- [41] S. Giddings, Princeton University preprint (Jan., 1986) to be published in Phys. Letts. B.
- [42] A. M. Polyakov, Phys. Lett. 103B (1981) 207, 211.
- [43] J. Scherk, Rev. of Mod. Phys. 47 (1975) 122.
- [44] H. Hata, K. Itoh, T. Kugo, H Kunitomo and K. Ogawa, Phys. Lett 172B (1986) 186; Kyoto University preprint, RIFP – 660 (May, 1986).
- [45] H. Hata, K. Itoh, T. Kugo, H Kunitomo and K. Ogawa, Phys. Lett 172B (1986) 195;
- [46] A. Neveu and P. C. West, CERN preprints TH.4315/85; CERN preprint TH.4358/86.
- [47] A. Jevicki, Phys. Lett. 169B (1986) 359.
- [48] T. Goto and S. Naka, Prog. Theor. Phys. 51 (1974) 299.
- [49] S. Fubini and G. Veneziano, Nuovo Cimento 67A (1970) 29.
- [50] J. Lykken and S. Raby, Los Alamos preprint LA – UR – 86 – 1334 (1986).
- [51] S. Giddings and E. Martinec, Princeton preprint (April, 1986).
- [52] S. Giddings, E. Martinec and E. Witten, Princeton preprint (May, 1986).
- [53] E. Witten, Princeton preprint (March, 1986).

TABLE CAPTION

1. The Neumann Functions for the Six String Process Through Fourth Level.

For m and $n > 0$, $\bar{N}_{mn} = \bar{N}_{nm}$ so it is sufficient to compute the Neumann functions with the first index fixed (fixed here to be the sixth string). For $m > 0$ a constant κ^r can be added to $\bar{N}_{m0}$ so that it too satisfies the cyclic relation. Because of momentum conservation this change does not effect couplings for the bosonic vertex. It also does not effect the X_{mn} since the additive constant cancels in the difference in Eq.(3).

r	$\bar{N}_{10}^{6r}$	$\bar{N}_{20}^{6r}$	$\bar{N}_{11}^{6r}$	$\bar{N}_{30}^{6r}$	$\bar{N}_{12}^{6r}$	$\bar{N}_{40}^{6r}$	$\bar{N}_{13}^{6r}$	$\bar{N}_{22}^{6r}$
1	$\frac{1}{\sqrt{3}}$	$\frac{7}{2 \cdot 3^3}$	$\frac{2^2}{3^2}$	$\frac{-5}{3^3 \cdot \sqrt{3}}$	$\frac{2^2}{3^2 \cdot \sqrt{3}}$	$\frac{-5 \cdot 29}{2^2 \cdot 3^6}$	$\frac{2^2}{3^5}$	$\frac{-2^2 \cdot 5}{3^4}$
2	$\frac{1}{3\sqrt{3}}$	$\frac{5}{2 \cdot 3^2}$	$\frac{2^2}{3^3}$	$\frac{-23}{3^5 \cdot \sqrt{3}}$	$\frac{2^2}{3^4 \cdot \sqrt{3}}$	$\frac{-107}{2^2 \cdot 3^5}$	$\frac{-2^2 \cdot 7}{3^6}$	$\frac{-2^2}{3^5}$
3	0	$\frac{2^3}{3^3}$	$\frac{1}{3^2}$	0	0	$\frac{-2^3 \cdot 11}{3^6}$	$\frac{-2^3}{3^5}$	$\frac{-1}{2 \cdot 3^4}$
4	$-\frac{1}{3\sqrt{3}}$	$\frac{5}{2 \cdot 3^2}$	$\frac{2^2}{3^3}$	$\frac{23}{3^5 \cdot \sqrt{3}}$	$\frac{-2^2}{3^4 \cdot \sqrt{3}}$	$\frac{-107}{2^2 \cdot 3^5}$	$\frac{-2^2 \cdot 7}{3^6}$	$\frac{-2^2}{3^5}$
5	$-\frac{1}{\sqrt{3}}$	$\frac{7}{2 \cdot 3^3}$	$\frac{2^2}{3^2}$	$\frac{5}{3^3 \cdot \sqrt{3}}$	$\frac{-2^2}{3^2 \cdot \sqrt{3}}$	$\frac{-5 \cdot 29}{2^2 \cdot 3^6}$	$\frac{2^2}{3^5}$	$\frac{-2^2 \cdot 5}{3^4}$
6	0	0	$\frac{-2^3}{3^3}$	0	0	0	$\frac{2^3 \cdot 7}{3^6}$	$\frac{2^3}{3^5}$

Table 1