QCD Results from ATLAS and CMS

Michael Leyton
(on behalf of the ATLAS and CMS collaborations)

30th International Workshop on High Energy Physics
IHEP, Protvino, Russia
24 June 2014

Outline

- Introduction
- QCD, Jets, MC modeling
- Measurements
  - Soft-QCD
  - Inclusive and di- jet cross sections
  - Color coherence
  - PDF sensitivity
  - Photon + jets cross sections
  - Multi-jet cross sections
  - Extraction of α_s
- Summary

Strong interactions are responsible for the bulk of production cross sections at the LHC

The Large Hadron Collider

ATLAS

&

CMS

CMS DETECTOR

Total weight : 14,000 tonnes
Overall diameter : 15.0 m
Overall length : 28.7 m
Magnetic field : 3.8 T

STEEL RETURN YOKE
12,500 tonnes

SILICON TRACKERS
Pixel (100x150 μm) $\sim 16\text{m}^2 \sim 66\text{M}$ channels
Microstrips (80x180 μm) $\sim 200\text{m}^2 \sim 9.6\text{M}$ channels

SUPERCONDUCTING SOLENOID
Niobium titanium coil carrying $\sim 18,000\text{A}$

MUON CHAMBERS
Barrel: 250 Drift Tube, 480 Resistive Plate Chambers
Endcaps: 468 Cathode Strip, 432 Resistive Plate Chambers

PRESHOWER
Silicon strips $\sim 16\text{m}^2 \sim 137,000$ channels

FORWARD CALORIMETER
Steel + Quartz fibres $\sim 2,000$ Channels

CRYSTAL ELECTROMAGNETIC CALORIMETER (ECAL)
 $\sim 76,000$ scintillating PbWO_4 crystals

HADRON CALORIMETER (HCAL)
Brass + Plastic scintillator $\sim 7,000$ channels

	ATLAS	CMS
Tracker:	$ \eta < 2.5$	$ \eta < 2.5$
EM calorimeter:	$ \eta < 3.2$	$ \eta < 3.2$
Hadronic calorimeter:	$ \eta < 4.9$	$ \eta < 5.0$
Muon system:	$ \eta < 2.7$	$ \eta < 2.4$

$$\eta = -\ln(\tan \theta/2)$$

Run-I	$\sqrt{s}$ (TeV)	$\langle \mu \rangle$	L_{int}
2010	7	2	$\sim 40 \text{ pb}^{-1}$
2011	7	9	$\sim 5 \text{ fb}^{-1}$
2011	2.76	~ 1	0.2 pb^{-1}
2012	8	21	$\sim 20 \text{ fb}^{-1}$

$\langle \mu \rangle$ = average number of interactions
per bunch crossing

QCD in pp collisions

A proton-proton collision consists of:

- Soft processes (non-perturbative QCD)
  - Underlying event (UE)
  - Fragmentation of spectator partons
  - Initial- and final-state radiation (ISR/FSR)
  - Multiple parton interactions (MPI)
- Hard processes (perturbative QCD)
  - $\sigma_{pp \rightarrow X}$
    - PDFs
    - Matrix element

Run Number: 201269, Event Number: 80898559

Date: 2012-04-14 22:30:13 CEST

Jet reconstruction

- Jet algorithms cluster a spray of particles into a single object
- This object can act as a proxy for the partons from the hard process
- **ATLAS** and **CMS** primarily use anti- k_T algorithm
  - Sequential recombination algorithm
  - Longitudinally invariant
  - Infrared and collinear safe to all orders in pQCD
  - Better control on shape and area (wrt other algorithms)
    - essential for calibrating jet energy

$R = 0.4, 0.6$

$R = 0.5, 0.7$

Modeling the data

Data is modeled using Monte Carlo generators:

- **Matrix element:** exact part up to the stated accuracy (e.g. LO or NLO)
- **Parton shower:** a good approximation to describe radiative QCD
- **Hadronization:** a model to describe non-perturbative effects
- **Underlying event:** describes how the rest of the colliding protons interact with the event and each other

Underlying Event (UE)

- Motivation: more accurate phenomenological modeling of soft and semi-hard parton interactions
- Classic definition of event topology, oriented by **highest p_T object** in the event (p_T^{lead})
  - Azimuthal segmentation of events
  - **Transverse regions** less affected by hard scatter process and most sensitive to UE
- Observables: ΣE_T , n_{ch} , Σp_T , $\langle p_T \rangle$ vs. n_{ch}
  - Studied as function of p_T^{lead} (inversely proportional to impact parameter)
- All data distributions corrected to particle level and compared to MC models

arXiv:1406.0392, sub. to EPJC

Underlying Event

arXiv:1406.0392, sub to EPJC

- Using the calorimeters allows the inclusion of neutral particles
- Relatively well described by the MC generators, but Pythia tuned to ATLAS UE data gives worse agreement

Inclusive (≥ 1 jet): Increase indicates contribution from wide-angle emissions from the hard scattering

Exclusive (= 2 jets with p_T balance): flat, slightly decreasing with p_T^{lead} , indicates rate of MPI reaches plateau

Underlying Event in Drell-Yan

- Dependence of UE on di-muon invariant mass and transverse momentum

- UE activity independent of $M_{\mu\mu}$
- Well described by Pythia and Herwig++
- Illustrates universality of UE

- Slow increase with increasing $p_T^{\mu\mu}$
- Accurately described by MadGraph, which simulates multiple hard emissions

Double parton interactions (DPI)

Production of $W + 2$
associated jets

vs.

DPI producing a W
and a dijet

New J. Phys. 15 (2013) 033038

$$\sigma_{\text{eff}} = 15 \pm 3 \text{ (stat.) } {}^{+5}_{-3} \text{ (syst.) mb}$$

$$\sigma_{\text{eff}} = 20.7 \pm 0.8 \text{ (stat.) } \pm 6.6 \text{ (syst.) mb}$$

arXiv: 1312.5729

Comparable to values measured at other CM
energies at other experiments

DPI-produced jets expected to be back-
to-back $\rightarrow$ require jet-jet p_T balance

DPI-sensitive kinematical distributions
studied at particle level and a template
method used to extract fraction of DPI
events

Cross section measurements

- General motivation:

- Test of pQCD
- Constrain proton PDFs
- Strong coupling measurement
- Constraints on physics beyond the SM

- Basic strategy:

- Select anti- k_T jets,
 $R = 0.4, 0.5, 0.6, \text{ or } 0.7$
- Unfold to particle level
- Compare to NLO predictions
- Dominant systematic uncertainty typically due to jet energy scale

EPJC (2013) 73 2509

Inclusive jet cross section

PRD 86 (2012) 014022

PRD 87 (2013) 112002

Inclusive jet cross section

2.76 TeV

EPJC (2013) 73 2509

8 TeV

CMS-PAS-FSQ-12-031

Via inclusive cross section measurements, QCD has been tested at 3 CM energies and over 7 orders of magnitude!

Dijet cross section

7 TeV

JHEP 05 (2014) 059

PRD 87 (2013) 112002

Measurements generally well described by NLO, except for high m_{12} and large y^*

Sensitivity to physics beyond the SM

- Implement contact interaction (CI) in CIJET
- Restrict study to where theory predicts largest effect:

$$m_{12} > 1.31 \text{ TeV}, y^* < 0.5, R = 0.6$$

- CI exclusion limit:
  - Create pseudo-experiments for both SM and SM + CI
  - Calculate generalized χ^2 , taking into account asymmetric uncertainties and their correlations
  - Scan CLs over compositeness scale Λ

Exclusion of $\Lambda < 6.9\text{--}7.7 \text{ TeV}$ at 95% CL, depending on PDF set used

Color coherence

- Event selection: $N_{\text{jet}} \geq 3$ and two back-to-back leading jets
- The emission of a colored object depends on other colored objects in the event $\rightarrow$ color coherence
- Look at direction of emission of third jet (quantified by β)
- None of the examined MC give an adequate description of the data

PDF sensitivity

- HERA data tightly constrain distributions for $x < 0.01$
- Combined fit using HERA-1 and **ATLAS**/**CMS** data leads to improved shape constraints
  - Both **ATLAS** and **CMS** data favor harder gluons (and softer sea quarks)

EPJC (2013) 73:2509

CMS-PAS-SMP-12-028

Inclusive isolated prompt photon cross section

- Motivation:
  - Precise test of pQCD
  - Photons provide a cleaner environment than jets
  - Sensitive to gluons: dominant process is $q g \rightarrow q \gamma$
- NLO Jetphox: good agreement up to highest photon energies
- LO parton shower MCs
  - Pythia describes data well
  - Herwig 10-20% lower than data

PRD 89, 052004 (2014)

Potential to constrain shape and uncertainty of gluon PDF in $0.03 < x < 0.3$ (see backup)

(Isolated) photon + jet production

- Measure cross section as functions of E_T , p_T , γ , $\Delta\phi$, $|\cos \theta^{\gamma j}|$
  - $\theta^{\gamma j}$ is polar angle between photon and jet in their rest frame
- Good test of dominance of the quark exchange channel
- Good agreement between data and Jetphox NLO prediction

Nucl. Phys, B 875 (2013) 483-535

arXiv:1311.6141 (accepted by JHEP)

$R_{3/2}$ • ratio of three-jet cross section ($\sim \alpha_s^3$) over two-jet cross section ($\sim \alpha_s^2$)
 • as a function of p_T^{lead}

ATLAS-CONF-2013-041

EPJC 73 (2013) 2604

$N_{3/2}$ • ratio of three-jet cross section ($\sim \alpha_s^3$) over two-jet cross section ($\sim \alpha_s^2$)
 • as a function of $p_T^{\text{all jets}}$

- $N_{3/2}$ has a smaller dependence on scale choice and theoretical prediction
- Dominant uncertainty comes from theory

ATLAS-CONF-2013-041

Measuring α_s

Extracted using $N_{3/2}$ since it is less sensitive to choice of renormalization and factorization scales

$$\alpha_s(M_Z) = 0.111 \pm 0.006 \text{ (exp.) } {}^{+0.016}_{-0.003} \text{ (theory)}$$

ATLAS-CONF-2013-041

Extracted using $R_{3/2}$, but also inclusive jet cross section spectrum up to 2 TeV

$$\alpha_s(M_Z) = 0.1185 \pm 0.0019 \text{ (exp.) } \pm 0.0028 \text{ (PDF)} \\ \pm 0.0004 \text{ (NP)} \pm 0.0019 \text{ (exp.) } {}^{+0.0055}_{-0.0022} \text{ (scale)}$$

CMS-PAS-SMP-12-028

Measuring α_s

Both are consistent with world average and renormalization group equation (RGE)

Summary

- QCD has been tested across a range of energies and phase spaces
  - Broadly speaking, it is performing remarkably well
  - Disagreement when looking into extreme areas of phase space
- Soft-QCD results help us to better understand the underlying event processes in pp collisions
- New pQCD measurements are providing powerful constraints on PDFs
- More results can be found here:
  - <https://twiki.cern.ch/twiki/bin/view/AtlasPublic/StandardModelPublicResults>
  - <https://twiki.cern.ch/twiki/bin/view/CMSPublic/PhysicsResultsSMP>
- 8 TeV analyses ongoing → expect new results soon!

BACKUP

Jet energy scale uncertainty

The jet energy scale uncertainty is the accuracy with which the calorimeters have been calibrated

- Determined by:
  - Looking for a jet recoiling off EM objects (photon or Z)
  - Test-beam data
  - MC simulation
- Typical range of systematic uncertainty (for $p_T > 20$ GeV):

1-3% (ATLAS)

1-5% (CMS)

ATLAS-CONF-2013-004

Jet energy resolution

The jet energy resolution is the inherent accuracy with which a jet can be measured

- Measured by looking for two objects with the same p_T and looking at the spread of their measured p_T
- Typical range of systematic uncertainty (for $p_T > 20$ GeV):

10-20%
(**ATLAS** and **CMS**)

CMS-PAS-JME-10-014

Underlying event (UE)

- Inclusive selection shows rising activity with p_T^{lead} , but exclusive selection gives flat behavior
- Relatively well described by the MC generators, but Pythia tuned to ATLAS UE data gives worse agreement

$\Phi(1020)$ cross section

- Measure cross section of $\Phi \rightarrow K^+K^-$ in MB events
- Probe the strangeness production and hadronization models at a scale of ~ 1 GeV
- Discriminating power among MC models: data best described by Pythia6 tune DW and EPOS-LHC

arXiv:1402.6162, sub to EPJC

Inclusive jet cross section

- Theory predicts a larger cross section than seen in data
- **ATLAS** 7 TeV data indicates the CT10 PDF description becomes worse at high p_T
- To some extent, MSTW 2008 predicts the trend seen in the data
- **CMS** 8 TeV data favors the high- p_T description from CT10

PRD 86 (2012) 014022

Ratio of inclusive jet cross sections

Many correlated uncertainties cancel when measuring ratio of cross sections at different CM energies

→ Strong constraint on the PDFs

Dijet cross section

8 TeV

CMS-PAS-SMP-14-002

Jet flavor in dijet events

- Unfolded fractions compared to MC at particle level
- Good data/MC agreement except for BU at high p_T

EPJC (2013) 73:2301

Jet substructure

JHEP 05 (2013) 090

- Boosted objects and jet substructure provide a new tool for separating QCD from the decay of heavy objects
- Several ways to examine jet substructure, which are all reasonably described in simulation
- In some cases, the more aggressive substructure techniques produce a better agreement between data and simulation

PDF sensitivity

- HERA data tightly constrain distributions for $x < 0.01$
- Combined fit using HERA-1 and **ATLAS**/**CMS** data leads to improved shape constraints
  - Both **ATLAS** and **CMS** data favor harder gluons (and softer sea quarks)

EPJC (2013) 73:2509

CMS-PAS-SMP-12-028

PDF sensitivity of the photon cross section

- Gluon PDF difference between different sets larger than quoted uncertainties
- NLO theory calculated using Jetphox and MCFM for different PDF sets
- χ^2 using all uncertainties from the experiment and theory (and correlations) using HERAFitter

ATL-PHYS-PUB-2013-018

Four-jet cross section

- First attempt at measuring $N_{\text{jet}} = 4$ double differential cross sections
- Softer jets poorly described by MC generators