On experimental neutrino physics

by **G. PLASS**, Nuclear Physics Apparatus Division

Neutrino physics is one of those fields of physics where, at present, we are at the limit of our possibilities in trying to extort from nature the answers to our many questions. This is primarily due to the very nature of neutrinos, which interact only 'weakly', with other matter and therefore extremely rarely. Nevertheless, the variety of the experiments that are done is surprising. Neutrino experiments* are carried out at accelerators (as at CERN), at reactors, and deep underground in coal mines or gold mines, where neutrinos of solar or cosmic origin can best be studied.

At the moment, these various approaches have two distinct aims: the accelerator and reactor experiments are designed to study the interaction mechanisms involved whilst those on neutrinos of extra-terrestrial origin are intended for the study of cosmological problems, such as the processes by which energy is produced in the sun and other stars or questions concerning the development of the universe.

Interactions induced by neutrinos were first observed in 1956, in an experiment carried out by Cowan, Reines *et al.* to detect neutrinos radiated by a nuclear reactor. This gave direct proof of the existence of these particles, superseding the previous indirect proof given by the momentum balance in radioactive beta decay (the particles radiated by a reactor arise from beta decay of the fission products and, to be exact, are now classed as antineutrinos), and thereby triggered off the whole field of experimental neutrino physics. Experiments similar to this pioneering one are still being carried on, as shown by two reports to the conference. One of these reports treated the interaction of neutrinos with protons and deuterons, whilst the other was concerned with what we may call a 'curiosity' — a detector of neutrino interactions with a volume of only two litres instead of the cubic metres that are normally used.

Neutrinos of natural origin

Nuclear processes involving neutrinos, of the kind we have recently been able to induce and detect on earth, are supposed to take place on a large scale in certain parts of the universe. Thus, an important field of neutrino physics, now in its initial stages of development, involves the search for neutrinos produced in these processes in the sun or in other celestial bodies. The measurement of the neutrino flux, of its variation with time, of its variation with direction, and so on, is of great interest for certain cosmological theories.

To obtain proper shielding against interactions induced by other cosmic particles, the detection

As noted in last month's CERN COURIER, an informal conference was held at CERN from 20-22 January 1965, on the subject of experimental neutrino physics. At this conference, which seems to have been a 'world première' and will probably turn out to be the first of a series, all aspects of neutrino physics were treated together for the first time. The accompanying article, based on the papers presented at that conference, summarizes the different kinds of neutrino experiment and gives information on various proposals for increasing the rates of observed neutrino events. This is followed by a discussion of the use of bubble chambers in neutrino experiments and of proposals for large new bubble chambers. The article ends with a brief summary of the present CERN plans for continuing experiments of this kind.

equipment in such experiments is most often installed in abandoned mine galleries. All strongly interacting particles are then completely absorbed by the rocks above, and even muons can only reach such depths if they arrive vertically from the zenith. A muon registered as travelling in a near-horizontal direction must be due to the interaction of a neutrino that has traversed the 'side-shield' formed by the earth itself. The deepest installation of this kind has been set up in the Kolar gold mines, in Southern India.

A similar experiment has also been proposed for setting up in the tunnel soon to be opened through the Mont Blanc massif, between Chamonix and Courmayeur. The equipment here would be located in a gallery pierced at right angles to the main tunnel, and the necessary shielding would be conveniently provided by the mountain range above.

The detectors developed for this kind of neutrino physics are of various kinds, but they all have the same basic characteristics: they must indicate the direction of flight of the particle registered and they must be simple and reliable, suited for 'runs' of many months duration. The Kolar experiment, for example uses large arrays of neon-filled flash tubes; at the University of Utah, an array of several hundred sonic spark counters, manufactured from iron pipes 15 cm in diameter and 10 m long, is being prepared.

Cherenkov detectors, registering the Cherenkov light emitted by fast-moving electrons from neutrino interactions, have also been proposed. It was even suggested that solar and cosmic neutrinos could be studied comparatively easily, just by lowering photomultipliers (perhaps combined with suitable light collectors) into the ocean depths and recording the Cherenkov light emitted by electrons and muons passing through the water. Liquid scintillators are another possibility for underground experiments, since they can be arranged rather easily in tanks to form a direction-sensitive 'telescope'. Radiochemistry can also be used, and a group at Brookhaven proposes to measure the flux of solar neutrinos by means of the reaction $\text{Cl} + \nu_e \rightarrow \text{A} + e^-$. Their 'detector' will be a tank containing some 400 000

* The term 'neutrino', in the context of this article, usually covers electron and muon neutrinos and electron and muon antineutrinos indiscriminately.

litres (100 000 gallons) of perchlorethylene, a dry-cleaning fluid. After several months a few of the chlorine atoms in the liquid are expected to be changed, as a result of neutrino interactions, into argon atoms and these could be extracted from the liquid and identified by their radioactivity.

Alternative search for lepton conservation law

It is interesting to note that at Brookhaven neutrino physics is being pursued by three different methods at the same time. Apart from this search for solar neutrinos with perchlorethylene and the continuation of the experiment at the AGS accelerator, confirmation of an important result of the high-energy neutrino experiments, namely the existence of a lepton conservation rule, is being sought for in a completely different way. In an experiment where extreme purity of materials and supreme patience are the essential ingredients (it takes two years to produce the necessary sample of the calcium isotope ^{48}Ca to begin the experiment) the possibility of so-called double beta decay is being examined. If this process, in which two electrons would be emitted from the nucleus without the emission of any corresponding neutrino, could be shown not to occur, the non-identity of neutrino and antineutrino would again be proved.

Neutrino experiments at accelerators

One may turn now to the field of neutrino physics at accelerators, where certainly most advances have been made in the past few years. The neutrinos here are produced in the decay of fast pions and kaons and have energies of around 1 GeV, that is between 100 and 1000 times greater than the energy of reactor-produced neutrinos. The interaction cross-section is perhaps 100 000 times higher and the event rates are of the order of 10 per day, per ton of detector, at present, probably reaching 100 per day in the not too distant future — much different to the few events per year that is all that can be anticipated in some of the experiments mentioned above. With these high rates and energies, the results of an interaction can actually be seen in spark chambers and bubble chambers and the processes can be analysed in detail; one does not have to be content just with the information that something has happened.

One of the main accents of the CERN neutrino experiments, in 1963 and 1964, particularly in the spark-chamber observations, was the question of the quantum of the field mediating the weak interactions, a boson usually known as the W. Though of very short life itself, it should produce decay products that would have been identified in both detectors — spark chambers and bubble chamber — of the CERN experiment, if the particle existed and had a mass small enough for it to be created by neutrinos of the energy available. An early paper in the conference was, in fact, concerned with the detailed proof that the mass of this particle must be greater than $1.9 \text{ GeV}/c^2$, that is, more than twice that of the proton. Unfortunately this leaves the question of the existence of such a particle completely open, since the theoretical upper limit of its mass is very much bigger.

Various other refinements to the results from the CERN heavy-liquid bubble chamber were treated in

* Slightly higher than the value deduced on the basis of the former incomplete analysis of the results.

Physics underground. Photograph exhibited by Dr. Reines at the conference, showing part of a detection system for cosmic neutrinos installed at a depth of over 3000 metres in a mine in South Africa. The aim is to detect muons arising from neutrino interactions in the earth on either side. Tanks of liquid scintillator are arranged in two rows about 40 metres long and 2 metres high, each row having an upper, middle and lower section, so that some information can be obtained on the angle to the horizontal made by the muon path.

a paper presented by the group concerned. For example, some of the so-called 'elastic' events had been reclassified, leading to a readjustment of the 'form factor' derived from their analysis. Concerning the fundamental question of the non-identity of the electron neutrino and the muon neutrino, the CERN results had of course amply confirmed the distinction, first established at Brookhaven in 1962**.

The progress of the related accelerator experiments at Brookhaven and Argonne was treated in several contributions. In both laboratories, much emphasis has been placed on the measurement of the neutrino energy spectrum, whereas at CERN, until recently, computed spectra served as the whole basis for the analysis (though in the end these were confirmed by experiment, within the limits of error). The essential result from the latest runs at Brookhaven was another determination of the lowest possible mass for the W boson, agreeing very well with the CERN results. At Argonne, the actual neutrino run had not yet started.

Facilities for accelerator experiments

An important part of the conference was reserved for a discussion of neutrino beams at accelerators and of neutrino detectors, for both existing and planned experiments, and to the question of the determination

** The reader may well ask what it means to establish sub-groups of a 'particle' that has none of the familiar qualities such as mass and electric charge and appears to be nothing but a package of directed energy. The neutrino was postulated because the mechanics of observed beta-decay processes could only be explained on the basis of a reaction involving three particles instead of the two that were seen. One 'neutrino' was then sufficient for all cases, but when the existence of antiparticles was confirmed for all other particles the question naturally arose whether there was an antineutrino distinguishable from a neutrino. On the basis of a conservation law for certain particles, a distinction between neutrino and antineutrino could be inferred and it was found that in fact certain reactions could only be caused by one, certain reactions only by the other, thus providing a good case for the distinction. In a similar way, since the electron and the muon appear to be identical apart from their mass, the question whether neutrinos (or antineutrinos) emitted with electrons (positive or negative) were the same as those emitted with muons was of obvious interest. We are thus faced with four entities, all of which have no electric charge and probably no rest mass but which are distinguishable by the interactions they induce. Unlike the photon (which also has no mass or charge), neutrinos cannot be identified as quanta of one of the familiar force fields. Because they obey Fermi-Dirac statistics, rather than Bose-Einstein statistics, they must be classed as 'particles' although their description can only be by sets of quantum-mechanical quantities.

of the neutrino spectrum. Methods included measurement of the muon flux in the shield, measurement of the flux of particles giving rise to the neutrinos — in auxiliary experiments using counter methods or a target immersed in a bubble chamber — and a proposal to measure this 'parent' flux directly, in the decay tunnel between target and shielding. The neutrino spectrum could also be determined from the energy measured in certain types of event; this could be the most reliable method, if the mechanism of weak interactions were already completely understood and if there were a greater number of events of the necessary kind.

Neutrino intensities in the accelerator experiments have already reached values a hundred or more times higher than expected when experiments without extracted beams were considered. However, the obtaining of a further increase in event rate is still one of the dominant problems in approaching a quantitative evaluation of the weak-interaction phenomena, and higher accelerator intensities would be of great value in this respect.

At Argonne, work is still directed mainly towards reaching the design goal of 10^{13} protons per pulse, but the possibility of replacing the present 15-MeV linac injector by one giving 200 MeV has already been considered as a further improvement. Both at Brookhaven and at CERN, substantial improvements are being planned in respect of the repetition rate, so that acceleration to top energy would become possible once every second (instead of once every 2.4 seconds or 3 seconds, respectively, as at present). Space-charge forces would still limit the intensity during each acceleration cycle to a value around 2×10^{12} protons per pulse, in both cases, and plans are therefore under consideration to install completely new linacs. At Brookhaven, this would provide injection into the synchrotron at 500 MeV, at CERN 200 MeV, instead of the present 50 MeV. Intensities of 10^{13} protons per pulse, or more, should then be possible, although serious problems of radiation damage and difficulties of servicing would in turn arise. These improvements

would take several years of materialize, in any case, and the necessary funds are not yet authorized.

Particularly at the present accelerated-proton intensities, efficient use of the available beam, by means of focusing devices for the neutrino parents and optimum design of shielding, are of primary importance. Talks were given on the 'plasma lens' developed at Brookhaven and on the 'magnetic horn' designed and used at CERN. The CERN device, which has also been developed at Argonne for use under somewhat different conditions, appears to give a rather higher gain than the plasma lens, though a fault that destroyed the power cables on the latter soon after it was put into operation prevented a detailed comparison.

As one of the highlights of the conference there came a proposal by R. Palmer, from Brookhaven, for a new type of focusing apparatus, the 'magnetic finger' or 'inverted horn'. An improvement of the flux to close on the theoretical limit is claimed for this device, which would mean an increase over present levels by a factor of two in most of the energy band and even more for the particles of highest energy. Detailed computations have been started at CERN in order to evaluate as soon as possible whether there would be any real advantage in replacing the existing installation with equipment of the new design.

New detectors for accelerator experiments

There is in principle another, very simple (though costly) way of increasing the event rate — just by increasing the mass of the detector. A first step in this direction is now being accomplished at CERN by increasing the visible volume of the CERN heavy-liquid bubble chamber to almost 1100 litres, the total volume being rather more than doubled. However, much useful information will still be lost, even with this increased volume, because of particles that pass out of the chamber without leaving a sufficient length of track to give full data for their identification. A much larger chamber of this type, with a diameter of 1.9 metres, is therefore being designed by a French

CERN/PI 145.2.64

A view of part of the spark-chamber installation for the second phase of the CERN neutrino experiment, now completed, taken during setting up of the equipment early in 1964. In the centre of the picture, the vertical spark-chamber modules can be identified by the white lines along the edges of the mirrors that enabled one camera to 'see' along all the gaps. Between the chambers are the magnetized iron plates that caused positive and negative particles to be deviated in opposite directions. On the right, the close-packed spark-chambers of the production region can be seen.

collaboration, with the hope that it can be built in the next few years for use at CERN. Since this chamber would contain 12 000 litres of liquid, with a visible volume of 8000-10 000 litres, the rate of analysable events would be increased at the same time; the actual improvement factor would depend on the properties of the beam and lie between the ratio of the chamber diameters and that of the chamber volumes.

Even so, there is still another important aspect of the picture. Certain data concerning the interactions can be extracted only with considerable error from events that occur in 'heavy liquid' or in the metal plates of spark chambers, because the primary interactions are obscured by secondary effects inside the large atomic nuclei. The metal plates of the spark chambers, generally aluminium or brass, or the freon liquid in the bubble chamber, as used so far, form the most suitable detectors in many respects, but have this disadvantage of being composed solely of large nuclei. There is hence considerable interest in studying at least some interactions with hydrogen nuclei, which of course consist of just single nucleons. One way of doing this is to fill the heavy-liquid chamber with propane. The density, and hence the event rate, would be only about one third of that for freon, but there would in fact be more hydrogen in the chamber than in a similar volume of pure liquid hydrogen. The carbon atoms in the propane would, unfortunately, still disturb certain experiments, but on the other hand they would permit the carrying out of others (polarization experiments, for example) that would be difficult to perform otherwise. For the study of processes occurring in almost free neutrons, propane containing deuterium instead of ordinary hydrogen could be used, though this would be expensive to manufacture.

An approach from the opposite angle has been made at Brookhaven, where early tests have shown that a hydrogen bubble chamber can be operated successfully with a high proportion of neon mixed with the hydrogen. This would enable the overall density of the liquid to be increased somewhat whilst still retaining a useful number of hydrogen nuclei in the target volume.

For completely unambiguous results on the interaction between neutrinos and protons, it remains true to say that liquid hydrogen by itself (or liquid deuterium if one wants to look specifically at neutrino reactions on neutrons) remains the ideal substance. Experiments in the 2-m hydrogen bubble chambers at both Brookhaven and CERN have therefore been considered, but the low density will give an event rate that will be below that needed for the study of many rare kinds of interactions. Partly because of this, but also for more general reasons, huge hydrogen chambers, with volumes between 25 000 and 40 000 litres, have been proposed at Argonne, Brookhaven and CERN.

Theoretical ideas

One session of the conference was devoted to theoretical contributions. In an invited review paper, also, the problems in weak-interaction physics that underlie the accelerator experiments were formulated. In particular, a summary was given of some recent suggestions that certain basic hypotheses in weak-interaction theory (conserved vector currents and partially

Although the conference dealt primarily with the experimental aspects of neutrino physics, theory could not be forgotten. The final talk, in fact, was given by Nobel Prize winner Prof. T. D. Lee, seen here as he resumed the basic problems in the theory of neutrino interactions.

conserved axial-vector currents) could be tested in polarization experiments with neutrinos. Unfortunately, there was no similar, concise review of the motivation for the experiments on extraterrestrial neutrinos.

Looking ahead

The field of experimental neutrino physics at accelerators for the next decade was thus sketched in. With higher accelerator intensities, improved neutrino facilities and enlarged detectors, one may expect neutrino physics after 1970 to be done with overall event rates that are customary now in strong-interaction experiments. One paper, a contribution from the Berkeley study group for a 200-GeV synchrotron, looked even further; for neutrino experiments at this accelerator a magnetic-lens channel about 1 km long is projected, going so far as to provide the possibility of selecting neutrinos in a certain momentum band at the end of it. Before this time the Russian 70-GeV accelerator at Serpukhov will go into operation — in 1968 according to present plans. A brief status report was given on this machine, but detailed studies on the experimental arrangements do not appear to have been carried out yet.

'Think big', in the words of Prof. Goldhaber, Director of Brookhaven National Laboratory and one of the two closing speakers, appeared to be the motto of the conference. But it is not only the future projects that seem so huge; our present-day installations would have appeared like science fiction only 25 years ago. To end, then, we might look at the more immediate plans for the CERN neutrino facility. A run with the enlarged heavy-liquid chamber filled with freon, in an antineutrino beam, is scheduled for the spring of this year. The chamber will then be used for an experiment on another problem, but plans are being made to prepare it after this for a propane filling. The possibilities of obtaining deuterated propane are also being investigated, with a view to the later use of this material, as already described. At the same time, the possibility of using spark chambers in polarization experiments, of the type mentioned in the report on the theoretical contributions, is also being looked into at CERN ●