EUROPEAN ORGANIZATION FOR NUCLEAR RESEARCH

CERN-EP/86-189
18 November 1986

THE BETA-DECAY HALF-LIFE OF $^{130}_{48}\text{Cd}_{82}$ AND ITS
IMPORTANCE FOR ASTROPHYSICAL r-PROCESS SCENARIOS

K.-L. Kratz¹, H. Gabelmann², W. Hillebrandt³, B. Pfeiffer¹,
H.L. Ravn², F.-K. Thielemann⁴,

¹ Institut für Kernchemie, Universität Mainz, D-6500 Mainz,
Federal Republic of Germany

² CERN-ISOLDE, CH-1211 Geneva, Switzerland

³ MPI für Physik und Astrophysik, D-8046 Garching, Federal
Republic of Germany

⁴ Department of Astronomy, Harvard University, Cambridge,
MA 02 138, U.S.A.

and

the ISOLDE Collaboration, CERN, Geneva, Switzerland

Submitted to Z. Phys. A - Atoms and Nuclei

Abstract: The β -decay half-life of ^{130}Cd has been measured and its importance as $N = 82$ 'waiting-point' nucleus for astrophysical rapid neutron capture (r-) process scenarios is discussed.

The investigation of β -decay properties of ^{130}Cd is of considerable interest for both nuclear and astrophysics. With regard to nuclear structure, experimental information on $J^\pi = 1^+$ two-quasi-particle (2QP) configurations in the daughter ^{130}In , for example, is of importance for the understanding of the basic shell-model states in the valence nuclei around the doubly-closed shell nucleus ^{132}Sn (see, e.g. [1-3]). As far as astrophysics is concerned, the β -decay half-life ($T_{1/2}$) of ^{130}Cd as a 'waiting-point' nucleus is of interest (see, for example [4-7]).

The solar-system isotopic abundances of nuclei produced in the r-process show two major maxima at $A = 130$ and $A = 196$ (see, e.g. Fig. 1 of [8]). These maxima are caused by the magic neutron numbers $N = 82$ and $N = 126$ where the r-process path comes closest to the valley of β -stability (see Fig. 1) and falls on nuclei with longer-than-average $T_{1/2}$. As a result, the abundances of these nuclei will build up as they wait to decay. These 'waiting points' govern the time scale of the r-process and give constraints on astrophysical scenarios. For example, until now it is not possible to decide whether the duration of the r-process is sufficiently long so that a steady-flow solution for the β -decay reaction equations is established in a $(n, \gamma) - (\gamma, n)$ equilibrium, or whether the neutron-exposure times are only of the order of individual β -decay times. The first alternative was originally suggested by Burbidge et al. (B²FH) in 1957 [4], whereas the second possibility applies, for example, to He-burning scena-

rios investigated more recently (see, e.g. [6,7]).

Since for the $N = 126$ magic number, the r-process must pass very-neutron-rich nuclei like ^{196}Yb which will be out of experimental reach in the near future, the only area accessible to experiment is that of the $N = 82$, $A = 128-130$ abundance maximum. In this region, the nucleus closest to the valley of β -stability with the presumably longest $T_{1/2}$ is ^{130}Cd which lies 14 neutrons away from the heaviest stable Cd isotope. The measurement of even this single $T_{1/2}$ in the r-process path may already be sufficient to determine the scenario in which the heavy elements in nature are formed.

Fig. 1: Illustration of the r-process path (solid line). At the numbers $N = 82$ and 126 the r-process path comes closest to the line of β -stability. For $N = 82$, ^{130}Cd is expected to be the longest-lived isotope in the r-process path.

The isotope ^{130}Cd was produced at the ISOLDE on-line mass separator facility connected to the CERN-SC which supplied a beam of 600 MeV protons at a current of 2.2 μA . The target consisted of uranium carbide in a graphite-cloth matrix, heated to 2,200 $^{\circ}\text{C}$. A plasma-discharge ion source was connected to the target via a heated quartz tube. This tube retained the isobar ^{130}In and suppressed the many orders of magnitude higher activity of ^{130}Cs , allowing preferential extraction of ^{130}Cd . After mass separation, the beam was guided by ion optics to a well shielded counting cabin where sources were collected on an aluminized tape at the detection position. For acquisition of multiscaling curves, the tape was moved periodically in order to remove the background of long-lived activities. Although the extraction of 29 min ^{130}Cs could be considerably suppressed, its activity was still too high to allow β -counting or γ -spectroscopy. Therefore, the only way of identifying ^{130}Cd proved to be the selective detection of β -delayed neutrons (βdn). This was done by using a neutron long-counter which consisted of eight parallel-coupled ^3He -filled proportional counters embedded in a paraffine matrix. The absolute efficiency of this neutron counter was determined to $(9.5 \pm 0.5) \%$. However, even with neutron counting, the occurrence of (a priori unexpected) $\text{CaBr}^+ \beta\text{dn}$ -activities - although having only per-mill yields of the Br^+ yields - severely complicated the experimental conditions. In consequence, very short collection times had to be chosen by use of pulsed-beam techniques in order to enhance ^{130}Cd and minimize the 1.9 s βdn -background from $^{40}\text{Ca}^{90}\text{Br}^+$ at $A = 130$. Data from about 36,000 growth (300 ms) and decay (900 ms) cycles were accumulated at 10 ms intervals. After subtraction of normal neutron background and the ^{90}Br βdn -component, the growth and decay curve was analyzed by least-squares fitting.

From the βdn -growth curve the $T_{1/2}$ of ^{130}Cd was determined to (195 ± 35) ms. The complex ^{130}Cd - ^{130}In decay curve yielded only an approximate $T_{1/2}$ of 230 ms. These values can be compared with a RPA shell model [9] prediction of 0.3 s for pure Gamow-Teller β -decay, implying that in addition to GT decay also fast ($\log ft \approx 6.0$) first-forbidden (near-) ground-state transitions must occur in ^{130}Cd . By relating the absolute βdn -saturation activity to the production yield of ^{130}Cd extrapolated from the measured β -activities of lighter-mass Cd isotopes, we obtain a βdn -emission probability (P_n) of the order of 4 %. This rather low P_n value is in reasonable agreement with our RPA prediction of 2.5 %. On the basis of the calculated $J^\pi = 1^+ 2\text{QP}$ level scheme, this implies that the neutron binding energy in ^{130}In should lie in the range 4.9 to 5.2 MeV, to be compared to (5.03 ± 0.37) MeV from the 1985 Wapstra-Audi mass evaluation [10].

From the measured $T_{1/2}$ of ^{130}Cd we now can derive an important and new constraint on the mode of operation of the r-process which relies on experimental data only. If we assume that the equilibrium steady-state solution is realized in

nature, the r-process equations reduce to [5]:

$$\frac{n(Z)}{n(Z-1)} = \frac{\tau_{\beta}(Z)}{\tau_{\beta}(Z-1)} \quad (1)$$

where $n(Z) = \sum_A n(Z,A)$ is the sum of the number densities in an isotopic chain Z and $\tau_{\beta} = [\sum_A \tau_{\beta}^{-1}(Z,A) \cdot n(Z,A)/n(Z)]^{-1}$ is the effective β -decay half-life. Near the neutron magic number $N = 82$ the abundances in the r-process path are strongly peaked at ^{130}Cd ($Z=48$) and at $^{131,132}\text{In}$ ($Z=49$). Since the P_n values are small for these nuclei [11], the observed abundances of ^{130}Te and $^{131,132}\text{Xe}$ will be almost equal to those of the r-process nuclei ^{130}Cd and $^{131,132}\text{In}$, respectively. The solar abundance of ^{130}Te and the r-process contributions to $^{131,132}\text{Xe}$ are fairly well known [8], and we obtain for the abundance ratio $n(^{130}\text{Cd})/[n(^{131}\text{In}) + n(^{132}\text{In})] \approx n(^{130}\text{Te})/[n(^{131}\text{Xe}) + n(^{132}\text{Xe})] \approx 0.91$. From the measured $T_{1/2}$ of $^{131,132}\text{In}$ [2,11] we find an effective $\tau_{\beta}(Z=49)$ of 225 ms. Therefore, from eq. (1), the $T_{1/2}$ of ^{130}Cd should be (205 ± 20) ms if the mass peak at $A = 130$ indeed was formed in a steady state. The main uncertainties in this argument come from the neglect of βdn -emission which, however, will not modify the value for $\tau_{\beta}(^{130}\text{Cd})$ by more than 10 %.

The good agreement with our measured $T_{1/2}$ leads us, therefore, to the conclusion that the neutron-exposure times for the r-process have been sufficiently long (≥ 5 to 10 s) so that a steady state was established. This rules out short-duration scenarios [5,6], unless the above agreement would be accidental. This possibility could, however, be excluded by an independent check, e.g. via the $N = 82$ isotone ^{129}Ag . From the same arguments given before, we predict a $T_{1/2}$ of about 150 ms for ^{129}Ag which should, in principle, be measurable.

We would like to acknowledge valuable discussions with J.W. Truran. This work was supported by the Deutsche Forschungsgemeinschaft through grant Kr 806/1-2.

References:

1. Blomqvist, J.: Proc. 4th Int. Conf. on Nuclei far from Stability, CERN 81-09, 536 (1981).
2. Björnstad, T. et al.: Z. Phys. **A306**, 95 (1982).
3. Fogelberg, B. and Blomqvist, J.: Nucl. Phys. **A429**, 205 (1984); and Refs. therein.
4. Burbidge, G.R. et al.: Rev. Mod. Phys. **29**, 547 (1957).
5. Hillebrandt, W.: Sp. Sci. Rev. **21**, 639 (1978).
6. Thielemann, F.-K. et al.: Z. Phys. **A309**, 301 (1983).
7. Cameron, A.G.W. et al., in: Nucleosynthesis, Challenges and New Developments; eds. Arnett, W.D. and Truran, J.W.; Univ. of Chicago Press, p. 190 (1985).
8. Käppeler, F. et al.: Ap. J. **257**, 821 (1982).
9. Krumlinde, J. and Möller, P.: Nucl. Phys. **A417**, 419 (1984).
10. Wapstra, A.H. and Audi, G.: Nucl. Phys. **A432**, 1 (1985).
11. Lund, E. et al.: Z. Phys. **A294**, 233 (1980).