***B*-hadrons spectroscopy and decay properties with the ATLAS experiment**

Tomáš Jakoubek

(Institute of Physics ASCR, Prague)
on behalf of the ATLAS Collaboration

tomas.jakoubek@cern.ch

QCD 2014, Montpellier, France

01. 07. 2014

Outline

- The ATLAS detector overview
- B -physics triggers
- Selection of related topics discussed today:
 - Measurement of the Λ_b^0 lifetime and mass [Phys. Rev. D 87, 032002 (2013)]
 - Measurement of the parity-violating asymmetry parameter α_b and the helicity amplitudes for the decay $\Lambda_b^0 \rightarrow J/\psi \Lambda^0$ with the ATLAS detector [Phys. Rev. D 89 092009 (2014)]
 - Observation of an excited $B_c^\pm$ meson state with the ATLAS detector [ATLAS-BPHY-2012-04-002, to be submitted to Phys. Rev. Lett.]

The ATLAS Detector

- Multipurpose detector at LHC, optimized for high p_T physics and instantaneous luminosity up to $10^{34} \text{ cm}^{-2}\text{s}^{-1}$
- Inner Detector
 - $|\eta| < 2.5$
 - Solenoid $B = 2 \text{ T}$
 - Si pixel, Si strips, Transition Radiation Tracker
 - Impact parameter res. $15 \mu\text{m}$ in $r\phi$, $100 \mu\text{m}$ in z
 - Momentum res. $\Delta p_T/p_T \sim 0.04 \%$
 $p_T [\text{GeV}] \otimes 1.5 \%$

- Muon Spectrometer

- $|\eta| < 2.7$
- Toroid average $B = 0.5 \text{ T}$ (barrel) or 1 T (end-caps)
- Momentum res. $\Delta p_T/p_T < 10 \%$ up to $\sim 1 \text{ TeV}$

B-physics triggers

- ATLAS has 3-level trigger system (L1 - hardware, L2 and Event Filter - computer farms)
- From 40 MHz to $O(200)$ Hz
- B-physics triggers are based on single and di-muons...
- Topological triggers process two L1 muon Rols and refine results in the HLT with a good vertex fit and mass cut
- TrigDiMuon triggers require one L1 muon Rol and then search for a second muon in ID tracks

Trigger name	Mass window	# of evts.
2mu4_Jpsimumu	2.5-4.3 GeV	14 M
2mu4_Upsimumu	8-12 GeV	9.1 M
2mu4_Bmumu	4-8.3 GeV	3.7 M

Example with 2.3 fb^{-1} data

Measurement of the Λ_b^0 lifetime and mass

- Λ_b^0 is the lightest baryon containing a b -quark (5620 MeV)
- Not produced at B -factories $\Rightarrow$ currently hadron colliders are the only facilities to study b -baryons properties
- Lifetime, although measured by many experiments, still suffers from a large experimental uncertainty
- Data from 2011 proton-proton 7 TeV collisions, used integrated luminosity 4.9 fb^{-1}
- Reconstructed via $\Lambda_b^0 \rightarrow J/\psi (\mu^-\mu^+) \Lambda^0 (p\pi^-)$
- Topological J/ψ trigger chain
- Candidate selection:
 - Global $\chi^2/\text{n.d.o.f} < 3$
 - $p_T(\Lambda^0) > 3.5 \text{ GeV}$
 - $L_{xy}(\Lambda^0) > 10 \text{ mm}$
 - $5.38 \text{ GeV} < m(J/\psi \Lambda^0) < 5.90 \text{ GeV}$

Measurement of the Λ_b^0 lifetime and mass

- Unbinned maximum likelihood fit to mass and lifetime:
 - Mass: signal – Gaussian, background – 1st order polynomial
 - Lifetime: signal – exponential function, prompt background – sum of δ -function and a symmetric exponential distribution, non-prompt background – exponential function
- $m = 5619.7 \pm 0.7$ (stat) ± 1.1 (syst) MeV
- $\tau = 1.449 \pm 0.036$ (stat) ± 0.017 (syst) ps

Measurement of the Λ_b^0 lifetime and mass

- Comparison with other experiments

Parity violation in the decay $\Lambda_b^0 \rightarrow J/\psi \Lambda^0$

- Parity violation – well-known feature of weak interactions – not maximal in decays of hadrons due to the presence of strongly bound spectator quarks
- Results of parity violation measurement can be used to test predictions made by different quark interaction models...
- **pQCD** predicts α_b (parity-violating asymmetry parameter) to be in the range **from -0.17 to -0.14**, while calculations based on **HQET** predicts a value of **~0.78**
- Decay described by 4 helicity amplitudes $A(\lambda_\Lambda, \lambda_{J/\psi})$ normalized to 1:
 - $a_+ = A(1/2, 0)$
 - $a_- = A(-1/2, 0)$
 - $b_+ = A(-1/2, -1)$
 - $b_- = A(1/2, 1)$
- Parameter $\alpha_b = |a_+|^2 - |a_-|^2 + |b_+|^2 - |b_-|^2$

Parity violation in the decay $\Lambda_b^0 \rightarrow J/\psi \Lambda^0$

- Full angular PDF:

$$w(\Omega, \vec{A}, P) = \frac{1}{(4\pi)^3} \sum_{i=0}^{19} f_{1i}(\vec{A}) f_{2i}(P, \alpha_\Lambda) F_i(\Omega)$$

f_{1i} ... bilinear combinations of the helicity amplitudes

f_{2i} ... $P\alpha_\Lambda$, P , α_Λ or 1

F_i ... moments of the angular distribution

- pp collisions symmetry + ATLAS η symmetry $\rightarrow$ overall polarization $P = 0 \rightarrow$ reducing PDF from 20 to 6 terms
- Helicity amplitudes and decay asymmetry parameter for Λ_b^0 extracted by least square fit from measured averages of each F_i :

$$\langle F_i \rangle = \frac{1}{N^{\text{data}}} \sum_{n=1}^{N^{\text{data}}} F_i(\Omega_n)$$

$$\chi^2 = \sum_i \sum_j (\langle F_i \rangle^{\text{expected}} - \langle F_i \rangle) V_{ij}^{-1} (\langle F_j \rangle^{\text{expected}} - \langle F_j \rangle)$$

V_{ij} ... covariance matrix of measured $\langle F_i \rangle$; $\langle F_i \rangle^{\text{expected}}$ is evaluated from models including detector effects and depends on the helicity amplitudes

Parity violation in the decay $\Lambda_b^0 \rightarrow J/\psi \Lambda^0$

- Background contribution in the signal region can be estimated as an average of the values in the two sidebands

- Results checked by comparing F_i distributions for data with weighted signal MC plus sideband background

Parity violation in the decay $\Lambda_b^0 \rightarrow J/\psi \Lambda^0$

- Helicity amplitudes:

$$|a_+| = 0.17_{-0.17}^{+0.12}(\text{stat}) \pm 0.09(\text{syst})$$

$$|a_-| = 0.59_{-0.07}^{+0.06}(\text{stat}) \pm 0.03(\text{syst})$$

$$|b_+| = 0.79_{-0.05}^{+0.04}(\text{stat}) \pm 0.02(\text{syst})$$

$$|b_-| = 0.08_{-0.08}^{+0.13}(\text{stat}) \pm 0.06(\text{syst})$$

- Decay asymmetry parameter:

$$\alpha_b = 0.30 \pm 0.16 (\text{stat}) \pm 0.06 (\text{syst})$$

- Large $|a_-|$ and $|b_+| \rightarrow$ negative helicity states for Λ^0 preferred
- Λ^0 and J/ψ are highly polarized in the direction of their momenta
- α_b consistent with LHCb (Phys. Lett. B 724 (2013) 27):
 $\alpha_b = 0.05 \pm 0.17 (\text{stat}) \pm 0.07 (\text{syst})$
- α_b differs from pQCD expectation (-0.17 to -0.14) by $\sim 2.6\sigma$ and from HQET expectation (0.78) by $\sim 2.8\sigma$
- Analysis using 2012 data is ongoing

Observation of an excited $B_c^\pm$ state

- Excited states of $B_c^\pm$ meson have not previously been observed
- Spectrum and properties predicted by non-relativistic potential models, pQCD, and lattice calculations
- Predicted $B_c^\pm(2S)$ mass is in the range 6835 – 6917 MeV, production cross-section should be 15–20 % of the ground state; production of P and D states is highly suppressed relative to that of $B_c^\pm(2S)$ state
- Data from 2011 (7 TeV) and 2012 (8 TeV) proton-proton collisions, used integrated luminosity 4.9 fb^{-1} (2011) + 19.2 fb^{-1} (2012)
- New state observed through its hadronic transition to the ground state, with $B_c^\pm \rightarrow J/\psi (\mu^-\mu^+) \pi^\pm$
- To minimize uncertainties due to detector resolution, peaks are sought in the distribution of the variable

$$Q = m(B_c^\pm \pi \pi) - m(B_c^\pm) - 2m(\pi^\pm)$$

Observation of an excited B_c meson state

- Invariant mass of $B_c^\pm \rightarrow J/\psi \pi^\pm$ (unbinned maximum likelihood fit)

- The $m(B_c^\pm \pi \pi) - m(B_c^\pm) - 2m(\pi^\pm)$ mass difference distribution, wrong-charge combinations also shown (unbinned maximum likelihood fit)

Observation of an excited B_c meson state

- Systematics:
 - Uncertainty coming from the statistical uncertainty of the ground state mass mostly cancels in Q -value distribution
 - Uncertainty from the fit, estimated with various fit models including toy MC studies basing on the background shape from the wrong-charge combinations (~ 3.4 MeV) and varying the fit mass range from 0 – 700 MeV to 0 – 1500 MeV (~ 1.2 MeV); these propagate to $m(B_c^\pm(2S))$ as ~ 3.6 MeV
- Mass of the peak
$$m(B_c^\pm(2S)) = 6842 \pm 7 \text{ (stat)} \pm 5 \text{ (syst) MeV}$$
consistent with the theory (6835 – 6917 MeV), as well as the decay mode
- 5.2 sigma observation for the combination of 7 and 8 TeV datasets

Summary and Outlook

- ATLAS has produced impressive and competitive results in heavy quark hadrons spectroscopy
- Presented today:
 - Λ_b^0 mass and lifetime measurement in agreement with other experiments
 - Parity violation parameter in decay $\Lambda_b^0 \rightarrow J/\psi \Lambda^0$ consistent with LHCb result, but lies between two theoretical predictions (pQCD and HQET); updates with 2012 data ongoing
 - Observation of a new excited $B_c^\pm$ meson state, $B_c^\pm(2S)$, mass consistent with the theory
- For RUN-2, insertion of Pixel Insertable B-Layer (IBL) should provide increasing vertexing performance
- More public results on ATLAS *B*-physics TWiki page:

<https://twiki.cern.ch/twiki/bin/view/AtlasPublic/BPhysPublicResults>

BACKUP

Observation of an excited B_c meson state

- Topological J/ψ trigger chain
 - Early 2011: both muons with $p_T > 4$ GeV
 - Later 2011 and 2012: $p_T(\mu_H) > 6$ GeV, $p_T(\mu_S) > 4$ GeV
- $B_c^\pm$ candidate selection:
 - J/ψ muons p_T cuts according to trigger
 - π track $p_T > 4$ GeV in both 2011 and 2012
 - $p_T(B_c^\pm) > 15$ GeV (2011) or > 18 GeV (2012)
 - Cut on π track impact parameter (XY-plane projection) to reduce combinatorial background: $d_{xy}^0/\sigma(d_{xy}^0) > 5$ (2011) or > 4.5 (2012)
 - J/ψ π vertex $\chi^2/\text{n.d.o.f} < 2$ (2011) or < 1.5 (2012)
- Excited $B_c^\pm$ candidate selection:
 - $B_c^\pm$ candidate mass in 3σ interval around its PDG mass
 - π tracks with $p_T > 400$ MeV; from the corresponding primary vertex (PV)
 - Refitted triplet of $B_c^\pm$ tracks and the pair of π tracks from PV must intersect in two separate $B_c^\pm$ and $B_c^\pm(2S)$ vertices
 - Taking a candidate with the best χ^2 of the combined cascade fit with the ground state