EUROPEAN ORGANIZATION FOR NUCLEAR RESEARCH

CERN-EP/87-184
September 30th, 1987

Production of Λ_c with Large- x_F at the ISR

(R608 Collaboration)

P. Chauvat¹, R. Cousins², K. Hayes³, A.M. Smith
CERN, 1211 Geneva 23, Switzerland.

R. Bonino⁴, A. Castellina⁵, S. Erhan, M. Medinnis,
P.E. Schlein, P. Sherwood⁶, S. Vernetto⁵, J.G. Zweizig
University of California^{*)}, Los Angeles, California 90024, USA.

J. Alitti, B. Bloch-Devaux, J.B. Cheze, J. Mallet,
A. Montag⁷, B. Pichard, J. Zsembery
Centre d'Etudes Nucleaires - Saclay, 91191 Gif-sur-Yvette, France.

L. Meritet, M. Reyrolle⁸, F. Vazeille
LPC-IN2P3, Universite de Clermont-Ferrand, 63 Aubiere, France.

Abstract

We report results of an inclusive measurement of the charmed baryon, Λ_c , in the forward direction of proton-proton interactions with $\sqrt{s} = 63$ GeV. The signal, seen in the final state $\Lambda_c \pi^+ \pi^+ \pi^-$, has 6.0σ significance with about 620 events on a large background and corresponds to a cross section times branching ratio of $2.84 \pm 0.50 \pm 0.72$ μbarn for Feynman- $x_F > 0.5$ in both hemispheres. Asymmetries are observed in the decay angular distributions which are characteristic of weak decays and have a combined 3.0σ significance.

Submitted to Physics Letters B

^{*)} Supported by U.S. National Science Foundation grants PHY79-24766 and PHY85-09175

¹⁾ present address: 29 Ave. General LeClerc, 78220 Viroflay, France

²⁾ now at University of California, Los Angeles, CA, USA

³⁾ now at Stanford Linear Accelerator Center, Stanford, CA, USA

⁴⁾ now at Istituto Nazionale de Fisica Nuclere, Rome, Italy

⁵⁾ now at Istituto di Cosmogeofisica del C.N.R., Torino, Italy

⁶⁾ now at University College, London, UK

⁷⁾ now at Weizmann Institute, Rehovoth, Israel

⁸⁾ now at CERN, Geneva, Switzerland

We report final results of an inclusive measurement of Λ_c production in the reaction

$$pp \rightarrow \Lambda_c^+ + X \rightarrow (\Lambda_s^0 \pi^+ \pi^+ \pi^-) + X \quad (1)$$

with $\sqrt{s} = 63$ GeV at the CERN Intersecting Storage Rings. These reinforce earlier results[1,2,3,4] and have higher statistical significance.

The apparatus used for these measurements was the forward multi-particle spectrometer[5,6,7] of Experiment R608. The apparatus had an aperture of ± 150 mrad measured from the outgoing beam direction and contained a two-threshold Cerenkov counter system with 16 cells and a 16,448 wire mini-drift chamber system for tracking. The trigger used for the data acquisition of Reaction 1 included multiplicity conditions in the wire chambers which demanded that five or more tracks traversed the spectrometer. No requirements were imposed on particles which were outside the aperture of the spectrometer, thus insuring that inclusive cross sections were measured for states observed in the spectrometer. Off-line analysis of the reconstructed tracks yielded events with $\Lambda_s \rightarrow p\pi^-$ about 5% of the time.

Because of the difficulty of finding Λ_c in an inclusive hadronic spectrum with a large combinatoric background, we have taken great care in determining chamber alignment, wire positions and magnetic field calibration. As part of this calibration program, we have analyzed our signals for Λ_s , K^0 and $\Lambda_s^*(1520) \rightarrow K^- p$, in addition to the exclusive final state with five forward tracks:

$$pp \rightarrow (p\pi^+\pi^-\pi^+\pi^-)p \quad (2)$$

in which there is a clear signal for $f_1(1285) \rightarrow 4\pi$. This state is especially interesting because the 726 MeV Q-value in its decay is comparable to that of the Λ_c in Reaction 1. Fig. 1(a) shows the invariant mass spectra for the Λ_s in the $\Lambda_s 3\pi$ data sample. Fig. 1(b) shows the $f_1(1285)$ in Reaction 2. The fitted curve through the data points in Fig. 1(b) is the sum of a second order polynomial and a Breit-Wigner function ($\Gamma = 25$ MeV) folded with a Gaussian resolution function of $\sigma = 12$ MeV. The fitted mass of the resonance is 1279.6 ± 1.6 MeV, which agrees well with the Particle Data Group[8] value of 1283 ± 5 MeV. The fitted mass of the $\Lambda_s^*(1520)$ shown in Fig. 1(c) is 1519.1 ± 0.3 MeV, compared with the world average of 1519.5 ± 1.0 MeV.

Fig. 2(a) shows the invariant mass spectrum of the $\Lambda_s \pi^+ \pi^+ \pi^-$ system in Reaction 1 when Feynman- x_F and transverse momentum of the $\Lambda_s \pi^+ \pi^+ \pi^-$ system are in the ranges, $0.5 - 0.9$ and $p_T < 1.1$ GeV/c, respectively. The fit shown in Fig. 2(a) is the sum of a third order polynomial and a Gaussian distribution with $\sigma = 14$ MeV, corresponding to the resolution in our invariant mass measurement. In performing the fit, the free parameters were the central mass of the Gaussian component, its fraction of the histogram's area and the three shape constants of the polynomial. The fitted number of signal events in the Gaussian component of Fig. 2(a) is 621 ± 103 , a 6.0σ (standard deviation) effect. The central mass of the best fit is $(2305 \pm 3 \pm 6)$ MeV (statistical and systematic uncertainties, respectively).

Our mass value for the Λ_c in Reaction 1 may be compared with the 1986 world average of 2281.2 ± 3.0 MeV quoted by the Particle Data Group[8] and several newer values: 2285.6 ± 1.1 MeV (statistical error only)[9], 2289 ± 2 MeV[10] and 2284.7 ± 2.3 MeV[11]. We have, in addition, refit the mass spectra of the R603 collaboration[1] and find a value of 2295 ± 6 MeV. Since a minimum bias trigger was used in Ref. [1], this latter value is interesting despite its larger error.

The signal seen in Fig. 2(a) is tracked as a function of x_F of the $\Lambda_s \pi^+ \pi^+ \pi^-$ system by performing independent fits of the type used in Fig. 2 on data samples in bins of x_F . The resulting signal magnitudes are plotted in Fig. 3. The solid and dashed curves in Fig. 3 show the expected (uncorrected) distributions if the Λ_c is produced with a cross section of the form $d\sigma/dx_F \propto (1 - x_F)^n$, with $n = 1$ and 2 , respectively.

Fig. 4 shows the acceptance-corrected differential cross section times branching ratio, $Bd\sigma/dx_F$, for Reaction 1 integrated over all p_T , for bins of $\Delta x_F = 0.1$ from 0.5 to 1.0 . The acceptance correc-

tions are model independent (except that they assume isotropic decay of the Λ_c) and account for all geometrical and other effects, such as pion decay, absorption, tails of cut distributions such as vertex χ^2 , etc., neutral Λ_s decay and include a factor of 2 for both arms in reaction 1. The solid circle points in Fig. 4 have the values: (15.4 ± 4.2) , (7.1 ± 2.2) , (4.3 ± 1.3) and (1.6 ± 0.7) μbarn . A fit of $Bd\sigma/dx_F = (1-x_F)^n$ to these points¹ yields $n = 2.1 \pm 0.3$. In a similar way, the dependence on transverse momentum is determined to be $d\sigma/dp_t^2 \propto \exp(-bt)$ with $b = (3.4 \pm 0.4) \text{ GeV}^{-2}$. Fig. 4 also shows $d\sigma/dx_F$ for inclusive Λ_s production reported by Erhan et al.[12]. The Λ_c is seen to have a somewhat steeper dependence on x_F than does the Λ_s .

The integrated cross section times branching ratio for the points with $x_F > 0.5$ in Fig. 4 is $B\sigma = 2.84 \pm 0.50 \pm 0.72 \mu\text{barn}$, where we estimate a 25% systematic uncertainty from the correction procedure. This may be compared with the earlier inclusive measurement of Reaction 1 made by Ref. [1] for $x_F > 0.75$. With our x_F -dependence presented above, $(15 \pm 3)\%$ of our measured cross section, or $0.42 \pm 0.18 \mu\text{barn}$, falls in the range $x_F > 0.75$. This is $\approx 2.5\sigma$ below the older (two hemisphere) value $B\sigma = 5.6 \pm 2.0 \mu\text{barn}$, reported in Ref. [1].

In order to obtain the inclusive cross section for $pp \rightarrow \Lambda_c + X$ from our measurements of $Bd\sigma/dx_F$, we use the branching ratio for $\Lambda_c \rightarrow \Lambda_s 3\pi$, $(2.8 \pm 0.7 \pm 1.1)\%$, reported by Bowcock et al.[13]. The results are shown as the solid squares in Fig. 4. It should be noted, however, that there is some uncertainty in this procedure. The ratio $(\Lambda_c \rightarrow \Lambda_s 3\pi)/(\Lambda_c \rightarrow K^- p \pi^+) = (0.61 \pm 0.16)$, reported by the ARGUS collaboration[14], when combined with the branching fraction, $(2.0 \pm 0.8)\%$, for $\Lambda_c \rightarrow K^- p \pi^+$ from the MARK II collaboration[15], leads to a smaller value for the $\Lambda_c \rightarrow \Lambda_s 3\pi$ branching fraction of $(1.2 \pm 0.6)\%$. However, the LEBE-EHS Collaboration[16] has reported a lower limit to the $\Lambda_c \rightarrow K^- p \pi^+$ branching fraction of 4.4% (at 90% confidence level) and it has recently been pointed out that a value as low as 2% for the $K^- p \pi^+$ branching ratio also leads to another difficulty.²

The curve in Fig. 4 is a prediction of Λ_c production based on the leading order QCD fusion processes, $g+g \rightarrow c+\bar{c}$ and $q+\bar{q} \rightarrow c+\bar{c}$ [17], and hadronization with the Lund string model[18], calculated using the program, TWISTER, of Ingelman[19]. The uncertainty of the charm quark mass and the choice of gluon structure function parameterization lead to an overall uncertainty of the perturbative QCD calculation of about a factor of 4. The curve, obtained with $m_c = 1.2 \text{ GeV}$ and the EHLQ set-1 structure functions[20] which enhance the cross section, is an approximate upper limit for Λ_c production from conventional first order QCD processes. The result also depends on the fragmentation model used, and in particular on the recombination with spectator beam partons that occurs in the string model[21]. Such a mechanism tends to increase the contribution of Λ_c production at large x_F , leading to a shape in Fig. 4 which is markedly different from the steep x_F distribution obtained for the c-quark itself. The importance of a final state interaction between a charmed quark and spectator quarks has also been pointed out by Bjorken[22] and by Brodsky et al.[23].

Finally, we investigate the decay angular distributions of the Λ_c and Λ_s in Reaction 1. In the Λ_c rest frame we study the direction of the Λ_s , $\hat{\Lambda}_s$, with respect to the normal to the production plane, $n \propto \hat{p}_{\text{beam}} \times \hat{\Lambda}_c$. If the Λ_c is polarized at production and if the direction of the Λ_s offers analyzing power in the decay of the Λ_c , an asymmetry would be observed. We also measure the helicity of the outgoing Λ_s by looking at the direction of the proton in its decay, $\hat{p}$, with respect to the Λ_s direction of motion. Naive V-A predicts that this helicity should be -1 . Table 1 contains the number of signal events in the two hemispheres for both angular distributions. These numbers are obtained by fitting the mass spectra for the data selected in each hemisphere. Also shown are the fraction of Monte Carlo events accepted by the apparatus when the two generated angular distributions are isotropic. This Monte Carlo calculation is used to obtain the acceptance-corrected asymmetries for both distributions

¹ A fit of the same function to the points plotted as invariant cross section $(x_F/\pi)d\sigma/dx_F$ yields $n = 1.7 \pm 0.3$.

² The ARGUS Collaboration observes that the MARK II value for the $K^- p \pi^+$ branching ratio, when combined with the ARGUS charmed particle data, leads to comparable cross sections for charmed hyperon and meson production in e^+e^- collisions. This unlikely result may mean that the $\Lambda_c \rightarrow K^- p \pi^+$ branching ratio is much larger than 2% [K. Schubert, private communication - 1987].

in Table 1. Using the well-known Λ_s decay distribution, $1 + \alpha P \cos \theta$, the values for $\hat{p} \cdot \hat{\Lambda}_s$ directly give a helicity of $-(1.2 \pm 0.5)$ for the Λ_s . The asymmetry in $\hat{\Lambda}_s \cdot \hat{n}$ is 2.2 standard deviations from zero and provides some evidence that the Λ_c is polarized at production and that the $\Lambda_s 3\pi$ decay is a good analyzer. We note that the two observed asymmetries are both characteristic of weak decay processes and together constitute a combined 3 standard deviation effect.

The observation of asymmetries in the decays of Λ_c and Λ_s allows us to display an improved signal/background ratio in Fig. 2(b). In that figure, only events which satisfy the following selections on decay angles are used: $(\hat{\Lambda}_s \cdot \hat{n}) < 0$ and $(\hat{p} \cdot \hat{\Lambda}_s) < 0$. The background decreases to 25% of the level in Fig. 2(a), whereas the signal decreases to 55% of its original value, as expected from the asymmetries shown in Table 1. The signal/background ratio increases from about 1/10 to about 1/5 after the angle selections are made. However, because of the loss in statistics, the significance of the signal (6.3σ) changes little. The Λ_c mass obtained from the fit in Fig. 2(b) is identical to that quoted above for Fig. 2(a).

Since the 4-body final state in Reaction 1 seems to possess significant analyzing power for polarization, it may be that the decay proceeds through a quasi-two-body intermediate state. In fact, the dominance of $a_1(1270)$ in the decay $\tau \rightarrow \nu 3\pi$ should have a parallel in $c \rightarrow s 3\pi$ in Reaction 1. To investigate this question, we show in Fig. 5 the distribution in invariant mass of the 3π system in Reaction 1. Although the statistics are limited, the points do show a peaking near the kinematic limit, which is not far from the central mass of the $a_1(1270)$. The solid curve is the result of a fit to the points of 3-body phase space added to a Gaussian-smeared Breit-Wigner distribution multiplied by phase space. The parameters of the Breit-Wigner distribution are fixed at the values found by the MARK II collaboration in fitting their τ decay data[24]. In the fit, the Breit-Wigner distribution is found to comprise $47 \pm 25\%$ of the total area. The dashed curve under the solid curve shows the area of background phase space obtained in the fit. The dot-dashed curve is the phase space component normalized to the data and has a $\chi^2/D.F.$ (degree of freedom) of 5.0/4, compared with 0.6/3 for the phase space plus a_1 fit. Clearly, a substantial contribution of $a_1(1270)$ can be accommodated by the data.

The presence of Λ_c polarization may mean that a significant component in the production of Λ_c is similar to that of Λ_s , for which a large polarization is observed at $\sqrt{s} = 63$ GeV[25]. Such polarization can be understood[26] as arising from local angular momentum conservation in the ss creation from a stretched color string. The possibility of a contribution to the Λ_c cross section other than that of parton-parton fusion, coupled with the branching ratio uncertainty referred to above, may be responsible, in part, for the discrepancy between the experimental results and the prediction in Fig. 4.

We wish to thank G. Ingelman for valuable discussions concerning the predictions for Λ_c production and, in particular, for help in the use of his Monte Carlo program, TWISTER. We also owe continuing gratitude to the ISR Operations and Experimental Support Groups for their invaluable efforts, and to the CERN administration for their support.

References

- [1] W. Lockman et al., Phys. Let. 85B (1979) 443.
- [2] D. Drijard et al., Phys. Let. 85B (1979) 452.
- [3] K. L. Giboni et al., Phys. Let. 85B (1979) 437.
- [4] M. Basile et al., Lett. Nuovo Cimento 30 (1981) 481, 487.
- [5] P. Chauvat et al., The Performance of a 16,000 Wire Mini Drift MWPC System; CERN-EP/82-198; Proceedings of the 1983 DPF Workshop on Collider Detectors (LBL-15973), p. 81; and contribution to XXI International Conference on High Energy Physics, Paris, July 1982.
- [6] P. Chauvat et al., Phys. Letters 127B (1983) 384.
- [7] P. Chauvat et al., Phys. Letters 163B (1985) 273.
- [8] M. Aguilar-Benitez et al. (Particle Data Group); Physics Letters 170B (1986) 1.
- [9] S. Barlag et al. (ACCMOR Collaboration), Phys. Letters 184 (1987) 283.
- [10] R. Morrison - Fermilab experiment E691 [private communication - 1987].
- [11] M. Aguilar-Benitez et al., Phys. Letters B189 (1987) 254.
- [12] S. Erhan et al., Phys. Letters 85B (1979) 447.
- [13] T. Bowcock et al., Phys. Rev. Letters 55 (1985) 923.
- [14] L. Albrecht et al. (ARGUS Collaboration), contribution to 1987 Uppsala Conference.
- [15] D. Hitlin, Rapporteur Talk at the 1987 International Symposium on Lepton and Photon Interactions at High Energies, Hamburg. This value is slightly different from the value, 2.2%, from the Mark II experiment quoted in Ref. [8] and is due to a reanalysis of the data by the authors.
- [16] M. Aguilar-Benitez et al., CERN/EP 87-126, Submitted to Physics Letters B (July 1987).
- [17] B.L. Combridge, Nucl. Phys. B151 (1979) 429.
- [18] B. Andersson, G. Gustafson, G. Ingelman, T. Sjostrand, Phys. Rep. 97 (1983) 31.
- [19] G. Ingelman, Computer Physics Communications 46 (1987) 217.
- [20] E. Eichten, I. Hinchliffe, K. Lane, C. Quigg, Rev. Mod. Phys. 56 (1984) 579, *ibid.* 58 (1986) 1047.
- [21] B. Andersson, H.-U. Bengtsson, G. Gustafsson, Lund preprint LU TP 83-4.
- [22] J.D. Bjorken, private communication (1987).
- [23] S.J. Brodsky, J.F. Gunion, D.E. Soper, OITS 359, invited talk by D.E. Soper at the Rencontre de Moriond, Les Arcs (March 1987).
- [24] W.B. Schmidke et al., Phys. Rev. Letters 547 (1986) 527.
- [25] A.M. Smith et al., Phys. Letters 185B (1987) 209.
- [26] B. Andersson et al., Phys. Letters 85B (1979) 417.

Table 1: Polarization Measurements

	Events	Isotropic M.C.(a)	Asymmetry(b)
$(\hat{\Lambda}_s \cdot \hat{n}) > 0$	196 ± 76	0.50	$-(0.39 \pm 0.17)$
< 0	437 ± 75	0.50	
$(\hat{p} \cdot \hat{\Lambda}_s) > 0$	176 ± 70	0.47	$-(0.38 \pm 0.17)$
< 0	451 ± 77	0.53	

(a) The fractions of isotropically decaying Monte Carlo Λ_c events that satisfy the selections shown.

(b) Asymmetry ratio is, e.g., (forward - backward)/(forward + backward)

Figure Captions

1. Calibration invariant mass signals: (a) $\Lambda_s \rightarrow p\pi^-$, mean value = 1115.6 ± 0.1 MeV compared with 1115.6 MeV[8]; (b) $f_1(1285) \rightarrow \pi^+\pi^-\pi^+\pi^-$, mean value = 1279.6 ± 1.6 MeV compared with 1283 ± 5 MeV[8]; (c) $\Lambda_s^*(1520) \rightarrow K^-p$, mean value = 1519.1 ± 0.3 MeV compared with 1519.5 ± 1.0 MeV[8]. Note the suppressed zero on the vertical scales in (b) and (c).
2. (a) Invariant mass of $\Lambda_s\pi^+\pi^+\pi^-$ system with $0.5 < x_F < 0.9$ and $p_t < 1.1$ GeV/c (15 MeV bins); the curve is the sum of a third order polynomial and a Gaussian resolution function with $\sigma = 14$ MeV. Note the suppressed zero on the vertical scale; (b) same as (a), but with additional selections on the decay angles of Λ_c and Λ_s , $(\hat{\Lambda}_s \cdot \hat{n}) < 0$ and $(\hat{p} \cdot \hat{\Lambda}_s) < 0$, as discussed in the text.
3. Number of observed (uncorrected) Λ_c signal events vs. x_F when $p_t < 1.1$ GeV/c; The solid and dashed curves are the expected distributions if the cross section $d\sigma/dx_F \propto (1-x_F)^n$ with $n = 2$ and 1, respectively.
4. Differential cross section for Reaction 1 vs. $1-x_F$; Solid circles are branching ratio times cross section, $Bd\sigma/dx_F$. The solid squares show the cross section after correcting for B (there is a 25% systematic uncertainty on all solid square points, in addition to the uncertainty due to the branching ratio used, as discussed in the text). The curve is the QCD and Lund Model prediction discussed in the text. The open squares show $d\sigma/dx_F$ for Λ_s [12]. All points on this figure contain a factor of 2 to account for both hemispheres.
5. Acceptance-corrected invariant mass distribution of $\pi^+\pi^+\pi^-$ in Reaction 1 for the Λ_c signal events in Fig. 2(a). The solid curve is the fit described in the text of $a_1(1270)$ plus phase space background; the dashed curve shows the phase space component of this fit. The dash-dot curve is the phase space component normalized to the data points.

FIG.1

FIG. 2

FIG. 3

FIG. 4

FIG.5