AD

PRE 306L9
c₁

University of Munich Preprint

-7 DEC. 1987

**The mixing angles in matter for three generations of neutrinos
and the MSW mechanism**

H.W. ZAGLAUER[†] and K.H. SCHWARZER

*Sektion Physik, University of Munich
Theresienstrasse 37, Munich (Fed. Rep. of Germany)*

ABSTRACT

Mikheyev and Smirnov have recently shown that oscillations between two species of neutrinos may be amplified in matter. We calculate the energy eigenvalues and mixing angles in matter for three generations of neutrinos using the Fritzsche parametrization for the flavor mixing matrix. For clearly separated neutrino masses $\Delta m_{31}^2 \gg \Delta m_{21}^2$ we find two MSW resonance effects – one for $\nu_e \leftrightarrow \nu_\mu$ and one for $\nu_e \leftrightarrow \nu_\tau$ conversions – which can each be approximated by a separate two neutrino treatment.

PLEASE
MAKE A
PHOTOCOPY
or check out as
NORMAL
LOAN

[†] Current address: Washington University, St. Louis, MO 63130, USA

1. Introduction

The question of the nature and the magnitude of the neutrino mass remains one of the most prominent problems of modern day physics. While there is no known principle that requires the neutrinos to be massless, their mass is certainly much smaller than that of any other fermion and may even be zero. The existence of massive neutrinos would not only lead to an extension of the standard model of electroweak interactions but might also help to solve the problem of the missing or dark matter [1] in the universe.

Due to the absence of righthanded neutrinos (and accordingly lefthanded antineutrinos) a standard Dirac mass term does not occur in the Glashow-Weinberg-Salam theory. A neutrino mass can only be introduced by enlarging the particle content of the theory either by heavy righthanded neutrinos or by a weak isovector Higgs boson. Both mechanisms generate Majorana mass terms in the Lagrangian for the various neutrino generations. Since there is no physical law that demands that the mass matrix is diagonal in the flavour and helicity eigenstates, there may be mixing between these states similar to the flavour mixing in the quark sector. Oscillations between the different neutrino flavours as well as between neutrinos and antineutrinos will result from such mixing. To account for the fact that so far no righthanded neutrino has been seen the righthanded neutrino components should be quite massive ([2],[3]). Thus the search for neutrino flavour oscillations seems to be much more promising than for the highly suppressed lepton number violating $\nu \leftrightarrow \bar{\nu}$ transitions. Since the concept of neutrino oscillations was devised about 20 years ago [4] a number of oscillation experiments have been performed at reactors and accelerators [5] — without conclusive evidence for or against oscillations. These experiments could so far only set limits on the range of the oscillation parameters, namely the mass-squared difference Δm^2 and the sine of the mixing angle $\sin \theta$. Vacuum neutrino oscillations have also been invoked as a possible solution for the solar neutrino puzzle [6]. In order to explain via neutrino oscillations the observed discrepancy for the ^{37}Cl - detector between measured ($2.1 \pm 0.3 \text{ SNU}$ [7]) and theoretically predicted ($5.8 \pm 2.2 \text{ SNU}$ [8]) solar neutrino flux the mixing angles have to be large. Recently Mikheyev and Smirnov [9] proposed a new approach to the solar neutrino puzzle where even small mixing angles would lead to a siz-

able reduction of the solar neutrino flux through resonant neutrino transformations in the interior of the sun. This mechanism derives from the different charged current contributions to the forward scattering amplitudes for the various flavours of neutrinos propagating through matter, as was first observed by Wolfenstein [10].

Several authors ([11],[12]) have thoroughly investigated this so called MSW effect concentrating mainly on oscillations between two neutrino generations and considering special cases for three neutrino families ([13],[14]). For weak eigenstates ν_e and ν_μ which are given as mixtures

$$\begin{pmatrix} |\nu_e\rangle \\ |\nu_\mu\rangle \end{pmatrix} = \begin{pmatrix} \cos \theta & \sin \theta \\ -\sin \theta & \cos \theta \end{pmatrix} \begin{pmatrix} |\nu_1\rangle \\ |\nu_2\rangle \end{pmatrix} \quad (1)$$

of the mass eigenstates ν_1 and ν_2 with masses m_1 and m_2 respectively, the evolution in time of a neutrino state

$$|\nu(t)\rangle = \psi_e(t) |\nu_e\rangle + \psi_\mu(t) |\nu_\mu\rangle \quad (2)$$

in vacuum is given in the ultrarelativistic limit $E \gg m_1, m_2$ through the Schrödinger-like equation

$$i \frac{d}{dt} \begin{pmatrix} \psi_e \\ \psi_\mu \end{pmatrix} = \frac{1}{2E} \begin{pmatrix} m_1^2 \cos^2 \theta + m_2^2 \sin^2 \theta & \Delta m^2 \sin \theta \cos \theta \\ \Delta m^2 \sin \theta \cos \theta & m_1^2 \sin^2 \theta + m_2^2 \cos^2 \theta \end{pmatrix} \begin{pmatrix} \psi_e \\ \psi_\mu \end{pmatrix} \quad (3)$$

From equation (3) we find for the transition probabilities the well known oscillation formulae

$$\begin{aligned} P(\nu_e \rightarrow \nu_e) &= 1 - \sin^2 2\theta \sin^2 \frac{\Omega t}{2} \\ P(\nu_e \rightarrow \nu_\mu) &= \sin^2 2\theta \sin^2 \frac{\Omega t}{2} \end{aligned} \quad (4)$$

with the frequency $\Omega = \Delta m^2 / 2E$. In matter the charged current contribution to the $\nu_e e^-$ forward scattering amplitude modifies the evolution equation

$$i \frac{d}{dt} \begin{pmatrix} \psi_e \\ \psi_\mu \end{pmatrix} = \frac{1}{2E} \begin{pmatrix} m_1^2 + \Delta m^2 \sin^2 \theta + D & \Delta m^2 \sin \theta \cos \theta \\ \Delta m^2 \sin \theta \cos \theta & m_1^2 + \Delta m^2 \cos^2 \theta \end{pmatrix} \begin{pmatrix} \psi_e \\ \psi_\mu \end{pmatrix} \quad (5)$$

with $D = 2\sqrt{2}G_F N_e E$, Fermi constant G_F and electron density N_e (the neutral current contributions for ν_e and ν_μ are equal and give therefore only an unobservable constant

phase). By diagonalizing the matrix in equation (5) we obtain the mass eigenvalues in matter

$$M_{1,2}^2 = \frac{1}{2}[D + m_1^2 + m_2^2 \mp \sqrt{(D - \Delta m^2 \cos 2\theta)^2 + (\Delta m^2 \sin 2\theta)^2}] \quad (6)$$

and the corresponding mass eigenstates with components given by

$$\begin{pmatrix} \psi_{1m} \\ \psi_{2m} \end{pmatrix} = \begin{pmatrix} \cos \theta_m & -\sin \theta_m \\ \sin \theta_m & \cos \theta_m \end{pmatrix} \begin{pmatrix} \psi_e \\ \psi_\mu \end{pmatrix} \quad (7)$$

where θ_m is the mixing angle in matter. In complete analogy to (3) the evolution equation (4) can be written in terms of the matter parameters as

$$i \frac{d}{dt} \begin{pmatrix} \psi_e \\ \psi_\mu \end{pmatrix} = \frac{1}{2E} \begin{pmatrix} M_1^2 + \Delta M^2 \sin^2 \theta_m & \Delta M^2 \sin \theta_m \cos \theta_m \\ \Delta M^2 \sin \theta_m \cos \theta_m & M_1^2 + \Delta m^2 \cos^2 \theta_m \end{pmatrix} \begin{pmatrix} \psi_e \\ \psi_\mu \end{pmatrix}. \quad (8)$$

A simple comparison of the off-diagonal matrix elements yields the matter mixing angle θ_m as a function of D

$$\begin{aligned} \sin^2 2\theta_m &= \left(\frac{\Delta m^2}{\Delta M^2} \right)^2 \sin^2 2\theta \\ &= \frac{(\Delta m^2)^2 \sin^2 2\theta}{(D - \Delta m^2 \cos 2\theta)^2 + (\Delta m^2 \sin 2\theta)^2} \end{aligned} \quad (9)$$

and

$$\sin^2 \theta_m = \frac{\Delta m^2(2 \sin^2 \theta - 1) + D + \sqrt{(D - \Delta m^2 \cos 2\theta)^2 + (\Delta m^2 \sin 2\theta)^2}}{2\sqrt{(D - \Delta m^2 \cos 2\theta)^2 + (\Delta m^2 \sin 2\theta)^2}}. \quad (10)$$

For small vacuum mixing angle θ , $\sin^2 2\theta_m$ shows a typical resonance profile as a function of D , while $\sin^2 \theta_m$ rises from its vacuum value to 1 as D increases through the resonance. As a consequence an electron neutrino ($\approx \nu_2$ since $\theta_m \approx \frac{\pi}{2}$) produced in matter where D is far beyond the resonance density $D_r = \Delta m^2 \cos 2\theta$ will in transversing adiabatically into the vacuum turn into a muon neutrino ($\approx \nu_1$ in vacuum since $\theta_m \approx \theta$ is small). Thus the MSW effect results from the resonance behaviour of the mixing angle.

2. The mass eigenvalues in matter

Unfortunately nature does not restrict itself to two fermion generations but presents us with at least three different fermion families. In general for neutrinos flavour and

vacuum mass eigenstates mix via a unitary $n \times n$ - matrix U that can be parametrized by $\frac{1}{2}n(n-1)$ real mixing angles and $\frac{1}{2}(n-1)(n-2)$ or $\frac{1}{2}n(n-1)$ CP-violating phases for Dirac and Majorana particles respectively. Fortunately the additional Majorana phases do not enter into neutrino oscillations ([3],[15]), so that we can use those parametrizations of the flavour mixing matrix which were devised for the quark sector. The three mixing angles and one CP-violating phase in three dimensions can be chosen in many different ways to parametrize U ; we use the form introduced by Chau and Keung and Fritzsch [16]

$$\begin{pmatrix} |\nu_e\rangle \\ |\nu_\mu\rangle \\ |\nu_\tau\rangle \end{pmatrix} = \begin{pmatrix} c_{12}c_{13} & s_{12}c_{13} & s_{13}e^{-i\delta} \\ -c_{23}s_{12} - c_{12}s_{23}s_{13}e^{i\delta} & c_{12}c_{23} - s_{12}s_{23}s_{13}e^{i\delta} & c_{13}s_{23} \\ s_{12}s_{23} - c_{12}c_{23}s_{13}e^{i\delta} & -c_{12}s_{23} - c_{23}s_{12}s_{13}e^{i\delta} & c_{13}c_{23} \end{pmatrix} \begin{pmatrix} |\nu_1\rangle \\ |\nu_2\rangle \\ |\nu_3\rangle \end{pmatrix} \quad (11)$$

with $c_{kj} = \cos \theta_{kj}$ and $s_{kj} = \sin \theta_{kj}$. The $|\nu_i\rangle$, $i=1,2,3$ are the mass eigenstates with masses m_i ; we shall assume $m_1 < m_2 < m_3$ while $\theta_{ij} \in [0, \frac{\pi}{2}]$ (this accounts for all possible effective masses of the flavour eigenstates, if no mass hierarchy were to be fixed we could restrict θ_{ij} to the interval $[0, \frac{\pi}{4}]$ but must then allow negative quantities $\Delta m_{ij}^2 = m_i^2 - m_j^2$). This particular parametrization was employed because the matter mixing angles s_{12}^m and s_{13}^m and therefore the ν_e - transition probabilities for this choice do not depend on the CP-phase δ as will be shown later.

The propagation of a neutrino in vacuum is given in the mass eigenbasis through

$$i \frac{d}{dt} \psi_j(t) = \frac{1}{2E} m_j^2 \psi_j(t) \quad (12)$$

and in the flavour eigenbasis through

$$i \frac{d}{dt} \psi_\alpha(t) = \frac{1}{2E} \sum_{\beta=e,\mu,\tau} \sum_{j=1}^n m_j^2 U_{\alpha j}^* U_{\beta j} \psi_\beta(t) \quad (13)$$

Just as in the case of two neutrinos the contribution from the interaction with matter gives for the ν_e an additional 'density'-dependent potential D which modifies equation (13) to

$$i \frac{d}{dt} \psi_\alpha(t) = \frac{1}{2E} \sum_{\beta=e,\mu,\tau} \left(\sum_{j=1}^n m_j^2 U_{\alpha j}^* U_{\beta j} + D \delta_{\alpha e} \delta_{\beta e} \right) \psi_\beta(t) \quad (14)$$

In the (vacuum) mass eigenbasis we then find through a unitary transformation

$$i \frac{d}{dt} \psi_j(t) = \frac{1}{2E} \sum_{k=1}^3 (m_j^2 \delta_{jk} + D U_{ej} U_{ek}^*) \psi_k(t) \quad (15)$$

As one can see only matrix elements involving the electron flavour appear in the Hamiltonian. With U as in (11) equation (15) can be written as

$$i \frac{d}{dt} \begin{pmatrix} \psi_1 \\ \psi_2 \\ \psi_3 \end{pmatrix} = \frac{1}{2E} \begin{pmatrix} m_1^2 + Dc_{12}^2 c_{13}^2 & Dc_{12}s_{12}c_{13}^2 & Dc_{12}c_{13}s_{13}e^{i\delta} \\ Dc_{12}s_{12}c_{13}^2 & m_2^2 + Ds_{12}^2 c_{13}^2 & Ds_{12}c_{13}s_{13}e^{i\delta} \\ Dc_{12}c_{13}s_{13}e^{-i\delta} & Ds_{12}c_{13}s_{13}e^{-i\delta} & m_3^2 + Ds_{13}^2 \end{pmatrix} \begin{pmatrix} \psi_1 \\ \psi_2 \\ \psi_3 \end{pmatrix}. \quad (16)$$

The characteristic polynomial of matrix (16)

$$\begin{aligned} X^3 - (m_1^2 + m_2^2 + m_3^2 + D)X^2 + [m_1^2 m_2^2 + m_2^2 m_3^2 + m_1^2 m_3^2 \\ + D(m_3^2 c_{13}^2 + m_2^2(c_{13}^2 c_{12}^2 + s_{13}^2) + m_1^2(c_{13}^2 s_{12}^2 + s_{13}^2)]X \\ - [m_1^2 m_2^2 m_3^2 + D(m_3^2 m_2^2 c_{13}^2 c_{12}^2 + m_3^2 m_1^2 c_{13}^2 s_{12}^2 + m_2^2 m_1^2 s_{13}^2)] = 0 \end{aligned} \quad (17)$$

leads to the mass eigenvalues in matter [17]

$$M_1^2 = m_1^2 + \frac{A}{3} - \frac{1}{3}\sqrt{A^2 - 3BS} - \frac{\sqrt{3}}{3}\sqrt{A^2 - 3B}\sqrt{1 - S^2} \quad (18)$$

$$M_2^2 = m_1^2 + \frac{A}{3} - \frac{1}{3}\sqrt{A^2 - 3BS} + \frac{\sqrt{3}}{3}\sqrt{A^2 - 3B}\sqrt{1 - S^2} \quad (19)$$

$$M_3^2 = m_1^2 + \frac{A}{3} + \frac{2}{3}\sqrt{A^2 - 3BS} \quad (20)$$

where we have introduced for convenience the abbreviations

$$A = \Delta m_{21}^2 + \Delta m_{31}^2 + D \quad (21)$$

$$B = \Delta m_{31}^2 \Delta m_{21}^2 + D[\Delta m_{31}^2 c_{13}^2 + \Delta m_{21}^2(c_{13}^2 c_{12}^2 + s_{13}^2)] \quad (22)$$

$$C = D\Delta m_{31}^2 \Delta m_{21}^2 c_{13}^2 c_{12}^2 \quad (23)$$

$$S = \cos \frac{1}{3} \arccos \frac{2A^3 - 9AB + 27C}{2\sqrt{A^2 - 3B}^3}. \quad (24)$$

The differences of the eigenvalues that determine the oscillation wavelengths are given by

$$\Delta M_{21}^2 = \frac{2\sqrt{3}}{3}\sqrt{A^2 - 3B}\sqrt{1 - S^2} \quad (25)$$

$$\Delta M_{31}^2 = \sqrt{A^2 - 3BS} + \frac{\sqrt{3}}{3}\sqrt{A^2 - 3B}\sqrt{1 - S^2} \quad (26)$$

$$\Delta M_{32}^2 = \sqrt{A^2 - 3BS} - \frac{\sqrt{3}}{3}\sqrt{A^2 - 3B}\sqrt{1 - S^2}; \quad (27)$$

they do only depend on the differences of the squared masses Δm_{21}^2 and Δm_{31}^2 and on the vacuum mixing angles θ_{12} and θ_{13} . In figures 1 and 2 we show in an exemplary way

the dependence of the mass eigenvalues and of their differences on the different vacuum parameters for firstly $\Delta m_{31}^2 = 10 \Delta m_{21}^2$ and secondly $\Delta m_{31}^2 = 1.1 \Delta m_{21}^2$. For the first choice of clearly separated vacuum masses we find two distinct 'densities' D where two mass eigenvalues approach each other closely, i.e. where one of the ΔM_{jk}^2 reaches a minimum (fig. 1 (a), (d)). These minima correspond to two separate resonance densities as will be shown later. The ΔM_{21}^2 - minimum is governed by the mixing angle θ_{12} and the ΔM_{32}^2 - minimum is governed by θ_{13} - an increase in a mixing angle leads to an increase in the mass splittings (fig. 1 (b),(c)). For the second choice of nearly degenerate neutrino masses $m_3 \approx m_2$ only ΔM_{31}^2 runs through a minimum corresponding to a single resonance density (fig. 2 (a), (d)) while M_2^2 is almost independent of D . A change in either mixing angle does affect M_1^2 and M_3^2 in the same way (fig. 2 (b), (c)). For large densities D the lower mass eigenvalues M_1^2 and M_2^2 approach constant asymptotic values while M_3^2 increases linearly with D .

3. The mixing angles in matter

We are now going to derive the matter mixing angles which requires us to determine the eigenstates of the Hamiltonian belonging to the different eigenvalues we obtained in the last section. Starting with the Hamiltonian in the vacuum mass eigenbasis (16) we can find a unitary matrix W such that

$$W \begin{pmatrix} m_1^2 + D c_{12}^2 c_{13}^2 & D c_{12} s_{12} c_{13}^2 & D c_{12} c_{13} s_{13} e^{i\delta} \\ D c_{12} s_{12} c_{13}^2 & m_2^2 + D s_{12}^2 c_{13}^2 & D s_{12} c_{13} s_{13} e^{i\delta} \\ D c_{12} c_{13} s_{13} e^{-i\delta} & D s_{12} c_{13} s_{13} e^{-i\delta} & m_3^2 + D s_{13}^2 \end{pmatrix} W^{-1} =$$

$$= \begin{pmatrix} M_1^2 & 0 & 0 \\ 0 & M_2^2 & 0 \\ 0 & 0 & M_3^2 \end{pmatrix} . \quad (28)$$

This matrix W transforms the vacuum mass eigenstates into matter mass eigenstates

$$\begin{pmatrix} \psi_{1m} \\ \psi_{2m} \\ \psi_{3m} \end{pmatrix} = W \begin{pmatrix} \psi_1 \\ \psi_2 \\ \psi_3 \end{pmatrix} \quad (29)$$

with the rows of W being the eigenvectors of the Hamiltonian in the vacuum mass eigenbasis. The flavour mixing matrix V in matter is thus obtained as

$$V = UW^T \quad (30)$$

where

$$\begin{pmatrix} |\nu_e\rangle \\ |\nu_\mu\rangle \\ |\nu_\tau\rangle \end{pmatrix} = V \begin{pmatrix} |\nu_{1m}\rangle \\ |\nu_{2m}\rangle \\ |\nu_{3m}\rangle \end{pmatrix} . \quad (31)$$

This procedure does not automatically guarantee that the matrix V will admit the parametrization (11). It could well be that some or all of the matrix elements $V_{11}, V_{12}, V_{23}, V_{33}$ were complex instead of real. In order to avoid this, a phase transformation

$$e^{i(aT_3+bY)} V e^{-i(cT_3+dY)} \quad (32)$$

with

$$T_3 = \begin{pmatrix} 1 & 0 & 0 \\ 0 & 0 & 0 \\ 0 & 0 & -1 \end{pmatrix} , \quad Y = \begin{pmatrix} 1 & 0 & 0 \\ 0 & -2 & 0 \\ 0 & 0 & 1 \end{pmatrix}$$

and real numbers a, b, c and d has to be applied to the matrix V such that we obtain the Fritzsche form (11). Fortunately we do not need all the matrix elements in order to calculate the behaviour of the mixing angles in matter. For the absolute values of the first row and the third column of the matter mixing matrix we find

$$s_{13}^2 = \frac{M_3^4 - \alpha M_3^2 + \beta}{\Delta M_{31}^2 \Delta M_{32}^2} \quad (34)$$

$$s_{12}^2 c_{13}^2 = \frac{\alpha M_2^2 - M_2^4 - \beta}{\Delta M_{21}^2 \Delta M_{32}^2} \quad (35)$$

$$c_{12}^2 c_{13}^2 = \frac{M_1^4 - \alpha M_1^2 + \beta}{\Delta M_{21}^2 \Delta M_{31}^2} \quad (36)$$

$$s_{23}^2 c_{13}^2 = \frac{E^2 s_{23}^2 + F^2 c_{23}^2 + 2EF c_{23} s_{23} c_\delta}{\Delta M_{31}^2 \Delta M_{32}^2 (M_3^4 - \alpha M_3^2 + \beta)} \quad (37)$$

$$c_{23}^2 c_{13}^2 = \frac{E^2 c_{23}^2 + F^2 s_{23}^2 - 2EF c_{23} s_{23} c_\delta}{\Delta M_{31}^2 \Delta M_{32}^2 (M_3^4 - \alpha M_3^2 + \beta)} \quad (38)$$

with the convenient abbreviations

$$\alpha = m_3^2 c_{13}^2 + m_2^2 (c_{13}^2 c_{12}^2 + s_{13}^2) + m_1^2 (c_{13}^2 s_{12}^2 + s_{13}^2) \quad (39)$$

$$\beta = m_3^2 c_{13}^2 (m_2^2 c_{12}^2 + m_1^2 s_{12}^2) + m_2^2 m_1^2 s_{13}^2 \quad (40)$$

$$E = [\Delta m_{31}^2 (M_3^2 - m_1^2 - \Delta m_{21}^2) - \Delta m_{21}^2 (M_3^2 - m_1^2 - \Delta m_{31}^2) s_{12}^2] c_{13} s_{13} \quad (41)$$

$$F = (M_3^2 - m_1^2 - \Delta m_{31}^2) \Delta m_{21}^2 c_{12} s_{12} c_{13} \quad (42)$$

The mixing angles in matter are then given by

$$s_{12}^2 = \frac{-(M_2^4 - \alpha M_2^2 + \beta) \Delta M_{31}^2}{\Delta M_{32}^2 (M_1^4 - \alpha M_1^2 + \beta) - \Delta M_{31}^2 (M_2^4 - \alpha M_2^2 + \beta)} \quad (43)$$

$$s_{13}^2 = \frac{M_3^4 - \alpha M_3^2 + \beta}{\Delta M_{31}^2 \Delta M_{32}^2} \quad (44)$$

$$s_{23}^2 = \frac{E^2 s_{23}^2 + F^2 c_{23}^2 + 2EF c_{23} s_{23} c_\delta}{E^2 + F^2} \quad (45)$$

and the CP-phase is determined as

$$e^{-i\delta^m} = \frac{(E^2 e^{-i\delta} - F^2 e^{i\delta}) s_{23} c_{23} + EF (c_{23}^2 - s_{23}^2)}{\sqrt{(E^2 s_{23}^2 + F^2 c_{23}^2 + 2EF c_{23} s_{23} c_\delta)(E^2 c_{23}^2 + F^2 s_{23}^2 - 2EF c_{23} s_{23} c_\delta)}} \quad (46)$$

The mixing angles and the CP-phase in matter depend in a quite complicated way on the vacuum mass-squared differences Δm_{21}^2 and Δm_{31}^2 , on the mixing angles s_{12}^2 , s_{13}^2 , s_{23}^2 and on the vacuum CP-phase δ . In figures 3 - 7 the angles are plotted as functions of D and their dependences on the various vacuum parameters are given in an exemplary fashion illustrating the characteristic features of the MSW effect for three species of neutrinos.

Considering the angles as functions of D they typically are nearly constant up to a certain resonance density D , where they sharply increase to an asymptotic value (fig. 3). The smaller the vacuum mixing angles and the larger Δm_{21}^2 are, the more pronounced the resonance is. While in the case of two families only the sine of the mixing angle tends towards 1 (for matter densities far beyond the resonance density) for three generations not all the angles necessarily have to be close to $\frac{\pi}{2}$ in highly dense matter. For certain values of the vacuum parameters, s_{12}^m does not even show the simple resonance behaviour (fig. 4). For small mass splittings $\Delta m_{31}^2 \approx \Delta m_{21}^2$, s_{12}^2 is no longer a strictly monotone function of

D; the asymptotic value for $D \rightarrow \infty$ is lower than its maximal value (fig. 4 (b),(c)). The larger the mass difference between the two heaviest neutrino mass states is the closer to 1 the asymptotic value for s_{12}^2 will be (fig. 4 (a)). This can also be seen by first expanding (43) in powers of $\frac{1}{D}$ and taking $D \rightarrow \infty$; for $\Delta m_{31}^2 \gg \Delta m_{21}^2$ we then obtain

$$s_{12}^2 \xrightarrow{D \rightarrow \infty} 1. \quad (47)$$

For θ_{13} we find that the asymptotic value is independent of the neutrino mass differences (fig. 5). Changes in the vacuum parameters only determine the density and the shape of the resonance. Just as for two neutrinos the resonance density is in this case directly proportional to the mass difference Δm_{31}^2 (fig. 5 (a)), while a rising s_{12}^2 tends to decrease the resonance density (fig. 5 (b)). An increase of s_{13}^2 tends to flatten the resonance peak, thereby making the mixing angle almost independent of D (fig. 5 (c)).

Whilst the matter angles θ_{12}^m and θ_{13}^m depend only on the four vacuum parameters Δm_{21}^2 , Δm_{31}^2 , s_{12}^2 and s_{13}^2 , both θ_{23}^m and δ^m do also depend on s_{23}^2 and on the vacuum phase δ . The matter mixing angle s_{23}^{m2} is displayed as a function of D in fig. 6. Only for nearly degenerate heavy neutrinos $m_3 \approx m_2$ do we find the typical resonance shape but with arbitrary asymptotic values. For $\Delta m_{31}^2 \gg \Delta m_{21}^2$ an expansion of (45) shows that

$$s_{23}^{m2} \xrightarrow{D \rightarrow \infty} s_{23}^2, \quad (48)$$

i.e. s_{23}^{m2} remains almost unchanged for varying density D (fig. 6(a)). A change in the vacuum angle θ_{12} and θ_{13} does affect the asymptotic value of s_{23}^{m2} in different ways: While enlarging s_{13}^2 means reducing the asymptotic value, the opposite is true for s_{12}^2 and s_{23}^2 (fig. 6 (b), (c), (d)). The CP-phase does depend markedly on the density only for small mass differences Δm_{32}^2 (fig 7 (a)), while for $\Delta m_{31}^2 \gg \Delta m_{21}^2$, δ^m remains very close to its vacuum value

$$e^{-\delta^m} \xrightarrow{D \rightarrow \infty} e^{-i\delta} \quad (49)$$

as is apparent from fig. 7 (b). Thus we find two distinct cases. If $\Delta m_{31}^2 \approx \Delta m_{21}^2$, only s_{13}^{m2} runs through a typical resonance with asymptotic value 1, and accordingly there is only one resonance density with D proportional to $\Delta m_{31}^2 (\approx \Delta m_{21}^2)$. For $\Delta m_{31}^2 \gg \Delta m_{21}^2$,

s_{12}^{m2} as well as s_{13}^{m2} go through a resonance so that there will be two different resonance regions – one governed by Δm_{21}^2 and the other by Δm_{31}^2 . Here s_{23}^{m2} and the CP-phase δ^m deviate only marginally from their vacuum values.

4. Oscillations with three neutrino generations

Now that we know how the mixing angles depend upon the electron density we can take a closer look at neutrino oscillations in matter. Here we will restrict ourselves to the physically most interesting case, i. e. the case where we expect a sizable resonance. We shall therefore assume small vacuum mixing angles θ_{12} and θ_{13} and clearly separated neutrino masses with $\Delta m_{31}^2 \gg \Delta m_{21}^2$. Large vacuum mixing angles were to lead to large neutrino oscillations, but they would be almost independent of the electron density implying that there were almost no difference between oscillations in vacuum and in matter. If the masses are nearly degenerate, only the angle θ_{13} shows the typical resonance behaviour – thus this case appears to be quite similar to the case of two neutrinos which has been treated at length elsewhere ([11],[12]). A further argument for small mixing angles and large mass splittings is that they also appear in the quark sector – and the idea of grand unification suggests an intimate connection between the lepton and quark masses.

Neutrino flavour oscillations occur when the flavour eigenstates are linear superpositions of mass eigenstates which evolve differently in time. The probability amplitude for finding a neutrino flavour ν_β at time t if a neutrino ν_α is produced at time 0 is given by

$$A(\nu_\alpha \rightarrow \nu_\beta) = \sum_{j=1}^3 V_{\alpha j} e^{-i \frac{m_j^2}{2E} t} V_{\beta j}^* \quad (50)$$

with the mixing matrix V in matter of constant density. The probability is then

$$\begin{aligned} P(\nu_\alpha \rightarrow \nu_\beta) &= |A(\nu_\alpha \rightarrow \nu_\beta)|^2 \\ &= \left| \sum_{j=1}^3 V_{\alpha j} e^{-i \frac{m_j^2}{2E} t} V_{\beta j}^* \right|^2 \\ &= \sum_{k,j=1}^3 |V_{\alpha j} V_{\alpha k}^* V_{\beta j}^* V_{\beta k}| - 4 \sum_{j < k} |V_{\alpha j} V_{\alpha k}^* V_{\beta j}^* V_{\beta k}| \sin^2 \left(\frac{\Omega_{kj} t + \varphi_{kj}^{\alpha\beta}}{2} \right) \end{aligned} \quad (51)$$

where $\Omega_{kj} = \frac{m_k^2 - m_j^2}{2E}$ is the oscillation frequency and the phase $\varphi_{kj}^{\alpha\beta}$ is defined by

$$V_{\alpha j} V_{\alpha k}^* V_{\beta j}^* V_{\beta k} = |V_{\alpha j} V_{\alpha k}^* V_{\beta j}^* V_{\beta k}| e^{i\varphi_{kj}^{\alpha\beta}}. \quad (52)$$

These phases reflect the fact that the elements of the mixing matrix V are, in general, complex, and only vanish if V contains no CP-violating phase. The phases also disappear for the diagonal transitions $\nu_\alpha \rightarrow \nu_\alpha$ and the probabilities for detecting at time t the same neutrino flavour as the one created at $t=0$ becomes

$$P(\nu_\alpha \rightarrow \nu_\alpha) = 1 - 4 \sum_{j < k} |U_{\alpha j}|^2 |U_{\alpha k}|^2 \sin^2 \frac{\Omega_{kj} t}{2}. \quad (53)$$

For three neutrino generations using the Fritzsch parametrization for the mixing matrix we find

$$\begin{aligned} P(\nu_e \rightarrow \nu_e) = & 1 - 4c_{12}^2 s_{12}^2 c_{13}^4 \sin^2 \frac{\Omega_{12} t}{2} \\ & - 4c_{12}^2 c_{13}^2 s_{13}^2 \sin^2 \frac{\Omega_{13} t}{2} - 4s_{12}^2 c_{13}^2 s_{13}^2 \sin^2 \frac{\Omega_{23} t}{2}. \end{aligned} \quad (54)$$

As a result of the D dependence of s_{12}^m and s_{13}^m we have five density regions with differing oscillation behaviour. If $0 \leq D \ll \Delta m_{21}^2$, both sines of the mixing angles, s_{12}^m and s_{13}^m , are close to their vacuum values, i.e. almost 0, and sizable oscillations can only occur between ν_μ and ν_τ , with amplitude $\sin^2 2\theta_{23}$ and frequency $\frac{\Omega_{23}}{2}$. For $D \approx \Delta m_{21}^2$, s_{13}^m is still approximately 0, while s_{12}^m runs through its resonance. Here we find $\nu_e \leftrightarrow \nu_\mu$ - oscillations with

$$P(\nu_e \rightarrow \nu_e) = 1 - \sin^2 2\theta_{12}^m \sin^2 \frac{\Omega_{12} t}{2}; \quad (55)$$

$\nu_\mu \leftrightarrow \nu_\tau$ - oscillations are also possible with the frequency changing smoothly from Ω_{23} to Ω_{13} over the resonance. In the intermediate zone $\Delta m_{21}^2 \ll D \ll \Delta m_{31}^2$ with $s_{12}^m \approx 1$ and $s_{13}^m \approx 0$, there could only be $\nu_\mu \leftrightarrow \nu_\tau$ - transitions with frequency Ω_{13} . In the s_{13}^m - resonance region with $D \approx \Delta m_{31}^2$, s_{12}^m is approximately 1 while s_{13}^m runs through its resonance. In this case $\nu_e \leftrightarrow \nu_\mu$ - transitions appear with

$$P(\nu_e \rightarrow \nu_e) = 1 - \sin^2 2\theta_{13}^m \sin^2 \frac{\Omega_{23} t}{2}. \quad (56)$$

For the $\nu_\mu \leftrightarrow \nu_\tau$ - oscillations the frequency smoothly interpolates from Ω_{13} for $s_{13}^{m^2} \approx 0$ to Ω_{12} for $s_{13}^{m^2} \approx 1$. Finally for very high densities $D \gg \Delta m_{31}^2$ oscillations of electron neutrinos are strongly suppressed because $s_{12}^{m^2} \approx 1$ and $s_{13}^{m^2} \approx 1$, while large $\nu_\mu \leftrightarrow \nu_\tau$ - oscillations are possible with frequency Ω_{12} .

Thus in matter with density $D \approx \Delta m_{21}^2$ or $D \approx \Delta m_{31}^2$ ν_e - oscillations are amplified. With respect to the solar neutrino puzzle we also need to know how the neutrinos behave in matter of varying density. For slowly varying D the adiabatic approximation was first used to establish the MSW effect([9],[11]). In complete analogy to the case of two neutrino generations, we find the mean adiabatic conversion probability for a neutrino originating at $\mathbf{r}_1$ and detected at $\mathbf{r}_2$

$$\langle P(\nu_\alpha \rightarrow \nu_\beta) \rangle = \sum_{j=1}^3 |V_{\beta j}^*(\mathbf{r}_2) V_{\alpha j}(\mathbf{r}_1)|^2 \quad (57)$$

This probability is averaged over the neutrino sample in order to eliminate the interference terms. Especially for the probability of detecting the original electron neutrino we have

$$\langle P(\nu_e \rightarrow \nu_e) \rangle = (c_{12}^2 s_{12}^{m^2} + s_{12}^2 s_{12}^{m^2}) c_{13}^2 c_{13}^{m^2} + s_{13}^2 s_{13}^{m^2} \quad (58)$$

Equation (58) resembles the case of two neutrinos [11] as can also be seen in figure 8. For densities above the s_{13} - resonance $P(\nu_e \rightarrow \nu_e)$ differs from this special case but only slightly so if θ_{13} is very close to θ_{12} while for low densities vacuum oscillations are enhanced by the additional neutrino flavour. With the help of equation (57) we can calculate the adiabatic transition probabilities between all the different neutrino flavours. Using the approximations for the mixing angles $s_{12}^{m^2}$ and $s_{13}^{m^2}$ in the different density regions as determined above, we find

$$P = (P_{\alpha\beta})_{\substack{\alpha=e,\mu,\tau \\ \beta=e,\mu,\tau}} = \begin{pmatrix} 0 & c_{23}^2 & s_{23}^2 \\ c_{23}^2 & s_{23}^4 & s_{23}^2 c_{23}^2 \\ s_{23}^2 & s_{23}^2 c_{23}^2 & c_{23}^4 \end{pmatrix} \quad (59)$$

with $P_{\alpha\beta} = \langle P(\nu_\beta \rightarrow \nu_\alpha) \rangle$ for adiabatic transitions between regions with $0 \leq D \ll \Delta m_{21}^2$ and with $\Delta m_{21}^2 \ll D \ll \Delta m_{31}^2$, and

$$P = \begin{pmatrix} 0 & s_{23}^2 & c_{23}^2 \\ s_{23}^2 & c_{23}^4 & s_{23}^2 c_{23}^2 \\ c_{23}^2 & s_{23}^2 c_{23}^2 & s_{23}^4 \end{pmatrix} \quad (60)$$

between regions with $\Delta m_{21}^2 \ll D \ll \Delta m_{31}^2$ and with $\Delta m_{31}^2 \ll D$. Therefore if we in addition assume that s_{12}^2 is small we can confirm the results of Petcov and Toshev [14]: For neutrinos propagating adiabatically through the s_{12}^2 - resonance there are enhanced $\nu_e \leftrightarrow \nu_\mu$ - conversions while through the s_{13}^2 - resonance $\nu_e \leftrightarrow \nu_\tau$ - conversions are amplified. With the two resonance densities clearly separated, the resonant conversions for both regions may each be described by the MSW treatment for two neutrinos. The results obtained above can be used to calculate the expected capture rate for the solar neutrino experiment (see fig. 9). Due to the possibility of two MSW resonance regions in the sun the Iso-SNU-plots may differ markedly from those of the two-dimensional model, as has already been discussed elsewhere [18].

5. Conclusions

The MSW effect between two neutrino flavours has been proposed as an elegant new solution to the solar neutrino puzzle. For small vacuum mixing angles a resonant amplification of neutrino oscillations in the sun accounts for the observed reduction of the solar neutrino flux. Since there exist at least three neutrino families in nature a full treatment of the case of three neutrinos becomes mandatory.

After a brief review of the case of two neutrino generations we have studied in detail the oscillation behaviour in matter for three neutrino generations. With the Fritzsch parametrization for the flavour mixing matrix the mass eigenvalues only depend on the two mixing angles θ_{12} and θ_{13} but not on θ_{23} or on the CP-violating phase δ that may appear in the mixing matrix.

We have derived the exact formulae for the mixing angles in matter and have also obtained an expression for the CP-violating phase in matter. If we have clearly separated neutrino masses, i.e. $\Delta m_{31}^2 \gg \Delta m_{21}^2$, the sines of the mixing angles s_{12} and s_{13} exhibit the typical MSW behaviour, independent of the vacuum parameters s_{23} and δ . We found two distinct resonance densities – one for $\nu_e \leftrightarrow \nu_\mu$ - transitions and one for $\nu_e \leftrightarrow \nu_\tau$ - transitions – which both can be approximated by a two neutrino treatment. The mixing angle s_{23} and the phase δ remain almost constant in matter, but since s_{23} does not influence the MSW behaviour of the electron neutrino there might be large $\nu_\mu \leftrightarrow \nu_\tau$ - oscillations

that could not be seen in ν_e experiments. For nearly degenerate neutrino mass differences $\Delta m_{31}^2 \approx \Delta m_{21}^2$ only the mixing angle θ_{13} runs through a resonance. Here the influence of the third generation becomes even more important with the mixing angle θ_{23} now changing considerably in matter. Also the CP-violating phase – a quantity that does not appear in the case of two neutrino flavours – now depends on the matter electron density.

Finally, for small vacuum mixing and large neutrino mass splittings we applied our formulae to oscillations of neutrinos in matter of constant densities and to adiabatic conversions in matter with slowly varying density. The adiabatic approximation was used to calculate the effect of the MSW mechanism with three neutrino flavours on the counting rate in the solar neutrino experiment.

5. Conclusions

The MSW effect between two neutrino flavours has been proposed as an elegant new solution to the solar neutrino puzzle. For small vacuum mixing angles a resonant amplification of neutrino oscillations in the sun accounts for the observed reduction of the solar neutrino flux. Since there exist at least three neutrino families in nature a full treatment of the case of three neutrinos becomes mandatory.

After a brief review of the case of two neutrino generations we have studied in detail the oscillation behaviour in matter for three neutrino generations. With the Fritsch parametrisation for the flavour mixing matrix the mass eigenvalues only depend on the two mixing angles θ_{12} and θ_{13} but not on θ_{23} or on the CP-violating phase δ that may appear in the mixing matrix.

We have derived the exact formulae for the mixing angles in matter and have also obtained an expression for the CP-violating phase in matter. If we have clearly separated neutrino masses, i.e. $\Delta m_{31}^2 \gg \Delta m_{21}^2$, the sizes of the mixing angles θ_{12} and θ_{13} exhibit the typical MSW behaviour, independent of the vacuum parameters θ_{23} and δ . We found two distinct resonance densities – one for $\nu_e \rightarrow \nu_\mu$ - transitions and one for $\nu_e \rightarrow \nu_\tau$ - transitions – which both can be approximated by a two neutrino treatment. The mixing angle θ_{12} and the phase δ remain almost constant in matter, but since θ_{13} does not influence the MSW behaviour of the electron neutrino there might be large $\nu_e \rightarrow \nu_\tau$ - oscillations

REFERENCES

- [1] S.M.Faber and J.S. Gallagher, *Annu.Rev.Astron.Astrophys.* **17** (1979) 135
- [2] M. Gell-Mann, P. Ramond and R. Slansky, in: *Supergravity*,
ed P. van Nieuwenhuizen and D.Freedmann, North Holland (1979)
- [3] P. Langacker, S.T. Petcov and G. Steigman, S. Toshev, *Nucl.Phys.* **B282** (1987) 589
- [4] B. Pontecorvo, *Sov.Phys.JETP* **26** (1968) 984;
V. Gribov and B. Pontecorvo, *Phys.Lett.* **28B** (1969) 493 ;
S.M. Bilenky and B. Pontecorvo, *Phys.Rep.* **41C** (1978) 225 ;
for a recent review see: J. Vergados, *Phys.Rep.* **133** (1986) 1
- [5] H. Blümer and K. Kleinknecht, *Phys.Lett.* **161B** (1985) 407
- [6] H. Fritzsch and P. Minkowski, *Phys.Lett.* **62B** (1976) 72;
V. Barger, K. Whisnant and R.J.N. Phillips, *Phys.Rev.* **D24** (1981) 538
- [7] R. Davis Jr., in: *Proc. of the Neutrino Mass Miniconference*, Telemark,
Wisconsin, ed. V.Barger and D. Cline (1980), 38
- [8] J.N. Bahcall et al., *Rev.Mod.Phys.* **54** (1982) 767;
J.N. Bahcall et al., *Astrophys.J.Lett.* **292** (1985), 179
- [9] S.P. Mikheyev and A.Yu. Smirnov, *Sov.J.Nucl.Phys.* **42** (1985) 913;
S.P. Mikheyev and A.Yu. Smirnov, *Nuovo Cimento* **9C** (1986) 17
- [10] L. Wolfenstein, *Phys.Rev.* **D17** (1978) 2369
- [11] V. Barger, R.J.N. Phillips and K. Whisnant, *Phys.Rev.* **D34** (1986) 980
- [12] H.A. Bethe, *Phys.Rev.Lett.* **56** (1986) 1305;
S.J. Parke, *Phys.Rev.Lett.* **57** (1986) 1275;
A. Halprin, *Phys.Rev.* **D34** (1986) 3462;
S.P. Rosen and J.M. Gelb, *Phys.Rev.* **D34** (1986) 969;
S.J. Parke and T.P. Walker, *Phys.Rev.Lett.* **57** (1986) 2322;
E.W. Kolb, S. Turner and T.P.Walker, *Phys.Lett.* **175B** (1986) 478;
W.C. Haxton, *Phys.Rev.Lett.* **57** (1986) 1271
- [13] A. Baldini and G.F. Guidice, *Phys.Lett.* **186B** (1987) 211;
C.W. Kim and W.K. Sze, *Phys.Rev.* **D35** (1987) 1404;

- C.W. Kim, S. Nussinov and W.K. Sze, *Phys.Lett.* **184B** (1987) 403;
T.K. Kuo and J. Pantaleone, *Phys.Rev.Lett.* **57** (1986) 1805
- [14] S. Toshev, *Phys.Lett.* **185B** (1987) 177;
S.T. Petcov and S. Toshev, *Phys.Lett.* **187B** (1987) 120
- [15] S.M. Bilenky, J. Hasek and S.T. Petcov, *Phys.Lett.* **94B** (1980) 495;
M.Doi et al., *Phys.Lett.* **102B** (1981) 323
- [16] L. Chau and K. Keung, *Phys.Rev.Lett.* **53** (1984) 1802;
H. Fritzsch, *Phys.Rev.* **D32** (1985) 3058
- [17] V. Barger, K. Whisnant, S. Pakvasa and R.J.N. Phillips, *Phys.Rev.* **D22** (1980) 271
- [18] H.W. Zaglauer and K.H. Schwarzer, *University of Munich preprint* (1987)

FIGURE CAPTIONS

Fig. 1 Mass eigenvalues in matter for three neutrino generations with clearly separated neutrino masses ($\Delta m_{31}^2/\Delta m_{21}^2 = 10.0$) and various vacuum mixing angles:

(a) $s_{12}^2 = 0.05, s_{13}^2 = 0.05$

(b) $s_{13}^2 = 0.05; s_{12}^2 = 0.01$ (solid line) and $s_{12}^2 = 0.2$ (dashed line)

(c) $s_{12}^2 = 0.05; s_{13}^2 = 0.01$ (solid line) and $s_{13}^2 = 0.2$ (dashed line)

In (d) the mass differences for the masses in matter are shown for $s_{12}^2 = 0.05$ and $s_{13}^2 = 0.05$

Fig. 2 Mass eigenvalues in matter for three neutrino generations with almost degenerate neutrino masses ($\Delta m_{31}^2/\Delta m_{21}^2 = 1.1$) and various vacuum mixing angles:

(a) $s_{12}^2 = 0.05, s_{13}^2 = 0.05$

(b) $s_{13}^2 = 0.05; s_{12}^2 = 0.01$ (solid line) and $s_{12}^2 = 0.2$ (dashed line)

(c) $s_{12}^2 = 0.05; s_{13}^2 = 0.01$ (solid line) and $s_{13}^2 = 0.2$ (dashed line)

In (d) the mass differences for the masses in matter are shown for $s_{12}^2 = 0.05$ and $s_{13}^2 = 0.05$

Fig. 3 The D dependence of the matter mixing angles $s_{12}^{m^2}$ (solid line), $s_{13}^{m^2}$ (dashed line) and $s_{23}^{m^2}$ (dotted line) for the following choices of vacuum parameters. In cases (a) and (b) for nearly degenerate neutrino masses

(a) $\Delta m_{31}^2/\Delta m_{21}^2 = 1.1, s_{12}^2 = s_{13}^2 = s_{23}^2 = 0.01$

(b) $\Delta m_{31}^2/\Delta m_{21}^2 = 1.1, s_{12}^2 = s_{13}^2 = s_{23}^2 = 0.1$

and in cases (c) and (d) likewise for clearly separated neutrino masses

(c) $\Delta m_{31}^2/\Delta m_{21}^2 = 10, s_{12}^2 = s_{13}^2 = s_{23}^2 = 0.01$

(d) $\Delta m_{31}^2/\Delta m_{21}^2 = 10, s_{12}^2 = s_{13}^2 = s_{23}^2 = 0.1$

Fig. 4 The D dependence of $\sin^2 \theta_{12}^m$ for illustrative vacuum parameters

- (a) $s_{12}^2 = 0.1$, $s_{13}^2 = 0.1$ and varying ratio of mass-squared differences $\Delta m_{31}^2/\Delta m_{21}^2$ (solid line: 1.1, dashed line: 1.5, dotted line: 10)
- (b) $\Delta m_{31}^2/\Delta m_{21}^2 = 2.0$, $s_{13}^2 = 0.1$ and for varying s_{12}^2 (solid line: 0.01, dashed line: 0.2, dotted line: 0.4)
- (c) $\Delta m_{31}^2/\Delta m_{21}^2 = 2.0$, $s_{12}^2 = 0.1$ and for varying s_{13}^2 (solid line: 0.01, dashed line: 0.2, dotted line: 0.4)

Fig. 5 The D dependence of $\sin^2 \theta_{13}^m$ for illustrative vacuum parameters

- (a) $s_{12}^2 = 0.1$, $s_{13}^2 = 0.1$ and varying ratio of mass-squared differences $\Delta m_{31}^2/\Delta m_{21}^2$ (solid line: 1.1, dashed line: 1.5, dotted line: 10)
- (b) $\Delta m_{31}^2/\Delta m_{21}^2 = 2.0$, $s_{13}^2 = 0.1$ and for varying s_{12}^2 (solid line: 0.01, dashed line: 0.2, dotted line: 0.4)
- (c) $\Delta m_{31}^2/\Delta m_{21}^2 = 2.0$, $s_{12}^2 = 0.1$ and for varying s_{13}^2 (solid line: 0.01, dashed line: 0.2, dotted line: 0.4)

Fig. 6 The D dependence of $\sin^2 \theta_{23}^m$ for illustrative vacuum parameters with vanishing CP-phase δ

- (a) $s_{12}^2 = 0.1$, $s_{13}^2 = 0.1$, $s_{23}^2 = 0.1$ and varying ratio of mass-squared differences $\Delta m_{31}^2/\Delta m_{21}^2$ (solid line: 1.1, dashed line: 1.5, dotted line: 10)
- (b) $\Delta m_{31}^2/\Delta m_{21}^2 = 2.0$, $s_{13}^2 = 0.1$, $s_{23}^2 = 0.1$ and for varying s_{12}^2 (solid line: 0.01,

dashed line: 0.2, dotted line: 0.4)

(c) $\Delta m_{31}^2/\Delta m_{21}^2 = 2.0$, $s_{12}^2 = 0.1$, $s_{23}^2 = 0.1$ and for varying s_{13}^2 (solid line: 0.01, dashed line: 0.2, dotted line: 0.4)

(d) $\Delta m_{31}^2/\Delta m_{21}^2 = 2.0$, $s_{12}^2 = 0.1$, $s_{13}^2 = 0.1$ and for varying s_{23}^2 (solid line: 0.01, dashed line: 0.2, dotted line: 0.4)

Fig. 7 The CP-phase δ^m in matter for vacuum phases $\frac{\pi}{4}$, $\frac{\pi}{2}$ and $\frac{3}{4}\pi$ with fixed $s_{12}^2 = 0.1$, $s_{13}^2 = 0.1$, $s_{23}^2 = 0.2$ and

(a) almost degenerate neutrino masses $\Delta m_{31}^2/\Delta m_{21}^2 = 1.1$

(b) clearly separated neutrino masses $\Delta m_{31}^2/\Delta m_{21}^2 = 10$

Fig. 8 Mean probability $\langle P(\nu_e \rightarrow \nu_e) \rangle$ for an adiabatic transition from matter of density D into the vacuum

Fig. 9 Exemplary iso-SNU-curves for the ^{37}Cl -experiment assuming a MSW effect for three neutrino flavours:

(a) $\Delta m_{31}^2 = 10^{-2} \text{ eV}^2$, $s_{13}^2 = 0.05$

(b) $\Delta m_{31}^2 = 10^{-4} \text{ eV}^2$, $s_{13}^2 = 0.05$

Figure 1
20

Figure 2
21

Figure 3
22

Figure 4

Figure 5

Figure 6

Figure 7

Figure 8
27

Figure 9